

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert Window: Last 30 Days · Generated: 2026-03-11 20:48:42

Clinical Summary

Recent developments in colorectal cancer research highlight several new clinical trials that may significantly impact treatment options for patients, particularly those with advanced or metastatic disease. Notably, trials are underway for Stage IV patients, including combinations of existing therapies like RC148 with chemotherapy and novel approaches such as dual-target CAR-NK cells for biomarker-selected patients. Additionally, studies are exploring the use of immunotherapy and chemotherapy combinations for both unresectable and locally advanced rectal cancer, which could enhance treatment effectiveness. Moving forward, it will be important to monitor the outcomes of these trials, as they may lead to new standard treatment protocols for various patient populations.

AI-generated from deterministic classification data · 2026-03-11 20:48:42

Situation Summary

Events Detected: 10
High Significance: 0
Top Category: New Trial
Primary Stage: Stage IV
Primary Target: General
Research Activity: Active Monitoring

Event Timeline

EVENT CLUSTER — Safety and Efficacy of Tegavivint in Patients With Metastatic Colorectal Carcinoma

MEDIUM **STAGE IV** **GENERAL** **NEW TRIAL** **VERIFIED** **PUBLISHED MAR 11**

Mar 11 19:09 · immediate · clinicaltrials.gov

A new trial is investigating the safety and effectiveness of Tegavivint for patients with advanced metastatic colorectal cancer, potentially offering a new treatment option for those with limited alternatives.

1 source(s) reporting · cluster 1 of 10

EVENT CLUSTER — A Phase II/III Clinical Trial of RC148 Plus Chemotherapy as 1L Therapy for Unresectable or Metastatic Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 10

Mar 11 19:09 · immediate · clinicaltrials.gov

A new clinical trial is testing the combination of RC148 and chemotherapy as a first-line treatment for patients with unresectable or metastatic colorectal cancer, potentially offering a new option for those facing advanced disease.

1 source(s) reporting · cluster 2 of 10

EVENT CLUSTER — Dual-Target CAR-NK Cells for Biomarker-Selected Advanced Colorectal Cancer

MEDIUM

STAGE IV

PD-1/PD-L1

NEW TRIAL

VERIFIED

PUBLISHED MAR 10

Mar 11 19:09 · immediate · clinicaltrials.gov

A new clinical trial is investigating the use of dual-target CAR-NK cells in patients with advanced colorectal cancer, potentially offering a promising treatment option for those whose tumors express specific biomarkers.

1 source(s) reporting · cluster 3 of 10

EVENT CLUSTER — Low-Dose Radiotherapy to Sensitize Pucotenlimab Plus CAPEOX for pMMR Locally Advanced Rectal Cancer

MEDIUM

STAGE III

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 4

Mar 11 19:09 · immediate · clinicaltrials.gov

A new clinical trial is exploring the use of low-dose radiotherapy combined with the immunotherapy drug pucotenlimab and chemotherapy for patients with locally advanced rectal cancer, which may enhance treatment effectiveness and improve outcomes.

1 source(s) reporting · cluster 4 of 10

EVENT CLUSTER — A Clinical Study of Iparomlimab and Tuvonralimab Combined With Bevacizumab and Alternating Triweekly CAPOX/mCAPIRI Regimen as First-line Treatment for Unresectable Advanced Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

VERIFIED

PUBLISHED MAR 3

Mar 11 19:09 · immediate · clinicaltrials.gov

A new clinical trial is exploring a combination treatment of Iparomlimab, Tuvonralimab, and Bevacizumab with a chemotherapy regimen for patients with advanced colorectal cancer, potentially offering a new first-line option for those with unresectable disease.

1 source(s) reporting · cluster 5 of 10

EVENT CLUSTER — Everolimus in CDK12-Deficient Metastatic Colorectal Cancer (EVER-RECODE)

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED FEB 27

Mar 11 19:09 · immediate · [clinicaltrials.gov](#)

A new trial is exploring the use of Everolimus for patients with Stage IV colorectal cancer that has CDK12 deficiencies, potentially offering a new treatment option for this specific group.

1 source(s) reporting · cluster 6 of 10

EVENT CLUSTER — Effect of Perioperative Lidocaine or High-Dose Dexamethasone on Immune Response in Colon Cancer Surgery (PILDI Study)

MEDIUM

STAGE III

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 6

Mar 11 19:09 · immediate · [clinicaltrials.gov](#)

The PILDI Study is investigating whether administering perioperative lidocaine or high-dose dexamethasone during colon cancer surgery can enhance the immune response, potentially improving outcomes for patients with Stage III colorectal cancer.

1 source(s) reporting · cluster 7 of 10

EVENT CLUSTER — FOLFOX Chemotherapy Combined With Fruquintinib and Serplulimab as First-Line Conversion Therapy for Initially Unresectable pMMR/MSS Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 3

Mar 11 19:09 · immediate · [clinicaltrials.gov](#)

A new clinical trial is exploring the combination of FOLFOX chemotherapy with Fruquintinib and Serplulimab as a promising first-line treatment option for patients with initially unresectable stage IV colorectal cancer, potentially improving their chances for surgery and better outcomes.

1 source(s) reporting · cluster 8 of 10

EVENT CLUSTER — Neoadjuvant Chemoradiotherapy Plus Tislelizumab With or Without Probio-M9 in pMMR/MSS Locally Advanced Rectal Cancer

MEDIUM

STAGE III

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 10

Mar 11 19:09 · immediate · [clinicaltrials.gov](#)

A new clinical trial is exploring the combination of neoadjuvant chemoradiotherapy with the immunotherapy drug Tislelizumab, potentially offering improved treatment options for patients with locally advanced rectal cancer.

1 source(s) reporting · cluster 9 of 10

EVENT CLUSTER — FOLFIRI and Bevacizumab With or Without Pelareorep for Second-Line Treatment of Metastatic RAS-Mutated, Microsatellite-Stable Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

VERIFIED

PUBLISHED MAR 4

Mar 11 19:09 · immediate · clinicaltrials.gov

A new clinical trial is exploring whether adding Pelareorep to the standard treatment of FOLFIRI and Bevacizumab can improve outcomes for patients with advanced colorectal cancer that has specific genetic characteristics.

1 source(s) reporting · cluster 10 of 10

Appendix A — Patient-Friendly Summary

Plain-language overview for patients, caregivers, and family members

What's happening

Recently, several important clinical trials have started that focus on new treatments for advanced colorectal cancer. One trial is looking at a drug called Tegavivint to see if it can help patients with metastatic colorectal cancer, which means the cancer has spread to other parts of the body. Another trial is testing a combination of a drug called RC148 with chemotherapy as a first treatment for patients with advanced disease. There are also trials exploring new combinations of existing treatments, like using low-dose radiotherapy with an immunotherapy drug called pucotenlimab, and a study on Everolimus for patients with specific genetic traits. These developments are exciting as they may provide new options for patients who have limited choices.

What this means for you

If you or a loved one is facing advanced colorectal cancer, these trials could be worth discussing with your oncologist. Ask about the specific trials that might be suitable based on your situation and whether you have any of the characteristics mentioned, like certain genetic markers. It's important to stay informed and engaged in your care. While these new options can be hopeful, they are still in the testing phase, so understanding what they mean for you is crucial. Remember, you are not alone in this journey, and there are people and resources to support you.

AI-generated in plain language from clinical data · 2026-03-11 20:48:42

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Colon & Colorectal Cancer Treatment Reference

Incidence & Trends

~150,000 new US cases/year · ~52,000 deaths/year · 4th most common US cancer

▲ Early-Onset Alert: CRC rates in adults under 50 have risen ~2% per year since the 1990s · Adults under 55 now account for ~20% of all new CRC diagnoses · Cause under active investigation (diet, microbiome, obesity, sedentary lifestyle)

Current Screening Guidelines (2025)

USPSTF / ACS / ACG — Average Risk: Begin screening at age 45 (lowered from 50 in 2021)

Colonoscopy every 10 years · FIT (fecal immunochemical test) annually · Cologuard every 1-3 years · CT colonography every 5 years

High Risk (family history / Lynch syndrome / IBD): Begin at age 40 or 10 years before youngest affected relative — whichever comes first · Colonoscopy every 1-5 years depending on risk

Symptoms at any age: Rectal bleeding, change in bowel habits >4 weeks, unexplained weight loss, iron deficiency anemia → immediate evaluation regardless of age

Key Biomarkers — Test All mCRC at Diagnosis

RAS (KRAS/NRAS) · BRAF V600E · MSI/MMR status · HER2 · NTRK fusion · TMB

Stage IV Targeted Options

KRAS G12C (~3-4%) → sotorasib + cetuximab · adagrasib + cetuximab

BRAF V600E (~8-10%) → encorafenib + cetuximab ± binimetinib

MSI-H/dMMR (~5%) → pembrolizumab (1st line preferred) · nivolumab ± ipilimumab

HER2 amplified (~2-3%) → trastuzumab + pertuzumab · trastuzumab deruxtecan

RAS/BRAF WT → FOLFOX or FOLFIRI + cetuximab or panitumumab (left-sided preferred)

1st Line mCRC Standard of Care

FOLFOX or FOLFIRI ± bevacizumab (RAS mutant / right-sided)

FOLFOX or FOLFIRI + cetuximab or panitumumab (RAS/BRAF WT, left-sided)

Pembrolizumab monotherapy (MSI-H/dMMR)

Key Resources

ClinicalTrials.gov · NCCN Guidelines (nccn.org) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org) · ACS (cancer.org)

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert classification fields are generated using deterministic rule-based analysis. Alert summaries are AI-generated from primary government sources including ClinicalTrials.gov, PubMed (NIH/NLM), and FDA.gov. This report aggregates publicly available data from U.S. government databases for situational awareness only. No article text is reproduced. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

© 2026 Randy Hinton. All rights reserved. · https://petroai1492.github.io/delta/timelines/coloncancer_timeline.pdf