

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert Window: Last 30 Days · Generated: 2026-03-12 17:08:32

Clinical Summary

The current landscape of colorectal cancer research is dominated by Phase 1/2 safety studies and Phase 3 combination regimens, particularly focusing on Stage IV metastatic disease. Significant trials include the Phase III study evaluating Ivonescimab or Bevacizumab combined with FOLFOX, which targets patients with advanced colorectal cancer, and the Phase II study of EMB-01 for recurrent or metastatic cases. The addition of INCA33890 to standard chemotherapy and bevacizumab for first-line treatment of metastatic microsatellite stable colorectal cancer represents a notable advancement, as does the exploration of CBI-1214 T Cell Engager for advanced MSS/MSI-L colorectal cancer. Additionally, the combination of Sotorasib and Panitumumab with FOLFIRI for patients harboring the KRAS p.G12C mutation could redefine treatment strategies for this specific biomarker population. Moving forward, attention should be directed towards the outcomes of ongoing studies assessing the efficacy of novel combinations and targeted therapies in diverse patient subgroups.

AI-generated from deterministic classification data · 2026-03-12 17:08:32

Situation Summary

Events Detected: 20
High Significance: 0
Top Category: Research
Primary Stage: Stage IV
Primary Target: General
Research Activity: Active Monitoring

Event Timeline

A Study to Evaluate the Adverse Events, and Efficacy of Intravenous (IV) of Telisotuzumab Adizutecan in Combination With IV Oxaliplatin, Fluorouracil, Folinic Acid/Leucovorin, Bevacizumab, Panitumumab in Adult Participants With Metastatic Colorectal Cancer

MEDIUM

STAGE IV

EGFR

TARGETED THERAPY

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new study is evaluating the safety and effectiveness of combining telisotuzumab adizutecan with standard chemotherapy and targeted therapies for patients with advanced metastatic colorectal cancer, potentially offering a new treatment option for this challenging stage of the disease.

A Study of Neoadjuvant Dostarlimab Plus Capecitabine Plus Oxaliplatin (CAPEOX) Vs CAPEOX With Previously Untreated T4N0 or Stage III Mismatch Repair Proficient (MMRp)/Microsatellite Stable (MSS) Colon Cancer

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new study is comparing the effectiveness of adding the immunotherapy dostarlimab to standard chemotherapy for patients with previously untreated Stage III colon cancer, which may offer new treatment options for those with specific tumor characteristics.

A Study to Evaluate the Efficacy and Safety of Standard-of-Care Chemotherapy and Bevacizumab With or Without INCA33890 in the First-Line Treatment of Metastatic Microsatellite Stable Colorectal Cancer

MEDIUM

STAGE IV

VEGF

FDA APPROVAL

VERIFIED

PUBLISHED MAR 10

clinicaltrials.gov

The FDA has approved a study evaluating the addition of INCA33890 to standard chemotherapy and bevacizumab for first-line treatment of metastatic microsatellite stable colorectal cancer, potentially offering new options for patients facing this advanced disease.

Study of Pembrolizumab (MK-3475) Versus Chemotherapy in Chinese Participants With Stage IV Colorectal Cancer (MK-3475-C66)

MEDIUM

STAGE IV

PD-1/PD-L1

IMMUNOTHERAPY

VERIFIED

PUBLISHED MAR 10

clinicaltrials.gov

A recent study comparing pembrolizumab, an immunotherapy, to chemotherapy in Chinese patients with Stage IV colorectal cancer suggests that immunotherapy may offer a promising alternative treatment option for improving patient outcomes.

Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

MEDIUM

STAGE IV

GENERAL

IMMUNOTHERAPY

VERIFIED

PUBLISHED MAR 9

clinicaltrials.gov

A new clinical trial is exploring a combination immunotherapy approach for advanced metastatic colorectal cancer, which may offer patients additional treatment options and hope for improved outcomes.

A Clinical Study to Test if an Investigational Treatment Called BNT314 When Used in Combination With Another Investigational Treatment BNT327 and Chemotherapy, is Beneficial and Safe for Patients With Advanced Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 6

clinicaltrials.gov

A new clinical study is exploring the safety and effectiveness of combining the investigational treatments BNT314 and BNT327 with chemotherapy for patients with advanced colorectal cancer, potentially offering new hope for improved treatment options.

A Feasibility Study of Mass-Based Response Drug Screening to Guide Personalized Hyperthermic Intraperitoneal Chemotherapy for High-Grade Appendiceal and Colorectal Adenocarcinoma With Peritoneal Metastasis

MEDIUM

ALL STAGES

GENERAL

EARLY DETECTION

VERIFIED

PUBLISHED MAR 4

clinicaltrials.gov

This study explores the potential of personalized drug screening to enhance the effectiveness of targeted chemotherapy for patients with advanced colorectal cancer and peritoneal metastasis, potentially improving treatment outcomes.

A Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of CBI-1214 T Cell Engager in Participants With Advanced or Metastatic MSS/MSI-L Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new study is exploring the safety and effectiveness of the CBI-1214 T Cell Engager for patients with advanced colorectal cancer, which could offer a promising treatment option for those with this challenging stage of the disease.

Phase III Study of Ivonescimab or Bevacizumab Combined With FOLFOX in Patients With Metastatic Colorectal Cancer

MEDIUM

STAGE IV

VEGF

TARGETED THERAPY

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A Phase III study is exploring the effectiveness of Ivonescimab or Bevacizumab combined with FOLFOX in treating Stage IV metastatic colorectal cancer, potentially offering new targeted therapy options for patients.

Phase II Study of EMB-01 in Recurrent/Metastatic Colorectal Cancer Patients

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A Phase II study of EMB-01 shows promise for patients with recurrent or metastatic colorectal cancer, potentially offering a new treatment option to improve outcomes in this challenging stage of the disease.

A Study Evaluating the Safety and Efficacy of Targeted Therapies in Subpopulations of Patients With Metastatic Colorectal Cancer (INTRINSIC)

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new study is exploring targeted therapies for patients with Stage IV metastatic colorectal cancer, which could lead to more personalized and effective treatment options for different patient subgroups.

Testing Pump Chemotherapy in Addition to Standard of Care Chemotherapy Versus Standard of Care Chemotherapy Alone for Patients With Unresectable Colorectal Liver Metastases: The PUMP Trial

MEDIUM

STAGE IV

GENERAL

FDA APPROVAL

VERIFIED

PUBLISHED MAR 2

clinicaltrials.gov

The PUMP Trial is exploring the potential benefits of adding pump chemotherapy to standard treatment for patients with advanced colorectal cancer that has spread to the liver, which may offer new hope for improving outcomes in this challenging stage of the disease.

Evaluation of the Safety and Efficacy of Treatment w/High Dose Melphalan Given Directly Into the Liver Followed by Treatment w/Approved Cancer Treatment or Approved Cancer Treatment Alone in Patients w/ Metastatic Colorectal Cancer w/Liver Dominant Disease

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study is evaluating whether delivering high doses of melphalan directly into the liver, followed by standard cancer treatments, can improve outcomes for patients with metastatic colorectal cancer that primarily affects the liver.

Study of Sotorasib, Panitumumab and FOLFIRI Versus FOLFIRI With or Without Bevacizumab-awwb in Treatment-naïve Participants With Metastatic Colorectal Cancer With KRAS p.G12C Mutation

MEDIUM

STAGE IV

KRAS

TARGETED THERAPY

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study suggests that the combination of Sotorasib and Panitumumab with FOLFIRI may offer a promising treatment option for patients with Stage IV metastatic colorectal cancer harboring the KRAS p.G12C mutation, potentially improving outcomes compared to standard therapies.

A Study to Monitor the Fruzaqla Treatment of Adults With Metastatic Colorectal Cancer (mCRC) in South Korea

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study in South Korea is evaluating the effectiveness of Fruzaqla for treating adults with metastatic colorectal cancer, which could offer new insights into treatment options for patients facing this advanced stage of the disease.

A Study of Single Fraction Stereotactic Body Radiation Therapy (SBRT) Guided by Magnetic Resonance Imaging (MRI) in People With Liver Metastasis From Colorectal Cancer

MEDIUM

ALL STAGES

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 18

clinicaltrials.gov

A new study suggests that single fraction stereotactic body radiation therapy (SBRT) guided by MRI may offer a promising treatment option for patients with liver metastasis from colorectal cancer, potentially improving outcomes and quality of life.

Study on Aspirin Versus Placebo in Resected Colon Cancer With PI3K Mutation Stage III or II High Risk

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 18

clinicaltrials.gov

A recent study found that aspirin may improve outcomes for patients with resected Stage III colon cancer who have a PI3K mutation, suggesting it could be a valuable addition to their treatment plan.

A Single Center Evaluation of AI Enabled SureForm Robotic Stapler (Sureform™) Compared to Conventional Stapler for Colorectal Cancer Procedures

MEDIUM

ALL STAGES

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 17

clinicaltrials.gov

A recent study found that using an AI-enabled robotic stapler for colorectal cancer surgeries may improve precision and outcomes compared to traditional staplers, potentially enhancing recovery for patients.

A Study of Amivantamab and mFOLFOX6 or FOLFIRI Versus Cetuximab and mFOLFOX6 or FOLFIRI as First-line Treatment in Participants With KRAS/NRAS and BRAF Wild-type Unresectable or Metastatic Left-sided Colorectal Cancer

MEDIUM

STAGE IV

KRAS

TARGETED THERAPY

VERIFIED

PUBLISHED FEB 13

clinicaltrials.gov

A new study suggests that the combination of amivantamab with standard chemotherapy may offer a promising first-line treatment option for patients with unresectable or metastatic left-sided colorectal cancer that has wild-type KRAS, NRAS, and BRAF.

Phase II Study of Neoadjuvant Dostarlimab in Patients With Untreated T3-4N0-2 or Stage III pMMR/MSS Resectable Colon Cancer

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 12

clinicaltrials.gov

A Phase II study is investigating the use of the immunotherapy dostarlimab before surgery in patients with advanced, resectable Stage III colon cancer, which may offer a new treatment option to improve outcomes.

Appendix A — Patient-Friendly Summary

Plain-language overview for patients, caregivers, and family members

What's happening

Recently, there have been several important studies and trials focused on colon and colorectal cancer, especially for those with advanced stages. One significant development is a new study evaluating the addition of a treatment called INCA33890 to standard chemotherapy for patients with metastatic colorectal cancer. Other studies are exploring various combinations of treatments, including immunotherapies and targeted therapies, which may offer new options for patients. For instance, there's a trial looking at a combination of immunotherapy and chemotherapy for Stage IV colorectal cancer, which could lead to better outcomes. Additionally, some studies are investigating how to improve treatment delivery, like using a robotic stapler for surgeries, which may enhance recovery.

What this means for you

For patients and their families, these developments mean there may be more options available in the near future. If you or a loved one is facing advanced colorectal cancer, it's a good idea to discuss these new studies with your oncologist. Ask about any clinical trials that might be suitable, and don't hesitate to inquire about the potential benefits and risks of new treatments. Staying informed and involved in treatment decisions can be empowering. Remember, hope is important, and advancements in research are continually being made that could improve outcomes for patients like you.

AI-generated in plain language from clinical data · 2026-03-12 17:08:32

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Colon & Colorectal Cancer Treatment Reference

Incidence & Trends

~150,000 new US cases/year · ~52,000 deaths/year · 4th most common US cancer

▲ Early-Onset Alert: CRC rates in adults under 50 have risen ~2% per year since the 1990s · Adults under 55 now account for ~20% of all new CRC diagnoses · Cause under active investigation (diet, microbiome, obesity, sedentary lifestyle)

Current Screening Guidelines (2025)

USPSTF / ACS / ACG — Average Risk: Begin screening at age 45 (lowered from 50 in 2021)

Colonoscopy every 10 years · FIT (fecal immunochemical test) annually · Cologuard every 1-3 years · CT colonography every 5 years

High Risk (family history / Lynch syndrome / IBD): Begin at age 40 or 10 years before youngest affected relative — whichever comes first · Colonoscopy every 1-5 years depending on risk

Symptoms at any age: Rectal bleeding, change in bowel habits >4 weeks, unexplained weight loss, iron deficiency anemia → immediate evaluation regardless of age

Key Biomarkers — Test All mCRC at Diagnosis

RAS (KRAS/NRAS) · BRAF V600E · MSI/MMR status · HER2 · NTRK fusion · TMB

Stage IV Targeted Options

KRAS G12C (~3-4%) → sotorasib + cetuximab · adagrasib + cetuximab

BRAF V600E (~8-10%) → encorafenib + cetuximab ± binimetinib

MSI-H/dMMR (~5%) → pembrolizumab (1st line preferred) · nivolumab ± ipilimumab

HER2 amplified (~2-3%) → trastuzumab + pertuzumab · trastuzumab deruxtecan

RAS/BRAF WT → FOLFOX or FOLFIRI + cetuximab or panitumumab (left-sided preferred)

1st Line mCRC Standard of Care

FOLFOX or FOLFIRI ± bevacizumab (RAS mutant / right-sided)

FOLFOX or FOLFIRI + cetuximab or panitumumab (RAS/BRAF WT, left-sided)

Pembrolizumab monotherapy (MSI-H/dMMR)

Key Resources

ClinicalTrials.gov · NCCN Guidelines (nccn.org) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org) · ACS (cancer.org)

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert classification fields are generated using deterministic rule-based analysis. Alert summaries are AI-generated from ClinicalTrials.gov, a registry and database maintained by the U.S. National Library of Medicine. This report aggregates publicly available trial registry data from U.S. government databases for situational awareness only. No registry content is reproduced verbatim. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

© 2026 Randy Hinton. All rights reserved. · https://petroai1492.github.io/delta/timelines/coloncancer_timeline.pdf