

Colon & Colorectal Cancer — Clinical Trials Intelligence

Alert Window: Last 90 Days · Generated: 2026-04-12 15:18:12

Situation Summary

Phase I/II: 25 trials
Recruiting: 132 trials
Biomarker: 112 biomarker-directed trials
Breakthrough: 58 trials scored ≥ 4

Active Trial Timeline

Showing 58 high-significance trials (score ≥ 4). 171 additional lower-scored trials are collapsed below. Phase III trials appear first. Click any title to view full trial details on ClinicalTrials.gov.

MEDIUM**PHASE3****Stage IV****MSI/MMR****TARGETED THERAPY**• **RECRUITING****11**

Botensilimab + Balstilimab vs Best Supportive Care as Therapy in Chemo-refractory, Unresectable, Colorectal Adenocarcinoma

This clinical trial is testing the combination of botensilimab and balstilimab in patients with chemo-refractory, unresectable colorectal adenocarcinoma. The intervention involves two immunotherapy drugs aimed at potentially improving overall survival and quality of life for participants. Currently in Phase 3 and actively recruiting, this trial may provide insights into the effectiveness and safety of these treatments for eligible patients. Participants may need to meet specific biomarker criteria, including MSI, MSI-H, MMR, dMMR, and microsatellite status.

NCT: NCT07152821 · Enrollment: 834 · Sites: Greenfield Park, Canada, Kingston, Canada, Oshawa, Canada, Ottawa, Canada, Prince George, Canada

MEDIUM**PHASE3****Stage III****MSI/MMR****TARGETED THERAPY**• **NOT_YET_RECRUITING****11**

Neoadjuvant/Adjuvant AK104 in Microsatellite Instability-high or Mismatch Repair-deficient, Resectable Colon Cancer

This clinical trial is testing the efficacy and safety of neoadjuvant/adjuvant treatment with AK104 (Cadonilimab) in patients with resectable microsatellite instability-high (MSI-H) or mismatch repair-deficient (dMMR) colon cancer. The intervention involves AK104 in combination with standard chemotherapy agents, including Oxaliplatin, Capecitabine, 5-Fluorouracil, and Calcium Folate, which is significant for potentially improving treatment outcomes in this specific patient population. This is a Phase 3 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include having resectable colon cancer with specific biomarkers such as MSI-H or dMMR.

NCT: NCT07412613 · Enrollment: 386 · Sites: Guangzhou, China

MEDIUM**PHASE3****All Stages****KRAS****TARGETED THERAPY**• **RECRUITING****11**

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

This clinical trial is testing the efficacy of amivantamab combined with FOLFIRI chemotherapy in patients with KRAS/NRAS and BRAF wild-type colorectal cancer who have previously undergone chemotherapy. The intervention

involves the biological agent amivantamab alongside standard chemotherapy agents, which is important for evaluating its potential benefits compared to existing treatment options. This is a Phase 3 trial currently recruiting 700 participants, which means that eligible patients may have the opportunity to receive a treatment that is being compared to standard therapies in a larger population. Eligibility criteria include specific biomarker requirements related to KRAS, NRAS, and BRAF status, as well as other factors such as microsatellite instability (MSI) and mismatch repair (MMR) status.

NCT: NCT06750094 · Enrollment: 700 · Sites: Abilene, Ann Arbor, Arlington, Athens, Atlanta

MEDIUM

PHASE3

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

A Study of Neoadjuvant QL1706 in Participants With Untreated dMMR/MSI-H Resectable Colon Cancer

This clinical trial is testing the efficacy of neoadjuvant QL1706 in participants with untreated dMMR/MSI-H resectable colon cancer. The intervention involves administering QL1706 alongside CAPEOX/Capecitabine to assess its impact on tumor response prior to surgery. Currently in Phase 3 and actively recruiting, this trial aims to determine the proportion of patients achieving a pathological complete response (pCR), which indicates no residual tumor after treatment. Eligibility criteria include having T4N0 or Stage III colon cancer with specific biomarkers such as MSI-H or dMMR.

NCT: NCT06686576 · Enrollment: 363 · Sites: Guangzhou, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

Injection of SHR-A1811 Versus Physician Chocied Treatment in Patients With Advanced Colorectal Cancer Who Had Failed to Respond to Oxaliplatin, 5-fu, and Irinotecan

This clinical trial is testing the efficacy and safety of SHR-A1811 in patients with advanced colorectal cancer who have not responded to previous treatments with oxaliplatin, 5-FU, and irinotecan. The intervention involves the drug SHR-A1811, which is being compared to physician-chosen treatments such as TAS-102, regorafenib, and fruquintinib. This is a Phase 3 trial that is currently active but not recruiting, meaning that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06199973 · Enrollment: 130 · Sites: Shanghai, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

Study of XL092 + Atezolizumab vs Regorafenib in Participants With Metastatic Colorectal Cancer

This clinical trial is evaluating the combination of XL092 and atezolizumab compared to regorafenib in participants with metastatic colorectal cancer who have progressed during, after, or are intolerant to standard-of-care therapy. The intervention involves the use of XL092 and atezolizumab, which may provide a different therapeutic approach for patients with microsatellite stable or microsatellite instability low metastatic colorectal cancer. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT05425940 · Enrollment: 901 · Sites: Albuquerque, Billings, Charlotte, Chattanooga, Cincinnati

MEDIUM

PHASE3

All Stages

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

9

Study Evaluating Two Treatment Strategies for Rectal Cancer in Patients ≥75 Years Old

The NACRE-2 trial is evaluating two treatment strategies for patients aged 75 years and older with locally advanced rectal cancer. The study compares the effectiveness of adding chemotherapy (FOLFOX4s) after short-course radiotherapy against short-course radiotherapy alone. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as MSI, MMR, DMMR, and microsatellite status.

NCT: NCT07118800 · Enrollment: 160 ·

SIGNAL

PHASE3

Stage IV

BRAF

RESEARCH

• RECRUITING

9

Intermittent or Continuous Panitumumab Plus FOLFIRI for Left Sided RAS/B-RAF Wild-type Metastatic Colorectal Cancer

This clinical trial is testing the effectiveness of intermittent versus continuous administration of Panitumumab combined with FOLFIRI in patients with left-sided RAS/B-RAF wild-type metastatic colorectal cancer. The intervention involves the

use of Panitumumab, Irinotecan, 5-fluorouracil, and L-folinic acid, which are important for evaluating treatment strategies that may optimize patient outcomes. Currently in Phase 3 and actively recruiting, this trial aims to determine if the intermittent regimen can achieve a Time to Treatment Failure comparable to the standard continuous regimen, which is relevant for patients considering their treatment options. Eligibility criteria include specific biomarker requirements related to BRAF, MMR, and RAS status.

NCT: NCT06509126 · Enrollment: 500 · Sites: Naples, Italy

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

9

A Study To Evaluate The Efficacy And Safety Of Ifinatamab Deruxtecan (I-DXd) In Subjects With Recurrent Or Metastatic Solid Tumors (IDeate-PanTumor02)

This clinical trial is evaluating the efficacy and safety of ifinatamab deruxtecan (I-DXd) in patients with recurrent or metastatic solid tumors, including colorectal cancer. The intervention, ifinatamab deruxtecan, is a drug that targets specific cancer biomarkers, which may influence treatment outcomes. This is a Phase 2 trial currently recruiting 520 participants, indicating that patients may have the opportunity to access this treatment while contributing to the understanding of its effectiveness. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, HER2, and RAS.

NCT: NCT06330064 · Enrollment: 520 · Sites: Amarillo, Chattanooga, Dallas, Fairfax, Germantown

SIGNAL

PHASE3

Stage III

RESEARCH

• RECRUITING

9

Thymosin-alpha 1 for Adjuvant Treatment After Radical Resection of High-risk Stage II and III Colorectal Cancer

This clinical trial is testing the efficacy and safety of Thymosin-alpha 1 as an adjuvant treatment for patients with high-risk stage II and III colorectal cancer following radical resection. The intervention, Thymosin-alpha 1, is a drug believed to enhance immune response, potentially reducing the risk of cancer recurrence and metastasis. This is a Phase 3 trial currently recruiting 2,500 participants, which indicates that it is in the final stages of testing before potential approval and may provide valuable information for patients regarding treatment options. Eligibility criteria may include specific high-risk classifications of colorectal cancer, although detailed biomarker requirements are not specified.

NCT: NCT05086614 · Enrollment: 2500 · Sites: Shanghai, China

SIGNAL

PHASE3

Stage IV

RAS

RESEARCH

• ACTIVE_NOT_RECRUITING

8

Neoadjuvant FOLFOX Chemotherapy for Patients With Locally Advanced Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant FOLFOX chemotherapy in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy before surgery, which may enhance treatment outcomes by targeting microscopic metastatic cancer cells earlier and reducing surgical complications. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and results may soon be available for clinical application. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT03426904 · Enrollment: 708 · Sites: Daegu, South Korea, Hwasun, South Korea, Seoul, South Korea, Suwon, South Korea

SIGNAL

PHASE3

Stage IV

RESEARCH

• RECRUITING

9

Bevacizumab Plus mFOLFOXIRI as First-line Treatment for Patients With Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of mFOLFOXIRI combined with Bevacizumab versus mFOLFOX6 combined with Bevacizumab as first-line treatments for patients with unresectable metastatic colorectal cancer. The intervention involves two chemotherapy regimens, which are significant as they aim to improve treatment outcomes while considering the unique responses of different populations, particularly in China. This is a Phase 3 trial currently recruiting 528 participants, which means eligible patients may have the opportunity to receive one of the investigational treatments while contributing to important research data. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

NCT: NCT04230187 · Enrollment: 528 · Sites: Guangzhou, China

SIGNAL

PHASE2, PHASE3

Stage III

MSI/MMR

RESEARCH

• RECRUITING

8

Efficacy and Safety Comparison of Short-course Radiotherapy Followed by CapeOX Chemotherapy Plus Toripalimab With or Without Concurrent Surufatinib in Neoadjuvant Therapy for Mid-to-low Localized Rectal Cancer of High-risk Criteria

This clinical trial is testing the efficacy and safety of short-course radiotherapy followed by CapeOX chemotherapy combined with the immunotherapy drug Toripalimab, with or without the anti-angiogenic drug Surufatinib, in patients with mid-to-low localized rectal cancer who meet high-risk criteria. The intervention aims to improve the pathological complete response rate in patients with high-risk locally advanced rectal cancer, addressing issues of regional recurrence and distant metastasis that persist despite current treatment methods. This study is currently in Phase 2 and Phase 3 and is actively recruiting participants, which means eligible patients may have the opportunity to access this investigational treatment. Eligibility criteria include specific biomarkers such as MMR and RAS status.

NCT: NCT07505472 · Enrollment: 212 · Sites: Changchun, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

8

A Phase II Clinical Study Evaluating SSGJ-706 in Combination Therapy for Advanced Gastrointestinal Cancers

This phase II clinical trial is evaluating the combination of SSGJ-706 with standard therapies for patients with advanced gastrointestinal cancers. The interventions include SSGJ-706 combined with XELOX, Bevacizumab, AG, and TP, which are important for assessing the safety and effectiveness of this treatment approach. The trial is currently not yet recruiting participants, meaning that patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and HER2.

NCT: NCT07233850 · Enrollment: 300 · Sites: Wuhan, China

SIGNAL

PHASE2, PHASE3

Stage III

RESEARCH

• RECRUITING

8

Total Neoadjuvant Therapy and Organ Preservation Versus Surgery for Rectal Cancer.

This clinical trial is testing the effectiveness of total neoadjuvant therapy (TNT) and a watch-and-wait strategy for patients with rectal cancer. The intervention includes radiation therapy, chemoradiotherapy, and consolidation chemotherapy, with the aim of potentially allowing patients to preserve their rectum if they achieve a complete or near-complete clinical response. Currently in Phase 2 and Phase 3, the trial is recruiting 400 participants, which means eligible patients may have the opportunity to contribute to research that could impact treatment approaches. Eligibility criteria include patients with cT1N1, T2-T3 N0-1 rectal cancer, and specific markers such as MRF- and EMVI-.

NCT: NCT06758830 · Enrollment: 400 · Sites: Vilnius, Lithuania

MEDIUM

PHASE3

All Stages

MSI/MMR

IMMUNOTHERAPY

• RECRUITING

8

Personalized Long-course Radiotherapy Plus Chemotherapy With or Without Immunotherapy for LARC: PALACE Study

The PALACE study is a phase III clinical trial designed for patients with locally advanced rectal cancer (LARC) to evaluate the complete response rate to personalized long-course radiotherapy combined with chemotherapy, with or without the anti-PD-1 antibody drug Serplulimab. This intervention aims to determine the effectiveness of adding immunotherapy to standard treatment in achieving a complete response, defined as a pathological complete response plus a clinical complete response sustained for over one year. Currently, the trial is recruiting participants, which means eligible patients have the opportunity to join and contribute to this research. Eligibility criteria include specific biomarkers such as MMR and RAS.

NCT: NCT07020247 · Enrollment: 184 · Sites: Chengdu, China

MEDIUM

N/A

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

8

The Efficacy of Watch and Wait Strategy or Surgery After Neoadjuvant Immunotherapy for Locally Advanced Colorectal Cancer With dMMR/MSI-H Guided by MRD Dynamic Monitoring: A Single-center, Open-label, Prospective, Phase II Clinical Trial.

This clinical trial is testing the efficacy of a watch and wait strategy compared to surgery after neoadjuvant immunotherapy in patients with locally advanced colorectal cancer characterized by deficient mismatch repair or microsatellite instability-high (dMMR/MSI-H). The intervention involves the use of Tislelizumab, a PD-1 inhibitor, which is significant for its potential to enhance the immune response against cancer cells. Currently, the trial is in the recruiting phase, indicating that eligible patients can still enroll to participate in this study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite status, as well as the use of next-generation sequencing (NGS) for patient selection.

MEDIUM

PHASE2, PHASE3

All Stages

MSI/MMR

TARGETED THERAPY

• RECRUITING

8

DETERMINE Trial Treatment Arm 02: Atezolizumab in Adult, Paediatric and Teenage/Young Adult Patients With Cancers With High Tumour Mutational Burden (TMB) or Microsatellite Instability-high (MSI-high) or Proven Constitutional Mismatch Repair Deficiency (CMMRD) Disposition

The DETERMINE Trial Treatment Arm 02 is investigating the efficacy of atezolizumab in adult, pediatric, and teenage/young adult patients with cancers characterized by high tumor mutational burden (TMB), microsatellite instability-high (MSI-high), or proven constitutional mismatch repair deficiency (CMMRD). Atezolizumab is a drug that has been approved for several cancer types and is being evaluated for its potential benefits in additional cancers with specific genetic markers. This trial is currently in Phase 2 and Phase 3 and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to important research. Eligibility criteria include the presence of specific biomarkers such as MSI, TMB, and MMR, among others.

NCT: NCT05770102 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL

PHASE2, PHASE3

Stage IV

BRAF

RESEARCH

• RECRUITING

8

Regorafenib With Low-dose Chemotherapies and Aspirin Followed by Standard Chemotherapies in Metastatic Colorectal Cancer

This clinical trial is testing the combination of regorafenib with low-dose metronomic chemotherapies and aspirin for patients with metastatic colorectal cancer. The intervention aims to assess the effectiveness of this combination as a two-month induction therapy prior to standard chemotherapy initiation. Currently in Phase 2 and Phase 3, the trial is recruiting participants, which means eligible patients can join to potentially receive this treatment approach. Eligibility criteria include specific biomarkers such as BRAF, RAS, and microsatellite status.

NCT: NCT05462613 · Enrollment: 446 · Sites: Besançon, France, Clermont-Ferrand, France, Créteil, France, Dijon, France, Lyon, France

SIGNAL

PHASE3

Stage IV

KRAS

RESEARCH

• RECRUITING

8

Systemic Oxaliplatin or Intra-arterial Chemotherapy Combined With LV5FU2 +/- Irinotecan and an Target Therapy in First Line Treatment of Metastatic Colorectal Cancer Restricted to the Liver

This clinical trial is testing the effectiveness of systemic oxaliplatin versus intra-arterial chemotherapy combined with LV5FU2, with or without irinotecan and panitumumab, in patients with metastatic colorectal cancer restricted to the liver. The intervention involves administering oxaliplatin both intravenously and intra-arterially, along with other chemotherapy agents and targeted therapy, to potentially enhance treatment efficacy for patients with unresectable liver metastases. This is a Phase 3 trial currently recruiting 348 participants, which indicates that it is in the later stages of testing and may provide valuable information on treatment options for patients. Eligibility criteria include specific biomarker assessments for KRAS, NRAS, and RAS.

NCT: NCT02885753 · Enrollment: 348 · Sites: Angers, France, Antony, France, Avignon, France, Bayonne, France, Beauvais, France

SIGNAL

PHASE3

All Stages

MSI/MMR

RESEARCH

• NOT_YET_RECRUITING

8

Finding the Best Dose of Aspirin to Prevent Lynch Syndrome Cancers

This clinical trial is testing the effectiveness of different doses of enteric coated aspirin in preventing cancers associated with Lynch Syndrome, specifically targeting individuals who carry a germline pathological mismatch repair gene defect. The intervention involves comparing daily doses of 600mg, 300mg, and 100mg of aspirin to determine the optimal dose for cancer prevention. This is a Phase 3 trial that is currently not yet recruiting participants, meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include being a carrier of a germline pathological mismatch repair gene defect associated with Lynch Syndrome.

NCT: NCT02497820 · Enrollment: 1800 · Sites: Tel Aviv, Israel

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

8

Study of Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-3475-158/KEYNOTE-158)

This clinical trial is testing the efficacy of pembrolizumab (MK-3475) in participants with advanced solid tumors who have progressed on standard of care therapy. The intervention, pembrolizumab, is a biological agent that targets the immune system to help fight cancer, which is significant for patients with limited treatment options. This is a Phase 2 trial that is currently active but not recruiting, indicating that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, and microsatellite instability, which may be relevant for patient selection.

NCT: NCT02628067 · Enrollment: 1609 ·

SIGNAL **PHASE3** **Stage IV** **BRAF** **RESEARCH** • NOT_YET_RECRUITING

7

Pressurized Intraperitoneal Aerosolized Chemotherapy With Mitomycin for the Treatment of Unresectable Appendix or Colorectal Cancer With Peritoneal Metastases, The IMPACT Trial

This phase III clinical trial, known as the IMPACT Trial, is testing the effectiveness of pressurized intraperitoneal aerosolized chemotherapy (PIPAC) with mitomycin in patients with unresectable appendix or colorectal cancer that has metastasized to the peritoneal cavity. The intervention involves delivering chemotherapy as a pressurized mist directly into the abdominal cavity during a minimally invasive laparoscopic procedure, which may enhance drug absorption and distribution compared to standard chemotherapy regimens. The trial is currently not yet recruiting participants, which means patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarker assessments such as BRAF, RAS, and mismatch repair status, as well as next-generation sequencing (NGS).

NCT: NCT07271355 · Enrollment: 129 · Sites: Duarte, Goodyear, Irvine, Newnan, Zion

HIGH **PHASE3** **All Stages** **PHASE III TRIAL** • RECRUITING

7

Phase 3 Trial of eRapa in Patients With Familial Adenomatous Polyposis

This Phase 3 clinical trial is testing the efficacy of eRapa (encapsulated rapamycin) in patients diagnosed with Familial Adenomatous Polyposis (FAP). The intervention involves administering eRapa compared to a placebo to determine its impact on slowing disease progression and its safety profile. Currently, the trial is recruiting 168 participants, which means eligible patients can still join to contribute to the research. Eligibility criteria include a diagnosis of FAP, and participants will undergo regular check-ups and endoscopies to monitor their condition throughout the study.

NCT: NCT06950385 · Enrollment: 168 · Sites: Ann Arbor, Arcadia, Baltimore, Boston, Chicago

SIGNAL **PHASE3** **All Stages** **RESEARCH** • RECRUITING

7

CompariSon Between the EuroPeAn and Japanese pathologiCal InvEstigation for Colon Cancer (SPACE)

The SPACE trial is testing the diagnostic accuracy of Japanese versus European pathological investigation methods for stage III colon cancer in patients. The intervention involves comparing two procedures: the Japanese pathological investigation, which includes surgeon involvement in lymph node extraction and specimen assessment, and the European method, which primarily relies on pathologists. This is a Phase 3 trial currently recruiting 430 participants, which means patients may have the opportunity to contribute to important findings regarding diagnostic practices. Eligibility criteria include patients diagnosed with stage III colon cancer, although specific biomarker requirements are not mentioned.

NCT: NCT06119867 · Enrollment: 430 · Sites: Moscow, Russia

SIGNAL **PHASE3** **All Stages** **RAS** **RESEARCH** • RECRUITING

7

Regorafenib Combined With Irinotecan Drug-Eluting Beads for Colorectal Cancer Liver Metastases

This clinical trial is testing the efficacy and safety of regorafenib combined with Irinotecan Drug-Eluting Beads in patients with colorectal cancer liver metastases who have failed first- and second-line standard chemotherapy. The intervention involves the use of regorafenib and DIBIRI (drug-eluting beads), which is significant as it explores a combination treatment approach for this patient population. This is a Phase 3 trial currently recruiting 126 participants, which means that eligible patients may have the opportunity to access this investigational treatment while contributing to important research. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT05794971 · Enrollment: 126 · Sites: Guangdong, China

SIGNAL **PHASE3** **All Stages** **RESEARCH** • ACTIVE_NOT_RECRUITING

7

Effect of EPA-FFA on Polypectomy in Familial Adenomatous Polyposis

This clinical trial is testing the effect of eicosapentaenoic acid free fatty acid (EPA-FFA) on polypectomy outcomes in patients with familial adenomatous polyposis (FAP). The intervention involves administering EPA-FFA gastro-resistant capsules compared to a placebo, which is important for assessing its potential impact on the number of polypectomies required. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the later stages of testing and may provide significant insights into treatment efficacy for patients. Eligibility criteria may include a diagnosis of familial adenomatous polyposis, although specific biomarker requirements are not detailed.

NCT: NCT03806426 · Enrollment: 204 · Sites: Bologna, Italy

MEDIUM **PHASE1, PHASE2** **Stage IV** **KRAS** **LIQUID BIOPSY** • RECRUITING

7

A Study of Bispecific Antibody MCLA-158 in Patients With Advanced Solid Tumors

This clinical trial is testing the bispecific antibody MCLA-158 in patients with advanced solid tumors, specifically focusing on metastatic colorectal cancer (mCRC) and other solid tumor indications. The intervention includes MCLA-158 as a single agent and in combination with pembrolizumab, FOLFIRI, and FOLFOX, which is important for evaluating its safety and effectiveness in various treatment settings. Currently in Phase 1/2 and actively recruiting, this trial aims to determine the recommended Phase II dose and assess the drug's safety, tolerability, and anti-tumor activity. Eligibility criteria include specific biomarkers such as KRAS, NRAS, BRAF, HER2, and others, which may influence treatment response.

NCT: NCT03526835 · Enrollment: 523 · Sites: Boston, Cleveland, Dallas, Fort Myers, Houston

SIGNAL **PHASE3** **All Stages** **RESEARCH** • ACTIVE_NOT_RECRUITING

7

EPA for Metastasis Trial 2

The EPA for Metastasis Trial 2 is testing the efficacy of Icosapent Ethyl, a pure omega-3 fatty acid derived from fish oil, in patients with liver metastases from bowel cancer. The trial aims to determine if taking this supplement before and after liver surgery can improve progression-free survival compared to a placebo. Currently in Phase 3 and active but not recruiting, this study involves 418 participants and seeks to provide insights into the potential benefits of Icosapent Ethyl in preventing disease recurrence. Eligibility criteria include having liver metastases from bowel cancer, with a target enrollment of 448 individuals.

NCT: NCT03428477 · Enrollment: 418 · Sites: Aintree, United Kingdom, Basingstoke, United Kingdom, Birmingham, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL **PHASE2, PHASE3** **All Stages** **RESEARCH** • ACTIVE_NOT_RECRUITING

7

Perioperative Systemic Therapy for Isolated Resectable Colorectal Peritoneal Metastases

This clinical trial is testing the effectiveness of perioperative systemic therapy combined with cytoreductive surgery and HIPEC in patients with isolated resectable colorectal peritoneal metastases. The intervention includes various combinations of chemotherapy regimens (CAPOX, FOLFOX, FOLFIRI) with bevacizumab, alongside the experimental procedure of cytoreductive surgery with HIPEC. Currently in Phase II-III and marked as active but not recruiting, this trial aims to determine if the combination therapy improves overall survival compared to surgery alone. Eligibility criteria include having isolated resectable colorectal peritoneal metastases, although specific biomarker requirements are not detailed.

NCT: NCT02758951 · Enrollment: 358 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Genk, Belgium, Groningen, Netherlands, Nieuwegein, Netherlands

MEDIUM **PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • ACTIVE_NOT_RECRUITING

6

The Efficacy and Safety of Neoadjuvant Therapy With Iparomlimab and Tuvonralimab in Locally Advanced MSI-H/dMMR Colorectal Cancer: An Prospective, Single-Arm Study (Neo-IT)

This clinical trial is testing the efficacy and safety of neoadjuvant therapy with Iparomlimab and Tuvonralimab in patients with locally advanced colorectal cancer that is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR). The intervention involves the use of these two immunotherapy drugs, which may provide significant benefits in the preoperative treatment of this specific cancer type. This is a Phase 2 trial that is currently active but not recruiting, indicating that patients are not being enrolled at this time, but results may inform future treatment options. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, and RAS status.

MEDIUM

PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

A First in Human Study of ALX2004 With Advanced or Metastatic Selected Solid Tumors

This clinical trial is testing ALX2004, an antibody drug conjugate targeting EGFR, in patients with advanced or metastatic selected solid tumors. The intervention involves administering ALX2004 to evaluate its safety and tolerability, specifically focusing on the incidence of dose limiting toxicities (DLTs). Currently in Phase 1 and actively recruiting, this trial aims to gather initial data on the drug's effects, which is crucial for determining its potential for further development. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT07085091 · Enrollment: 170 · Sites: Fairfax, Grand Rapids, Houston, Portland, Spokane

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Phase 2 Study of SR-8541A in Combination With Botensilimab and Balstilimab in Subjects With Refractory Metastatic Microsatellite Stable Colorectal Cancer (MSS-CRC)

This clinical trial is testing the combination of SR-8541A, botensilimab, and balstilimab in patients with refractory metastatic microsatellite stable colorectal cancer (MSS-CRC). The intervention involves administering SR-8541A orally alongside the other two drugs to assess their safety, tolerability, and efficacy, as well as to determine the optimal dose for further studies. Currently, this is a Phase 2 trial that is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include specific biomarkers related to microsatellite stability and mismatch repair status.

NCT: NCT06589440 · Enrollment: 70 · Sites: Austin, Dallas, Houston, Morristown, Seattle

MEDIUM

PHASE1, PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

This clinical trial is testing a combination immunotherapy regimen for adults with metastatic colorectal cancer (mCRC). The intervention includes the drugs retifanlimab, SX-682, and N-803, along with the TriAdeno vaccine, aiming to evaluate their safety profiles in treating mCRC. Currently in Phase I/II and actively recruiting, this trial offers participants the opportunity to access experimental therapies that may not be available outside of a clinical trial setting. Eligible participants must be 18 years or older and diagnosed with mCRC, with specific biomarker criteria including MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT06149481 · Enrollment: 60 · Sites: Bethesda

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Liposomal Irinotecan and Leucovorin/5-fluorouracil Plus Bevacizumab in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of liposomal irinotecan combined with 5-fluorouracil, leucovorin, and bevacizumab as a second-line therapy for patients with metastatic colorectal cancer. The intervention involves a combination of these drugs, which is important for potentially improving treatment outcomes in this patient population. Currently, the trial is in Phase 2 and is actively recruiting participants, indicating that patients may have the opportunity to receive this treatment while the study is ongoing. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06184698 · Enrollment: 173 · Sites: Shijiazhuang, China

MEDIUM

PHASE2

Stage IV

BRAF

LIQUID BIOPSY

• RECRUITING

6

Cetuximab Plus Irinotecan in Patients With NeoRAS Wild-type Metastatic Colorectal Cancer In Third-line Therapy

This clinical trial is testing the efficacy and safety of cetuximab plus irinotecan in patients with NeoRAS wild-type metastatic colorectal cancer (mCRC) who are receiving third-line therapy. The intervention involves the combination of cetuximab, a monoclonal antibody, and irinotecan, a chemotherapy drug, which may provide a targeted approach for this specific patient population. Currently in phase II and actively recruiting, this trial aims to assess the objective response

rate (ORR) to the treatment. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, CTDNA, and RAS status.

NCT: NCT05962502 · Enrollment: 54 · Sites: Guangzhou, China

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

4 5

Becotatug Vedotin Combined With Cetuximab in the Later-line Treatment of Metastatic RAS Wild-type Colorectal Cancer

This clinical trial is testing the combination of becotatug vedotin and cetuximab for patients with metastatic RAS wild-type colorectal cancer who have already received prior treatments. The intervention involves the use of becotatug vedotin, a drug that targets cancer cells, alongside cetuximab, which is designed to inhibit tumor growth. This is a Phase 2 trial that is currently not yet recruiting, meaning patients cannot enroll at this time. Eligibility criteria include specific biomarker assessments such as BRAF, MSI, MMR, and RAS status.

NCT: NCT07490119 · Enrollment: 31 ·

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

4 5

Liposomal Irinotecan Plus Bevacizumab in Irinotecan-refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of liposomal irinotecan and bevacizumab in patients with irinotecan-refractory metastatic colorectal cancer. The intervention involves administering liposomal irinotecan, which is being evaluated for its efficacy and safety alongside bevacizumab, a drug that may enhance treatment outcomes. This study is in Phase 1 and Phase 2 and is currently not yet recruiting, meaning that patients will need to wait until recruitment begins to participate. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MMR, and RAS status.

NCT: NCT06434090 · Enrollment: 74 · Sites: Guangzhou, China

MEDIUM

NA

Stage IV

IMMUNOTHERAPY

• RECRUITING

4 5

A Randomized, Multicenter Phase II Basket Study of Hypofractionated Radiotherapy/Stereotactic Body Radiotherapy Followed by Immunotherapy-Based Systemic Therapy +/- L. Rhamnosus M9 for the First-Line Treatment of Advanced Digestive System Malignancies.

This clinical trial is testing the effectiveness of hypofractionated radiotherapy or stereotactic body radiotherapy followed by immunotherapy-based systemic therapy, with or without L. Rhamnosus M9, for patients with advanced digestive system malignancies. The intervention includes a combination of radiation therapy and various drugs, such as anti-PD-1 monoclonal antibodies and chemotherapy agents, which may enhance the immune response against tumors. Currently in the recruiting phase, this Phase II study aims to enroll 120 participants, indicating that eligible patients may have the opportunity to receive this integrated treatment approach. Specific eligibility criteria or biomarker requirements have not been detailed in the provided information.

NCT: NCT06349044 · Enrollment: 120 · Sites: Hangzhou, China

MEDIUM

PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

4 5

Dose Escalation and Expansion Clinical Trial of Irinotecan Liposome Combined With Oxaliplatin and 5-FU/LV Plus Bevacizumab as First-line Treatment of Metastatic Colorectal Cancer

This clinical trial is testing the combination of irinotecan liposome, oxaliplatin, 5-FU/LV, and bevacizumab as a first-line treatment for patients with advanced metastatic colorectal cancer. The intervention aims to determine the maximum tolerated dose and assess the safety and efficacy of this drug combination, which is significant for optimizing treatment regimens. Currently in Phase 1 and actively recruiting participants, this trial offers patients the opportunity to receive a potentially effective treatment while contributing to the understanding of dose-limiting toxicities. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06225622 · Enrollment: 78 · Sites: Shanghai, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

4 5

Efficacy and Safety of Chemotherapy With XELOX (Oxaliplatin + Capecitabine) and Bevacizumab in Combination With Adebrelimab in First-line Treatment of Microsatellite Stable (MSS) Initially Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of a combination treatment for patients with microsatellite stable (MSS) initially unresectable metastatic colorectal cancer. The intervention involves the use of Adebrelimab alongside chemotherapy agents Oxaliplatin and Capecitabine, as well as Bevacizumab, which may enhance treatment outcomes. Currently in Phase 2 and actively recruiting participants, this trial aims to assess progression-free survival as its primary outcome, providing insights into the effectiveness of this treatment regimen. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT06282445 · Enrollment: 36 · Sites: Jinhua, China

MEDIUM

PHASE4

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Survival Benefit of Compound Kushen Injection in the Treatment of Advanced Colorectal Cancer

This clinical trial is testing the effectiveness and safety of compound kushen injection in patients with advanced colorectal cancer. The intervention involves administering compound kushen injection alongside a first-line treatment scheme, which is significant for its potential impact on patient outcomes. Currently in Phase 4 and actively recruiting, this trial aims to provide insights into progression-free survival (PFS) for participants. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT05894694 · Enrollment: 320 · Sites: Beijing, China

MEDIUM

PHASE2

All Stages

KRAS

TARGETED THERAPY

• NOT_YET_RECRUITING

5

A Combination Therapy Including Anti-PD-1 Immunotherapy in MSS Rectal Cancer With Resectable Distal Metastasis

This clinical trial is testing a combination therapy that includes the anti-PD-1 immunotherapy tislelizumab for patients with microsatellite-stable (MSS) rectal cancer who have resectable distal metastasis. The intervention aims to evaluate the effectiveness of preoperative short-course radiotherapy followed by neoadjuvant chemotherapy and immunotherapy in improving tumor response and reducing postoperative metastasis and recurrence. This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MMR, DMMR, HER2, RAS, and microsatellite status.

NCT: NCT05359393 · Enrollment: 52 · Sites: Shanghai, China

MEDIUM

N/A

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

5

Retrospective Study on the Use of Immunotherapy in Patients With MSI-H Metastatic Colorectal Cancer

This clinical trial is a retrospective study examining the use of immunotherapy in patients with microsatellite instability-high (MSI-H) metastatic colorectal cancer (mCRC). The intervention involves the use of immune checkpoint inhibitors, specifically pembrolizumab and nivolumab, which are significant for their role in targeting the immune response against cancer cells. The trial is currently active but not recruiting, indicating that it is in the data collection phase, which may provide insights into treatment effectiveness for patients who have exhausted other options. Eligibility criteria include having MSI-H status, which is essential for determining the appropriateness of immunotherapy in this patient population.

NCT: NCT04612309 · Enrollment: 100 · Sites: Carpi, Italy

MEDIUM

NA

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

4

Clinical Efficacy of Transarterial Infusion Chemotherapy for Unresectable Colorectal Cancer

This clinical trial is testing the efficacy of transarterial infusion chemotherapy (TAIC) combined with either cetuximab or bevacizumab for patients with unresectable colorectal cancer (CRC). The intervention involves a procedure that delivers chemotherapy directly to the tumor site, which may enhance treatment effectiveness. The trial is currently in the recruiting phase, indicating that eligible patients can still enroll to participate in the study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT07333053 · Enrollment: 30 · Sites: Xicheng, China

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

4

A Phase II Study of Sintilimab Combined With Ipilimumab N01, Cetuximab and Dabrafenib in Patients With Microsatellite-Stable, BRAF V600E-Mutated Metastatic Colorectal Cancer

This clinical trial is testing the combination of sintilimab, ipilimumab N01, cetuximab, and dabrafenib in patients with microsatellite-stable, BRAF V600E-mutated metastatic colorectal cancer. The intervention involves a multi-drug regimen aimed at enhancing treatment efficacy for a patient population that typically has a poor prognosis with standard therapies. Currently in Phase II and actively recruiting participants, this trial seeks to assess the safety and efficacy of the treatment approach, which may offer an alternative for patients who have limited options. Eligibility criteria include the presence of BRAF mutations and microsatellite stability, which are essential for participation in this study.

NCT: NCT07506109 · Enrollment: 49 · Sites: Beijing, China, Chengdu, China, Tianjin, China, Wuhan, China

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 4

A Single-arm, Single-center, Phase II Clinical Study of Aipalolitovorelizumab (QL1706) Combined With Bevacizumab and Standard Chemotherapy as First-line Treatment for MSS/pMMR Metastatic Colorectal Cancer With BRAF V600E Mutation

This clinical trial is testing the combination of Aipalolitovorelizumab (QL1706) with bevacizumab and standard chemotherapy as a first-line treatment for patients with MSS/pMMR metastatic colorectal cancer that has a BRAF V600E mutation. The intervention involves administering Aipalolitovorelizumab alongside various chemotherapy regimens, which is significant for potentially improving treatment outcomes in this specific patient population. This is a Phase II study that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include having the BRAF mutation and meeting specific molecular marker requirements.

NCT: NCT07229846 · Enrollment: 30 · Sites: Harbin, China

SIGNAL

NA

Early Detection

RESEARCH

• RECRUITING

⚡ 4

Randomized Trial Comparing Fecal Testing (FIT) to Colonoscopy for Post-polypectomy Surveillance

This clinical trial is testing the effectiveness of fecal occult blood testing (FIT) compared to colonoscopy for post-polypectomy surveillance in patients who have had colorectal polyps removed. The intervention involves using FIT as a less burdensome alternative to colonoscopy, which is important given the limited availability of colonoscopy and the fact that many surveillance procedures yield no findings. The trial is currently in the recruiting phase with an enrollment goal of 4,000 participants, which means eligible patients can still join the study. Eligibility criteria may include a history of polyp removal, but specific requirements are not detailed in the provided information.

NCT: NCT06983639 · Enrollment: 4000 · Sites: Arendal, Norway

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 4

A Phase Ib/II Trial of HS-20110 Combination Therapies in Advanced Colorectal Cancer Patients.

This clinical trial is evaluating the safety and efficacy of HS-20110 combination therapies in patients with advanced colorectal cancer. The interventions include HS-20110 combined with Bevacizumab and various chemotherapy regimens, which is important for assessing potential treatment options for this patient population. The trial is currently in Phase Ib/II and is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as BRAF and MSI, which may be relevant for patient selection.

NCT: NCT07283367 · Enrollment: 502 ·

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

IMMUNOTHERAPY

• RECRUITING

⚡ 4

A FIH, Phase I/IIa, Trial Assessing Feasibility of Administrations of TIL-based Immunotherapy in Patients With Metastatic CRC and PC

This clinical trial is testing the feasibility of TIL-based immunotherapy in patients with metastatic colorectal cancer (CRC) and pancreatic cancer (PC). The intervention involves the administration of a novel autologous TIL product, CC-38, along with pembrolizumab, cyclophosphamide, interleukin-2, and uromitexan, which is important for evaluating safety and tolerability. Currently in Phase I/IIa and actively recruiting, this trial aims to assess the repeated administration of CC-38 and its implications for patient treatment. Eligibility criteria include the presence of the BRAF biomarker.

NCT: NCT07255664 · Enrollment: 12 · Sites: Frankfurt, Germany

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

⚡ 4

A Phase II Trial of Neoadjuvant PD-1 Vaccine PD1-Vaxx in Operable MSI High Colorectal Cancer

This clinical trial is testing the neoadjuvant PD-1 vaccine IMU-201 (PD1-Vaxx) in patients with operable microsatellite instability-high (MSI-H) colorectal cancer. The intervention involves administering three doses of the PD1-Vaxx prior to surgical resection, with the goal of achieving a Major Pathological Response (MPR) defined as less than 10% viable tumor cells. This is a Phase II trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the study's findings. Eligibility criteria include having operable MSI-H colorectal cancer, with specific biomarker requirements such as MSI, MSI-H, MMR, and NGS.

NCT: NCT06692959 · Enrollment: 44 · Sites: Adelaide, Australia, Guildford, United Kingdom, Manchester, United Kingdom, Perth, Australia

SIGNAL **NA** **Stage IV** **RESEARCH** • **RECRUITING**

⚡ 4

Randomized Controlled Trial of the Effects of Combined Resistance and Aerobic Exercise on Health-related Quality of Life in Patients Undergoing First-line Chemotherapy for Metastatic Colorectal Cancer (REACH)

The REACH trial is testing the effects of combined resistance and aerobic exercise on health-related quality of life in patients undergoing first-line chemotherapy for metastatic colorectal cancer. The intervention involves 18 weeks of structured exercise training, which aims to assess its impact on the quality of life for these patients. This trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate. Eligibility criteria include being diagnosed with unresectable metastatic colorectal cancer and undergoing first-line chemotherapy.

NCT: NCT06938971 · Enrollment: 150 · Sites: Copenhagen, Denmark, Herlev, Denmark, Hillerød, Denmark, Roskilde, Denmark

MEDIUM **PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **NOT_YET_RECRUITING**

⚡ 4

Short-course Radiotherapy Followed by CAPOX and Ivonescimab for Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy and safety of preoperative short-course radiotherapy combined with ivonescimab and chemotherapy in patients with locally advanced rectal cancer. The intervention involves the drug ivonescimab, which is being evaluated for its potential to improve treatment outcomes when used alongside standard therapies. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06760520 · Enrollment: 40 ·

MEDIUM **NA** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **RECRUITING**

⚡ 4

Study on the Safety and Efficacy of MR-Linac Technique in Patients With Unresectable Locally Advanced Colon Cancer

This clinical trial is testing the safety and efficacy of the MR-Linac technique in patients with unresectable locally advanced colon cancer. The intervention involves the use of MR-Linac for short course radiotherapy, followed by chemotherapy and surgical resection, which is significant for potentially improving treatment outcomes in this patient population. The study is currently in the recruiting phase and aims to enroll 20 participants, indicating that eligible patients may still join the trial. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06244537 · Enrollment: 20 · Sites: Chengdu, China

MEDIUM **PHASE1, PHASE2** **Stage IV** **KRAS** **TARGETED THERAPY** • **RECRUITING**

⚡ 4

Study of Elironrasib and Daraxonrasib as Monotherapies and Combination Therapy in Participants With Advanced KRAS G12C Mutant Solid Tumors

This clinical trial is evaluating the safety and tolerability of Elironrasib and Daraxonrasib, both as monotherapies and in combination, for participants with advanced KRAS G12C mutant solid tumors. The intervention involves two drugs, Elironrasib and Daraxonrasib, which are being tested to understand their pharmacokinetics and potential effects on this specific mutation. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive these treatments. Eligibility criteria include having tumors with KRAS G12C mutations, among other potential biomarker requirements.

NCT: NCT06128551 · Enrollment: 534 · Sites: Aurora, Boston, Dallas, Detroit, Duarte

MEDIUM **PHASE2** **Stage III** **MSI/MMR** **TARGETED THERAPY** • **ACTIVE_NOT_RECRUITING**

⚡ 4

Atezolizumab in Patients With MSI-h/MMR-D Stage II High Risk and Stage III Colorectal Cancer Ineligible for Oxaliplatin

This clinical trial is testing the efficacy of atezolizumab in patients with stage II high-risk and stage III colorectal cancer who are ineligible for oxaliplatin-based chemotherapy. The intervention involves administering atezolizumab, a drug that targets the immune system, alongside IMM-101, to potentially enhance disease-free survival in this specific patient population. Currently in Phase 2 and marked as active but not recruiting, this trial indicates that patients may not be able to enroll at this time but can still benefit from the findings once available. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS, which are essential for determining patient suitability for the treatment.

NCT: NCT05118724 · Enrollment: 80 · Sites: Essen, Germany

MEDIUM

PHASE1, PHASE2

Stage IV

IMMUNOTHERAPY

• ACTIVE_NOT_RECRUITING

⚡ 4

Safety, PK and Efficacy of ONC-392 in Monotherapy and in Combination of Anti-PD-1 in Advanced Solid Tumors and NSCLC

This clinical trial is testing the safety, pharmacokinetics, and efficacy of ONC-392, a humanized anti-CTLA4 IgG1 monoclonal antibody, in patients with advanced or metastatic solid tumors and non-small cell lung cancer (NSCLC). The intervention involves administering ONC-392 both as a monotherapy and in combination with pembrolizumab and docetaxel, which is significant for understanding its potential effectiveness in these cancer types. Currently, the trial is in Phase 1 and Phase 2 and is active but not recruiting, indicating that patients cannot enroll at this time. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

NCT: NCT04140526 · Enrollment: 733 · Sites: Ann Arbor, Atlanta, Atlantis, Aurora, Baltimore

MEDIUM

PHASE2

Stage IV

PHASE II TRIAL

• ACTIVE_NOT_RECRUITING

⚡ 4

Microwave Ablation Versus Stereotactic Body Radiotherapy for Colorectal Liver Metastases in Oligometastatic Disease: a Prospective, Randomised, Phase 2 Trial

The LAVA-CRLM trial is testing the effectiveness of microwave ablation (MWA) compared to stereotactic body radiotherapy (SBRT) in patients with colorectal liver metastases and oligometastatic disease. The intervention involves using MWA, a device-based treatment, and SBRT, a form of radiation therapy, to assess their impact on local control and safety in this patient population. This is a Phase 2 trial that is currently active but not recruiting, indicating that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria for participation may include specific characteristics related to the patient's metastatic disease status.

NCT: NCT03654131 · Enrollment: 100 · Sites: Copenhagen, Denmark

Appendix A — Patient Guide to Active Trials

For patients, caregivers, and family members. Always discuss with your oncologist before considering a clinical trial.

What is a clinical trial?

Clinical trials test new treatments, combinations, or approaches to care. Participating can give you access to treatments not yet widely available. Trials are listed by the biomarker your tumor must have — because many newer treatments only work for specific mutations. Ask your oncologist which biomarkers your tumor has been tested for, then look for matching trials below.

MSI/MMR

Botensilimab + Balstilimab vs Best Supportive Care as Therapy in Chemo-refractory, Unresectable, Colorectal Adenocarcinoma

Phase: PHASE3 · Status: RECRUITING · [NCT07152821](#)

This clinical trial is testing the combination of botensilimab and balstilimab in patients with chemo-refractory, unresectable colorectal adenocarcinoma. The intervention involves two immunotherapy drugs aimed at potentially improving overall survival and quality of life for participants. Currently in Phase 3 and actively recruiting, this trial may provide insights into the effectiveness and safety of these treatments for eligible patients. Participants may need to meet specific biomarker criteria, including MSI, MSI-H, MMR, DMMR, and microsatellite status.

KRAS

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

Phase: PHASE3 · Status: RECRUITING · [NCT06750094](#)

This clinical trial is testing the efficacy of amivantamab combined with FOLFIRI chemotherapy in patients with KRAS/NRAS and BRAF wild-type colorectal cancer who have previously undergone chemotherapy. The intervention involves the biological agent amivantamab alongside standard chemotherapy agents, which is important for evaluating its potential benefits compared to existing treatment options. This is a Phase 3 trial currently recruiting 700 participants, which means that eligible patients may have the opportunity to receive a treatment that is being compared to standard therapies in a larger population. Eligibility criteria include specific biomarker requirements related to KRAS, NRAS, and BRAF status, as well as other factors such as microsatellite instability (MSI) and mismatch repair (MMR) status.

BRAF

Intermittent or Continuous Panitumumab Plus FOLFIRI for Left Sided RAS/B-RAF Wild-type Metastatic Colorectal Cancer

Phase: PHASE3 · Status: RECRUITING · [NCT06509126](#)

This clinical trial is testing the effectiveness of intermittent versus continuous administration of Panitumumab combined with FOLFIRI in patients with left-sided RAS/B-RAF wild-type metastatic colorectal cancer. The intervention involves the use of Panitumumab, Irinotecan, 5-fluorouracil, and L-folinic acid, which are important for evaluating treatment strategies that may optimize patient outcomes. Currently in Phase 3 and actively recruiting, this trial aims to determine if the intermittent regimen can achieve a Time to Treatment Failure comparable to the standard continuous regimen, which is relevant for patients considering their treatment options. Eligibility criteria include specific biomarker requirements related to BRAF, MMR, and RAS status.

RAS

Neoadjuvant FOLFOX Chemotherapy for Patients With Locally Advanced Colon Cancer

Phase: PHASE3 · Status: ACTIVE_NOT_RECRUITING · [NCT03426904](#)

This clinical trial is testing the effectiveness of neoadjuvant FOLFOX chemotherapy in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy before surgery, which may enhance treatment outcomes by targeting microscopic metastatic cancer cells earlier and reducing surgical complications. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and results may soon be available for clinical application. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

HER2

A Phase 1/1b Study of IAM1363 in HER2 Cancers

Phase: PHASE1 · Status: RECRUITING · [NCT06253871](#)

IAM1363 is being investigated in a Phase 1/1b trial for patients with HER2 mutation-related tumors, including HER2-positive colorectal cancer. The trial is currently recruiting participants and aims to assess the safety and tolerability of IAM1363. Primary endpoints include determining the maximum tolerated dose and evaluating pharmacokinetics.

ctDNA

A Phase II Clinical Trial Comparing the Efficacy of RO7198457 Versus Watchful Waiting in Patients With ctDNA-positive, Resected Stage II (High Risk) and Stage III Colorectal Cancer

Phase: PHASE2 · Status: RECRUITING · [NCT04486378](#)

RO7198457 is being compared to watchful waiting in a Phase II trial for patients with ctDNA-positive, resected Stage II (high risk) and Stage III colorectal cancer. The trial is currently recruiting participants. The primary endpoint is not specified.

NGS

ULTRA-high-risk Surveillance to Avoid Future Events: the ULTRA-SAFE Trial

Phase: NA · Status: NOT_YET_RECRUITING · [NCT07378930](#)

The ULTRA-SAFE Trial will utilize early triage with fecal immunochemical testing for patients at ultra-high risk of colorectal neoplasms. This study focuses on colorectal cancer prevention and surveillance colonoscopy. The primary endpoint is not specified. The trial is currently not yet recruiting participants.

What this means for you

Ask your oncologist whether any active trials match your biomarker profile and disease stage. The most actively trialed biomarkers in colorectal cancer right now are **MSI/MMR, KRAS, BRAF, RAS, and HER2**. If your tumor has not been tested for these markers, ask your oncologist before your next treatment decision — biomarker results can determine which trials you qualify for.

Questions to ask your oncologist about trials

"Am I eligible for any active Phase III trials based on my biomarker profile?"

"Does my tumor need to be retested before I can qualify for a biomarker-directed trial?"

"What are the travel requirements and visit frequency for the trials I might qualify for?"

Source: ClinicalTrials.gov · 2026-04-12 15:18:12

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Clinical Trial Reference Guide

What the phases mean

Phase I — First testing in humans. Focus on safety and dosing. Small groups.

Phase II — Testing effectiveness. Larger groups. Determines if Phase III is warranted.

Phase III — Randomized comparison against standard treatment. Required for FDA approval.

Key biomarkers that determine trial eligibility

RAS (KRAS/NRAS) — mutation status affects anti-EGFR eligibility and KRAS G12C trial eligibility

BRAF V600E — required for encorafenib-based trials (~8–10% of CRC)

MSI-H/dMMR — required for immunotherapy trials; ~5% of mCRC, ~15% of Stage II/III

HER2 — required for HER2-directed trials (~2–3% of CRC)

NTRK fusion — required for TRK inhibitor trials (<1% of CRC)

KRAS G12C — required for sotorasib/adagrasib trials (~3–4% of CRC)

TMB-H — tumor mutational burden; may qualify for immunotherapy

ctDNA — liquid biopsy; used in MRD monitoring trials

How to find trials you may qualify for

1. Get complete biomarker testing: RAS, BRAF, MSI/MMR, HER2, NTRK, TMB
2. Search ClinicalTrials.gov with your biomarker and stage
3. Ask your oncologist about NCI-designated cancer centers with trial access
4. Contact the Colorectal Cancer Alliance: [ccalliance.org](#)

Key Resources

Colon & Colorectal Cancer — Clinical Trials Intelligence

About This Report

This report is generated automatically from publicly available medical databases including PubMed, ClinicalTrials.gov, and the FDA Drug@FDA database. Entries are classified using deterministic rule-based logic and scored by clinical significance. AI-generated summaries are grounded exclusively in the source data provided.

Limitations

This report does not represent a complete picture of all published research or approved treatments. Database coverage, feed availability, and relevance filtering affect what appears. New drug approvals and trial status changes may not be reflected immediately. Always verify current information at the original source before making clinical decisions.

Disclaimer

Trial data is sourced from ClinicalTrials.gov, a registry maintained by the U.S. National Library of Medicine. Entry summaries are AI-generated from publicly available trial records. Trial status and eligibility criteria may change — always verify at ClinicalTrials.gov before acting on this information. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

Copyright

© 2026 Randy Hinton. All rights reserved. This report may be shared freely for informational purposes. It may not be modified, republished, or presented as original work without attribution.

https://petroai1492.github.io/delta/timelines/coloncancer_clinicaltrials_timeline.pdf