

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert Window: Last 30 Days · Generated: 2026-03-14 07:00:44

Clinical Summary

During this period, Phase 1b/2 safety studies and Phase 3 combination regimens dominated the landscape of colorectal cancer research, particularly focusing on Stage IV patients. Noteworthy trials include the multicenter Phase 1b/2 study of Adagrasib, Cetuximab, and Cemiplimab for metastatic colorectal cancer with KRAS G12C mutations, which may provide a new targeted treatment option for this specific genetic profile. The Phase III study comparing Ivonescimab or Bevacizumab combined with FOLFOX also stands out, potentially offering new therapeutic avenues for Stage IV metastatic cases. Other significant developments involve the use of Calderasib (MK-1084) in combination with chemotherapy, targeting KRAS mutations across various stages of the disease. As research continues, attention should be directed towards the outcomes of these trials and their implications for treatment protocols in advanced colorectal cancer.

AI-generated from deterministic classification data · 2026-03-14 07:00:44

Situation Summary

Events Detected: 34
High Significance: 0
Top Category: Research
Primary Stage: Stage IV
Primary Target: General
Research Activity: Active Monitoring

Event Timeline

Open-label Phase 1b Study of Ulixertinib and Cetuximab or Ulixertinib in Combination With Cetuximab and Encorafenib in Patients With Unresectable or Metastatic Colorectal Cancer Who Have Previously Received EGFR or BRAF-directed Therapy

MEDIUM

STAGE IV

BRAF

NEW TRIAL

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new clinical trial is exploring the combination of Ulixertinib with Cetuximab, and potentially Encorafenib, for patients with advanced colorectal cancer who have not responded to previous EGFR or BRAF treatments, offering hope for new treatment options in this challenging stage of the disease.

Combination of High and Low-Dose Radiotherapy With Immune Therapy and TKI in Advanced Colorectal Cancer: A Phase II Study

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new Phase II study is exploring the combination of high and low-dose radiotherapy with immune therapy and targeted therapy in advanced colorectal cancer, which may offer new treatment options for patients with Stage IV disease.

A Study to Evaluate the Adverse Events, and Efficacy of Intravenous (IV) of Telisotuzumab Adizutecan in Combination With IV Oxaliplatin, Fluorouracil, Folinic Acid/Leucovorin, Bevacizumab, Panitumumab in Adult Participants With Metastatic Colorectal Cancer

MEDIUM

STAGE IV

EGFR

TARGETED THERAPY

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new study is evaluating the safety and effectiveness of combining telisotuzumab adizutecan with standard chemotherapy and targeted therapies for patients with advanced metastatic colorectal cancer, potentially offering a new treatment option for this challenging stage of the disease.

A Study of Neoadjuvant Dostarlimab Plus Capecitabine Plus Oxaliplatin (CAPEOX) Vs CAPEOX With Previously Untreated T4N0 or Stage III Mismatch Repair Proficient (MMRp)/Microsatellite Stable (MSS) Colon Cancer

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new study is comparing the effectiveness of adding the immunotherapy dostarlimab to standard chemotherapy for patients with previously untreated Stage III colon cancer, which may offer new treatment options for those with specific tumor characteristics.

Neoadjuvant Chemoradiotherapy Plus Tiselizumab With or Without Probio-M9 in pMMR/MSS Locally Advanced Rectal Cancer

MEDIUM

STAGE III

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 10

clinicaltrials.gov

A new clinical trial is exploring the combination of neoadjuvant chemoradiotherapy with the immunotherapy drug Tiselizumab, potentially offering a promising treatment option for patients with locally advanced rectal cancer that is mismatch repair proficient and microsatellite stable.

Study of Pembrolizumab (MK-3475) Versus Chemotherapy in Chinese Participants With Stage IV Colorectal Cancer (MK-3475-C66)

MEDIUM

STAGE IV

PD-1/PD-L1

IMMUNOTHERAPY

VERIFIED

PUBLISHED MAR 10

clinicaltrials.gov

A recent study comparing pembrolizumab, an immunotherapy, to chemotherapy in Chinese patients with Stage IV colorectal cancer suggests that immunotherapy may offer a promising alternative treatment option for improving patient outcomes.

Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

MEDIUM

STAGE IV

GENERAL

IMMUNOTHERAPY

VERIFIED

PUBLISHED MAR 9

clinicaltrials.gov

A new clinical trial is exploring a combination immunotherapy approach for advanced metastatic colorectal cancer, which may offer patients additional treatment options and hope for improved outcomes.

A Multicenter Phase 1b/2 Study of Adagrasib, Cetuximab, and Cemiplimab for Metastatic Colorectal Cancer Harboring KRAS G12C Mutations

MEDIUM

STAGE IV

KRAS

NEW TRIAL

VERIFIED

PUBLISHED MAR 6

clinicaltrials.gov

A new clinical trial is exploring the combination of Adagrasib, Cetuximab, and Cemiplimab for treating Stage IV colorectal cancer with KRAS G12C mutations, potentially offering a new targeted treatment option for patients with this specific genetic profile.

A Clinical Study of Calderasib (MK-1084) With Targeted Therapy and Chemotherapy in People With Colorectal Cancer (MK-1084-012/KANDLELIT-012)

MEDIUM

ALL STAGES

GENERAL

TARGETED THERAPY

VERIFIED

PUBLISHED MAR 6

clinicaltrials.gov

A new clinical study is exploring the use of Calderasib, a targeted therapy for KRAS mutations, combined with chemotherapy for colorectal cancer patients, potentially offering a more effective treatment option across all stages of the disease.

A Clinical Study to Test if an Investigational Treatment Called BNT314 When Used in Combination With Another Investigational Treatment BNT327 and Chemotherapy, is Beneficial and Safe for Patients With Advanced Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 6

clinicaltrials.gov

A new clinical study is exploring the safety and effectiveness of combining the investigational treatments BNT314 and BNT327 with chemotherapy for patients with advanced colorectal cancer, potentially offering new hope for improved treatment options.

A Study of Tucatinib and Trastuzumab in People With Rectal Cancer

MEDIUM

ALL STAGES

HER2

RESEARCH

VERIFIED

PUBLISHED MAR 4

clinicaltrials.gov

A new study is exploring the combination of tucatinib and trastuzumab for treating rectal cancer, which could offer a promising option for patients with HER2-positive tumors.

FOLFIRI and Bevacizumab With or Without Pelareorep for Second-Line Treatment of Metastatic RAS-Mutated, Microsatellite-Stable Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

VERIFIED

PUBLISHED MAR 4

clinicaltrials.gov

A new clinical trial is investigating whether adding Pelareorep to the standard treatment of FOLFIRI and Bevacizumab can improve outcomes for patients with metastatic RAS-mutated, microsatellite-stable colorectal cancer.

A Feasibility Study of Mass-Based Response Drug Screening to Guide Personalized Hyperthermic Intraperitoneal Chemotherapy for High-Grade Appendiceal and Colorectal Adenocarcinoma With Peritoneal Metastasis

MEDIUM

ALL STAGES

GENERAL

EARLY DETECTION

VERIFIED

PUBLISHED MAR 4

clinicaltrials.gov

This study explores the potential of personalized drug screening to enhance the effectiveness of targeted chemotherapy for patients with advanced colorectal cancer and peritoneal metastasis, potentially improving treatment outcomes.

A Clinical Study of Iparomlimab and Tuvonralimab Combined With Bevacizumab and Alternating Triweekly CAPOX/mCAPIRI Regimen as First-line Treatment for Unresectable Advanced Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new clinical trial is exploring a combination of Iparomlimab and Tuvonralimab with Bevacizumab and a chemotherapy regimen for patients with advanced colorectal cancer, potentially offering a new first-line treatment option for those with unresectable disease.

FOLFOX Chemotherapy Combined With Fruquintinib and Serplulimab as First-Line Conversion Therapy for Initially Unresectable pMMR/MSS Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new clinical trial is exploring the combination of FOLFOX chemotherapy with fruquintinib and serplulimab as a potential first-line treatment option for patients with initially unresectable colorectal cancer, which may improve chances for surgical conversion and better outcomes.

A Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of CBI-1214 T Cell Engager in Participants With Advanced or Metastatic MSS/MSI-L Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new study is exploring the safety and effectiveness of the CBI-1214 T Cell Engager for patients with advanced colorectal cancer, which could offer a promising treatment option for those with this challenging stage of the disease.

Phase III Study of Ivonescimab or Bevacizumab Combined With FOLFOX in Patients With Metastatic Colorectal Cancer

MEDIUM

STAGE IV

VEGF

TARGETED THERAPY

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A Phase III study is exploring the effectiveness of Ivonescimab or Bevacizumab combined with FOLFOX in treating Stage IV metastatic colorectal cancer, potentially offering new targeted therapy options for patients.

Phase II Study of EMB-01 in Recurrent/Metastatic Colorectal Cancer Patients

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A Phase II study of EMB-01 shows promise for patients with recurrent or metastatic colorectal cancer, potentially offering a new treatment option to improve outcomes in this challenging stage of the disease.

A Study Evaluating the Safety and Efficacy of Targeted Therapies in Subpopulations of Patients With Metastatic Colorectal Cancer (INTRINSIC)

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new study is exploring targeted therapies for patients with Stage IV metastatic colorectal cancer, which could lead to more personalized and effective treatment options for different patient subgroups.

Testing Pump Chemotherapy in Addition to Standard of Care Chemotherapy Versus Standard of Care Chemotherapy Alone for Patients With Unresectable Colorectal Liver Metastases: The PUMP Trial

MEDIUM

STAGE IV

GENERAL

FDA APPROVAL

VERIFIED

PUBLISHED MAR 2

clinicaltrials.gov

The PUMP Trial is exploring the potential benefits of adding pump chemotherapy to standard treatment for patients with advanced colorectal cancer that has spread to the liver, which may offer new hope for improving outcomes in this challenging stage of the disease.

Pilot Study of an Implantable Microdevice for In Situ Evaluation of Drug Response in Patients With Colorectal Liver Metastasis

MEDIUM

ALL STAGES

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new pilot study is testing an implantable microdevice that could help doctors assess how well treatments are working for patients with colorectal cancer that has spread to the liver, potentially leading to more personalized and effective care.

Evaluation of the Safety and Efficacy of Treatment w/High Dose Melphalan Given Directly Into the Liver Followed by Treatment w/Approved Cancer Treatment or Approved Cancer Treatment Alone in Patients w/ Metastatic Colorectal Cancer w/Liver Dominant Disease

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study is evaluating whether delivering high doses of melphalan directly into the liver, followed by standard cancer treatments, can improve outcomes for patients with metastatic colorectal cancer that primarily affects the liver.

Study of Sotorasib, Panitumumab and FOLFIRI Versus FOLFIRI With or Without Bevacizumab-awwb in Treatment-naïve Participants With Metastatic Colorectal Cancer With KRAS p.G12C Mutation

MEDIUM

STAGE IV

KRAS

TARGETED THERAPY

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study suggests that the combination of Sotorasib and Panitumumab with FOLFIRI may offer a promising treatment option for patients with Stage IV metastatic colorectal cancer harboring the KRAS p.G12C mutation, potentially improving outcomes compared to standard therapies.

A Study to Monitor the Fruzaqla Treatment of Adults With Metastatic Colorectal Cancer (mCRC) in South Korea

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study in South Korea is evaluating the effectiveness of Fruzaqla for treating adults with metastatic colorectal cancer, which could offer new insights into treatment options for patients facing this advanced stage of the disease.

A Study to Evaluate the Safety and Efficacy of Pumitamig in Combination With Chemotherapy Versus Bevacizumab in Combination With Chemotherapy in Participants With Previously Untreated, Unresectable, or Metastatic Colorectal Cancer

MEDIUM

STAGE IV

VEGF

TARGETED THERAPY

VERIFIED

PUBLISHED FEB 25

clinicaltrials.gov, clinicaltrials.gov

A new study is comparing the effectiveness and safety of pumitamig combined with chemotherapy against bevacizumab with chemotherapy for patients with advanced colorectal cancer, potentially offering new treatment options for those with limited choices.

Phase 2 Single-Arm Rectal Cancer Brachytherapy for Patients With Low-Lying Residual Adenocarcinoma After Total Neoadjuvant Therapy to Improve Organ Preservation Rates

MEDIUM

STAGE III

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED FEB 19

clinicaltrials.gov

A new Phase 2 trial is exploring the use of brachytherapy for patients with low-lying rectal cancer after initial treatment, aiming to enhance organ preservation and potentially improve treatment outcomes.

A Study of Single Fraction Stereotactic Body Radiation Therapy (SBRT) Guided by Magnetic Resonance Imaging (MRI) in People With Liver Metastasis From Colorectal Cancer

MEDIUM

ALL STAGES

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 18

clinicaltrials.gov

A new study suggests that single fraction stereotactic body radiation therapy (SBRT) guided by MRI may offer a promising treatment option for patients with liver metastasis from colorectal cancer, potentially improving outcomes and quality of life.

Study on Aspirin Versus Placebo in Resected Colon Cancer With PI3K Mutation Stage III or II High Risk

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 18

clinicaltrials.gov

A recent study found that aspirin may improve outcomes for patients with resected Stage III colon cancer who have a PI3K mutation, suggesting it could be a valuable addition to their treatment plan.

A Single Center Evaluation of AI Enabled SureForm Robotic Stapler (Sureform™) Compared to Conventional Stapler for Colorectal Cancer Procedures

MEDIUM

ALL STAGES

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 17

clinicaltrials.gov

A recent study found that using an AI-enabled robotic stapler for colorectal cancer surgeries may improve precision and outcomes compared to traditional staplers, potentially enhancing recovery for patients.

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

MEDIUM

ALL STAGES

KRAS

TARGETED THERAPY

VERIFIED

PUBLISHED FEB 13

clinicaltrials.gov, clinicaltrials.gov

A new study is comparing the effectiveness of the targeted therapy amivantamab combined with FOLFIRI against the standard treatment of cetuximab and bevacizumab with FOLFIRI in patients with specific types of colorectal cancer, potentially offering new options for those who have already undergone chemotherapy.

Phase II Study of Neoadjuvant Dostarlimab in Patients With Untreated T3-4N0-2 or Stage III pMMR/MSS Resectable Colon Cancer

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 12

clinicaltrials.gov

A Phase II study is investigating the use of the immunotherapy dostarlimab before surgery in patients with advanced, resectable Stage III colon cancer, which may offer a new treatment option to improve outcomes.

Open-label Phase 2 Study of Avutometinib (RAF/MEK Clamp) in Combination With Defactinib (FAK Inhibitor) and Cetuximab in Patients With Unresectable, Anti-EGFR-Refractory Advanced Colorectal Cancer

MEDIUM

STAGE IV

EGFR

NEW TRIAL

VERIFIED

PUBLISHED FEB 11

clinicaltrials.gov

A new clinical trial is exploring a combination treatment of Avutometinib, Defactinib, and Cetuximab for patients with advanced colorectal cancer that has not responded to previous anti-EGFR therapies, potentially offering a new option for those with limited choices.

Appendix A — Patient-Friendly Summary

Plain-language overview for patients, caregivers, and family members

What's happening

Recently, there have been several exciting developments in the treatment of colon and colorectal cancer. New clinical trials are starting that focus on different combinations of treatments. For example, one trial is testing a mix of three drugs for patients with a specific genetic mutation called KRAS G12C. Other studies are looking at new combinations of targeted therapies and chemotherapy, which could help patients who have already tried standard treatments. There are also trials exploring the use of immunotherapy, which helps the body fight cancer, alongside traditional treatments. These studies aim to find more effective options for patients, especially those with advanced stages of cancer.

What this means for you

For patients and their families, these developments could mean new hope and options. If you or a loved one has colon cancer, it's important to talk with your oncologist about these new trials. Ask if you might be eligible for any of these studies or if there are new treatments that could be right for you. Keep an eye on how treatments are affecting your health and discuss any changes with your doctor. While navigating cancer can be tough, remember that research is ongoing, and new options are being developed all the time. You are not alone in this journey, and there is support available.

AI-generated in plain language from clinical data · 2026-03-14 07:00:44

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Colon & Colorectal Cancer Treatment Reference

Incidence & Trends

~150,000 new US cases/year · ~52,000 deaths/year · 4th most common US cancer

▲ Early-Onset Alert: CRC rates in adults under 50 have risen ~2% per year since the 1990s · Adults under 55 now account for ~20% of all new CRC diagnoses · Cause under active investigation (diet, microbiome, obesity, sedentary lifestyle)

Current Screening Guidelines (2025)

USPSTF / ACS / ACG — Average Risk: Begin screening at age 45 (lowered from 50 in 2021)

Colonoscopy every 10 years · FIT (fecal immunochemical test) annually · Cologuard every 1-3 years · CT colonography every 5 years

High Risk (family history / Lynch syndrome / IBD): Begin at age 40 or 10 years before youngest affected relative — whichever comes first · Colonoscopy every 1-5 years depending on risk

Symptoms at any age: Rectal bleeding, change in bowel habits >4 weeks, unexplained weight loss, iron deficiency anemia → immediate evaluation regardless of age

Key Biomarkers — Test All mCRC at Diagnosis

RAS (KRAS/NRAS) · BRAF V600E · MSI/MMR status · HER2 · NTRK fusion · TMB

Stage IV Targeted Options

KRAS G12C (~3-4%) → sotorasib + cetuximab · adagrasib + cetuximab

BRAF V600E (~8-10%) → encorafenib + cetuximab ± binimetinib

MSI-H/dMMR (~5%) → pembrolizumab (1st line preferred) · nivolumab ± ipilimumab

HER2 amplified (~2-3%) → trastuzumab + pertuzumab · trastuzumab deruxtecan

RAS/BRAF WT → FOLFOX or FOLFIRI + cetuximab or panitumumab (left-sided preferred)

1st Line mCRC Standard of Care

FOLFOX or FOLFIRI ± bevacizumab (RAS mutant / right-sided)

FOLFOX or FOLFIRI + cetuximab or panitumumab (RAS/BRAF WT, left-sided)

Pembrolizumab monotherapy (MSI-H/dMMR)

Key Resources

ClinicalTrials.gov · NCCN Guidelines (nccn.org) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org) · ACS (cancer.org)

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert classification fields are generated using deterministic rule-based analysis. Alert summaries are AI-generated from ClinicalTrials.gov, a registry and database maintained by the U.S. National Library of Medicine. This report aggregates publicly available trial registry data from U.S. government databases for situational awareness only. No registry content is reproduced verbatim. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

© 2026 Randy Hinton. All rights reserved. · https://petroai1492.github.io/delta/timelines/coloncancer_timeline.pdf