Colon & Colorectal Cancer — Clinical Trials Intelligence

Alert Window: Last 30 Days · Generated: 2026-11-09 07:30:00

Situation Summary

Phase Iii: 240 trials
Recruiting: 1309 trials
Biomarker: 1137 biomarker-directed trials
Breakthrough: 568 trials scored ≥ 4

Active Trial Timeline

Showing 568 high-significance trials (score ≥ 4). 1685 additional lower-scored trials are collapsed below. Phase III trials appear first. Click any title to view full trial details on ClinicalTrials.gov.

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 11

Anti-PCSK9 Antibody Tafolecimab and Anti-PD-1 Antibody Sintilimab Combined With Neoadjuvant Chemoradiotherapy for pMMR/ MSS Locally Advanced Rectal Cancer : A Prospective, Multicenter, Randomized, Open-Label, Parallel-Controlled Trial

NCT: NCT07686796 · Enrollment: 148 · Sites: Guangzhou, China

MEDIUM NA Stage IV MSI/MMR LIQUID BIOPSY • NOT_YET_RECRUITING 11

A Multicenter Randomized Open-Label Trial Evaluating ctDNA-Guided Interruption Versus Standard of Care Immune-Checkpoint Inhibitor (ICI) Therapy In Patients With Advanced / Metastatic Solid Tumors.

This clinical trial is evaluating the effectiveness of ctDNA-guided interruption of immune-checkpoint inhibitor (ICI) therapy compared to standard of care continuous ICI therapy in patients with advanced or metastatic solid tumors, including colorectal cancer. The intervention involves the Signatera Genome ultra-sensitive ctDNA blood test, which is significant for its potential to personalize treatment based on tumor dynamics. The trial is currently not yet recruiting participants and aims to enroll 920 patients, which means that patients interested in this study will need to wait until recruitment begins. Eligibility criteria include patients with histologically confirmed advanced/metastatic non-small cell lung cancer, melanoma, MSI-High/dMMR colorectal cancer, renal cell carcinoma, and other solid tumors.

NCT: NCT07689812 · Enrollment: 920 ·

MEDIUM PHASE3 All Stages MSI/MMR LIQUID BIOPSY • RECRUITING 11

Zanzalintinib Alone or in Combination With Pembrolizumab and Berahyaluronidase Alfa in Participants With Colorectal Cancer With Molecular Residual Disease

NCT: NCT07750158 · Enrollment: 600 · Sites: Bridgewater, Council Bluffs, East Brunswick, Edison, Grand Island

MEDIUM PHASE3 Stage IV KRAS TARGETED THERAPY • RECRUITING 11

Study of Petosemtamab Plus Chemotherapy Versus Cetuximab Plus Chemotherapy in RAS and BRAF Wild-type, Unresectable or Metastatic, Left-sided Colorectal Cancer (LiGeR-CRC1)

NCT: NCT07702032 · Enrollment: 960 · Sites: Hot Springs, Indianapolis, Los Angeles, Olympia, Savannah

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Efficacy of the Use of Neoadjuvant With/Without Hyperthermic Intraperitoneal Chemotherapy in the Treatment of Locally Advanced Colon Cancer

This clinical trial is testing the efficacy of neoadjuvant treatment strategies, with or without hyperthermic intraperitoneal chemotherapy (HIPEC), in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy followed by cytoreductive surgery and may include mitomycin C as part of the HIPEC procedure, aiming to improve disease-free survival rates. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to receive a potentially more effective treatment approach compared to standard care. Eligibility criteria include patients with locally advanced colon adenocarcinoma (cT4, cT3 with invasion >5mm) Nx and no metastases, with specific biomarker assessments for MMR, DMMR, and microsatellite status.

NCT: NCT06783491 · Enrollment: 1083 · Sites: Córdoba, Spain

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

A Clinical Trial Comparing Long-Course Versus Short-Course Radiotherapy Followed by Immunotherapy Combined With Total Neoadjuvant Therapy (TNT) to Long-Course Radiotherapy Followed by TNT in Locally Advanced Rectal Cancer

NCT: NCT07113275 · Enrollment: 444 · Sites: Wuhan, China

MEDIUM PHASE2, PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

A Phase II/III Clinical Trial of RC148 Plus Chemotherapy as 1L Therapy for Unresectable or Metastatic Colorectal Cancer

NCT: NCT07462143 · Enrollment: 80 · Sites: Beijing, China, Changchun, China, Chengdu, China, Chongqing, China, Fuzhou, China

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 11

Neoadjuvant/Adjuvant AK104 in Microsatellite Instability-high or Mismatch Repair-deficient, Resectable Colon Cancer

This clinical trial is testing the efficacy and safety of neoadjuvant/adjuvant treatment with AK104 (Cadonilimab) in patients with resectable microsatellite instability-high (MSI-H) or mismatch repair-deficient (dMMR) colon cancer. The intervention involves AK104 in combination with standard chemotherapy agents, including Oxaliplatin, Capecitabine, 5-Fluorouracil, and Calcium Folinate, to assess its impact on treatment outcomes. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to receive a potentially effective treatment option while contributing to important research. Eligibility criteria include having resectable MSI-H or dMMR colon cancer, with specific biomarker requirements related to microsatellite instability and mismatch repair status.

NCT: NCT07412613 · Enrollment: 386 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Botensilimab + Balstilimab vs Best Supportive Care as Therapy in Chemo-refractory, Unresectable, Colorectal Adenocarcinoma

This clinical trial is testing the combination of botensilimab and balstilimab in patients with chemo-refractory, unresectable colorectal adenocarcinoma. The intervention involves two immunotherapy drugs aimed at potentially improving overall survival and quality of life for participants. Currently in Phase 3 and actively recruiting, this trial may provide insights into treatment options for patients who have limited alternatives. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and microsatellite status.

NCT: NCT07152821 · Enrollment: 834 · Sites: Angers, France, Avignon, France, Besançon, France, Bordeaux, France, Caen, France

MEDIUM PHASE3 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 11

JSKN003 Versus Physician Choiced Treatment in Patients With HER2-positive and Advanced Colorectal Cancer Who Had Failed to Respond to Oxaliplatin, 5-Fu, and Irinotecan

NCT: NCT07384377 · Enrollment: 123 ·

MEDIUM PHASE3 Stage IV BRAF TARGETED THERAPY • RECRUITING 11

Symbiotic-GI-03: A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Chemotherapy in Adult Participants With Metastatic Colorectal Cancer

NCT: NCT07222800 · Enrollment: 800 · Sites: Aberdeen, Abilene, Albany, Albuquerque, Amarillo

HIGH PHASE3 Stage IV BRAF TARGETED THERAPY • RECRUITING 11

Phase III Study of Ivonescimab or Bevacizumab Combined With FOLFOX in Patients With Metastatic Colorectal Cancer

This Phase III clinical trial is evaluating the efficacy of Ivonescimab compared to Bevacizumab when combined with FOLFOX in patients with metastatic colorectal cancer. The intervention involves administering either Ivonescimab or Bevacizumab alongside the chemotherapy regimen mFOLFOX6, which is significant for assessing treatment options in this patient population. Currently, the trial is in the recruiting phase, meaning eligible patients can still enroll to participate in the study. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and microsatellite status.

NCT: NCT07228832 · Enrollment: 600 · Sites: Akron, Atlanta, Baltimore, Beverly Hills, Billings

HIGH PHASE3 Stage IV PHASE III TRIAL • RECRUITING 11

A Randomized, Double-blind, Placebo-controlled, Parallel-group, Comparative, Phase III Study to Evaluate the Efficacy and Safety of Nuvastatic® 300mg Capsule in Reducing Cancer-Tumor in Patients With Metastatic Colorectal Cancer Receiving Standard Chemotherapy.

NCT: NCT07669454 · Enrollment: 250 · Sites: Erode, India, Jalgaon, India, Kanpur, India, Mysore, India, Pune, India

MEDIUM PHASE3 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 11

The Study of JMT101 Combined With Irinotecan as a ≥3rd-Line Treatment in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of JMT101 combined with irinotecan for patients with metastatic colorectal cancer who have received at least two prior lines of treatment. The intervention involves the use of JMT101, a drug being evaluated alongside irinotecan, to determine its effectiveness in improving patient outcomes. This is a phase III trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, and RAS status.

NCT: NCT07134205 · Enrollment: 252 ·

MEDIUM PHASE3 Stage IV KRAS TARGETED THERAPY • RECRUITING 11

Continuation of Cetuximab Beyond First-Line Progression in Metastatic Colorectal Cancer

NCT: NCT07389265 · Enrollment: 480 · Sites: Ancona, Italy, Avellino, Italy, Aviano, Italy, Barcelona, Spain, Bari, Italy

HIGH PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 11

Neoadjuvant Radiotherapy for Rectal Adenocarcinoma With Capecitabine Versus TAS-102 (Neo-REACT): A Multi-center, Randomized, Phase III Trial

This clinical trial, titled Neo-REACT, is testing the efficacy of TAS-102 compared to capecitabine in patients with locally advanced rectal adenocarcinoma undergoing neoadjuvant treatment. The intervention involves administering TAS-102, an oral anti-tumor formulation, alongside radiotherapy, which may offer improved outcomes over the standard capecitabine

regimen. Currently in Phase III and actively recruiting 210 participants, this trial aims to assess the complete response rate, providing critical data for treatment decisions. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06850090 · Enrollment: 210 · Sites: Jinan, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 11

Irinotecan Liposomes in Total Neoadjuvant Therapy in Locally Advanced Rectal Cancer

NCT: NCT07074353 · Enrollment: 360 · Sites: Hangzhou, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

NALIRIFOX Plus Targeted Therapy Versus FOLFOX Plus Targeted Therapy as First-line Treatment for Metastatic Colorectal Cancer

This clinical trial is testing the safety and efficacy of NALIRIFOX plus targeted therapy compared to FOLFOX plus targeted therapy as first-line treatment for patients with metastatic colorectal cancer. The intervention involves two different drug combinations, which is important for determining the most effective treatment approach for this patient population. Currently in Phase 3 and actively recruiting participants, this trial aims to assess the 18-month progression-free survival rate, providing valuable insights for future treatment options. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT07309289 · Enrollment: 144 · Sites: Shanghai, China

MEDIUM PHASE2, PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Node-Sparing Short-Course Radiotherapy Plus Chemotherapy, Bevacizumab and PD-1 Inhibitor in Metastatic pMMR/MSS Colorectal Cancer (MODIFI-CRC)

This clinical trial, known as MODIFI-CRC, is testing the effectiveness of node-sparing short-course radiotherapy combined with chemotherapy, bevacizumab, and a PD-1 inhibitor in patients with metastatic pMMR/MSS colorectal cancer. The intervention aims to enhance tumor response by utilizing radiotherapy to improve the efficacy of immunotherapy and targeted treatments, addressing the limitations of current standard therapies. Currently in Phase 2/3 and actively recruiting participants, this trial may offer patients an opportunity to receive a potentially more effective treatment regimen. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and NGS.

NCT: NCT06958419 · Enrollment: 286 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

PRODIGE 90

The PRODIGE 90 trial is investigating the combination of neoadjuvant Dostarlimab and short course radiotherapy for patients with microsatellite unstable or mismatch repair-deficient locally advanced rectal cancer. The intervention involves administering Dostarlimab, an anti-PD1 immunotherapy, alongside radiotherapy, which may offer a nonoperative management strategy for patients who do not show detectable residual cancer after treatment. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to access this investigational approach while contributing to the research. Eligibility criteria include specific molecular characteristics such as MSI, MMR, and RAS status, which are important for determining the suitability of patients for this treatment strategy.

NCT: NCT06762405 · Enrollment: 68 · Sites: Dijon, France

MEDIUM PHASE3 All Stages KRAS TARGETED THERAPY • RECRUITING 11

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

NCT: NCT06750094 · Enrollment: 700 · Sites: Abilene, Ann Arbor, Arlington, Arlington Heights, Athens

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 11

A Study of Neoadjuvant QL1706 in Participants With Untreated dMMR/MSI-H Resectable Colon Cancer

NCT: NCT06686576 · Enrollment: 363 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 11

Adjuvant PD-1 Blockade for High-risk Stage-II DMMR/MSI-H Colorectal Cancer

This phase III clinical trial is investigating the efficacy of Tislelizumab, a PD-1 blockade drug, as adjuvant therapy for patients with high-risk stage II dMMR/MSI-H colorectal cancer. The trial aims to assess how two cycles of Tislelizumab combined with standard adjuvant chemotherapy compare to standard care in improving disease-free survival (DFS). Currently, the trial is recruiting 180 participants, which means eligible patients may have the opportunity to receive this investigational treatment while contributing to important research. Eligibility criteria include the presence of specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06520683 · Enrollment: 180 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage IV KRAS TARGETED THERAPY • RECRUITING 11

A Study of Amivantamab and mFOLFOX6 or FOLFIRI Versus Cetuximab and mFOLFOX6 or FOLFIRI as First-line Treatment in Participants With KRAS/NRAS and BRAF Wild-type Unresectable or Metastatic Left-sided Colorectal Cancer

This clinical trial is testing the effectiveness of amivantamab in combination with standard chemotherapy regimens (mFOLFOX6 or FOLFIRI) compared to cetuximab with the same chemotherapy in adult patients with KRAS, NRAS, and BRAF wild-type unresectable or metastatic left-sided colorectal cancer. The intervention involves the biological agents amivantamab and cetuximab, along with standard chemotherapy drugs, to evaluate their impact on progression-free survival (PFS). This is a Phase 3 trial currently recruiting 1,000 participants, which indicates that it is in the later stages of testing and may provide critical information on treatment efficacy for patients. Eligibility criteria include specific biomarker requirements related to KRAS, NRAS, BRAF, and other molecular characteristics.

NCT: NCT06662786 · Enrollment: 1000 · Sites: Abilene, Albuquerque, Ann Arbor, Arlington, Arlington Heights

HIGH PHASE3 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 11

First-line Treatment of MCapOX + Cetuximab Vs. MFOLFOX6 + Cetuximab for RAS/BRAF Wild-type, MSS, Unresectable Left-Sided MCRC: a Multicenter, Randomized, Controlled, Phase III Study

NCT: NCT06616259 · Enrollment: 452 · Sites: Chengdu, China

SIGNAL PHASE3 Stage III ctDNA LIQUID BIOPSY • RECRUITING 11

ctDNA-guided Selection of Adjuvant Chemotherapy Regimens for Elderly Colon Cancer Patients After Surgery: A Single-center, Randomized, Controlled Study

This clinical trial is testing the effectiveness of ctDNA-guided selection of adjuvant chemotherapy regimens in elderly patients with high-risk stage II and stage III colon cancer following surgery. Participants will receive either 6 months of 5-FU monotherapy or an intensive XELOX treatment regimen, which is significant for tailoring chemotherapy based on ctDNA detection. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into disease-free survival outcomes for eligible patients. Eligibility criteria include being an elderly patient with high-risk stage II or stage III colon cancer and undergoing ctDNA testing post-surgery.

NCT: NCT06609551 · Enrollment: 312 · Sites: Hangzhou, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Personalising and Refining Neo-adjuvant Chemotherapy in Locally Advanced but Resecable Colon Cancer in the Elderly of 70 Years Old or More

This clinical trial is testing the effectiveness of neoadjuvant chemotherapy using Folfox in elderly patients aged 70 years or older with locally advanced but resectable colon cancer. The intervention involves administering Folfox prior to surgery to potentially improve disease-free survival outcomes. Currently in Phase 3 and actively recruiting participants, this trial

aims to refine treatment approaches for older patients who may be at risk of not receiving adjuvant chemotherapy due to frailty. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06293625 · Enrollment: 150 · Sites: Dijon, France

MEDIUM PHASE2, PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Sacituzumab Govitecan in Metastatic Colorectal Cancer

NCT: NCT06243393 · Enrollment: 80 · Sites: Heidelberg, Germany

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

Injection of SHR-A1811 Versus Physician Choiced Treatment in Patients With Advanced Colorectal Cancer Who Had Failed to Respond to Oxaliplatin, 5-fu, and Irinotecan

NCT: NCT06199973 · Enrollment: 130 · Sites: Shanghai, China

MEDIUM PHASE3 Stage IV KRAS TARGETED THERAPY • RECRUITING 11

Cadonilimab (AK104) Combined With Chemotherapy and Bevacizumab as First-line Treatment for RAS Mutated or Right Sided-metastatic MSS Colorectal Cancer

NCT: NCT06566755 · Enrollment: 30 · Sites: Shenyang, China

MEDIUM PHASE2, PHASE3 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

A Phase II/III Study of HR070803 in Combination With Oxaliplatin, 5-fluorouracil, Calcium Folinate and Bevacizumab Versus FOLFOX in Combination With Bevacizumab for First-line Treatment of Advanced Colorectal Cancer

NCT: NCT05945901 · Enrollment: 669 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 11

Study of Perioperative Dostarlimab in Participants With Untreated T4N0 or Stage III dMMR/MSI-H Resectable Colon Cancer

This clinical trial is testing the efficacy of perioperative dostarlimab in participants with untreated T4N0 or Stage III dMMR/MSI-H resectable colon cancer. The intervention involves dostarlimab, a biological therapy, in combination with standard chemotherapy regimens CAPEOX and FOLFOX, which is significant for its potential to improve event-free survival outcomes. This is a Phase 3 trial currently recruiting 892 participants, indicating that it is in the later stages of testing and may provide valuable information for treatment options. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT05855200 · Enrollment: 892 · Sites: Akron, Ann Arbor, Appleton, Baton Rouge, Bethesda

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

A Clinical Trial Evaluating the Efficacy and Safety of IBI310 in Combination With Sintilimab, for Neoadjuvant Treatment of MSI-H/dMMR Resectable Colon Cancer

This clinical trial is evaluating the efficacy and safety of IBI310 in combination with Sintilimab for patients with MSI-H/dMMR resectable colon cancer. The intervention involves the use of IBI310, a CTLA-4 antibody, alongside Sintilimab, which is significant for its potential to enhance treatment outcomes in this specific cancer type. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients may not be able to enroll at this time. Eligibility criteria include the presence of specific biomarkers such as MSI, MSI-H, MMR, and dMMR.

NCT: NCT05890742 · Enrollment: 453 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Preoperative Short-course Radiation Followed by Envafolimab Plus CAPEOX for MSS Locally Advanced Rectal Adenocarcinoma

This clinical trial is testing the combination of preoperative short-course radiation followed by Envafolimab and CAPEOX chemotherapy in patients with microsatellite stable (MSS) locally advanced rectal adenocarcinoma. The intervention involves the use of Envafolimab, a PD-L1 monoclonal antibody, which is being evaluated for its potential to enhance the effectiveness of short-course radiotherapy and chemotherapy. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to access this investigational treatment while contributing to the understanding of its efficacy and safety. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT05752136 · Enrollment: 108 · Sites: Hangzhou, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Total Neoadjuvant Treatment ±Immunotherapy for High Risk Locally Advanced Rectal Cancer (TNTi)

NCT: NCT06229041 · Enrollment: 472 · Sites: Beijing, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

A Study of HX008 Compared to Chemotherapy in the First-Line Treatment of Subjects With MSI-H/dMMR Metastatic Colorectal Cancer

NCT: NCT05652894 · Enrollment: 190 · Sites: Beijing, China, Bengbu, China, Changchun, China, Changde, China, Changsha, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

TNT of SCRT+CAPOX vs SCRT+CAPOXIRI for Locally Advanced Rectal Cancer

NCT: NCT05646511 · Enrollment: 608 · Sites: Chiba, Japan, Ehime, Japan, Fukuoka, Japan, Gifu, Japan, Hirosaki, Japan

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

Study of XL092 + Atezolizumab vs Regorafenib in Participants With Metastatic Colorectal Cancer

NCT: NCT05425940 · Enrollment: 901 · Sites: Albuquerque, Billings, Charlotte, Chattanooga, Cincinnati

MEDIUM PHASE2, PHASE3 Stage III MSI/MMR LIQUID BIOPSY • RECRUITING 11

Colon Adjuvant Chemotherapy Based on Evaluation of Residual Disease

NCT: NCT05174169 · Enrollment: 1912 · Sites: Abbeville, Aberdeen, Akron, Albany, Albuquerque

MEDIUM PHASE2, PHASE3 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

A Study of a Patient-Specific Neoantigen Vaccine in Combination With Immune Checkpoint Blockade for Patients With Metastatic Colorectal Cancer

This clinical trial is investigating the effectiveness of a patient-specific neoantigen vaccine in combination with immune checkpoint blockade for patients with metastatic colorectal cancer. The intervention includes GRT-C901, GRT-R902, Atezolizumab, Ipilimumab, and fluoropyrimidine plus leucovorin, aiming to enhance treatment outcomes by targeting specific tumor characteristics. Currently, the trial is in Phase 2 and Phase 3 and is active but not recruiting, meaning that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as BRAF, MSI, and MMR status, which may influence treatment response.

NCT: NCT05141721 · Enrollment: 700 · Sites: Austin, Bethesda, Boca Raton, Charlottesville, Chicago

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 11

PACE: PD-1 Antibody For dMMR/MSI-H Stage III Colorectal Cancer

The PACE trial is investigating the efficacy of the PD-1 antibody Sintilimab in patients with dMMR/MSI-H Stage III colorectal cancer. Participants will receive either Sintilimab alone or a combination of oxaliplatin and capecitabine, which are standard chemotherapy agents, to assess their impact on disease-free survival. This is a Phase 3 study currently recruiting 323 participants, which means eligible patients may have the opportunity to receive a potentially effective treatment while contributing to important research. Eligibility criteria include having local advanced colon cancer with specific biomarker characteristics such as dMMR or MSI-H.

NCT: NCT05236972 · Enrollment: 323 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 11

Neoadjuvant FOLFOXIRI Versus Immediate Surgery for Stage II and III Colon Cancers

NCT: NCT05194878 · Enrollment: 840 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

A Study of Anlotinib Hydrochloride Capsule Combined With Chemotherapy as First-line Treatment in Subjects With RAS/BRAF Wild Metastatic Colorectal Cancer

This clinical trial is testing the combination of Anlotinib hydrochloride capsules with chemotherapy as a first-line treatment for patients with RAS/BRAF wild-type unresectable metastatic colorectal cancer. The intervention involves administering Anlotinib alongside standard chemotherapy agents Bevacizumab, Oxaliplatin, and Capecitabine, which is significant for evaluating potential improvements in treatment outcomes. Currently, the trial is in Phase 3 and is active but not recruiting, indicating that it is in the final stages of testing before potential approval. Eligibility criteria include specific biomarker requirements related to BRAF, MSI, MMR, and RAS status.

NCT: NCT04854668 · Enrollment: 748 · Sites: Beijing, China, Changchun, China, Changsha, China, Changzhi, China, Changzhou, China

MEDIUM PHASE2, PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

A Clinical Study to Evaluate Efficacy and Safety of Serplulimab (HLX10) Combined With Bevacizumab (HLX04) and Chemotherapy (XELOX) in Patients With Metastatic Colorectal Cancer (mCRC)

This clinical trial is evaluating the efficacy and safety of Serplulimab (HLX10) combined with Bevacizumab (HLX04) and chemotherapy (XELOX) in patients with metastatic colorectal cancer (mCRC). The intervention involves the use of HLX10, a drug that is being tested against a placebo to determine its effectiveness in improving patient outcomes. Currently, the trial is in Phase 2 and Phase 3 and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment as part of the study. Eligibility criteria include the presence of specific biomarkers, such as MSI or MSI-H.

NCT: NCT04547166 · Enrollment: 568 · Sites: Guangzhou, China, Hangzhou, China, Kashiwa, Japan, Linyi, China, Shanghai, China

MEDIUM PHASE3 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

A Study of Encorafenib Plus Cetuximab With or Without Chemotherapy in People With Previously Untreated Metastatic Colorectal Cancer

NCT: NCT04607421 · Enrollment: 841 · Sites: Aventura, Basking Ridge, Berkeley Heights, Chicago, City of Saint Peters

MEDIUM PHASE3 Stage III MSI/MMR LIQUID BIOPSY • RECRUITING 11

Circulating Tumour DNA Based Decision for Adjuvant Treatment in Colon Cancer Stage II Evaluation

The CIRCULATE study is evaluating adjuvant therapy for patients with UICC stage II colon cancer. The intervention involves the use of capecitabine to assess its impact on disease-free survival in patients who test positive for postoperative circulating tumor DNA. This is a Phase 3 trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive capecitabine as part of their treatment. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT04089631 · Enrollment: 4812 · Sites: Dresden, Germany

MEDIUM PHASE3 Stage IV BRAF LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 11

Identification and Treatment Of Micrometastatic Disease in Stage III Colon Cancer

NCT: NCT03803553 · Enrollment: 400 · Sites: Boston, New York

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 11

Neoadjuvant mFOLFOXIRI Plus Bevacizumab in Patients With High-Risk Locally Advanced Rectal Cancer

NCT: NCT04215731 · Enrollment: 582 · Sites: Guangzhou, China

HIGH PHASE3 Stage IV PHASE III TRIAL • ACTIVE_NOT_RECRUITING 11

A Randomized Phase III Study Comparing Maintenance Treatment With Fluoropyrimidine + Bevacizumab Versus Fluoropyrimidine After Induction Chemotherapy for a Metastatic Colorectal Cancer

This clinical trial, titled BEVAMAINT, is testing the effectiveness of maintenance treatment with a combination of fluoropyrimidine and bevacizumab compared to fluoropyrimidine alone after induction chemotherapy in patients with metastatic colorectal cancer. The intervention involves the use of fluoropyrimidine, a chemotherapy drug, and bevacizumab, a targeted therapy, to assess their impact on treatment outcomes. Currently, the trial is in Phase III and is active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients are not being enrolled at this time. Eligibility criteria have not been specified, but the study focuses on patients who have undergone initial chemotherapy for metastatic colorectal cancer.

NCT: NCT04188145 · Enrollment: 400 · Sites: Dijon, France

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

A Study of Nivolumab, Nivolumab Plus Ipilimumab, or Investigator's Choice Chemotherapy for the Treatment of Participants With Deficient Mismatch Repair (dMMR)/Microsatellite Instability High (MSI-H) Metastatic Colorectal Cancer (mCRC)

This clinical trial is testing the effectiveness of nivolumab, both alone and in combination with ipilimumab, compared to investigator's choice chemotherapy in patients with deficient mismatch repair (dMMR) or microsatellite instability high (MSI-H) metastatic colorectal cancer (mCRC). The intervention involves the use of nivolumab, a monoclonal antibody, and ipilimumab, which may enhance the immune response against cancer cells, making it a significant focus for treatment in this patient population. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the advanced stages of testing and may provide valuable insights into treatment options for patients. Eligibility criteria include the presence of specific biomarkers such as MSI-H or dMMR.

NCT: NCT04008030 · Enrollment: 839 · Sites: Arlington Heights, Dallas, Denver, Los Angeles, New York

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 11

Prospectively Randomized Control Clinical Trial of FOLFOXIRI Preoperative Chemotherapy Alone on Rectal Cancer in Local Advance Comparing to Oral Capecitabine Combined With Long-term Radiation

NCT: NCT03671252 · Enrollment: 776 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 11

Testing the Addition of Atezolizumab to Combination Chemotherapy or Atezolizumab Alone for Metastatic Colon or Rectal Cancer, the COMMIT Study

The COMMIT study is a phase III clinical trial evaluating the effectiveness of adding atezolizumab to combination chemotherapy or administering atezolizumab alone in patients with metastatic colorectal cancer characterized by deficient DNA mismatch repair. The intervention involves the use of atezolizumab, a monoclonal antibody that may enhance the

immune system's ability to combat cancer, alongside standard chemotherapy agents like fluorouracil and bevacizumab, which target tumor growth and blood vessel formation. Currently, the trial is active but not recruiting new participants, indicating that it is in the process of collecting data from enrolled patients. Eligibility criteria include specific biomarkers such as MSI, MMR, and RAS status, which may determine suitability for participation in the study.

NCT: NCT02997228 · Enrollment: 120 · Sites: Aberdeen, Akron, Albany, Albuquerque, Allentown

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 10

A Study of the Duration of Dostarlimab in Untreated Mismatch Repair Deficient (dMMR)/ Microsatellite Instability-high (MSI-H) Locally Advanced Rectal Cancer

NCT: NCT07740720 · Enrollment: 90 ·

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 10

Personalizing Chemotherapy Selection After Surgery for Patients With Stage III Colorectal Cancer Using a Blood Test

NCT: NCT07340567 · Enrollment: 2450 · Sites: Gothenburg, Sweden, Lund, Sweden, Stockholm, Sweden, Uppsala, Sweden, Villejuif, France

SIGNAL PHASE3 Stage IV BRAF LIQUID BIOPSY • NOT_YET_RECRUITING 10

De-scalation or swItch of Treatment According to Circulating tuMOr DNA Variation After 2 Cycles of Doublet Chemotherapy Plus Targeted Agent in Metastatic Unresectable Colorectal Cancer

NCT: NCT06446557 · Enrollment: 408 · Sites: Rouen, France

MEDIUM PHASE3 All Stages IMMUNOTHERAPY • NOT_YET_RECRUITING 10

Long-Term Extension Study for Participants Previously Enrolled in an Exelixis-Sponsored Study

This clinical trial is a long-term extension study for participants who have previously enrolled in an Exelixis-sponsored study and are experiencing clinical benefit. The intervention involves multiple drugs, including Cabozantinib, Zanzalintinib, Atezolizumab, Nivolumab, and Abiraterone, which are being provided to ensure continued access to treatment for those without local options. This is a Phase 3 study that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria may include prior participation in an Exelixis-sponsored study and the ability to demonstrate clinical benefit from the treatment.

NCT: NCT07620574 · Enrollment: 5000 ·

SIGNAL PHASE3 Stage III ctDNA LIQUID BIOPSY • RECRUITING 10

Investigate the Efficacy of Chemotherapy in Patients With Positive ctDNA After Surgery and Adjuvant Chemotherapy for a Stage III Colorectal Cancer

This clinical trial is investigating the efficacy of chemotherapy in patients with positive circulating tumor DNA (ctDNA) following surgery and adjuvant chemotherapy for stage III colorectal cancer. The intervention includes FOLFIRI and Trifluridine drug regimens, along with biological assessments and imaging studies, to evaluate their impact on disease recurrence. This is a Phase III study currently recruiting 1,660 participants, which means eligible patients may have the opportunity to contribute to research that could influence future treatment protocols. Eligibility criteria include the presence of ctDNA, as well as specific biomarkers such as RAS and NGS.

NCT: NCT06197425 · Enrollment: 1660 · Sites: Dijon, France

SIGNAL PHASE2, PHASE3 All Stages RESEARCH • RECRUITING 10

Antegrade Intestinal Fluid Reinfusion for Prevention of Low Anterior Resection Syndrome After Low Anterior Resection: a Single-Center, Prospective Randomized Controlled Trial.

This clinical trial is testing the effectiveness of antegrade intestinal fluid reinfusion in preventing low anterior resection syndrome (LARS) in patients with a prophylactic ileal stoma. The intervention involves administering either intestinal fluid or potable water through the ileal stoma prior to ileostomy reversal surgery, which is significant for managing post-surgical

complications. Currently in Phase 2 and Phase 3, the trial is actively recruiting 160 participants, which means eligible patients can join to contribute to the research. Participants will be required to maintain a diary of their symptoms during the study period.

NCT: NCT07537998 · Enrollment: 160 · Sites: Guangzhou, China

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 10

Patient Preference for Subcutaneous vs. Intravenous Immune Therapy

This clinical trial is testing patient and healthcare professional preferences for subcutaneous versus intravenous administration of immune therapies nivolumab and pembrolizumab in individuals with colorectal cancer. The intervention involves comparing the two methods of drug delivery to determine which is preferred by patients, which may impact treatment adherence and quality of life. Currently in Phase 2 and actively recruiting, this trial aims to enroll 880 participants, providing valuable insights into patient preferences that could influence future treatment protocols. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, and tumor mutational burden.

NCT: NCT07223424 · Enrollment: 880 · Sites: Pittsburgh

HIGH PHASE3 Stage III NGS PHASE III TRIAL • ACTIVE_NOT_RECRUITING 10

A Phase III Trial to Evaluate the Efficacy, Immunogenicity and Safety of a Recombinant Human Papillomavirus 9-Valent (Types 6, 11, 16, 18, 31, 33, 45, 52, 58) Vaccine (E.Coli) in Chinese Males

NCT: NCT06866574 · Enrollment: 9300 · Sites: Chengdu, China, Fuzhou, China, Hangzhou, China, Kunming, China

MEDIUM N/A Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 10

Prospective National Cohort Evaluating Predictive Biomarkers of Resistance to Immunotherapy in Patients With MSI/dMMR Metastatic Colorectal Cancer (CORESIM)

The CORESIM trial is evaluating predictive biomarkers of resistance to immunotherapy in patients with metastatic colorectal cancer characterized by microsatellite instability (MSI) or mismatch repair deficiency (dMMR). The intervention focuses on identifying biomarkers such as MSI, MMR, and DMMR to determine which patients may not respond to pembrolizumab, an anti-PD1 monoclonal antibody, in first-line treatment. This study is currently in the recruiting phase, which means eligible patients can still enroll to participate in the trial. Eligibility criteria include having MSI or dMMR metastatic colorectal cancer, with a target enrollment of 600 participants.

NCT: NCT06353854 · Enrollment: 600 · Sites: Bayonne, France, Beauvais, France, Besançon, France, Boujan-sur-Libron, France, Boulogne-sur-Mer, France

MEDIUM PHASE3 Stage IV BRAF TARGETED THERAPY • RECRUITING 10

Efficacy and Safety of Tunlametinib Plus Vemurafenib in Patients With BRAF V600E-mutant Metastatic Colorectal Cancer

NCT: NCT06008119 · Enrollment: 165 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR LIQUID BIOPSY • RECRUITING 10

FOG-001 in Locally Advanced or Metastatic Solid Tumors

NCT: NCT05919264 · Enrollment: 619 · Sites: Aurora, Baltimore, Boston, Charlottesville, Cleveland

MEDIUM PHASE3 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 10

COLLISION RELAPSE Trial

The COLLISION RELAPSE Trial is testing the effectiveness of neoadjuvant systemic therapy followed by repeat local treatment in patients with at least one locally treatable recurrent colorectal liver metastasis (CRLM) without extrahepatic disease. The intervention involves various combinations of chemotherapy regimens (CAPOX, FOLFOX, FOLFIRI) and aims to improve overall survival compared to upfront repeat local treatment alone. This is a Phase 3 trial currently recruiting 360 participants, which means eligible patients may have the opportunity to receive a potentially beneficial

treatment option while contributing to important research. Eligibility criteria include specific biomarkers such as MSI, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT05861505 · Enrollment: 360 · Sites: Amsterdam, Netherlands

MEDIUM PHASE3 Stage IV HER2 TARGETED THERAPY • RECRUITING 10

A Study of Tucatinib With Trastuzumab and mFOLFOX6 Versus Standard of Care Treatment in First-line HER2+ Metastatic Colorectal Cancer

NCT: NCT05253651 · Enrollment: 400 · Sites: Anaheim, Atlanta, Aurora, Aventura, Baldwin Park

MEDIUM PHASE2 Stage IV MSI/MMR LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 10

Fecal Microbiota Transplant and Re-introduction of Anti-PD-1 Therapy (Pembrolizumab or Nivolumab) for the Treatment of Metastatic Colorectal Cancer in Anti-PD-1 Non-responders

This phase II clinical trial is investigating the combination of fecal microbiota transplantation and the re-introduction of anti-PD-1 therapy (pembrolizumab or nivolumab) for patients with metastatic colorectal cancer who have not responded to previous anti-PD-1 treatments. The intervention includes procedures such as fecal microbiota transplantation and the use of specific drugs, which aim to enhance the immune response against cancer cells. Currently, the trial is active but not recruiting new participants, indicating that it is ongoing with enrolled patients receiving treatment. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, CTDNA, microsatellite status, and mismatch repair status.

NCT: NCT04729322 · Enrollment: 15 · Sites: Houston

HIGH PHASE3 Stage IV RAS IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 10

Phase 3 Clinical Trial of Carboplatin and Paclitaxel +/- Nivolumab in Metastatic Anal Cancer Patients

The EA2176 trial is testing the effectiveness of adding nivolumab to a chemotherapy regimen of carboplatin and paclitaxel in patients with metastatic anal cancer. The intervention involves the use of nivolumab, an immunotherapy drug, alongside standard chemotherapy, which may enhance the immune response against cancer cells while also targeting tumor growth through traditional chemotherapy mechanisms. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients are not currently being enrolled. Eligibility for participation includes specific biomarker requirements related to RAS.

NCT: NCT04444921 · Enrollment: 205 · Sites: Aberdeen, Albemarle, Allentown, Altamonte Springs, Alton

HIGH PHASE3 Stage IV RAS PHASE III TRIAL • ACTIVE_NOT_RECRUITING 10

Phase III Study in First-line Treatment of Patients With Metastatic Colorectal Cancer Who Are Not Candidate for Intensive Therapy.

This Phase III clinical trial is testing the efficacy of trifluridine/tipiracil hydrochloride (S95005) in combination with bevacizumab compared to capecitabine with bevacizumab in patients with metastatic colorectal cancer who are not candidates for intensive therapy. The intervention involves the use of S95005, which is being evaluated for its potential to improve progression-free survival (PFS) in this patient population. Currently, the trial is active but not recruiting, indicating that while no new participants are being enrolled, the study is ongoing and data is being collected. Eligibility for participation includes specific biomarker requirements related to RAS status.

NCT: NCT03869892 · Enrollment: 856 · Sites: Aalborg, Denmark, Aberdeen, United Kingdom, Alicante, Spain, Almada, Portugal, Arkhangelsk, Russia

MEDIUM PHASE2, PHASE3 Stage IV RAS IMMUNOTHERAPY • RECRUITING 10

Trans-Artery/Intra-Tumor Infusion of Checkpoint Inhibitors Plus Chemodrug for Immunotherapy of Advanced Solid Tumors

This clinical trial is testing the combination of checkpoint inhibitors, such as Pembrolizumab, with chemotherapy for patients with advanced solid tumors, including colorectal cancer. The intervention involves a trans-artery/intra-tumor infusion of these therapies, which aims to enhance the effectiveness of treatment by targeting the tumor directly. Currently in Phase 2 and Phase 3, the trial is actively recruiting 200 participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include specific biomarker requirements, such as RAS status.

NCT: NCT03755739 · Enrollment: 200 · Sites: Guanzhou, China

SIGNAL PHASE3 Stage IV KRAS LIQUID BIOPSY • NOT_YET_RECRUITING 9

Continuous or Intermittent Cetuximab Plus FOLFIRI as First-line Treatment in RAS/BRAF Wild-type mCRC Patients

This clinical trial is testing the effectiveness of continuous versus intermittent administration of FOLFIRI combined with cetuximab in adults with RAS/BRAF wild-type metastatic colorectal cancer (mCRC). The intervention involves the established chemotherapy regimen FOLFIRI alongside cetuximab, with the aim of determining whether treatment breaks can improve progression-free survival and quality of life compared to continuous treatment. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include being an adult with mCRC and having specific biomarkers such as KRAS, NRAS, and BRAF assessed through liquid biopsy.

NCT: NCT07004413 · Enrollment: 267 · Sites: Berlin, Germany

MEDIUM N/A All Stages KRAS TARGETED THERAPY • NOT_YET_RECRUITING 9

NOMAD-GI: Molecular-guided Non-operative Management and Organ Preservation Strategy After Precision Therapy in Gastrointestinal Cancers

The NOMAD-GI study is investigating the safety and efficacy of molecular biomarker-guided non-operative management and organ preservation strategies in patients with gastrointestinal cancers. The intervention involves assessing patients with actionable molecular biomarkers after receiving precision therapy, which is significant for determining individualized treatment approaches. This study is currently not yet recruiting participants, meaning that patients will need to wait until enrollment begins to participate. Eligible patients must have specific biomarkers, including dMMR/MSI-H, PD-L1 high expression, and others, to qualify for the study.

NCT: NCT07775859 · Enrollment: 200 ·

MEDIUM PHASE3 Stage II IMMUNOTHERAPY • NOT_YET_RECRUITING 9

A Multicenter Study of Active Specific Immunotherapy With OncoVax® in Patients With Stage II Colon Cancer

NCT: NCT02448173 · Enrollment: 550 · Sites: Port Orange

MEDIUM PHASE3 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 9

Study Evaluating Two Treatment Strategies for Rectal Cancer in Patients ≥75 Years Old

NCT: NCT07118800 · Enrollment: 160 ·

SIGNAL PHASE3 Stage IV KRAS RESEARCH • RECRUITING 9

Comparing Efficacy and Safety of Enlituo® Versus Erbitux® Plus FOLFOX in Locally Advanced/Metastatic Colorectal Cancer With RAS/BRAF Wild-type and MSS/pMMR Status.

NCT: NCT07595276 · Enrollment: 608 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 9

A Study to Access Intravenous (IV) Telisotuzumab Adizutecan in Combination With IV Bevacizumab Compared to Standard of Care IV Bevacizumabin Combination With Oral Trifluridine and Tipiracil in Adult Participants With Refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of intravenous (IV) telisotuzumab adizutecan and IV bevacizumab against the standard of care, which includes IV bevacizumab combined with oral trifluridine and tipiracil, in adult participants with refractory metastatic colorectal cancer (mCRC). The intervention involves the investigational drug telisotuzumab adizutecan, which is being evaluated for its potential effectiveness in altering disease activity and managing adverse events in this patient population. Currently in Phase 3 and actively recruiting, this trial aims to enroll approximately 990 participants across 125 sites globally, which may provide eligible patients with access to a potentially different treatment

approach. Eligibility criteria include being an adult with refractory mCRC, although specific biomarker requirements are not detailed in the provided information.

NCT: NCT07525206 · Enrollment: 990 · Sites: Abilene, Aurora, Blacksburg, Chicago, Cincinnati

SIGNAL PHASE3 Stage III MSI/MMR RESEARCH • RECRUITING 9

SCRT + Chemo Targeted Immuno-neoadjuvant Therapy for High-risk pMMR/MSS RC

NCT: NCT07549399 · Enrollment: 204 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 9

A Phase II Randomized Trial of Serplulimab With Second-Line Chemo/Targeted Therapy for Early Relapse Colorectal Cancer After Adjuvant Chemotherapy

NCT: NCT07604909 · Enrollment: 40 · Sites: Shanghai, China

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 9

A Study of Precemtabart Tocentecan With or Without Bevacizumab Compared to Trifluridine/Tipiracil Plus Bevacizumab in Participants With Previously Treated Metastatic Colorectal Cancer (PROCEADE-CRC-03)

This clinical trial is testing the efficacy of precemtabart tocentecan, both as a single agent and in combination with bevacizumab, in participants with previously treated metastatic colorectal cancer (mCRC). The intervention involves comparing these treatments to trifluridine/tipiracil plus bevacizumab, aiming to demonstrate an improvement in overall survival for patients. Currently in Phase 3 and actively recruiting, this trial offers eligible patients the opportunity to participate in research that may lead to better treatment options. Participants must have previously received treatment with irinotecan, oxaliplatin, a fluoropyrimidine, and bevacizumab to qualify for enrollment.

NCT: NCT07549412 · Enrollment: 1020 · Sites: Abilene, Arlington Heights, Austin, Bedford, Columbia

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • RECRUITING 9

Neoadjuvant CAPOX With or Without Pucotenlimab Plus Selective Radiotherapy for Locally Advanced Rectal Cancer

NCT: NCT07528209 · Enrollment: 556 · Sites: Guangzhou, China

MEDIUM PHASE3 Stage IV HER2 TARGETED THERAPY • NOT_YET_RECRUITING 9

A Clinical Study to Evaluate Injection TQB2102 for the Treatment of Patients With HER2 IHC3+ Advanced Colorectal Cancer Who Progressed After Treatment With Oxaliplatin, Irinotecan and Fluoropyrimidine-Based Drugs

This clinical trial is evaluating the efficacy and safety of Injection TQB2102 for patients with HER2 IHC3+ advanced colorectal cancer who have progressed after treatment with oxaliplatin, irinotecan, and fluoropyrimidine-based drugs. The intervention, TQB2102, is being compared to the investigator's choice of treatment, which may include Trifluridine/Tipiracil (TAS-102), Fruquintinib, or Regorafenib, to determine its effectiveness in improving patient outcomes. This is a Phase III trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligible participants must have a HER2 IHC3+ biomarker status and have previously failed standard treatments.

NCT: NCT07483684 · Enrollment: 142 · Sites: Beijing, China, Changchun, China, Changsha, China, Chengdu, China, Chongqing, China

MEDIUM NA Stage IV MSI/MMR LIQUID BIOPSY • NOT_YET_RECRUITING 9

Neoadjuvant CAPEOX Versus Upfront Surgery for Locally Advanced Colon Cancer With Elevated CEA: A Single-Center, Open-Label, Randomized Controlled Trial

This clinical trial is testing the efficacy of neoadjuvant chemotherapy using CAPOX compared to upfront surgery in adults aged 18-70 years with stage II (high-risk)-III, non-MSI-H colon adenocarcinoma and elevated baseline CEA levels. The intervention includes ctDNA-based molecular residual disease (MRD) monitoring and Immunoscore assessment, which

are important for evaluating treatment response and predicting disease-free survival. This study is currently not yet recruiting participants and aims to enroll 100 individuals, meaning patients will need to wait for recruitment to begin before they can participate. Eligibility criteria include specific biomarkers such as MSI, MMR, and RAS, which will help identify suitable candidates for the trial.

NCT: NCT07413146 · Enrollment: 100 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 9

Neoadjuvant Immunotherapy ± Radiotherapy in MSI-H/dMMR Locally Advanced Colorectal Cancer

NCT: NCT07403877 · Enrollment: 114 · Sites: Shanghai, China

MEDIUM PHASE3 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 9

Prophylactic PIPAC in Patients With High-risk Colorectal Cancer

NCT: NCT06839092 · Enrollment: 160 ·

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 9

A Randomized Phase 2 Trial of Fruquintinib and TAS-102 as Compared to Fruquintinib in Patients With Refractory Advanced/Metastatic Colorectal Cancer

This clinical trial is testing the combination of Fruquintinib and TAS-102 (trifluridine and tipiracil) against Fruquintinib alone in patients with refractory advanced or metastatic colorectal cancer. The intervention aims to evaluate the potential progression-free survival benefit of the combination therapy. Currently in Phase 2 and actively recruiting, this trial may provide insights into treatment options for patients who have not responded to previous therapies. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status.

NCT: NCT06992258 · Enrollment: 120 · Sites: Charleston, Fairfax, Lancaster, Miami Beach, Nashville

SIGNAL PHASE3 Stage III ctDNA LIQUID BIOPSY • RECRUITING 9

ctDNA-Based Adjuvant Chemotherapy for High-Risk Rectal Cancer

This clinical trial is testing the effectiveness of adjuvant chemotherapy in preventing disease recurrence in patients with high-risk rectal cancer who have detectable ctDNA after surgery. The intervention involves administering either 4 cycles of CAPOX or 6 cycles of FOLFOX chemotherapy within 12 weeks post-surgery, which is significant for potentially improving patient outcomes. Currently in Phase 3 and actively recruiting participants, this trial aims to provide insights into disease-free survival for eligible patients. Participants must have detectable ctDNA following their surgical treatment to qualify for the study.

NCT: NCT07188025 · Enrollment: 103 · Sites: 's-Hertogenbosch, Netherlands, Almelo, Netherlands, Amersfoort, Netherlands, Amsterdam, Netherlands, Breda, Netherlands

MEDIUM PHASE2, PHASE3 Stage IV MSI/MMR IMMUNOTHERAPY • NOT_YET_RECRUITING 9

Neoadjuvant Chemoradiotherapy Combined With Immunotherapy and Anti-angiogenesis in Treating Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy and safety of neoadjuvant chemoradiotherapy combined with the PD-1 monoclonal antibody serplulimab and the anti-angiogenic agent bevacizumab in patients with locally advanced rectal cancer. The intervention involves a combination of these drugs with short-course chemoradiation therapy, which may enhance treatment outcomes for patients. This is a multicenter, prospective, randomized phase II and phase III trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin. Eligibility criteria include patients with previously untreated pMMR/MSS middle and low locally advanced rectal cancer.

NCT: NCT07134218 · Enrollment: 180 · Sites: Shanghai, China

SIGNAL PHASE2, PHASE3 Stage IV KRAS RESEARCH • NOT_YET_RECRUITING 9

A Clinical Study to Evaluate the Efficacy and Safety of Envafolimab Combined With Cetuxima-βand mFOLFOX6 in Patients With MSS, RAS/BRAF Wild-Type Metastatic Colorectal Cancer (mCRC)

This clinical trial is evaluating the efficacy and safety of Envafolimab combined with Cetuximab-β and mFOLFOX6 in patients with microsatellite stable (MSS), RAS/BRAF wild-type metastatic colorectal cancer (mCRC). The intervention involves a combination of three drugs, which is significant for potentially improving treatment outcomes in this specific patient population. The trial is in Phase 2/3 and is currently not yet recruiting, indicating that patients will need to wait for enrollment to begin before participating. Eligibility criteria include the presence of specific biomarkers such as KRAS, NRAS, BRAF, and RAS.

NCT: NCT06959693 · Enrollment: 590 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 9

AK112 and Chemotherapy in First-line Metastatic Colorectal Cancer

NCT: NCT06951503 · Enrollment: 560 · Sites: Guanzhou, China

SIGNAL PHASE3 All Stages ctDNA LIQUID BIOPSY • RECRUITING 9

tislelizUMaB in canceR Patients With molEcuLar residuaL Disease

This clinical trial is testing the efficacy of tislelizumab in patients with colorectal cancer who have molecular residual disease (MRD+) after surgery and perioperative treatments. The intervention involves administering tislelizumab, an immunotherapy drug, compared to a placebo, with the aim of improving disease-free survival (DFS) in this patient population. This is a Phase 3 trial currently recruiting 717 participants, which indicates that it is in the later stages of testing and may provide critical information on the effectiveness of tislelizumab for patients with detectable tumor DNA in their blood. Eligibility criteria include having detectable circulating tumor DNA and specific biomarker requirements related to RAS status.

NCT: NCT06332274 · Enrollment: 717 · Sites: Villejuif, France

HIGH PHASE3 Stage IV RAS PHASE III TRIAL • NOT_YET_RECRUITING 9

Short-course Radiation (SCRT) Followed by 6 Cycles of Cadonilimab Plus MFOLFOX6 As Neoadjuvant Therapy for Patients with Locally Advanced Rectal Cancer (LARC): a Multicenter, Two-arm Parallel, Open-label, Randomised Phase III Trial (NeoCaCRT-III)

This clinical trial is testing the effectiveness of neoadjuvant short-course radiation (SCRT) followed by 6 cycles of cadonilimab plus mFOLFOX6 in patients with locally advanced rectal cancer (LARC). The intervention involves combining a dual immunotherapy (cadonilimab) with chemotherapy (mFOLFOX6) to assess its impact on disease-free survival and safety compared to chemotherapy alone. This is a Phase III trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarker assessments, such as RAS and next-generation sequencing (NGS).

NCT: NCT06832917 · Enrollment: 238 ·

MEDIUM PHASE1 Stage IV KRAS TARGETED THERAPY • RECRUITING 9

Open-Label Study of BBO-10203 in Subjects With Advanced Solid Tumors

This clinical trial is testing BBO-10203, a PI3Kα:RAS breaker, in patients with advanced solid tumors. The intervention involves administering BBO-10203 both as a single agent and in combination with other drugs, including Trastuzumab, Fulvestrant, Ribociclib, and FOLFOX, to evaluate its safety and pharmacokinetics. Currently in Phase 1 and actively recruiting, this trial aims to determine the maximum tolerated dose and/or recommended Phase 2 dose of BBO-10203, which is essential for guiding future treatment options. Eligibility criteria include specific biomarkers such as KRAS, BRAF, MSI, MSI-H, MMR, DMMR, HER2, and NTRK.

NCT: NCT06625775 · Enrollment: 392 · Sites: Boston, Dallas, Duarte, Houston, Indianapolis

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 9

Stereotactic Ablative Radiotherapy (SABR) for the Treatment of Patients With Metastatic Cancer, ID-COMET Trial

The ID-COMET trial is testing the effectiveness of Stereotactic Ablative Radiotherapy (SABR) in patients with newly diagnosed metastatic colorectal cancer. The intervention involves administering SABR compared to best practice treatment, which is significant for assessing its impact on overall survival. This is a Phase 3 trial currently recruiting 800 participants, indicating that eligible patients may have the opportunity to receive this treatment while contributing to important research. Specific eligibility criteria or biomarker requirements have not been detailed in the provided information.

NCT: NCT06563388 · Enrollment: 800 · Sites: Buffalo

HIGH PHASE3 All Stages PHASE III TRIAL • ACTIVE_NOT_RECRUITING 9

A Phase III Study to Evaluate the Efficacy, Immunogenicity and Safety of the 9-valent HPV Vaccine in Chinese Males

NCT: NCT06465914 · Enrollment: 9000 · Sites: Changsha, China, Chengdu, China, Kunming, China, Nanning, China, Taiyuan, China

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • RECRUITING 9

DOSAGE Study: Upfront Dose-Reduced Chemotherapy in Older Patients with Metastatic Colorectal Cancer

NCT: NCT06275958 · Enrollment: 587 · Sites: 's-Hertogenbosch, Netherlands, Alkmaar, Netherlands, Amstelveen, Netherlands, Amsterdam, Netherlands, Arnhem, Netherlands

SIGNAL PHASE3 Stage IV BRAF RESEARCH • RECRUITING 9

Intermittent or Continuous Panitumumab Plus FOLFIRI for Left Sided RAS/B-RAF Wild-type Metastatic Colorectal Cancer

NCT: NCT06509126 · Enrollment: 500 · Sites: Naples, Italy

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 9

A Study to Evaluate the Efficacy and Safety of Ifinatamab Deruxtecan (I-DXd) in Subjects With Recurrent or Metastatic Solid Tumors (IDeate-PanTumor02)

NCT: NCT06330064 · Enrollment: 680 · Sites: Amarillo, Chattanooga, Dallas, Fairfax, Germantown

MEDIUM PHASE3 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 9

Testing Pump Chemotherapy in Addition to Standard of Care Chemotherapy Versus Standard of Care Chemotherapy Alone for Patients With Unresectable Colorectal Liver Metastases: The PUMP Trial

NCT: NCT05863195 · Enrollment: 408 · Sites: Allentown, Ann Arbor, Atlanta, Aurora, Aventura

SIGNAL PHASE3 Stage III ctDNA LIQUID BIOPSY • RECRUITING 9

Circulating Tumor DNA Methylation Guided Postoperative Adjuvant Chemotherapy for High-risk Stage II/III Colorectal Cancer

NCT: NCT05954078 · Enrollment: 340 · Sites: Shanghai, China

SIGNAL PHASE3 Stage III RESEARCH • RECRUITING 9

Adjuvant Chemotherapy Combined With Targeted Therapy or Not in the T3-4N2 Colorectal Cancer Patients

This clinical trial is testing the effectiveness of adjuvant chemotherapy combined with targeted therapy in patients with resectable stage T3-4N2 colorectal cancer. The intervention involves FOLFOX chemotherapy regimens with or without

Bevacizumab, which is significant for evaluating potential improvements in disease-free survival. This is a Phase 3 trial currently recruiting 366 participants, which means eligible patients may have the opportunity to contribute to important findings regarding treatment options. Specific eligibility criteria or biomarker requirements have not been detailed in the provided information.

NCT: NCT05797467 · Enrollment: 366 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage III ctDNA LIQUID BIOPSY • RECRUITING 9

ctDNA-guided Adjuvant Chemotherapy in Liver Metastasis of Colorectal Cancer

NCT: NCT05815082 · Enrollment: 490 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage III ctDNA LIQUID BIOPSY • RECRUITING 9

Platform Study of Circulating Tumor DNA Directed Adjuvant Chemotherapy in Colon Cancer (KCSG CO22-12)

NCT: NCT05534087 · Enrollment: 236 · Sites: Busan, South Korea, Daegu, South Korea, Goyang, South Korea, Hwasun, South Korea, Seongnam, South Korea

SIGNAL PHASE3 Stage III RESEARCH • RECRUITING 9

Xian-Lian-Jie-Du Optimization Decoction As an Adjuvant Treatment for Prevention of Recurrence of Colon Cancer

NCT: NCT05709249 · Enrollment: 730 · Sites: Nanjing, China

SIGNAL PHASE3 Stage III ctDNA LIQUID BIOPSY • NOT_YET_RECRUITING 9

Adjuvant mFOLFOXIRI vs. mFOLFOX6 in MRD Positive Stage II-III Colorectal Cancer (AFFORD)

NCT: NCT05427669 · Enrollment: 340 ·

MEDIUM PHASE1, PHASE2 Stage IV KRAS LIQUID BIOPSY • RECRUITING 9

A Study of Amivantamab Monotherapy and in Addition to Standard-of-Care Chemotherapy in Participants With Advanced or Metastatic Colorectal Cancer

NCT: NCT05379595 · Enrollment: 225 · Sites: Ann Arbor, Baltimore, Birmingham, Grand Rapids, Hattiesburg

SIGNAL PHASE3 Stage IV RAS RESEARCH • RECRUITING 9

Post-resection/Ablation Chemotherapy in Patients With Metastatic Colorectal Cancer (FIRE-9

NCT: NCT05008809 · Enrollment: 507 · Sites: Amberg, Germany, Bad Saarow, Germany, Bayreuth, Germany, Berlin, Germany, Bochum, Germany

SIGNAL PHASE3 Stage III RAS RESEARCH • ACTIVE_NOT_RECRUITING 9

Comparison of the Pathological Effect Between 2 and 4 Cycles Neoadjuvant CAPOX for Low/Intermediate Risk II/III Rectal Cancer

NCT: NCT04922853 · Enrollment: 554 · Sites: Chengdu, China, Kunming, China, Wuxi, China

HIGH PHASE3 Stage IV KRAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 9

Phase 3 Study of MRTX849 With Cetuximab vs Chemotherapy in Patients With Advanced Colorectal Cancer With KRAS G12C Mutation (KRYSTAL-10)

NCT: NCT04793958 · Enrollment: 461 · Sites: Alabaster, American Fork, Atlanta, Auburn, Aurora

SIGNAL PHASE3 Stage III RESEARCH • RECRUITING 9

Thymosin-alpha 1 for Adjuvant Treatment After Radical Resection of High-risk Stage II and III Colorectal Cancer

This clinical trial is testing the efficacy and safety of Thymosin-alpha 1 as an adjuvant treatment for patients with high-risk stage II and III colorectal cancer following radical resection. The intervention, Thymosin-alpha 1, is a drug that is believed to enhance immune response and may help reduce the risk of cancer recurrence and metastasis. Currently in Phase 3 and actively recruiting, this trial aims to enroll 2,500 participants to evaluate the 3-year disease-free survival rate. Eligibility criteria may include specific high-risk characteristics associated with stage II and III colorectal cancer.

NCT: NCT05086614 · Enrollment: 2500 · Sites: Shanghai, China

SIGNAL PHASE3 Stage IV RAS RESEARCH • ACTIVE_NOT_RECRUITING 9

Neoadjuvant FOLFOX Chemotherapy for Patients With Locally Advanced Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant FOLFOX chemotherapy in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy before surgery, which may enhance treatment outcomes by targeting microscopic metastatic cancer cells earlier and reducing surgical complications. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and results may soon be available for clinical application. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT03426904 · Enrollment: 708 · Sites: Daegu, South Korea, Hwasun, South Korea, Seoul, South Korea, Suwon, South Korea

SIGNAL PHASE3 Stage IV RAS RESEARCH • RECRUITING 9

Preoperative FOLFOX Versus Postoperative Risk-adapted Chemotherapy in Patients With Locally Advanced Rectal Cancer

This clinical trial is testing the effectiveness of preoperative FOLFOX compared to postoperative risk-adapted chemotherapy in patients with locally advanced rectal cancer who are at low risk for local failure. The intervention involves various chemotherapy regimens, including mFOLFOX and XELOX, administered either before or after surgery, which is important for determining the optimal timing of treatment to improve disease-free survival. This is a Phase 3 trial currently recruiting 550 participants, indicating that it is in a critical stage of evaluating the intervention's efficacy and safety for patients. Eligibility criteria include the presence of RAS biomarkers, which may influence treatment decisions.

NCT: NCT04495088 · Enrollment: 550 · Sites: Mannheim, Germany

SIGNAL PHASE3 Stage IV RESEARCH • ACTIVE_NOT_RECRUITING 9

Bevacizumab Plus mFOLFOXIRI as First-line Treatment for Patients With Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of the triplet regimen mFOLFOXIRI combined with bevacizumab compared to mFOLFOX6 plus bevacizumab as a first-line treatment for patients with unresectable metastatic colorectal cancer. The intervention is significant as it aims to improve treatment outcomes while addressing the high incidence of adverse events associated with existing regimens in the Asian population. This is a Phase 3 trial that is currently active but not recruiting, indicating that the study is ongoing and data is being collected, but no new participants are being enrolled at this time. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

NCT: NCT04230187 · Enrollment: 528 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage III MSI/MMR RESEARCH • RECRUITING 9

CIRCULATING TUMOR DNA BASED DECISION FOR ADJUVANT TREATMENT IN COLON CANCER STAGE II

NCT: NCT04120701 · Enrollment: 1980 · Sites: Dijon, France

SIGNAL PHASE2, PHASE3 Stage II ctDNA LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 9

Circulating Tumor DNA Testing in Predicting Treatment for Patients With Stage IIA Colon Cancer After Surgery

NCT: NCT04068103 · Enrollment: 635 · Sites: Aberdeen, Abilene, Akron, Albany, Albemarle

SIGNAL PHASE3 Stage IV RESEARCH • NOT_YET_RECRUITING 9

Triplet Combination or Doublet Regimen Versus Chemoradiation as Neoadjuvant Therapy for Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy of chemotherapy alone using the FOLFOXIRI or mFOLFOX protocols compared to standard chemoradiation as neoadjuvant therapy for patients with locally advanced rectal cancer. The intervention involves two chemotherapy regimens, which may reduce surgical complications and improve quality of life compared to traditional chemoradiation. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria have not been specified, but the study focuses on patients with locally advanced rectal cancer.

NCT: NCT03975049 · Enrollment: 933 ·

MEDIUM PHASE3 Stage II RAS IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 9

Nivolumab After Combined Modality Therapy in Treating Patients With High Risk Stage II-IIIB Anal Cancer

NCT: NCT03233711 · Enrollment: 344 · Sites: Aberdeen, Abington, Aitkin, Albemarle, Albuquerque

SIGNAL PHASE3 Stage III RESEARCH • ACTIVE_NOT_RECRUITING 9

Adjuvant Low Dose Aspirin in Colorectal Cancer

NCT: NCT02647099 · Enrollment: 626 · Sites: Aarhus, Denmark, Espoo, Finland, Falun, Sweden, Gothenburg, Sweden, Helsinki, Finland

HIGH NA All Stages PHASE III TRIAL • RECRUITING 9

Magnetic Resonance Tumour Regression Grade (mrTRG) as a Novel Biomarker

The TRIGGER trial is testing the effectiveness of using magnetic resonance tumour regression grade (mrTRG) as a biomarker to determine whether patients with locally advanced rectal cancer can avoid surgery after pre-operative treatment. The intervention involves high-resolution MRI scans and mrTRG assessments to evaluate treatment response, which is significant as it may allow for a watch-and-wait approach for patients showing a good response. This is a Phase III non-CTIMP trial currently recruiting 441 participants, which means eligible patients can potentially contribute to important findings while receiving standard care. Eligibility includes patients undergoing any pre-operative treatment for locally advanced rectal cancer.

NCT: NCT02704520 · Enrollment: 441 · Sites: Aberdeen, United Kingdom, Basingstoke, United Kingdom, Bristol, United Kingdom, Colchester, United Kingdom, East Kilbride, United Kingdom

SIGNAL PHASE3 Stage III RESEARCH • RECRUITING 9

Randomised Study Evaluating Adjuvant Chemotherapy After Resection of Stage III Colonic Adenocarcinoma in Patients of 70 and Over

This clinical trial is evaluating the effectiveness of adjuvant chemotherapy in patients aged 70 and over who have undergone resection for stage III colonic adenocarcinoma. The interventions being tested include LV5FU2, capecitabine, FOLFOX4, XELOX, and observation, which are important for understanding the potential benefits of chemotherapy in this age group. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into disease-free survival outcomes for elderly patients, which may influence treatment decisions. Eligibility criteria include being 70 years or older and having a diagnosis of stage III colonic adenocarcinoma.

NCT: NCT02355379 · Enrollment: 774 · Sites: Abbeville, France, Agen, France, Aix-en-Provence, France, Albi, France, Alès, France

SIGNAL PHASE3 Stage IV RESEARCH • ACTIVE_NOT_RECRUITING 9

Octreotide Acetate and Recombinant Interferon Alfa-2b or Bevacizumab in Treating Patients With Metastatic or Locally Advanced, High-Risk Neuroendocrine Tumor

NCT: NCT00569127 · Enrollment: 427 · Sites: Adrian, Akron, Alamosa, Albuquerque, Alexandria

MEDIUM PHASE2, PHASE3 All Stages MSI/MMR IMMUNOTHERAPY • NOT_YET_RECRUITING 8

Yang et al. Anti-PD-1/CTLA-4 Dual Immunotherapy for LARC

NCT: NCT07596290 · Enrollment: 342 ·

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 8

A Single-Arm, Phase II, Multicenter Clinical Study of Neoadjuvant Encorafenib Plus Cetuximab N01 and mFOLFOX6 in Patients With Locally Advanced BRAF V600E-Mutant Colorectal Cancer With or Without Resectable Metastases

This clinical trial is testing the safety and efficacy of neoadjuvant therapy combining encorafenib, cetuximab N01, and mFOLFOX6 in patients with locally advanced BRAF V600E-mutant colorectal cancer, with or without resectable metastases. The intervention involves a targeted drug regimen aimed at improving tumor control and long-term outcomes in this specific molecular subgroup. Currently in Phase II and actively recruiting participants, this trial offers patients the opportunity to receive a novel treatment approach before surgery, with the primary outcome focused on the 18-month disease-free survival rate. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MSI, and MMR status.

NCT: NCT07789210 · Enrollment: 25 · Sites: Hangzhou, China

SIGNAL PHASE2, PHASE3 Stage IV BRAF RESEARCH • NOT_YET_RECRUITING 8

A Clinical Study of Liposomal Irinotecan for Second-Line Therapy in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of various treatment regimens involving liposomal irinotecan for patients with metastatic colorectal cancer who are receiving second-line therapy. The interventions include liposomal irinotecan combined with 5-FU/LV and bevacizumab, with one group also receiving Enlonstobart, which is significant for evaluating potential improvements in treatment outcomes. The trial is currently in Phase II and Phase III stages and is not yet recruiting participants, meaning patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarker assessments such as BRAF, RAS, microsatellite instability, and mismatch repair status.

NCT: NCT07585279 · Enrollment: 408 · Sites: Shijiazhuang, China

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • NOT_YET_RECRUITING 8

SCRT Based iTNT vs. LCRT Based TNT for MSS Locally Advanced Rectal Cancer

This phase III clinical trial is investigating the effectiveness of short-course radiotherapy combined with Serplulimab versus long-course chemoradiotherapy in patients with locally advanced rectal cancer (LARC) who have at least one high-risk feature. The intervention involves two treatment regimens: the TNT group receives long-course chemoradiotherapy with capecitabine, while the iTNT group receives short-course radiotherapy followed by Serplulimab and capecitabine. Currently, the trial is not yet recruiting participants, which means patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include having locally advanced rectal cancer with specific high-risk features and may involve biomarker assessments for MSI and MMR.

NCT: NCT07551479 · Enrollment: 612 · Sites: Shanghai, China

SIGNAL PHASE2, PHASE3 All Stages RAS RESEARCH • NOT_YET_RECRUITING 8

Injection of IP-001 Into Thermally Ablated Hepatic Tumors in Patients With Colorectal Liver Metastases

This clinical trial is testing the efficacy of the injectable drug IP-001 in patients with colorectal cancer that has metastasized to the liver, following standard microwave tumor ablation. The intervention involves administering either a high-dose or low-dose solution of IP-001 after the ablation procedure to assess its potential in prolonging progression-free

survival compared to ablation alone. This study is in Phase 2 and Phase 3 and is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include being at least 18 years old and having RAS biomarker status relevant to the study.

NCT: NCT07321847 · Enrollment: 717 ·

SIGNAL PHASE3 Early Detection RESEARCH • NOT_YET_RECRUITING 8

EFFECTIVENESS OF EARLY INFORMATION AND ENDORSMENT BY GP TO INCREASE COLONOSCOPY FOLLOW-UP AND REDUCE DELAYS IN PEOPLE WITH A POSITIVE COLORECTAL CANCER SCREENING TEST. (SUIVICOLO)

The SUIVICOLO study is testing strategies to increase the rate of colonoscopy follow-up in individuals with a positive colorectal cancer screening test. The intervention includes providing an information leaflet about colonoscopy follow-up and endorsement of follow-up letters from general practitioners, which may enhance patient engagement and adherence to necessary diagnostic procedures. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria have not been specified in the provided information.

NCT: NCT07520578 · Enrollment: 2000 · Sites: Tours, France

SIGNAL PHASE2, PHASE3 Stage III MSI/MMR RESEARCH • RECRUITING 8

Efficacy and Safety Comparison of Short-course Radiotherapy Followed by CapeOX Chemotherapy Plus Toripalimab With or Without Concurrent Surufatinib in Neoadjuvant Therapy for Mid-to-low Localized Rectal Cancer of High-risk Criteria

This clinical trial is testing the efficacy and safety of short-course radiotherapy followed by CapeOX chemotherapy combined with Toripalimab, with or without concurrent Surufatinib, in patients with mid-to-low localized rectal cancer who meet high-risk criteria. The intervention involves a combination of radiotherapy, chemotherapy, and immunotherapy, aiming to enhance the pathological complete response rate in this patient population. Currently in Phase 2 and Phase 3, the trial is recruiting participants, which means eligible patients may have the opportunity to access this investigational treatment. Eligibility criteria include specific biomarkers such as MMR and RAS.

NCT: NCT07505472 · Enrollment: 212 · Sites: Changchun, China

SIGNAL PHASE3 Stage IV RAS RESEARCH • NOT_YET_RECRUITING 8

Fruquintinib Versus Bevacizumab Plus Chemotherapy in Second-Line RAS-Mutant Metastatic Colorectal Cancer (FRU-RAS)

NCT: NCT07362836 · Enrollment: 224 ·

MEDIUM PHASE2 All Stages MSI/MMR LIQUID BIOPSY • NOT_YET_RECRUITING 8

Phase II Basket Trial: Zanidatamab Plus Tislelizumab in HER2-Positive GI Tumors (UNION-HER2-BASKET)

NCT: NCT07243938 · Enrollment: 70 ·

SIGNAL PHASE3 Stage III MSI/MMR RESEARCH • NOT_YET_RECRUITING 8

PD-1 + FOLFOXIRI vs CAPOX as Total Neoadjuvant Therapy for pMMR Low Rectal Cancer

This clinical trial is testing the effectiveness of combining a PD-1 antibody (serplulimab) with FOLFOXIRI chemotherapy and radiotherapy compared to CAPOX chemotherapy with radiotherapy for adults with pMMR locally advanced low rectal cancer. The intervention aims to determine if this combination improves 3-year event-free survival and enhances the likelihood of clinical complete response while minimizing the need for a permanent stoma. This is a Phase 3 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include having pMMR low rectal cancer, and the trial will assess biomarkers such as MSI and MMR.

NCT: NCT07277842 · Enrollment: 382 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage IV RAS PHASE I TRIAL • RECRUITING 8

A Phase Ib/III Study of Suvemcitug Plus FTD/TPI in Participants With Refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of Suvemcitug and trifluridine/tipiracil tablets in participants with refractory metastatic colorectal cancer. The intervention involves administering Suvemcitug injections alongside trifluridine/tipiracil tablets to assess its safety and efficacy compared to a placebo. Currently in Phase III and actively recruiting, this trial aims to determine if the combination treatment is more effective than the standard treatment alone for patients. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT07361003 · Enrollment: 464 · Sites: Fuzhou, China, Hangzhou, China, Harbin, China, Jinan, China, Nanjing, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 8

A Phase II Clinical Study Evaluating SSGJ-706 in Combination Therapy for Advanced Gastrointestinal Cancers

NCT: NCT07233850 · Enrollment: 300 · Sites: Wuhan, China

SIGNAL PHASE3 All Stages NGS RESEARCH • RECRUITING 8

Evaluating Whether an Educational Website Called Current Together After Cancer (CTAC) Improves Follow-up Care for Colorectal Cancer Survivors

This phase III clinical trial is evaluating the effectiveness of an educational website called Current Together After Cancer (CTAC) in improving follow-up care for colorectal cancer survivors who have undergone surgical resection. The intervention involves access to the CTAC website, which aims to enhance patients' understanding of surveillance care and boost their confidence in managing their follow-up, thereby promoting adherence to guideline-concordant surveillance practices. The trial is currently recruiting participants, with an enrollment goal of 1,057 individuals, which means eligible patients can still join the study. Eligibility criteria include having a history of surgically removed colorectal cancer and may involve specific biomarker assessments such as next-generation sequencing (NGS).

NCT: NCT07018869 · Enrollment: 1057 · Sites: Aberdeen, Aitkin, Allentown, Alma, Alpena

SIGNAL PHASE2, PHASE3 All Stages RESEARCH • NOT_YET_RECRUITING 8

Impact of Comprehensive Geriatric Management on Morbidity and Quality of Life in Elderly Patients Undergoing Major Hepatectomy and Pancreaticoduodenectomy for Cancer

NCT: NCT06999512 · Enrollment: 526 · Sites: Clermont-Ferrand, France, Le Kremlin-Bicêtre, France, Lille, France, Lyon, France, Marseille, France

MEDIUM PHASE2 Stage IV KRAS LIQUID BIOPSY • NOT_YET_RECRUITING 8

5-fluorouracil Plus Panitumumab (Anti-EGFR) and Sotorasib (KRAS G12C Inhibitor) in First-line Treatment of Patients Non-eligible for a Doublet/Triplet Chemotherapy With Advanced Unresectab

NCT: NCT07124884 · Enrollment: 300 · Sites: Angers, France, Annecy, France, Antony, France, Aurillac, France, Aviano, Italy

SIGNAL PHASE3 Stage III MSI/MMR RESEARCH • NOT_YET_RECRUITING 8

Neoadjuvant Chemoradiotherapy Followed by Chemotherapy With or Without Tislelizumab for Resectable Ultra-low Rectal Cancer: The RELIEVE-02 Study

The RELIEVE-02 study is a Phase 3 clinical trial designed for patients with pathologically confirmed, previously untreated, resectable ultra-low rectal adenocarcinoma, specifically those with MSI-L or MSS/pMMR tumors. The trial investigates the effectiveness of long-course chemoradiotherapy followed by chemotherapy with or without the addition of Tislelizumab, an immunotherapy drug, to assess its impact on treatment outcomes. Currently, the trial is not yet recruiting participants, which means patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarker requirements such as MSI, MMR, and NGS status.

NCT: NCT07132463 · Enrollment: 154 ·

SIGNAL PHASE3 All Stages KRAS RESEARCH • RECRUITING 8

A Clinical Study of Calderasib (MK-1084) With Targeted Therapy and Chemotherapy in People With Colorectal Cancer (MK-1084-012/KANDLELIT-012)

NCT: NCT06997497 · Enrollment: 477 · Sites: Atlanta, Baltimore, Billings, Boise, Centerville

SIGNAL PHASE3 Stage IV RESEARCH • NOT_YET_RECRUITING 8

QL1706 Injection Plus Bevacizumab and XELOX vs Placebo Plus Bevacizumab and XELOX as First-Line Treatment of Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of QL1706 Injection combined with bevacizumab and XELOX in patients with unresectable metastatic colorectal cancer (mCRC). The intervention involves the use of QL1706, a drug being evaluated against a placebo, both in conjunction with bevacizumab and the chemotherapy regimen XELOX, which is significant for its potential impact on treatment outcomes. This is a Phase 3 trial that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Specific eligibility criteria or biomarker requirements have not been detailed in the provided information.

NCT: NCT07025239 · Enrollment: 430 ·

SIGNAL PHASE3 Stage III BRAF RESEARCH • RECRUITING 8

Adjuvant FOLFIRI Based-chemotherapy After Resection of CLM Responding to Preoperative FOLFIRI

This clinical trial is testing the effectiveness of postoperative reintroduction of FOLFIRI-based chemotherapy in patients who have undergone resection of colorectal liver metastases and have shown a good response to preoperative FOLFIRI treatment. The intervention involves administering FOLFIRI chemotherapy after surgery, which may provide benefits for patients who typically do not receive postoperative treatment. This is a Phase III trial currently recruiting participants, which means eligible patients may have the opportunity to contribute to research that could influence future treatment standards. Eligibility criteria include having a contraindication to preoperative oxaliplatin-based chemotherapy and demonstrating an objective response to prior treatment, with specific biomarker assessments for BRAF and RAS.

NCT: NCT06501482 · Enrollment: 254 · Sites: Paris, France

SIGNAL PHASE3 All Stages RESEARCH • NOT_YET_RECRUITING 8

The Use of Indocyanine Green Near-infrared Fluorescence for Bowel Perfusion Quantitative Assessment in Order to Prevent Anastomotic Leakage in Colorectal Surgery

This clinical trial is testing the use of indocyanine green (ICG) near-infrared fluorescence imaging for bowel perfusion assessment in patients undergoing colorectal surgery to prevent anastomotic leakage. The intervention involves ICG-guided bowel perfusion assessment compared to a conventional bowel anastomosis group, which is significant as anastomotic leakage is a serious complication that can increase mortality rates. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria have not been specified in the provided information.

NCT: NCT06845306 · Enrollment: 1268 · Sites: Vsevolozhsk District, Russia

MEDIUM PHASE2 Stage IV KRAS LIQUID BIOPSY • RECRUITING 8

Safety and Efficacy of IB-FOLFIRI in BRAF V600E-Mutant Metastatic Colorectal Cancer

This clinical trial is testing the safety and efficacy of a treatment regimen called IB-FOLFIRI, which includes Iparomlimab, Tuvonralimab, bevacizumab, and FOLFIRI, for adults with BRAF V600E-mutant metastatic colorectal cancer (mCRC). The intervention aims to determine if this combination improves clinical outcomes compared to historical data and to evaluate its safety profile in this specific patient population. Currently in Phase 2 and actively recruiting participants, this trial offers an opportunity for patients to receive a potentially beneficial treatment while contributing to research. Eligibility criteria include the presence of specific biomarkers such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, DMMR, and ctDNA.

NCT: NCT07150247 · Enrollment: 20 · Sites: Guangzhou, China

MEDIUM NA All Stages KRAS LIQUID BIOPSY • RECRUITING 8

Clinical Utility of ctDNA in Detecting Resistance Mechanisms and Delivering Precision Medicine to Cancer Patients

NCT: NCT07673861 · Enrollment: 100 · Sites: London, United Kingdom

SIGNAL PHASE3 Stage IV RAS RESEARCH • RECRUITING 8

Trifluridine/Tipiracil Plus Regorafenib vs Trifluridine/Tipiracil Plus Bevacizumab for Refractory Metastatic Colorectal Cancer

NCT: NCT06992648 · Enrollment: 302 · Sites: Hangzhou, China

SIGNAL PHASE3 All Stages RAS RESEARCH • NOT_YET_RECRUITING 8

Pre-Operative Treatment in REseCTable COlon CanceR

This clinical trial is testing the efficacy of preoperative treatments for patients with advanced colon cancer, including those with tumors in the upper third of the rectum, who are clinically staged as cT3-4 and/or cN+. The intervention involves administering mFOLFOXIRI, mFOLFOX-6, or CAPOX prior to surgery, which is significant for evaluating potential improvements in disease-free survival and patient quality of life compared to the standard care approach. This is a phase III trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific staging and biomarker assessments, such as RAS and microsatellite status.

NCT: NCT06899477 · Enrollment: 714 · Sites: Berlin, Germany, Berlin-Spandau, Germany, Chemnitz, Germany, Cologne, Germany, Dessau, Germany

SIGNAL PHASE2, PHASE3 Stage III RESEARCH • RECRUITING 8

Total Neoadjuvant Therapy and Organ Preservation Versus Surgery for Rectal Cancer.

NCT: NCT06758830 · Enrollment: 400 · Sites: Vilnius, Lithuania

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 8

Combination of IV Ascorbic Acid and Adebrelimab in Metastatic Colorectal Cancer

This clinical trial is testing the combination of intravenous ascorbic acid, Adebrelimab, and the FOLFOX protocol in patients with metastatic colorectal cancer who have high expression of the GLUT3 biomarker. The intervention involves administering ascorbic acid alongside a PD-L1 antibody and standard chemotherapy, which may enhance treatment efficacy based on previous preclinical findings. This is a Phase 3 trial currently recruiting 400 participants, indicating that it is in the later stages of testing and may provide valuable information on the effectiveness of this combination therapy for patients. Eligibility criteria include having metastatic colorectal cancer with high GLUT3 expression.

NCT: NCT04516681 · Enrollment: 400 · Sites: Shanhai, China

MEDIUM PHASE3 All Stages MSI/MMR IMMUNOTHERAPY • RECRUITING 8

Personalized Long-course Radiotherapy Plus Chemotherapy With or Without Immunotherapy for LARC: PALACE Study

The PALACE study is a phase III clinical trial designed for patients with locally advanced rectal cancer (LARC) to evaluate the complete response rate to personalized long-course radiotherapy combined with chemotherapy, with or without the anti-PD-1 antibody drug Serplulimab. This intervention is significant as it aims to determine the effectiveness of adding immunotherapy to standard treatment protocols. Currently, the trial is recruiting participants, which means eligible patients have the opportunity to join and contribute to the research. Eligibility criteria include specific biomarkers such as MMR and RAS.

NCT: NCT07020247 · Enrollment: 184 · Sites: Chengdu, China

MEDIUM PHASE3 Stage IV MSI/MMR IMMUNOTHERAPY • RECRUITING 8

FOLFOXIRI Plus Bevacizumab With or Without Atezolizumab as 1st Line Treatment of pMMR and IS IC-High Metastatic Colorectal Cancer Patients.

NCT: NCT06733038 · Enrollment: 238 · Sites: Aviano, Italy, Brescia, Italy, Catania, Italy, Florence, Italy, Livorno, Italy

SIGNAL PHASE3 Stage III MSI/MMR RESEARCH • RECRUITING 8

Planed Organ Preservation for Low Rectal Cancer After Neoadjuvant Chemotherapy (OPLAR)

The OPLAR trial is testing the effectiveness of total neoadjuvant therapy (TNT) versus immunotherapy combined with TNT (iTNT) in patients with low and intermediate-risk rectal cancer. The intervention involves administering two cycles of XELOX chemotherapy followed by either long-course chemoradiotherapy with consolidation chemotherapy or the same regimen with added immunotherapy, aiming to improve organ preservation rates. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to receive cutting-edge treatment while contributing to important research. Eligibility criteria include patients with effective neoadjuvant chemotherapy, as determined by high-definition MRI showing a significant reduction in tumor size.

NCT: NCT07070622 · Enrollment: 186 · Sites: Chengdu, China

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 8

The Sagittarius Trial

The Sagittarius Trial is testing the effectiveness of personalized therapy based on liquid biopsy results in patients with high-risk stage II and stage III colorectal cancer. The intervention involves a combination of drugs, including Oxaliplatin, Capecitabine, Folinic acid, Fluorouracil, and Temozolomide, which aims to improve outcomes and quality of life compared to conventional therapy. This is a Phase 3 trial currently recruiting 700 participants, which means eligible patients may have the opportunity to receive either standard or customized treatment based on their tumor genetics. Eligibility criteria include being diagnosed with high-risk stage II or stage III colon cancer and undergoing liquid biopsy testing.

NCT: NCT06490536 · Enrollment: 700 · Sites: Barcelona, Spain, Berlin, Germany, Biella, Italy, Brescia, Italy, Candiolo, Italy

SIGNAL PHASE3 Stage III RESEARCH • NOT_YET_RECRUITING 8

Neoadjuvant Chemoradiotherapy Versus Total Neoadjuvant Therapy in the Treatment of T3 Rectal Cancer

NCT: NCT06097416 · Enrollment: 100 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 8

Node-Sparing Short-Course Radiation With CAPOX and Sintilimab for MSS Locally Advanced Colon Cancer: A Randomized, Prospective, Multicenter Study

This clinical trial is testing the efficacy of node-sparing short-course radiotherapy combined with CAPOX chemotherapy and Sintilimab for patients with MSS locally advanced colon cancer. The intervention aims to evaluate the potential benefits of this combination therapy as neoadjuvant treatment, focusing on the role of regional lymph nodes in tumor immunotherapy. Currently in Phase 2 and not yet recruiting, this trial will provide insights into treatment options for eligible patients once enrollment begins. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06594692 · Enrollment: 140 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 8

Dostarlimab for Locally Advanced or Metastatic Cancer (Non-colorectal/Non-endometrial) With Tumor dMMR/MSI

This clinical trial is testing the efficacy of dostarlimab, an immunotherapy drug, for adult patients with locally advanced or metastatic cancers that are not colorectal or endometrial and have deficient mismatch repair (dMMR) or microsatellite instability (MSI). The intervention involves comparing dostarlimab to standard chemotherapy as a first-line treatment, which is significant for understanding alternative therapeutic options for these specific cancer types. This is a Phase 2 trial currently recruiting 120 participants, which means eligible patients may have the opportunity to receive either treatment while contributing to research on their effectiveness. Eligibility criteria include adults aged 18 and older with specific cancer types, including duodenum and small bowel adenocarcinoma, gastric adenocarcinoma, and others with documented dMMR/MSI status.

NCT: NCT06333314 · Enrollment: 120 · Sites: Angers, France, Avignon, France, Besançon, France, Brest, France, Caen, France

MEDIUM PHASE2 Stage IV KRAS LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 8

Phase II Study of ctDNA-guided Encorafenib Plus Cetuximab Retreatment in Patients BRAF V600E Mutated mCRC

This clinical trial is testing the effectiveness of ctDNA-guided retreatment with encorafenib plus cetuximab in patients with BRAF V600E mutated metastatic colorectal cancer (mCRC). The intervention involves the combination of the drugs encorafenib and cetuximab, which is significant for patients who have previously benefited from these treatments. This is a Phase II study that is currently active but not recruiting, indicating that it is ongoing and assessing the treatment's efficacy in a defined patient population. Eligibility criteria include having specific genetic mutations and statuses, such as BRAF V600E mutation and wild-type KRAS, NRAS, and MAP2K1, as well as ctDNA assessments at the time of entry.

NCT: NCT06578559 · Enrollment: 16 · Sites: Meldola, Italy, Milan, Italy, Monserrato, Italy, Naples, Italy, Padova, Italy

SIGNAL PHASE3 Stage IV KRAS RESEARCH • RECRUITING 8

Study of Sotorasib, Panitumumab and FOLFIRI Versus FOLFIRI With or Without Bevacizumab-awwb in Treatment-naïve Participants With Metastatic Colorectal Cancer With KRAS p.G12C Mutation

NCT: NCT06252649 · Enrollment: 450 · Sites: Atlanta, Bridgeton, Chandler, Chicago, Detroit

SIGNAL PHASE3 Stage IV RESEARCH • ACTIVE_NOT_RECRUITING 8

Borderline of CME/D3 Dissection for Local Advanced Right-sided Colon Cancer

This clinical trial is testing the effectiveness of SMA lymph node dissection compared to SMV lymph node dissection in patients with locally advanced right-sided colon cancer. The intervention involves two surgical procedures aimed at improving outcomes by addressing the lymph node dissection boundaries in this specific cancer type. Currently in Phase 3 and marked as active but not recruiting, this trial is designed to provide insights that could influence surgical practices for patients with this condition. Eligibility criteria include having advanced right-sided colon cancer, with a total enrollment of 896 participants.

NCT: NCT06709144 · Enrollment: 896 · Sites: Hangzhou, China

SIGNAL PHASE3 Stage III MSI/MMR RESEARCH • RECRUITING 8

Neoadjuvant Chemotherapy, Excision And Observation vs Chemoradiotherapy For Rectal Cancer

NCT: NCT06205485 · Enrollment: 250 · Sites: Albuquerque, Atlanta, Aurora, Basking Ridge, Bellevue

SIGNAL PHASE2, PHASE3 Stage IV MSI/MMR RESEARCH • NOT_YET_RECRUITING 8

Neoadjuvant Fruquintinib Plus Tislelizumab Combined With mCapeOX Versus CapeOX for Mid-high pMMR/MSS Locally Advanced Rectal Cancer

NCT: NCT06443671 · Enrollment: 132 · Sites: Beijing, China

SIGNAL PHASE3 Stage III MSI/MMR RESEARCH • RECRUITING 8

Neoadjuvant Chemotherapy for Obstructive Colon cancER First Treated by cOlostomy

NCT: NCT06107920 · Enrollment: 232 · Sites: Amiens, France, Beauvais, France, Besançon, France, Bobigny, France, Caen, France

SIGNAL PHASE3 Stage IV BRAF RESEARCH • NOT_YET_RECRUITING 8

Nimotuzumab Combined With Trifluridine/Tipiracil in the Treatment of Refractory Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of nimotuzumab combined with trifluridine/tipiracil in patients with refractory metastatic colorectal cancer (mCRC). The intervention involves the administration of nimotuzumab, a monoclonal antibody, alongside trifluridine/tipiracil, a chemotherapy regimen, to assess its impact on overall survival. This

is a Phase 3 trial that is currently not yet recruiting participants, indicating that it is in the preparatory stages before patient enrollment begins. Eligibility criteria include specific biomarker requirements related to BRAF, MMR, and RAS.

NCT: NCT06343116 · Enrollment: 420 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 8

Neoadjuvant Low-dose Immunotherapy in Locally Advanced MMR-deficient Colorectal Cancer

NCT: NCT07523763 · Enrollment: 30 · Sites: Moscow, Russia

HIGH PHASE3 Stage IV PHASE III TRIAL • RECRUITING 8

Phase III Study Evaluating Induction Chemotherapy Followed by Chemoradiotherapy Compared to Standard Chemoradiotherapy for Locally Advanced SCCA

This Phase III clinical trial is evaluating the effectiveness of induction chemotherapy using the modified DCF protocol (mDCF) followed by chemoradiotherapy compared to standard chemoradiotherapy for patients with locally advanced squamous cell carcinoma of the anus (SCCA). The intervention involves administering mDCF, followed by concomitant chemotherapy with Capecitabine and Mitomycin-C, along with radiotherapy, aiming to improve disease-related event-free survival (DREFS) at two years. The trial is currently recruiting 310 participants, which means eligible patients can enroll to potentially access this treatment approach. Eligibility criteria may include specific tumor stages and the presence of biomarkers related to HPV.

NCT: NCT06207981 · Enrollment: 310 · Sites: Agen, France, Aix-en-Provence, France, Amiens, France, Antony, France, Argenteuil, France

SIGNAL PHASE2, PHASE3 Stage III BRAF RESEARCH • RECRUITING 8

Neoadjuvant Serplulimab & Bevacizumab With FOLFOX vs. FOLFOX Alone in RAS/BRAF WT, pMMR/MSS CRC Patients

This clinical trial is testing the effectiveness of adding Serplulimab and Bevacizumab to standard FOLFOX chemotherapy in patients with RAS/BRAF wild-type, pMMR/MSS colorectal cancer liver metastases. The intervention involves the combination of Serplulimab, a PD-1 inhibitor, and Bevacizumab, an anti-angiogenesis agent, to potentially enhance the immune response and improve postoperative outcomes. This study is currently in Phase 2 and Phase 3 and is actively recruiting participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include specific biomarker requirements related to BRAF, MMR, and RAS status.

NCT: NCT06280495 · Enrollment: 156 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage IV KRAS RESEARCH • RECRUITING 8

Other Oncogene Mutations for Anti-EGFR Efficacy in Patients With Left-sided RAS-wild Type Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of cetuximab in patients with left-sided RAS-wild type metastatic colorectal cancer. The intervention involves administering cetuximab alongside FOLFOX chemotherapy, with patients undergoing extensive molecular genetic profiling to determine their eligibility for anti-EGFR therapy based on specific oncogene mutations. This is a Phase 3 trial currently recruiting 355 participants, which means eligible patients may have the opportunity to receive a treatment that is being evaluated for its effectiveness in improving progression-free survival. Eligibility criteria include the presence of specific biomarkers such as KRAS, NRAS, BRAF, and RAS.

NCT: NCT06226857 · Enrollment: 355 · Sites: Moscow, Russia, Reutov, Russia

SIGNAL PHASE3 Stage IV RAS RESEARCH • ACTIVE_NOT_RECRUITING 8

Adaptive Boost Radiotherapy to Primary Lesions and Positive Nodes in the Neoadjuvant Treatment of Locally Advanced Rectal Cancer

This clinical trial is testing the effectiveness of online adaptive radiotherapy (ART) with a sequential boost compared to conventional non-adaptive intensity-modulated radiotherapy (IMRT) in patients with high-risk, node-positive locally advanced rectal cancer. The intervention involves ART aimed at improving pathologic complete response (pCR) rates, which is significant for treatment outcomes in this patient population. This is a Phase 3 trial that is currently active but not recruiting, meaning that while no new participants are being enrolled, the study is ongoing for those already involved. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT06246344 · Enrollment: 128 · Sites: Jinan, China

MEDIUM N/A Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 8

The Efficacy of Watch and Wait Strategy or Surgery After Neoadjuvant Immunotherapy for Locally Advanced Colorectal Cancer With dMMR/MSI-H Guided by MRD Dynamic Monitoring: A Single-center, Open-label, Prospective, Phase II Clinical Trial.

This clinical trial is testing the efficacy of a watch and wait strategy compared to surgery after neoadjuvant immunotherapy in patients with locally advanced colorectal cancer characterized by deficient mismatch repair or microsatellite instability-high (dMMR/MSI-H). The intervention involves the use of Tireilizumab, a PD-1 inhibitor, which is significant for its potential to enhance the immune response against cancer cells. Currently, the trial is in the recruiting phase, indicating that eligible patients can still enroll to participate in this study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite status, as well as the use of next-generation sequencing (NGS) for patient selection.

NCT: NCT06477991 · Enrollment: 22 · Sites: Kunming, China

MEDIUM PHASE2, PHASE3 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 8

DETERMINE Trial Treatment Arm 02: Atezolizumab in Adult, Paediatric and Teenage/Young Adult Patients With Cancers With High Tumour Mutational Burden (TMB) or Microsatellite Instability-high (MSI-high) or Proven Constitutional Mismatch Repair Deficiency (CMMRD) Disposition

NCT: NCT05770102 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL PHASE3 Stage IV BRAF RESEARCH • RECRUITING 8

Testing the Addition of Total Ablative Therapy to Usual Systemic Therapy Treatment for Limited Metastatic Colorectal Cancer, The ERASur Study

The ERASur Study is a phase III clinical trial evaluating the addition of total ablative therapy to usual systemic therapy in patients with limited metastatic colorectal cancer, specifically those with cancer that has spread to up to four body sites. The intervention includes stereotactic ablative radiotherapy, surgical resection, and microwave ablation, alongside chemotherapy, aiming to eliminate cancer at metastatic sites while managing systemic disease. Currently, the trial is in the recruiting phase, meaning eligible patients can still enroll to participate in this study. Eligibility criteria include specific biomarkers such as BRAF, MSI, RAS, microsatellite status, and next-generation sequencing (NGS) results.

NCT: NCT05673148 · Enrollment: 364 · Sites: Albuquerque, Ankeny, Ann Arbor, Antigo, Appleton

MEDIUM PHASE2 All Stages KRAS TARGETED THERAPY • RECRUITING 8

Pre-operative Targeted Treatments in Molecularly Selected Resectable Colorectal Cancer (UNICORN)

The UNICORN trial is testing pre-operative targeted treatments in patients with non-metastatic resectable colorectal cancer who have specific targetable molecular alterations. The intervention involves a combination of drugs, including Trastuzumab deruxtecan, Durvalumab, Panitumumab, Botensilimab, and Balstilimab, which are administered as a short-course pre-operative strategy to assess their effectiveness. This is a Phase 2 trial currently recruiting participants, which means eligible patients may have the opportunity to receive these targeted therapies before standard surgical treatment. Eligibility criteria include the presence of biomarkers such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, DMMR, and HER2.

NCT: NCT05845450 · Enrollment: 197 · Sites: Milan, Italy

SIGNAL PHASE2, PHASE3 Stage IV BRAF RESEARCH • RECRUITING 8

Regorafenib With Low-dose Chemotherapies and Aspirin Followed by Standard Chemotherapies in Metastatic Colorectal Cancer

This clinical trial is testing the combination of regorafenib with low-dose metronomic chemotherapies and aspirin for patients with metastatic colorectal cancer. The intervention aims to assess the effectiveness of this combination as a two-month induction therapy prior to standard chemotherapy initiation. Currently in Phase 2 and Phase 3, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment approach. Eligibility criteria include specific biomarkers such as BRAF, RAS, and microsatellite status.

NCT: NCT05462613 · Enrollment: 446 · Sites: Besançon, France, Clermont-Ferrand, France, Créteil, France, Dijon, France, Lyon, France

MEDIUM PHASE2, PHASE3 All Stages HER2 TARGETED THERAPY • RECRUITING 8

DETERMINE Trial Treatment Arm 04: Trastuzumab in Combination With Pertuzumab in Adult, Paediatric and Teenage/Young Adult Patients With Cancers With HER2 Amplification or Activating Mutations

The DETERMINE Trial Treatment Arm 04 is investigating the efficacy of trastuzumab in combination with pertuzumab in adult, pediatric, and teenage/young adult patients with cancers that exhibit HER2 amplification or activating mutations. This combination of drugs is already approved as a standard treatment for adult breast cancer, highlighting its established role in targeting HER2-related malignancies. Currently in Phase 2 and Phase 3, the trial is actively recruiting participants, which means eligible patients may have the opportunity to access this treatment option during the study. Eligibility criteria include the presence of HER2 amplification or mutations, as determined by next-generation sequencing or other biomarker assessments.

NCT: NCT05786716 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL PHASE3 All Stages ctDNA LIQUID BIOPSY • RECRUITING 8

Postoperation Maintenance Therapy for Resectable Liver Metastases of Colorectal Cancer Guided by ctDNA

This clinical trial is testing the effectiveness of a combination of adjuvant chemotherapy and maintenance therapy in patients with resectable liver metastases from colorectal cancer. The intervention includes colorectal resection surgery followed by the FOLFOX chemotherapy regimen and Capecitabine, which are important for assessing their impact on patient outcomes. Currently in Phase 3 and actively recruiting, this trial aims to determine if the combination therapy improves 3-year progression-free survival compared to chemotherapy alone. Eligible participants must have positive ctDNA after surgery, which will guide the treatment approach.

NCT: NCT05797077 · Enrollment: 346 · Sites: Guangzhou, China

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 8

Intraoperative Indocyanine Green Fluorescence Angiography in Colorectal Surgery to Prevent Anastomotic Leakage

The FLUOCOL-1 study is testing the effectiveness of intraoperative fluorescence angiography (IOFA) with indocyanine green (ICG) in preventing anastomotic leakage in patients undergoing colorectal surgery for cancer. The intervention involves the use of ICG to enhance visualization during surgery, which may help reduce the incidence of this serious complication. This is a Phase 3 clinical trial currently recruiting 1,010 participants, which means that eligible patients may have the opportunity to contribute to research that could influence future surgical practices. Eligibility criteria include patients undergoing left-sided or low anterior resection with anastomosis for colorectal cancer.

NCT: NCT05168839 · Enrollment: 1010 · Sites: Amiens, France, Annecy, France, Besançon, France, Bourgoin, France, Dijon, France

SIGNAL PHASE2, PHASE3 Stage IV RAS RESEARCH • NOT_YET_RECRUITING 8

Short-course Radiotherapy Followed by Chemotherapy and PD-1 Inhibitor for Locally Advanced Rectal Cancer

NCT: NCT05484024 · Enrollment: 588 · Sites: Beijing, China, Shenzhen, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 8

A Study of Encorafenib Plus Cetuximab Taken Together With Pembrolizumab Compared to Pembrolizumab Alone in People With Previously Untreated Metastatic Colorectal Cancer

This clinical trial is testing the combination of encorafenib and cetuximab alongside pembrolizumab in individuals with previously untreated metastatic colorectal cancer that exhibits specific genetic characteristics. The intervention involves administering encorafenib orally and cetuximab via intravenous infusion, in addition to pembrolizumab, to evaluate their effects on progression-free survival. This is a Phase 2 trial that is currently active but not recruiting new participants,

indicating that it is in the process of gathering data on the treatment's efficacy and safety. Eligibility criteria include the presence of BRAF mutations and conditions related to DNA mismatch repair, such as MSI-H or DMMR.

NCT: NCT05217446 · Enrollment: 108 · Sites: Germantown, Houston, Miami Beach, Phoenix, Scottsdale

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 8

A Study to Evaluate the Immunogenicity and Safety of HPV Vaccine in Healthy Female Participants Aged 9-26 Years in China

This clinical trial is evaluating the immunogenicity and safety of a recombinant nonavalent HPV vaccine in healthy female participants aged 9-26 years in China. The intervention involves comparing the immune response of the study vaccine to that of GARDASIL® 9, which is significant for understanding the effectiveness of the new vaccine formulation. This is a Phase 3 trial that is currently active but not recruiting, indicating that the study is in its final stages of testing before potential approval. Eligibility criteria include being a healthy female within the specified age range, with specific focus on different dosing schedules for various age groups.

NCT: NCT05662020 · Enrollment: 2750 · Sites: Kunming, China

SIGNAL PHASE3 All Stages MSI/MMR RESEARCH • ACTIVE_NOT_RECRUITING 8

Study Comparing the Standard Administration of IO Versus the Same IO Administered Each 3 Months in Patients in Response After 6 Months of Standard IO

This clinical trial is testing the effectiveness of a reduced dose intensity of immunotherapy (IO) in patients with colorectal cancer who have shown a response after six months of standard IO treatment. The intervention involves administering the same IO drug every three months instead of the standard schedule, which may help maintain disease control while potentially reducing treatment costs and toxicity. This is a Phase 3 trial that is currently active but not recruiting new participants, indicating that it is in the later stages of evaluation and may soon provide important insights for patient care. Eligibility criteria include specific biomarkers such as microsatellite instability (MSI).

NCT: NCT05078047 · Enrollment: 646 · Sites: Angers, France, Avignon, France, Bayonne, France, Bordeaux, France, Boulogne-sur-Mer, France

SIGNAL PHASE3 Stage IV BRAF RESEARCH • ACTIVE_NOT_RECRUITING 8

Combination With Sintilimab and XELOX+Bevacizumab as 1st Line Therapy in RAS-mutant Metastatic Colorectal Cancer

NCT: NCT05171660 · Enrollment: 446 · Sites: Hangzhou, China

SIGNAL PHASE2, PHASE3 Stage IV RAS RESEARCH • RECRUITING 8

Study Evaluating the Tailored Management of Locally-advanced Rectal Carcinoma

This clinical trial is evaluating the tailored management of locally advanced rectal carcinoma for patients who may benefit from personalized treatment approaches. The intervention includes a modified FOLFIRINOX regimen for induction chemotherapy, early tumor response evaluation using MRI volumetry, radiochemotherapy with Cap 50, and radical proctectomy with total mesorectal excision, aiming to improve surgical outcomes and reduce metastatic risks. Currently in Phase 2 and Phase 3, the trial is actively recruiting participants, which means eligible patients have the opportunity to contribute to research that may influence future treatment protocols. Eligibility criteria include the presence of RAS biomarkers, which are being assessed to help predict tumor response to treatment.

NCT: NCT04749108 · Enrollment: 1075 · Sites: Amiens, France, Annecy, France, Besançon, France, Bordeaux, France, Clermont-Ferrand, France

SIGNAL PHASE3 Stage IV RAS RESEARCH • RECRUITING 8

Neoadjuvant FOLFOXIRI Versus CapeOX Chemotherapy for Local Advanced Rectal Cancer

NCT: NCT05201430 · Enrollment: 300 · Sites: Shanghai, China

SIGNAL PHASE3 All Stages ctDNA LIQUID BIOPSY • RECRUITING 8

FOLICOLOR TRIAL: Following Therapy Response Through Liquid Biopsy in Metatstatic Colorectal Cancer Patients

The FOLICOLOR trial is testing a liquid biopsy-guided follow-up strategy for adults with unresectable, metastatic colorectal cancer (mCRC) who are beginning first-line systemic treatment. The intervention involves evaluating therapy response through NPY methylation-based circulating tumor DNA (ctDNA) analysis from blood samples, which may enhance clinical decision-making and improve patient outcomes. This is a Phase 3 trial currently recruiting 150 participants, indicating that patients may have the opportunity to contribute to research that could influence future treatment approaches. Eligible patients must be adults diagnosed with unresectable, metastatic colorectal cancer and starting first-line treatment.

NCT: NCT07558083 · Enrollment: 150 · Sites: Brasschaat, Belgium, Bruges, Belgium, Charleroi, Belgium, Edegem, Belgium, Ghent, Belgium

SIGNAL PHASE3 Stage IV RAS RESEARCH • ACTIVE_NOT_RECRUITING 8

Short-Course Radiotherapy Followed by Neoadjuvant Chemotherapy and Camrelizumab in Locally Advanced Rectal Cancer (UNION)

This clinical trial is testing the effectiveness of short-course radiotherapy followed by neoadjuvant chemotherapy and camrelizumab in patients with locally advanced rectal cancer (LARC). The intervention involves a combination of short-term radiotherapy, camrelizumab, and the chemotherapy regimen CAPOX, which is significant for its potential to improve treatment outcomes. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients may not be able to enroll at this time. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT04928807 · Enrollment: 231 · Sites: Wuhan, China

SIGNAL PHASE2, PHASE3 Stage III RAS RESEARCH • RECRUITING 8

Adjuvant mFOLFIRINOX for High-risk Stage III Colon Cancer

NCT: NCT05179889 · Enrollment: 308 · Sites: Daejeon, South Korea

SIGNAL PHASE3 Stage IV BRAF LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 8

Study in mCRC Patients RAS/BRAF wt Tissue and RAS Mutated LIquid Blopsy to Compare FOLFIRI Plus CetuxiMAb or BevacizumaB

This clinical trial is testing the effectiveness of FOLFIRI combined with either cetuximab or bevacizumab in patients with metastatic colorectal cancer (mCRC) who have RAS/BRAF wild type tissue and RAS mutations detected in liquid biopsy. The intervention involves comparing the anti-VEGF drug bevacizumab with the anti-EGFR drug cetuximab, both used alongside chemotherapy, to determine which combination leads to better progression-free survival (PFS). This is a phase III trial that is currently active but not recruiting new participants, indicating that it is in the later stages of evaluation and may provide insights into treatment options for eligible patients. Eligibility criteria include having RAS/BRAF wild type tissue and RAS mutations identified through liquid biopsy.

NCT: NCT04776655 · Enrollment: 280 · Sites: Coppito, Italy, Guastalla, Italy, Mirano, Italy, Reggio Emilia, Italy

SIGNAL PHASE3 Stage IV BRAF RESEARCH • RECRUITING 8

Conversion Therapy of RAS/BRAF Wild-Type Colorectal Cancer Patients With Initially Unresectable Liver Metastases

This clinical trial is testing the effectiveness of two treatment combinations for patients with RAS/BRAF wild-type colorectal cancer who have initially unresectable liver metastases. The interventions being compared are modified FOLFOXIRI (mFOLFOXIRI) combined with cetuximab and mFOLFOXIRI combined with bevacizumab, as both combinations may improve the response and resectability rates of liver metastases. This is a Phase 3 trial currently recruiting 508 participants, which means that eligible patients may have the opportunity to receive one of the treatment regimens while contributing to important research on their effectiveness. Eligibility criteria include having RAS/BRAF wild-type colorectal cancer and initially unresectable liver-only metastases, confirmed by a local multidisciplinary team.

NCT: NCT04687631 · Enrollment: 508 · Sites: Shanghai, China

MEDIUM N/A All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 8

National Cancer Institute "Cancer Moonshot Biobank"

The "Cancer Moonshot Biobank" is a clinical trial designed for cancer patients, focusing on the collection of biospecimens and associated clinical data. The intervention involves various procedures, including biospecimen collection, computed tomography, magnetic resonance imaging, and medical chart review, which are essential for understanding cancer progression and treatment responses. This study is currently active but not recruiting, indicating that while it is ongoing, new participants cannot be enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, HER2, microsatellite status, and mismatch repair status.

NCT: NCT04314401 · Enrollment: 1600 · Sites: Ames, Ann Arbor, Arroyo Grande, Atlanta, Augusta

SIGNAL PHASE3 All Stages RAS RESEARCH • ACTIVE_NOT_RECRUITING 8

Short RT Versus RCT,Followed by Chemo and Organ Preservation for Interm and High-risk Rectal Cancer Patients

NCT: NCT04246684 · Enrollment: 702 · Sites: Amberg, Germany, Bad Saarow, Germany, Bayreuth, Germany, Berlin, Germany, Bielefeld, Germany

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 8

Non Inferiority Study of Preoperative Chemotherapy Without Pelvic Irradiation for Rectal Cancer

NCT: NCT03875781 · Enrollment: 540 · Sites: Le Kremlin-Bicêtre, France

MEDIUM PHASE2 All Stages KRAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 8

QUILT-3.055: A Study of Combination Immunotherapies in Patients Who Have Previously Received Treatment With Immune Checkpoint Inhibitors

NCT: NCT03228667 · Enrollment: 40 · Sites: Anchorage, Billings, Boston, Buffalo, Charleston

SIGNAL PHASE3 Stage IV KRAS RESEARCH • ACTIVE_NOT_RECRUITING 8

Study to Evaluate the Efficacy of FOLFOX + Panitumumab Followed by FOLFIRI + Bevacizumab (Sequence 1) Versus FOLFOX + Bevacizumab Followed by FOLFIRI + Panitumumab (Sequence 2) in Untreated Patients With Wild-type RAS Metastatic, Primary Left-sided, Unresectable Colorectal Cancer

NCT: NCT03635021 · Enrollment: 419 · Sites: Madrid, Spain

SIGNAL PHASE3 Stage IV RAS RESEARCH • RECRUITING 8

Perioperative Versus Postoperative CapOX Chemotherapy for Locally Advanced Colon Cancer

This clinical trial is testing the effectiveness of perioperative versus postoperative capecitabine plus oxaliplatin (CapOX) chemotherapy in patients with locally advanced colon cancer. The intervention involves administering CapOX both before and after surgery compared to administering it only after surgery, which may enhance tumor suppression and improve surgical outcomes. This is a Phase 3 trial currently recruiting 1,370 participants, indicating that it is in a critical stage of evaluating the treatment's efficacy and safety for broader patient use. Eligibility criteria include the presence of RAS biomarkers, which may help identify suitable candidates for this study.

NCT: NCT03125980 · Enrollment: 1370 · Sites: Shanghai, China

SIGNAL PHASE3 Stage IV RESEARCH • ACTIVE_NOT_RECRUITING 8

Consolidation Chemotherapy for Locally Advanced Mid or Low Rectal Cancer After Neoadjuvant Concurrent Chemoradiotherapy

This clinical trial is testing the efficacy and feasibility of consolidation chemotherapy for patients with locally advanced mid or low rectal cancer following neoadjuvant concurrent chemoradiotherapy. The intervention involves administering chemotherapy to evaluate its impact on achieving a pathologic complete response. Currently, the trial is in Phase 3 and is active but not recruiting, indicating that it is ongoing but not accepting new participants at this time. Eligibility criteria may include specific characteristics related to the cancer stage and treatment history.

NCT: NCT02843191 · Enrollment: 358 · Sites: Busan, South Korea, Cheonan, South Korea, Chuncheon, South Korea, Daegu, South Korea, Daejeon, South Korea

SIGNAL PHASE3 Stage IV KRAS RESEARCH • RECRUITING 8

Systemic Oxaliplatin or Intra-arterial Chemotherapy Combined With LV5FU2 +/- Irinotecan and an Target Therapy in First Line Treatment of Metastatic Colorectal Cancer Restricted to the Liver

This clinical trial is testing the effectiveness of systemic oxaliplatin versus intra-arterial chemotherapy combined with LV5FU2, with or without irinotecan and targeted therapy, in patients with metastatic colorectal cancer restricted to the liver. The intervention involves administering oxaliplatin both intravenously and intra-arterially, along with other chemotherapy agents and panitumumab, to potentially enhance treatment efficacy for patients with unresectable liver metastases. This is a Phase 3 trial currently recruiting 348 participants, which means patients may have the opportunity to access a treatment regimen that could improve progression-free survival. Eligibility criteria include specific biomarker assessments for KRAS, NRAS, and RAS mutations.

NCT: NCT02885753 · Enrollment: 348 · Sites: Angers, France, Antony, France, Avignon, France, Bayonne, France, Beauvais, France

SIGNAL PHASE3 All Stages MSI/MMR RESEARCH • NOT_YET_RECRUITING 8

Finding the Best Dose of Aspirin to Prevent Lynch Syndrome Cancers

NCT: NCT02497820 · Enrollment: 1800 · Sites: Tel Aviv, Israel

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 8

Study of Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-3475-158/KEYNOTE-158)

NCT: NCT02628067 · Enrollment: 1609 ·

SIGNAL PHASE3 Stage III RESEARCH • ACTIVE_NOT_RECRUITING 8

Safety of a Boost (CXB or EBRT) in Combination With Neoadjuvant Chemoradiotherapy for Early Rectal Adenocarcinoma

NCT: NCT02505750 · Enrollment: 148 · Sites: Besançon, France, Geneva, Switzerland, Guildford, United Kingdom, Hull, United Kingdom, Liverpool, United Kingdom

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 8

A Trial of Aspirin on Recurrence and Survival in Colon Cancer Patients

NCT: NCT02301286 · Enrollment: 770 · Sites: Almelo, Netherlands, Amersfoort, Netherlands, Assen, Netherlands, Breda, Netherlands, Capelle aan den IJssel, Netherlands

SIGNAL PHASE3 Early Detection RAS RESEARCH • RECRUITING 8

The Cancer of the Pancreas Screening-5 CAPS5)Study

The CAPS5 study is a Phase 3 clinical trial aimed at evaluating early cancer markers in pancreatic juice for individuals at risk of pancreatic cancer. The intervention includes the use of secretin, MRI, and a tumor marker gene test with CA19-9, which are important for detecting pancreatic cancer at an earlier stage. Currently, the trial is in the recruiting phase, indicating that eligible participants can still enroll to contribute to this research. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT02000089 · Enrollment: 9000 · Sites: Ann Arbor, Baltimore, Boston, Cleveland, New Haven

SIGNAL PHASE3 Stage IV RESEARCH • ACTIVE_NOT_RECRUITING 8

Neoadjuvant Chemotherapy Versus Standard Treatment in Patients With Locally Advanced Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant chemotherapy compared to standard treatment in patients with locally advanced colon cancer. The intervention involves administering three cycles of capecitabine and oxaliplatin before surgery, which may reduce the need for additional chemotherapy after the operation. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and may provide important insights for future treatment options. Eligibility criteria may include specific biomarkers, although details on these are not provided.

NCT: NCT01918527 · Enrollment: 250 · Sites: Aalborg, Denmark, Bergen, Norway, Copenhagen, Denmark, Gothenburg, Sweden, Herlev, Denmark

SIGNAL PHASE3 Stage III RESEARCH • ACTIVE_NOT_RECRUITING 8

Early Commencement of Adjuvant Chemotherapy for Colon Cancer

NCT: NCT01460589 · Enrollment: 303 · Sites: Daegu, South Korea

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 8

Rectal Cancer And Pre-operative Induction Therapy Followed by Dedicated Operation. The RAPIDO Trial

NCT: NCT01558921 · Enrollment: 920 · Sites: St Louis

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 8

The Northern-European Initiative on Colorectal Cancer

The Northern-European Initiative on Colorectal Cancer (NordICC) is a Phase 3 clinical trial designed to evaluate the effectiveness of colonoscopy screening in reducing colorectal cancer (CRC) incidence and mortality among individuals aged 55-64 years. The intervention involves a colonoscopy procedure, which is significant as it can detect and remove precursor lesions that may lead to CRC, potentially preventing the disease. Currently, the trial is active but not recruiting, indicating that it is ongoing with enrolled participants, and results will be assessed after 15 years of follow-up. Eligibility criteria include being part of the specified age group and drawn randomly from population registries in the participating Nordic countries, the Netherlands, and Poland.

NCT: NCT00883792 · Enrollment: 95000 · Sites: Boston, New York

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 7

FOLFOX8 Versus mFOLFOX6 With Bevacizumab or Cetuximab for First-Line Unresectable Metastatic Colorectal Cancer (Phase II)

NCT: NCT07621497 · Enrollment: 229 · Sites: Guangzhou, China, Shanghai, China

MEDIUM NA Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 7

A Prospective, Single-center, Phase II Study of Neoadjuvant Serplulimab Plus XELOX and Bevacizumab in Locally Advanced Rectal Cancer

NCT: NCT07854964 · Enrollment: 48 ·

SIGNAL PHASE3 Stage IV BRAF RESEARCH • NOT_YET_RECRUITING 7

Pressurized Intraperitoneal Aerosolized Chemotherapy With Mitomycin for the Treatment of Unresectable Appendix or Colorectal Cancer With Peritoneal Metastases, The IMPACT Trial

This phase III clinical trial, known as the IMPACT Trial, is investigating the effectiveness of pressurized intraperitoneal aerosolized chemotherapy (PIPAC) with mitomycin in patients with unresectable appendix or colorectal cancer that has metastasized to the peritoneal cavity. The intervention involves delivering chemotherapy as a pressurized mist directly into the abdominal cavity during a minimally invasive procedure, which may enhance drug absorption and distribution compared to standard chemotherapy regimens. Currently, the trial is not yet recruiting participants, which means patients

will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarker assessments such as BRAF, RAS, mismatch repair status, and next-generation sequencing (NGS).

NCT: NCT07271355 · Enrollment: 129 · Sites: Duarte, Goodyear, Irvine, Newnan, Zion

SIGNAL PHASE3 All Stages MSI/MMR RESEARCH • NOT_YET_RECRUITING 7

Preventive Dendritic Cell Vaccination for Lynch Syndrome Carriers

NCT: NCT07609901 · Enrollment: 372 · Sites: Nijmegen, Netherlands

SIGNAL PHASE2, PHASE3 All Stages RAS RESEARCH • RECRUITING 7

Study to Evaluate the Diagnostic Performance of GEH300079 (68Ga) Injection PET/CT for Detection of PC in Patients With Colorectal, Gastric, Ovarian, or Pancreatic Cancers (PERISCOPE)

This clinical trial is evaluating the diagnostic performance of GEH300079 (68Ga) Injection PET/CT for detecting peritoneal carcinomatosis (PC) in patients with colorectal, gastric, ovarian, or pancreatic cancers. The intervention involves the use of GEH300079 (68Ga) PET/CT imaging, which is significant for improving the accuracy of PC detection compared to standard imaging methods. This is a Phase 2/3 trial currently recruiting participants, which means eligible patients may have the opportunity to contribute to research that could enhance diagnostic practices. Eligibility criteria include having primary colorectal, gastric, ovarian, or pancreatic ductal adenocarcinoma with known or suspected PC, and biomarkers such as RAS and NGS may be relevant for participation.

NCT: NCT07219238 · Enrollment: 175 · Sites: Grand Rapids

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 7

A Study of IOV-5001 in Adults With Advanced Solid Tumors

NCT: NCT07743723 · Enrollment: 106 · Sites: New York

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 7

A Clinical Study of SHR-A1811 in Combination With Chemotherapy and Bevacizumab Versus Standard Therapy as First-Line Treatment for Advanced Colorectal Cancer

NCT: NCT07676162 · Enrollment: 300 · Sites: Beijing, China

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • RECRUITING 7

Short-Course Radiotherapy Followed by CAPOX With or Without Iparomlimab and Tuvonralimab in pMMR/MSS Locally Advanced Rectal Cancer

NCT: NCT07686640 · Enrollment: 180 · Sites: Jinan, China

MEDIUM NA Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 7

To Explore a Single-arm Phase II Study on Intestinal Microbiota Transplantation Combined With Neoadjuvant Immunotherapy and Chemoradiotherapy for Locally Advanced Rectal Cancer Under MSS/pMMR

This clinical trial is testing the effectiveness of intestinal microbiota transplantation combined with neoadjuvant immunotherapy and chemoradiotherapy for patients with locally advanced rectal cancer characterized by microsatellite stable (MSS) and proficient mismatch repair (pMMR). The intervention includes the drug sintilimab, the chemotherapy regimen CAPOX (oxaliplatin and capecitabine), short-course radiotherapy, and intestinal bacteria transplantation, aiming to evaluate its impact on the pathological complete response rate. This is a single-arm Phase II study that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status.

NCT: NCT07738042 · Enrollment: 46 · Sites: Tianjin, China

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • NOT_YET_RECRUITING 7

FOLFIRI Plus Bevacizuamb and QL1706 in Second-Line Treatment for mCRC

NCT: NCT07729605 · Enrollment: 270 · Sites: Guangzhou, China

MEDIUM NA All Stages MSI/MMR LIQUID BIOPSY • RECRUITING 7

Research on Colorectal Cancer Recurrence Monitoring and Individualized Treatment Based on MRD

NCT: NCT07775950 · Enrollment: 100 · Sites: Guangzhou, China

SIGNAL PHASE2, PHASE3 Stage III RAS RESEARCH • RECRUITING 7

Neoadjuvant Chemoradiotherapy Combined With QL1706 for MSS-Type Low-Rectal Cancer

NCT: NCT07866313 · Enrollment: 60 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR IMMUNOTHERAPY • NOT_YET_RECRUITING 7

Neoadjuvant Moderately Hypofractionated Radiotherapy Combined With Chemotherapy and Immunotherapy for High-risk pMMR/MSS Locally Advanced Rectal Cancer: A Prospective, Multi-center Randomized Control Phase II Trial

NCT: NCT07527520 · Enrollment: 165 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 7

Neoadjuvant Treatment of Fulzerasib Plus Cetuximab N01 in KRAS G12C Mutated Locally Advanced Colorectal Cancer With or Without Resectable Metastases

This clinical trial is testing the combination of fulzerasib and cetuximab N01 in patients with locally advanced colorectal cancer that has a KRAS G12C mutation, with or without resectable metastases. The intervention involves administering these targeted therapies to evaluate their effectiveness in improving the overall response rate in this specific patient population. This is a Phase 2 trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include individuals aged 18-80 years with histologically confirmed colorectal adenocarcinoma at specific stages and an ECOG performance status of 0-1.

NCT: NCT07581912 · Enrollment: 40 · Sites: Hangzhou, China

SIGNAL PHASE2, PHASE3 Stage IV RAS RESEARCH • NOT_YET_RECRUITING 7

FMT Combined With Standard First-Line Therapy in Initially Unresectable Colorectal Cancer

This clinical trial is evaluating the efficacy and safety of adding fecal microbiota transplantation (FMT) to standard first-line therapy for patients with initially unresectable colorectal cancer (CRC). The intervention involves combining FMT, which aims to restore intestinal microbiome balance, with standard chemotherapy, with or without targeted therapy. This study is in Phase 2 and Phase 3 and is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT07509398 · Enrollment: 220 ·

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 7

Sintilimab Versus Mitomycin in Combination With Capecitabine and IMRT for Limited-stage Anal Squamous Cell Carcinoma

This clinical trial is testing the efficacy and safety of sintilimab compared to mitomycin in combination with capecitabine and intensity-modulated radiotherapy (IMRT) for patients with limited-stage anal squamous cell carcinoma. The intervention involves replacing the traditional chemotherapeutic drug mitomycin with the PD-1 inhibitor sintilimab, which may offer a different approach to treatment. This is a Phase 3 trial that is currently recruiting 350 participants, which

means eligible patients may have the opportunity to receive either the standard treatment or the experimental regimen. Eligibility criteria include being previously untreated and having a diagnosis of limited-stage anal squamous cell carcinoma, with a focus on patients with specific RAS biomarkers.

NCT: NCT07692152 · Enrollment: 350 · Sites: Beijing, China, Shenzhen, China

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 7

Neoadjuvant Immunotherapy and Organ-sparing Treatment in Patients With Stage I-III dMMR Colon Cancer

NCT: NCT07409844 · Enrollment: 152 · Sites: Køge, Denmark

MEDIUM EARLY_PHASE1 Stage III MSI/MMR LIQUID BIOPSY • RECRUITING 7

COLONYVAQ™, a Quantum-Classical Guided Personalized Neoantigen Vaccine for MSS Stage III Colorectal Cancer

This clinical trial is testing the safety and tolerability of COLONYVAQ-CRC, a personalized neoantigen vaccine, in patients with completely resected stage III microsatellite-stable (MSS) colorectal cancer. The intervention involves administering COLONYVAQ-CRC alongside standard chemotherapy (mFOLFOX6 or CAPOX) and nivolumab, which is significant for its potential to enhance treatment outcomes in this patient population. Currently in the early phase I stage and actively recruiting, the trial aims to monitor treatment-emergent adverse events and immune-related adverse events in an initial cohort of 12 patients. Eligibility criteria include patients with MSS/proficient mismatch repair (pMMR) colorectal cancer, and specific biomarkers such as MSI, MMR, and ctDNA will be assessed.

NCT: NCT07328087 · Enrollment: 12 · Sites: Thessaloniki, Greece

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 7

A Clinical Research Study Evaluating EIK1005, a Werner Helicase Inhibitor, as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced Solid Tumors Including Microsatellite Instability High (MSI-H) Tumors

The EIK1005-002 clinical trial is evaluating the efficacy and safety of EIK1005, a Werner Helicase inhibitor, both as a monotherapy and in combination with pembrolizumab in participants with advanced solid tumors, including those with microsatellite instability high (MSI-H) tumors. The intervention involves administering EIK1005 to determine the optimal safe dosage and assess its tolerability alone and alongside pembrolizumab, which is significant for understanding potential treatment options for advanced cancer. This study is currently in Phase 1 and Phase 2 and is actively recruiting participants, indicating that eligible patients may have the opportunity to access this investigational treatment. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT07262619 · Enrollment: 160 · Sites: Houston, Morristown, New York

MEDIUM PHASE1 Stage IV BRAF TARGETED THERAPY • RECRUITING 7

A Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of CBI-1214 T Cell Engager in Participants With Advanced or Metastatic MSS/MSI-L Colorectal Cancer

This clinical trial is investigating the safety, tolerability, pharmacokinetics, and anti-tumor activity of CBI-1214 T Cell Engager in participants with advanced or metastatic Microsatellite Stable (MSS) or Microsatellite Instability Low (MSI-L) colorectal cancer. The intervention, CBI-1214, is a biological agent designed to engage T cells to target cancer cells, which is important for potentially improving treatment outcomes in this patient population. Currently in Phase 1 and actively recruiting, this trial aims to assess the safety and tolerability of CBI-1214 at various dose levels, which is crucial for determining the appropriate dosage for future studies. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and HER2.

NCT: NCT07321106 · Enrollment: 80 · Sites: Atlanta, Duarte, Fairfax, Grand Rapids, Jacksonville

SIGNAL PHASE3 Early Detection RESEARCH • RECRUITING 7

Improving Functional Outcomes and Quality of Life in Patients With Rectal Cancer After Surgery With Intensified Follow-up & Surveillance

NCT: NCT06936774 · Enrollment: 140 · Sites: Basel, Switzerland, Bern, Switzerland, Lucerne, Switzerland, Winterthur, Switzerland

MEDIUM PHASE1 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 7

Phase I Study of [177Lu]Lu-DFC413 in Patients With Solid Tumors

NCT: NCT07261631 · Enrollment: 180 · Sites: Princeton

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • RECRUITING 7

Megestrol Acetate in Improving Neoadjuvant Chemotherapy-Related Weight Loss in Locally Advanced CRC

This clinical trial is testing the effectiveness of Megestrol Acetate in improving weight loss associated with neoadjuvant chemotherapy in patients with locally advanced colorectal cancer (LACRC). The intervention, Megestrol Acetate, is a drug that aims to enhance appetite and promote weight gain, addressing the nutritional challenges faced by patients undergoing chemotherapy. This is a Phase 3 trial currently recruiting 60 participants, which indicates that it is in the later stages of testing and may provide valuable information on the drug's efficacy and safety for patients. Eligibility criteria include specific biomarkers such as MSI and MMR status.

NCT: NCT06998758 · Enrollment: 60 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage IV RAS RESEARCH • NOT_YET_RECRUITING 7

Neoadjuvant CAPOX Plus Tislelizumab vs CAPOX in MSS High-Risk Locally Advanced Colon Cancer

NCT: NCT07132008 · Enrollment: 94 · Sites: Nanning, China

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 7

Interest of Early Erectile Rehabilitation With Sildenafil After Radiotherapy and Proctectomy for Rectal Cancer

This clinical trial is testing the efficacy of early erectile rehabilitation with Sildenafil in men who have undergone neoadjuvant proctectomy and radiotherapy for rectal cancer. The intervention involves administering Sildenafil compared to a placebo, which is important for assessing its potential to prevent long-term erectile dysfunction following treatment. This is a Phase 3 trial that is currently recruiting participants, indicating that it is in the later stages of testing and may provide valuable information for patient care. Eligibility criteria include being a male patient who has received the specified treatments for rectal cancer.

NCT: NCT06123156 · Enrollment: 188 · Sites: Amiens, France, Beauvais, France, Besançon, France, Bicêtre, France, Bordeaux, France

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 7

Modulated Electro-Hyperthermia in Combination With Multimodal Therapy for Locally Advanced Rectal Cancer

This clinical trial is testing the effectiveness of modulated electro-hyperthermia (mEHT) in combination with Total Neoadjuvant Therapy (TNT) for adult patients with locally advanced rectal adenocarcinoma. The intervention involves the use of the Oncotherm EHY-2030 device to apply hyperthermia, which may enhance tumor down-staging and improve pathological responses. This is a Phase 3 trial currently recruiting 126 participants, which means eligible patients may have the opportunity to receive a potentially beneficial treatment while contributing to important research outcomes. Eligibility criteria include adults aged 20 years and above with specific stages of rectal cancer, including cT3N0M0 with high risk of recurrence, cT3N1-2M0, or cT4N0-2M0.

NCT: NCT07372300 · Enrollment: 126 · Sites: Chiayi City, Taiwan

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 7

Neoadjuvant Pembrolizumab With a Watch-and-wait Strategy for dMMR/MSI-H Localized Colon Cancer: PREMICES Study.

The PREMICES study is investigating the effectiveness of neoadjuvant pembrolizumab combined with a watch-and-wait strategy for patients with localized deficient mismatch repair/high microsatellite instability (dMMR/MSI-H) colorectal cancer. The intervention involves administering pembrolizumab, a drug that targets the immune system, alongside a non-operative management approach, which may reduce the need for immediate surgery. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligible participants will be randomized to either the experimental arm receiving pembrolizumab with the watch-and-wait strategy or the standard treatment involving surgical resection and possible adjuvant chemotherapy.

NCT: NCT06646445 · Enrollment: 60 · Sites: Avignon, France, Besançon, France, Levallois-Perret, France, Lyon, France, Paris, France

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 7

Endoscopic Therapy Or Surgery for Early Colon Cancer

This clinical trial is testing the effectiveness of endoscopic full-thickness resection (eFTR) compared to standard surgical procedures for patients with early colon cancer. The intervention, eFTR, is a minimally invasive endoscopic treatment that may offer different benefits and risks compared to traditional surgery. Currently in Phase 3 and actively recruiting participants, this trial aims to assess the rate of severe adverse events, re-admissions, and deaths within 30 days post-procedure. Specific eligibility criteria or biomarker requirements have not been detailed in the provided information.

NCT: NCT06940947 · Enrollment: 434 · Sites: Gdansk, Poland

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 7

Evaluation of NuvastaticTM in Reducing Cancer-Related Fatigue in Colon Cancer Patients Undergoing First-Line Chemotherapy

NCT: NCT07669519 · Enrollment: 180 · Sites: Belagavi, India, Lucknow, India, Mysuru, India, Satara, India, Srikakulam, India

MEDIUM PHASE2 Stage IV BRAF LIQUID BIOPSY • RECRUITING 7

Phase II Study of Anti-PD-1/VEGF Bispecific Antibody Ivonescimab in Patients With Previously Treated Metastatic Colorectal Cancer

NCT: NCT06959550 · Enrollment: 90 · Sites: Houston

HIGH PHASE3 All Stages PHASE III TRIAL • RECRUITING 7

Phase 3 Trial of eRapa in Patients With Familial Adenomatous Polyposis

This Phase 3 clinical trial is testing the efficacy of eRapa (encapsulated rapamycin) in patients diagnosed with Familial Adenomatous Polyposis (FAP). The intervention involves administering eRapa compared to a placebo to determine its impact on slowing disease progression and its safety profile. Currently, the trial is recruiting 168 participants, which means eligible patients can still join to contribute to the research. Eligibility criteria include a diagnosis of FAP, and participants will undergo regular check-ups and endoscopies to monitor their condition throughout the study.

NCT: NCT06950385 · Enrollment: 168 · Sites: Ann Arbor, Arcadia, Baltimore, Boston, Chicago

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 7

Study of Orally Administered MOMA-341 in Participants With Advanced or Metastatic Solid Tumors

NCT: NCT06974110 · Enrollment: 132 · Sites: Dallas, Detroit, Houston, Phoenix, Portland

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • NOT_YET_RECRUITING 7

Dual EGFR/HER2 Blockade Combined With Irinotecan for the Treatment of HER2-Positive Metastatic Colorectal Cancer

This clinical trial is testing the combination of trastuzumab, cetuximab beta, and irinotecan for patients with HER2-positive metastatic colorectal cancer (mCRC) who have not responded to previous chemotherapy. The intervention aims to target both HER2 and EGFR pathways, which may help overcome resistance to HER2-targeted therapies and improve

treatment outcomes. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, DMMR, and HER2 status.

NCT: NCT07059338 · Enrollment: 34 · Sites: Guangzhou, China

MEDIUM NA Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 7

Reducing the Resection Range After Neoadjuvant Therapy for Locally Advanced Upper Rectal and Rectosigmoid Junction Cancers

NCT: NCT07038122 · Enrollment: 874 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 7

Efficacy and Safety of Trastuzumab Biosimilars and Pertuzumab Biosimilars Combined With Chemotherapy for Neoadjuvant Treatment of Patients With Locally Advanced HER2-positive Rectal Cancer

NCT: NCT07108127 · Enrollment: 24 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 7

AK129 Combination Therapy for Advanced Solid Tumors

This clinical trial is testing the combination therapy of AK129 with Pemetrexed, Paclitaxel, and Carboplatin for patients with advanced solid tumors, including colorectal adenocarcinoma. The intervention involves administering different doses of AK129 alongside established chemotherapy agents to assess safety and efficacy. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to research. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, NTRK, and microsatellite status.

NCT: NCT06943820 · Enrollment: 230 · Sites: Shenyang, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 7

TL938 and Trastuzumab for Patients With HER2-positive Metastatic Colorectal Cancer

This Phase II clinical trial is testing the combination of TL938 capsules and trastuzumab for patients with HER2-positive metastatic colorectal cancer that is inoperable or has recurred. The intervention aims to determine the optimal dose while evaluating the effectiveness of these drugs in this specific patient population. Currently, the trial is in the recruiting status, meaning eligible patients can still enroll to participate. Eligibility criteria include specific biomarkers such as HER2, MSI, and MMR status.

NCT: NCT06589830 · Enrollment: 80 · Sites: Beijing, China

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 7

Bovine Reinforcement in Stomach, Colorectal and Lung Operation

NCT: NCT06968221 · Enrollment: 172 · Sites: Shanghai, China

SIGNAL PHASE2, PHASE3 All Stages MSI/MMR RESEARCH • RECRUITING 7

Node-Sparing Short-Course Radiotherapy Sequential Chemotherapy and PD-1 Inhibitor for Mid/Low pMMR/MSS Rectal Cancer (MODIFI-RC-II)

The MODIFI-RC-II trial is testing the effectiveness of node-sparing short-course radiotherapy followed by sequential chemotherapy and a PD-1 inhibitor in patients with mid-low pMMR/MSS rectal cancer. This intervention is significant as it aims to enhance tumor response to PD-1 blockade while preserving critical lymphatic functions that may be compromised by conventional radiotherapy. Currently in Phase 2 and Phase 3, the trial is recruiting 430 participants, which means eligible patients may have the opportunity to contribute to research that could influence future treatment approaches. Eligibility criteria include specific biomarkers such as MMR, RAS, and microsatellite status.

NCT: NCT06958432 · Enrollment: 430 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • NOT_YET_RECRUITING 7

An Open-label, Single-arm, Multicenter Phase II Study to Evaluate the Efficacy of Amivantamab in Combination With FOLFIRI as a Second-line Treatment in Patients With RAS/BRAF Wild-type Advanced Colorectal Cancer Progressing on Prior Anti-EGFR Based Treatment.

This clinical trial is evaluating the efficacy of amivantamab in combination with FOLFIRI as a second-line treatment for patients with RAS/BRAF wild-type advanced colorectal cancer who have progressed on prior anti-EGFR-based treatment. The intervention involves administering amivantamab alongside the FOLFIRI regimen, which includes folic acid, 5-FU, and irinotecan, to assess its impact on overall response rates. This is a Phase II study that is currently not yet recruiting participants, meaning that patients will not be able to enroll until the trial opens. Eligibility criteria include specific biomarker requirements such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, DMMR, and HER2 status.

NCT: NCT06855849 · Enrollment: 85 · Sites: Seoul, South Korea

MEDIUM N/A All Stages KRAS LIQUID BIOPSY • RECRUITING 7

Precision-MRD: Prospective Observational Study of Biomarker-directed Systemic Therapy for Colorectal Cancer Patients With Minimal Residual Disease

The Precision-MRD study is investigating the effects of biomarker-directed systemic therapy in colorectal cancer patients with minimal residual disease (MRD). The intervention involves monitoring participants receiving therapy based on specific biomarkers, such as KRAS, BRAF, and ctDNA, which is important for understanding treatment responses. This study is currently in the recruiting phase, meaning eligible patients can enroll to contribute to the research while receiving standard care. Eligibility criteria include the presence of specific biomarkers related to colorectal cancer.

NCT: NCT06878131 · Enrollment: 40 · Sites: Houston

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 7

The Effect of Discontinuation of Renin Angiotensin System Inhibitors on the Perioperative Myocardial Injury in Adult Patients Undergoing Colorectal Cancer Surgery

This clinical trial is testing the effect of discontinuing Renin Angiotensin System (RAS) inhibitors in adult patients undergoing colorectal cancer surgery. The intervention involves the use of ACE inhibitors and/or angiotensin receptor blockers (ARBs) to assess their impact on perioperative myocardial injury, which is significant for managing cardiovascular risks during surgery. This is a Phase 3 trial currently recruiting 156 participants, indicating that patients may have the opportunity to contribute to important findings that could influence clinical practices. Eligibility criteria include the presence of specific biomarkers related to the RAS.

NCT: NCT06862115 · Enrollment: 156 · Sites: Cairo, Egypt

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 7

DEBIRI Plus Chemotherapy vs. Chemotherapy Alone in Colorectal Cancer Liver Metastases

This clinical trial is testing the effectiveness of DEBIRI combined with chemotherapy compared to chemotherapy alone in patients with colorectal cancer liver metastases. The intervention involves the use of Drug-Eluting Embolic Beads (DEBIRI) alongside systemic chemotherapy, which aims to enhance tumor shrinkage and potentially convert unresectable tumors to resectable ones. This is a Phase 3 trial currently recruiting 116 participants, which means eligible patients may have the opportunity to receive either treatment while contributing to important research outcomes. Eligibility criteria include being chemotherapy-naïve for metastatic disease and having specific RAS biomarker status.

NCT: NCT06555003 · Enrollment: 116 · Sites: Tehran, Iran

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • RECRUITING 7

mFOLFOX6 + Bevacizumab + PD-1 Monoclonal Antibody Vs. mFOLFOX6 in Locally Advanced pMMR/MSS CRC

This clinical trial is testing the combination of mFOLFOX6, Bevacizumab, and a PD-1 monoclonal antibody in patients with locally advanced mismatch repair-proficient or microsatellite stable colorectal cancer (pMMR/MSS CRC). The intervention aims to enhance the efficacy of immunotherapy in this patient population, which has shown limited response to PD-1 monoclonal antibodies alone. Currently in Phase 3 and actively recruiting, this trial seeks to determine the three-year

disease-free survival (DFS) rate, providing critical insights for future treatment strategies. Eligibility criteria include specific biomarker assessments such as MMR, RAS, and microsatellite status, which are essential for patient selection.

NCT: NCT06791512 · Enrollment: 166 · Sites: Guangzhou, China

SIGNAL PHASE3 All Stages KRAS RESEARCH • RECRUITING 7

A Study of Tucidinostat in Combination With Sintilimab and Bevacizumab in MSS/pMMR Colorectal Cancer Patients

This clinical trial is testing the combination of tucidinostat, sintilimab, and bevacizumab in patients with MSS/pMMR colorectal cancer. The intervention aims to evaluate the efficacy and safety of this combination compared to fruquintinib monotherapy, which is significant for understanding potential treatment options for this patient population. Currently in Phase 3 and actively recruiting, this trial may provide insights into overall survival outcomes for participants. Eligibility criteria include specific biomarkers such as KRAS, MSI, MMR, and RAS.

NCT: NCT06497985 · Enrollment: 430 · Sites: Guangzhou, China

SIGNAL PHASE3 Stage IV BRAF RESEARCH • RECRUITING 7

Induction Treatment for Initially Unresectable Colorectal Liver Metastases: Combined Hepatic Arterial Infusion Pump Therapy With Systemic Therapy

NCT: NCT06857773 · Enrollment: 306 · Sites: Amsterdam, Netherlands

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 7

HDAC Inhibitor Combination With Chemoimmunotherapy in the Neoadjuvant Treatment of pMMR Locally Advanced Colon Cancer

NCT: NCT06709885 · Enrollment: 100 · Sites: Chongqing, China

MEDIUM PHASE1 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 7

Phase I Study of [177Lu]Lu-NNS309 in Patients With Pancreatic, Lung, Breast and Colorectal Cancers

NCT: NCT06562192 · Enrollment: 162 · Sites: Birmingham, Boston, Grand Rapids, Houston, Jacksonville

SIGNAL PHASE2, PHASE3 All Stages RESEARCH • RECRUITING 7

Colon Cancer Diagnosis With FAPI-PET Imaging

This clinical trial is testing the diagnostic efficacy of 18F-FAPI-74 PET/CT imaging in patients with primary colon cancer. The intervention involves the use of a novel PET tracer, 18F-FAPI-74, which targets fibroblast activation protein overexpressed by cancer-associated fibroblasts, potentially improving the detection of local lymph node and distant metastases. Currently in Phase 2 and Phase 3, the trial is recruiting 100 patients, which means eligible participants may contribute to understanding the effectiveness of this imaging tool in clinical practice. Specific eligibility criteria have not been detailed, but the study focuses on patients diagnosed with colon cancer.

NCT: NCT07272772 · Enrollment: 100 · Sites: Turku, Finland

SIGNAL PHASE3 All Stages RESEARCH • NOT_YET_RECRUITING 7

Preoperative Chemoradiotherapy and Transanal Endoscopic Microsurgery Versus Ttransanal Endoscopic Microsurgery in T1 N0, M0 Rectal Cancer (TAUTEM-T1 Study)

the T1 N0, M0 rectal cancer stage. This trial is testing the combination of preoperative chemoradiotherapy using Capecitabine and radiation therapy, followed by Transanal Endoscopic Microsurgery (TEM), against TEM alone. As a Phase 3 trial that is not yet recruiting, it aims to evaluate rectal preservation rates in patients with early-stage rectal cancer, which may influence treatment decisions and outcomes. Eligibility criteria include patients diagnosed with T1, N0, M0 rectal adenocarcinoma without poor prognostic factors.

NCT: NCT06450574 · Enrollment: 106 ·

SIGNAL PHASE3 All Stages MSI/MMR RESEARCH • ACTIVE_NOT_RECRUITING 7

Node-sparing Short-Course Radiation Combined With CAPOX and Tislelizumab for MSS Rectal Cancer

NCT: NCT06507371 · Enrollment: 170 · Sites: Hanzhou, China

MEDIUM NA Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 7

Targeted Treatment Plus Tislelizumab and HAIC for Advanced CRCLM Failed From Standard Systemic Treatment

This clinical trial is testing the efficacy and safety of a combination of hepatic arterial infusion chemotherapy (HAIC), targeted therapy, and the PD-1 inhibitor tislelizumab for patients with advanced colorectal cancer liver metastasis (CRCLM) who have not responded to standard systemic treatment. The intervention aims to provide an individualized treatment regimen based on genotyping or ctDNA genotyping, which is significant for optimizing therapy in microsatellite stable (MSS) CRCLM cases. The trial is currently active but not recruiting new participants, indicating that it is in the process of evaluating its outcomes with the enrolled patients. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, HER2, and RAS.

NCT: NCT06199232 · Enrollment: 47 · Sites: Beijing, China

SIGNAL PHASE3 All Stages NGS RESEARCH • RECRUITING 7

Localized Treatment Versus Palliative Chemotherapy in CRC Patients With 10 or More CRLM

This clinical trial is testing the effectiveness of localized interventions versus palliative chemotherapy in colorectal cancer patients with 10 or more liver metastases. The intervention includes surgical resection, ablation therapy, and/or stereotactic body radiotherapy (SBRT), which may provide a different approach to treatment compared to standard palliative chemotherapy. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to contribute to significant findings regarding treatment options. Participants must have achieved disease control through chemotherapy and meet specific eligibility criteria related to their condition.

NCT: NCT06208371 · Enrollment: 117 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 7

Efficacy and Safety of Combined With Immunotherapy After Induction Therapy With Chemotherapy and Targeted Therapy in the First-line Treatment of Microsatellite Stable (MSS) Initially Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of combining camrelizumab with chemotherapy and bevacizumab as a first-line treatment for patients with microsatellite stable (MSS) initially unresectable metastatic colorectal cancer. The intervention involves the use of camrelizumab, an immunotherapy drug, which is significant as it may enhance treatment outcomes in this patient population. Currently in Phase 2 and actively recruiting, this trial aims to provide insights into disease-free survival for participants. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT06176885 · Enrollment: 36 · Sites: Jinhua, China

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 7

CompariSon Between the EuroPeAn and Japanese pathologiCal InvEstigation for Colon Cancer (SPACE)

The SPACE trial is testing the diagnostic accuracy of Japanese versus European pathological investigation methods for stage III colon cancer in patients. The intervention involves comparing two procedures: the Japanese pathological investigation, which includes surgeon involvement in lymph node extraction and specimen assessment, and the European method, which primarily relies on pathologists. This is a Phase 3 trial currently recruiting 430 participants, which means patients may have the opportunity to contribute to important findings regarding diagnostic practices. Eligibility criteria have not been specified in the provided information.

NCT: NCT06119867 · Enrollment: 430 · Sites: Moscow, Russia

MEDIUM N/A Stage IV MSI/MMR LIQUID BIOPSY • RECRUITING 7

Circulating Tumor DNA (ctDNA) as a Predictive Biomarker for Immunotherapy in Advanced or Locally Advanced dMMR/MSI-H Colorectal Patients

This clinical trial is investigating the use of circulating tumor DNA (ctDNA) as a predictive biomarker for immunotherapy in patients with advanced or locally advanced dMMR/MSI-H colorectal cancer. The intervention focuses on assessing the correlation between changes in ctDNA and the rates of pathological complete response (pCR) and the duration of clinical complete response (cCR), which is important for evaluating treatment efficacy. The trial is currently in the recruiting phase and aims to enroll 60 participants, indicating that eligible patients may still join the study. Eligibility criteria include having dMMR/MSI-H colorectal cancer, with specific biomarkers such as MSI, MSI-H, MMR, and DMMR being relevant for participation.

NCT: NCT06098560 · Enrollment: 60 · Sites: Beijing, China, Shanghai, China

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 7

Neoadjuvant Chemotherapy With PD-1 Inhibitors Combined With SIB-IMRT in the Treatment of Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy and safety of neoadjuvant chemotherapy with PD-1 inhibitors, specifically tislelizumab, combined with simultaneous integrated boost intensity-modulated radiotherapy (SIB-IMRT) in patients with locally advanced rectal cancer. The intervention includes the drugs capecitabine and oxaliplatin alongside SIB-IMRT, aiming to improve treatment outcomes for this patient population. Currently in Phase 3 and actively recruiting, this trial may provide patients with access to a potentially effective treatment approach while contributing to the understanding of neoadjuvant therapies in rectal cancer. Eligibility criteria may include specific biomarkers, although details on these are not provided.

NCT: NCT06017583 · Enrollment: 48 · Sites: Nanning, China

SIGNAL PHASE3 Stage IV RAS RESEARCH • RECRUITING 7

Sulfasalazine in Patients With Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of sulfasalazine in patients with metastatic colorectal cancer. The intervention involves the drug sulfasalazine, which is being evaluated for its potential impact on treatment outcomes. This is a Phase 3 trial currently recruiting 50 participants, which indicates that it is in the later stages of testing and may provide important information about the drug's effectiveness and safety for patients. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT06134388 · Enrollment: 50 · Sites: Tanta, Egypt

SIGNAL PHASE2, PHASE3 All Stages BRAF RESEARCH • ACTIVE_NOT_RECRUITING 7

Testing the Use of BRAF-Targeted Therapy After Surgery and Usual Chemotherapy for BRAF-Mutated Colon Cancer

NCT: NCT05710406 · Enrollment: 1 · Sites: Allentown, Anaconda, Ankeny, Ann Arbor, Antigo

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 7

Regorafenib Combined With Irinotecan Drug-Eluting Beads for Colorectal Cancer Liver Metastases

This clinical trial is testing the efficacy and safety of regorafenib combined with Irinotecan Drug-Eluting Beads as a third-line treatment for patients with colorectal cancer liver metastases. The intervention involves the use of regorafenib and DIBIRI (drug-eluting beads), which is significant as it explores a combination therapy for patients who have not responded to first- and second-line chemotherapy. This is a Phase 3 trial currently recruiting 126 participants, which means eligible patients may have the opportunity to access this treatment option while contributing to important research. Eligibility criteria include patients who have failed prior standard chemotherapy and may also involve RAS biomarker considerations.

NCT: NCT05794971 · Enrollment: 126 · Sites: Guangdong, China

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 7

Prophylactic Mesh Placement During Stoma Closure After Low Anterior Resection

This clinical trial is testing the efficacy of polypropylene mesh for hernia prevention during stoma closure in patients with colorectal cancer. The intervention involves comparing outcomes between patients receiving mesh and those undergoing non-mesh repair, which is significant for understanding potential benefits in hernia prevention. Currently in Phase 3 and actively recruiting, this trial aims to enroll 142 participants who will be followed for two years to assess hernia rates. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT05939687 · Enrollment: 142 · Sites: Moscow, Russia

MEDIUM PHASE3 All Stages ctDNA LIQUID BIOPSY • RECRUITING 7

Evaluating Novel Therapies in ctDNA Positive GI Cancers

This clinical trial is evaluating the effectiveness of the drugs atezolizumab and bevacizumab in patients with ctDNA positive gastrointestinal (GI) cancers, specifically colorectal cancer, gastric cancer, pancreatic ductal adenocarcinoma (PDAC), and gallbladder adenocarcinomas. The intervention involves administering intravenous atezolizumab and bevacizumab to patients who have completed standard of care treatments but still test positive for ctDNA, which may indicate residual disease. This is a Phase 3 trial currently recruiting participants, meaning eligible patients may have the opportunity to access these therapies while contributing to the research. Eligibility criteria include having a histologically confirmed GI cancer, a positive SignateraTM ctDNA test, and being radiographically NED within a specified timeframe after completing curative-intent therapy.

NCT: NCT05482516 · Enrollment: 20 · Sites: Hackensack, Washington D.C.

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • RECRUITING 7

Comparing Chidamide+Sintilimab+Bev With Standard Second-line FOLFIRI+Bev in Advanced MSS/pMMR mCRC

This clinical trial is testing the combination of chidamide, sintilimab, and bevacizumab against the standard second-line treatment of FOLFIRI plus bevacizumab in patients with advanced microsatellite stable (MSS) and proficient mismatch repair (pMMR) metastatic colorectal cancer (mCRC) who have not responded to first-line oxaliplatin-based therapy. The intervention involves an experimental drug regimen that may offer a different therapeutic approach compared to the standard treatment, which is important for improving patient outcomes. This is a phase 3 trial currently recruiting participants, indicating that it is in the later stages of testing efficacy and safety before potential approval for broader use. Eligibility criteria include specific biomarkers such as MSI, MMR, RAS, and microsatellite status.

NCT: NCT05768503 · Enrollment: 176 · Sites: Chengdu, China, Foshan, China, Guangzhou, China, Hangzhou, China, Harbin, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 7

A Study of Evorpacept (ALX148) With Cetuximab and Pembrolizumab for Refractory Microsatellite Stable Metastatic Colorectal Cancer

This Phase 2 clinical trial is evaluating the combination of evorpacept (ALX148), cetuximab, and pembrolizumab for patients with refractory microsatellite stable metastatic colorectal cancer. The intervention involves administering these three drugs together to determine the recommended dose of evorpacept in this specific patient population. Currently, the trial is active but not recruiting new participants, which means that it is ongoing with enrolled patients receiving treatment. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status.

NCT: NCT05167409 · Enrollment: 48 · Sites: Aurora, Fairfax, New Brunswick, Tucson

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 7

Evaluation of Co-formulated Pembrolizumab/Quavonlimab (MK-1308A) Versus Other Treatments in Participants With Microsatellite Instability-High (MSI-H) or Mismatch Repair Deficient (dMMR) Stage IV Colorectal Cancer (CRC) (MK-1308A-008/KEYSTEP-008).

NCT: NCT04895722 · Enrollment: 302 · Sites: Athens, Chicago, Dallas, Germantown, Nashville

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 7

Study of Cabozantinib and Nivolumab in Refractory Metastatic Microsatellite Stable (MSS) Colorectal Cancer

NCT: NCT04963283 · Enrollment: 48 · Sites: Aurora, Colorado Springs, Fort Collins, Greeley, Loveland

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 7

Efficacy of Hyperthermic Intraperitoneal Chemotherapy

NCT: NCT04861558 · Enrollment: 213 · Sites: Ahmedabad, India, Gothenburg, Sweden, Malmö, Sweden, Stockholm, Sweden, Uppsala, Sweden

SIGNAL PHASE3 All Stages NGS RESEARCH • RECRUITING 7

Induction Chemotherapy for Locally Recurrent Rectal Cancer

This clinical trial is testing the effectiveness of induction chemotherapy followed by neoadjuvant chemoradiotherapy and surgery in patients with locally recurrent rectal cancer. The intervention involves a combination of drugs, radiation, and surgical procedures, which may improve surgical outcomes by increasing the likelihood of achieving clear resection margins. Currently in Phase III and actively recruiting participants, this trial aims to enroll 364 patients, which means eligible individuals may have the opportunity to contribute to important findings in treatment approaches. Eligibility criteria include the requirement for next-generation sequencing (NGS) biomarkers.

NCT: NCT04389086 · Enrollment: 364 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Ghent, Belgium, Gothenburg, Sweden, Groningen, Netherlands

MEDIUM NA All Stages TARGETED THERAPY • ACTIVE_NOT_RECRUITING 7

Targeted Agent Evaluation in Digestive Cancers in China Based on Molecular Characteristics

This clinical trial is evaluating the effectiveness of various targeted agents in patients with previously treated and refractory gastrointestinal tumors, including colorectal cancer. The intervention involves multiple inhibitors, such as FGFR, IDH1, HER2, PARP, BRAF, MEK, and others, aimed at providing individualized treatment based on tumor DNA sequencing. The trial is currently active but not recruiting participants, indicating that while the study is ongoing, no new patients are being enrolled at this time. Eligibility criteria may include specific molecular characteristics identified through tumor DNA sequencing, which will guide the selection of targeted therapies for each patient.

NCT: NCT04584008 · Enrollment: 600 · Sites: Beijing, China

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 7

Self-Management Survivorship Care in Stage I-III Non-small Cell Lung Cancer or Colorectal Cancer

This phase III clinical trial is testing the effectiveness of a telehealth self-management program for survivors of stage I-III non-small cell lung cancer or colorectal cancer. The intervention includes quality-of-life assessments, questionnaire administration, survivorship care plans, and telemedicine to enhance care coordination and patient outcomes. Currently, the trial is active but not recruiting new participants, which means that patients cannot enroll at this time. Eligibility criteria may include being a survivor of stage I-III non-small cell lung cancer or colorectal cancer.

NCT: NCT04428905 · Enrollment: 403 · Sites: Duarte

SIGNAL PHASE2, PHASE3 Stage II RAS RESEARCH • ACTIVE_NOT_RECRUITING 7

Duloxetine to Prevent Oxaliplatin-Induced Peripheral Neuropathy in Patients With Stage II-III Colorectal Cancer

NCT: NCT04137107 · Enrollment: 220 · Sites: Aberdeen, Akron, Albany, Albuquerque, Allentown

SIGNAL PHASE3 All Stages NGS RESEARCH • RECRUITING 7

Standard Dose Versus High Dose of Radiotherapy in Rectal Preservation With Chemo-radiotherapy in Rectal Cancer Patients

This clinical trial is testing the effectiveness of standard versus high doses of radiotherapy in rectal cancer patients who are undergoing chemoradiotherapy with the goal of rectal preservation. The intervention involves administering either 50.4 Gy or 62 Gy of radiation, along with Capecitabine, to evaluate which dosage leads to better outcomes in terms of organ preservation. This is a Phase 3 trial currently recruiting 162 participants, which means that eligible patients may have the opportunity to contribute to important findings that could influence treatment protocols. Eligibility criteria include specific biomarkers identified through next-generation sequencing (NGS).

NCT: NCT04095299 · Enrollment: 162 · Sites: Aalborg, Denmark, Copenhagen, Denmark, Næstved, Denmark, Roskilde, Denmark, Vejle, Denmark

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 7

Surgery Plus Chemo Versus Chemoradiotherapy Followed by Surgery Plus Chemo for Locally Recurrent Rectal Cancer

NCT: NCT04288999 · Enrollment: 110 · Sites: Chiba, Japan, Gifu, Japan, Hidaka, Japan, Hirakata, Japan, Hiroshima, Japan

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 7

Comparison of Transanal Irrigation and Glycerol Suppositories in Treatment of Low Anterior Resection Syndrome

This clinical trial is testing the effectiveness of transanal irrigation versus glycerol suppositories in patients experiencing major Low Anterior Resection Syndrome (LARS). The intervention involves the Qufora Irrisedo Cone System as a device and glycerol "OBA" as a drug, which are being evaluated for their impact on bowel function. Currently in Phase 3 and actively recruiting participants, this trial aims to provide insights that could inform treatment options for patients with LARS. Eligibility criteria may include specific symptoms related to bowel function, although detailed biomarker requirements are not specified.

NCT: NCT04087421 · Enrollment: 114 · Sites: Aarhus, Denmark

SIGNAL PHASE3 All Stages RAS RESEARCH • RECRUITING 7

Performance of SGM-101 for the Delineation of Primary and Recurrent Tumor and Metastases in Patients Undergoing Surgery for Colorectal Cancer

This clinical trial is testing the performance of SGM-101, an intraoperative imaging agent, in patients undergoing surgery for colorectal cancer. The intervention involves using SGM-101 to enhance the visualization of primary and recurrent tumors and metastases during surgical procedures, which may improve surgical outcomes. This is a Phase 3 trial currently recruiting 300 participants, indicating that it is in the later stages of testing and may provide valuable information for clinical practice. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT03659448 · Enrollment: 300 · Sites: Boston, Duarte, La Jolla, Philadelphia, Weston

SIGNAL PHASE3 Stage III RESEARCH • RECRUITING 7

Mebendazole as Adjuvant Treatment for Colon Cancer

This clinical trial is testing the effectiveness of mebendazole as an adjuvant treatment for patients with colon cancer. The intervention involves administering mebendazole alongside the standard treatment of Folfox and Avastin, which may enhance tumor response. Currently in Phase 3 and actively recruiting, this trial aims to determine the impact of mebendazole on treatment outcomes for participants. Eligibility criteria may include specific biomarkers, although details are not provided.

NCT: NCT03925662 · Enrollment: 40 · Sites: Cairo, Egypt

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 7

Effect of EPA-FFA on Polypectomy in Familial Adenomatous Polyposis

This clinical trial is testing the effect of eicosapentaenoic acid free fatty acid (EPA-FFA) on polypectomy outcomes in patients with familial adenomatous polyposis (FAP). The intervention involves administering EPA-FFA gastro-resistant capsules compared to a placebo, which is important for assessing its potential impact on the number of polypectomies

required. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the later stages of testing and may provide significant insights into treatment efficacy for patients. Eligibility criteria may include a diagnosis of familial adenomatous polyposis, although specific biomarker requirements are not detailed.

NCT: NCT03806426 · Enrollment: 204 · Sites: Bologna, Italy

MEDIUM PHASE1, PHASE2 Stage IV KRAS LIQUID BIOPSY • RECRUITING 7

A Study of Bispecific Antibody MCLA-158 in Patients With Advanced Solid Tumors

NCT: NCT03526835 · Enrollment: 560 · Sites: Atlanta, Boston, Cleveland, Columbus, Dallas

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 7

EPA for Metastasis Trial 2

The EPA for Metastasis Trial 2 is testing the efficacy of Icosapent Ethyl, a pure omega-3 fatty acid derived from fish oil, in patients with liver metastases from bowel cancer. The trial aims to determine if taking this supplement before and after liver surgery can improve progression-free survival compared to a placebo. Currently in Phase 3 and active but not recruiting, this study involves 418 participants and seeks to provide insights into the potential benefits of Icosapent Ethyl in preventing disease recurrence. Eligibility criteria include having liver metastases from bowel cancer, with a target enrollment of 448 individuals.

NCT: NCT03428477 · Enrollment: 418 · Sites: Aintree, United Kingdom, Basingstoke, United Kingdom, Birmingham, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL PHASE3 Early Detection RESEARCH • ACTIVE_NOT_RECRUITING 7

Second and Third Look Laparoscopy in pT4 Colon Cancer Patients for Early Detection of Peritoneal Metastases

NCT: NCT03413254 · Enrollment: 389 · Sites: Almere Stad, Netherlands, Amsterdam, Netherlands, Eindhoven, Netherlands, Groningen, Netherlands, Nieuwegein, Netherlands

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 7

ASPIRIN Trial Belgium

NCT: NCT03464305 · Enrollment: 148 · Sites: Antwerp, Belgium, Bornem, Belgium, Bouge, Belgium, Brasschaat, Belgium, Bruges, Belgium

SIGNAL PHASE2, PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 7

Aspirin in Colorectal Cancer Liver Metastases

The ASAC trial is investigating the efficacy of low-dose acetylsalicylic acid (ASA) as an adjuvant treatment for patients who have undergone resection for colorectal cancer liver metastases (CRCLM). The intervention involves administering ASA 160 mg once daily compared to a placebo for three years, with the aim of improving disease-free survival (DFS). This is a Phase 2/Phase 3 trial that is currently active but not recruiting, meaning that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include patients who have been operated on for CRCLM, with a total enrollment of 466 participants planned for the study.

NCT: NCT03326791 · Enrollment: 466 · Sites: Aarhus, Denmark, Bergen, Norway, Copenhagen, Denmark, Gothenburg, Sweden, Linköping, Sweden

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 7

iRECIST Evaluation's Relevance for DCR in MMR/MSI Metastatic Colorectal Cancer Patients on Nivolumab and Ipilimumab

This clinical trial is evaluating the disease control rate (DCR) in patients with metastatic colorectal cancer who have specific biomarkers, including MSI, MSI-H, MMR, DMMR, and RAS, while receiving treatment with the drugs nivolumab and ipilimumab. The intervention involves administering ipilimumab and nivolumab to assess their effectiveness in controlling the disease over a 12-week period. This is a Phase 2 trial that is currently active but not recruiting new

participants, indicating that the study is ongoing and data is being collected from the enrolled patients. Eligibility criteria include specific biomarker requirements related to microsatellite instability and mismatch repair status.

NCT: NCT03350126 · Enrollment: 57 · Sites: Avignon, France, Besançon, France, Créteil, France, Levallois-Perret, France, Lyon, France

SIGNAL PHASE2, PHASE3 All Stages RAS RESEARCH • ACTIVE_NOT_RECRUITING 7

Can we Save the Rectum by Watchful Waiting or TransAnal Surgery Following (Chemo)Radiotherapy Versus Total Mesorectal Excision for Early REctal Cancer?

NCT: NCT02945566 · Enrollment: 380 · Sites: Birmingham, United Kingdom, Leuven, Belgium, Nijmegen, Netherlands, Odense, Denmark, Stockholm, Sweden

SIGNAL PHASE2, PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 7

Perioperative Systemic Therapy for Isolated Resectable Colorectal Peritoneal Metastases

This clinical trial is testing the effectiveness of perioperative systemic therapy combined with cytoreductive surgery and HIPEC in patients with isolated resectable colorectal peritoneal metastases. The intervention includes various combinations of chemotherapy regimens (CAPOX, FOLFOX, FOLFIRI) with bevacizumab, alongside the experimental procedure of cytoreductive surgery with HIPEC. Currently in Phase II-III and marked as active but not recruiting, this trial aims to determine if the combination therapy improves overall survival compared to surgery alone. Eligibility criteria include having isolated resectable colorectal peritoneal metastases, although specific biomarker requirements are not detailed.

NCT: NCT02758951 · Enrollment: 358 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Genk, Belgium, Groningen, Netherlands, Nieuwegein, Netherlands

SIGNAL PHASE2, PHASE3 All Stages RAS RESEARCH • RECRUITING 7

Randomizing Two Radiotherapy Boost Options to Avoid Rectal Cancer Surgery

NCT: NCT03051464 · Enrollment: 131 · Sites: Dallas

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 7

S0820, Adenoma and Second Primary Prevention Trial

NCT: NCT01349881 · Enrollment: 354 · Sites: Aberdeen, Abilene, Abington, Adrian, Aitkin

SIGNAL PHASE3 Stage IV RESEARCH • RECRUITING 7

A Study of 2nd-line FOLFIRI ± Bevacizumab vs. Irinotecan ± Bevacizumab in mCRC

NCT: NCT03303495 · Enrollment: 280 · Sites: Guangzhou, China

MEDIUM PHASE1 Stage IV TARGETED THERAPY • NOT_YET_RECRUITING 6

Phase 1 Study of STRO-006 in Adults With Refractory Solid Tumors That Are Recurrent or Metastatic

This clinical trial is testing the safety and efficacy of STRO-006 in adults with refractory solid tumors that are recurrent or metastatic. The intervention involves administering STRO-006, both as a monotherapy and in combination with pembrolizumab, to evaluate its safety, tolerability, and preliminary anti-tumor activity. This is a Phase 1 study that is not yet recruiting participants, meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include the expression of the HER2 biomarker and the presence of specific tumor types that commonly express ITGB6.

NCT: NCT07807345 · Enrollment: 285 · Sites: Boston, Fairfax, Irving, La Jolla, Lubbock

MEDIUM PHASE2 All Stages MSI/MMR LIQUID BIOPSY • RECRUITING 6

ctDNA-Guided Immunotherapy for dMMR/MSI-H Colon Cancer

NCT: NCT07841236 · Enrollment: 23 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 6

Radiotherapy Combined With Capeox Chemotherapy, Sintilimab and Suvemcitug in Locally Advanced pMMR Rectal Cancer:a Prospective Clinical Trial

This clinical trial is investigating the safety and efficacy of a combination of neoadjuvant chemoradiotherapy with sintilimab and suvemcitug in patients with locally advanced rectal cancer that has a proficient mismatch repair (pMMR) status. The intervention includes long-course chemoradiotherapy with capecitabine, followed by a Capeox regimen combined with sintilimab and suvemcitug, which is significant for its potential to enhance treatment outcomes in this patient population. This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include having locally advanced rectal cancer with specific biomarker statuses such as pMMR.

NCT: NCT07774754 · Enrollment: 18 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 6

HSK42360-Na Tablets Combined With Cetuximab With or Without Chemotherapy in BRAF V600-Mutant Metastatic Colorectal Cancer

This clinical trial is testing the combination of HSK42360-Na tablets and cetuximab, with or without chemotherapy, in patients with BRAF V600-mutant metastatic colorectal cancer. The intervention involves HSK42360-Na, an investigational drug, alongside cetuximab, an EGFR inhibitor, to assess their safety, tolerability, and preliminary efficacy. Currently in Phase I/II and recruiting participants, this trial aims to establish the maximum tolerated dose and evaluate the treatment's effectiveness for eligible patients. Patients may need to meet specific biomarker criteria, including BRAF mutation status and other genetic markers such as MSI and RAS.

NCT: NCT07790510 · Enrollment: 180 · Sites: Beijing, China

SIGNAL PHASE2, PHASE3 All Stages RESEARCH • RECRUITING 6

Protective Role of Alpha-Lipoic Acid in Chemotherapy-Induced Mucositis

NCT: NCT07736495 · Enrollment: 50 · Sites: Cairo, Egypt

MEDIUM PHASE1 Stage IV KRAS TARGETED THERAPY • NOT_YET_RECRUITING 6

EMB-01 in Combination With Chemotherapy for Unresectable or Metastatic Colorectal Cancer

This clinical trial is testing the safety and preliminary efficacy of EMB-01 in combination with various chemotherapy regimens for patients with unresectable or metastatic colorectal cancer. The intervention includes EMB-01 along with irinotecan, TAS-102, mFOLFOX6, and FOLFIRI, which is important for assessing potential new treatment combinations. This is a Phase Ib trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility may include specific biomarker requirements such as KRAS, NRAS, BRAF, HER2, NTRK, and RAS.

NCT: NCT07649655 · Enrollment: 120 ·

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 6

A Clinical Study of TQB2930 Injection and Chemotherapy With or Without Bevacizumab in the Treatment of Advanced Colorectal Cancer

This clinical trial is testing the safety and preliminary efficacy of TQB2930 injection in combination with chemotherapy, with or without bevacizumab, for patients with HER2-positive unresectable locally advanced or metastatic colorectal cancer. The intervention involves administering TQB2930 alongside oxaliplatin and capecitabine, with the addition of bevacizumab for some participants, which is important for evaluating potential treatment options for this patient population. The trial is currently in Phase 1 and Phase 2 and is not yet recruiting, meaning that patients will need to wait until recruitment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, HER2, and microsatellite status.

NCT: NCT07653685 · Enrollment: 71 · Sites: Beijing, China, Changchun, China, Changsha, China, Harbin, China, Jingzhou, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 6

An Exploratory Study of Zanidatamab in HER2-positive Advanced Tumor After at Least One Line of Standard Therapy

NCT: NCT07631871 · Enrollment: 10 ·

MEDIUM NA Stage III MSI/MMR TARGETED THERAPY • RECRUITING 6

Active Surveillance vs Adjuvant Chemoradiotherapy for Locally Resected Intermediate-Risk T1 Rectal Cancer

NCT: NCT07669298 · Enrollment: 480 · Sites: Gdansk, Poland, Limoges, France, Warsaw, Poland

MEDIUM PHASE2 Stage IV BRAF LIQUID BIOPSY • NOT_YET_RECRUITING 6

SBRT Followed by PD-1 Inhibitor, Bevacizumab and TAS-102 as Third-Line Therapy for Recurrent/Metastatic Colorectal Cancer

This phase II clinical trial is investigating the effectiveness of stereotactic body radiotherapy (SBRT) followed by a combination of sintilimab, bevacizumab, and TAS-102 as a third-line treatment for patients with recurrent or metastatic colorectal cancer (mCRC) that has progressed after at least two prior lines of systemic therapy. The intervention involves administering SBRT to lung or liver metastases, followed by the immune checkpoint inhibitor sintilimab, the anti-VEGF drug bevacizumab, and the chemotherapy agent TAS-102, which may enhance treatment outcomes for this patient population. Currently, the trial is not yet recruiting participants, meaning that patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MSI, and MMR status, which may help identify suitable candidates for this treatment approach.

NCT: NCT07535632 · Enrollment: 58 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

Single-arm, Phase II Clinical Study Evaluating the Safety and Efficacy of Sintilimab Combined With IBI-310 and Chemotherapy as Neoadjuvant Therapy for Locally Advanced Rectal Cancer

NCT: NCT07750951 · Enrollment: 34 · Sites: Xi'an, China

MEDIUM PHASE2 Stage IV RAS IMMUNOTHERAPY • RECRUITING 6

Short-Course Radiotherapy Followed by Immunotherapy Combined With Chemotherapy for Microsatellite Stable Locally Advanced Rectal Cancer: A Prospective, Randomized, Controlled Phase II Clinical Trial

NCT: NCT07594925 · Enrollment: 130 · Sites: Tianjin, China

MEDIUM PHASE2 All Stages MSI/MMR LIQUID BIOPSY • RECRUITING 6

A Phase II Platform Study to Evaluate Treatment With Cemiplimab Monotherapy or Cemiplimab Plus Fianlimab or Other Novel Combinations in Patients With Colorectal Cancer With Minimal Residual Disease Following Definitive Surgery and Chemotherapy (EMPIRE)

The EMPIRE trial is evaluating the effectiveness of cemiplimab monotherapy and cemiplimab in combination with fianlimab or other novel therapies in patients with colorectal cancer who have minimal residual disease following definitive surgery and chemotherapy. The intervention involves immunotherapeutic drugs, which aim to enhance the immune system's ability to target and eliminate cancer cells, particularly in patients who are ctDNA-positive after treatment. This is a Phase II study currently recruiting participants, which means eligible patients may have the opportunity to receive these investigational treatments while contributing to the understanding of their effectiveness. Eligibility criteria include specific biomarkers such as MSI, MMR, and ctDNA status.

NCT: NCT07058012 · Enrollment: 79 · Sites: Gainesville, Iowa City, Kansas City, Kingsport, Midlothian

SIGNAL PHASE2, PHASE3 All Stages BRAF RESEARCH • RECRUITING 6

DETERMINE Trial Treatment Arm 07: Dabrafenib in Combination With Trametinib in Adult, Paediatric and Teenage/Young Adult Patients With BRAF V600 Mutation-Positive Cancers.

NCT: NCT07440290 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

MEDIUM PHASE2 Stage IV KRAS LIQUID BIOPSY • RECRUITING 6

ROMANCE: "Irinotecan Plus Cetuximab Rechallenge Versus Trifluridine/Tipiracil Plus Bevacizumab in Molecularly Selected Metastatic Colorectal Cancer"

The ROMANCE trial is testing the effectiveness of irinotecan plus cetuximab rechallenge versus trifluridine/tipiracil plus bevacizumab in patients with metastatic colorectal cancer who have experienced disease progression after standard therapies. The intervention involves two different drug combinations, which is significant for determining the best third-line treatment option for this patient population. This is a phase II trial currently recruiting 150 participants, which means eligible patients may have the opportunity to access these treatment options while contributing to the research. Enrollment requires specific molecular characteristics of the cancer, identified through circulating tumor DNA analysis, including biomarkers such as KRAS, NRAS, BRAF, HER2, and RAS.

NCT: NCT07381764 · Enrollment: 150 · Sites: Alessandria, Italy, Ancona, Italy, Aviano, Italy, Caserta, Italy, Castellana Grotte, Italy

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 6

Testing Ivonescimab in Combination With Chemotherapy in Patients With Metastatic Colorectal Cancers Without Liver Metastases

This clinical trial is testing the efficacy of ivonescimab in combination with the FOLFIRI chemotherapy regimen in patients with metastatic colorectal cancer (mCRC) who do not have liver metastases. The intervention involves administering ivonescimab alongside FOLFIRI, comparing its effectiveness to the standard treatment of FOLFIRI combined with bevacizumab, with a focus on progression-free survival (PFS). This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status, which will determine patient inclusion in the study.

NCT: NCT07359456 · Enrollment: 130 · Sites: Auxerre, France, Beuvry, France, Caen, France, Calais, France, Compiègne, France

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

Modified Short-Course Radiotherapy Combined With Immunochemotherapy and Targeted Therapy as First-Line Treatment for MSS Metastatic Colorectal Cancer

NCT: NCT07512674 · Enrollment: 33 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 6

Lubiprostone Combined With Maintenance Therapy for Prevention of Postoperative Recurrence in Peritoneal Metastatic Colorectal Cancer

NCT: NCT07405736 · Enrollment: 124 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

A Single-arm Phase II Clinical Study of Camrelizumab Combined With Long-course Chemoradiotherapy for Total Neoadjuvant Therapy in Locally Advanced Low pMMR/MSS Rectal Cancer

This clinical trial is testing the combination of camrelizumab with long-course chemoradiotherapy as a total neoadjuvant therapy for patients with locally advanced low pMMR/MSS rectal cancer. The intervention involves the use of camrelizumab, a drug that may enhance the effectiveness of standard treatment by targeting the immune system. This is a Phase II trial currently recruiting participants, which means that eligible patients may have the opportunity to receive this treatment while contributing to the understanding of its efficacy. Eligibility criteria include specific biomarker requirements related to microsatellite instability and mismatch repair status.

NCT: NCT07527026 · Enrollment: 44 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 6

A Phase Ib/II Study to Evaluate HLX43 Combined With HLX07 or Serplulimab in Patients With Advanced or Metastatic Colorectal Cancer

This clinical trial is testing the efficacy, safety, and tolerance of HLX43 in combination with either HLX07 or Serplulimab in patients with advanced or metastatic colorectal cancer who have failed or are intolerant to standard first-line therapy. The interventions include various doses of HLX43 paired with HLX07 or Serplulimab, which are being evaluated for their potential benefits in this patient population. This is a Phase Ib/II study that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT07358585 · Enrollment: 126 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 6

Phase II Study of EMB-01 in Recurrent/Metastatic Colorectal Cancer Patients

NCT: NCT07314294 · Enrollment: 54 · Sites: Beijing, China, Guangzhou, China

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 6

Radiotherapy Plus CAPOX, and Iparomlimab and Tuvonralimab (QL1706) as Neoadjuvant Therapy for LARC

NCT: NCT07291401 · Enrollment: 108 · Sites: Wuhan, China

MEDIUM PHASE1, PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 6

A Study of ABT-301 Plus Tislelizumab With Bevacizumab in pMMR/Non-MSI-H Locally Advanced or mCRC

This clinical trial is testing the combination of ABT-301, Tislelizumab, and Bevacizumab in patients with locally advanced or metastatic colorectal cancer (mCRC) who have proficient mismatch repair (pMMR) and are non-MSI-H. The intervention involves escalating doses of ABT-301 alongside fixed doses of Tislelizumab and Bevacizumab, which is important for assessing safety and tolerability in this patient population. Currently, the trial is in Phase 1 and Phase 2 and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include specific biomarker assessments such as KRAS, BRAF, and MMR status.

NCT: NCT07244705 · Enrollment: 66 · Sites: Clayton, Australia, Greenslopes, Australia, Heidelberg, Australia, Kaohsiung City, Taiwan, Liverpool, Australia

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 6

Neoadjuvant Therapy for Locally Advanced Low Rectal Cancer (SMARTi-RC01)

The SMARTi-RC01 clinical trial is investigating the effectiveness of combining serplulimab, a PD-1 inhibitor, with bevacizumab, short-course radiotherapy, and chemotherapy in adults with locally advanced mid-to-low rectal cancer. This trial aims to determine if the addition of bevacizumab enhances the complete remission rate compared to serplulimab combined with chemotherapy and radiotherapy alone, while also assessing the safety of these treatments. Currently in Phase 2 and not yet recruiting, this trial will enroll 138 participants, which means patients will need to wait until recruitment begins to potentially participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and microsatellite status.

NCT: NCT07134101 · Enrollment: 138 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

The Efficacy and Safety of Neoadjuvant Therapy With Iparomlimab and Tuvonralimab in Locally Advanced MSI-H/dMMR Colorectal Cancer: An Prospective, Single-Arm Study (Neo-IT)

This clinical trial is testing the efficacy and safety of neoadjuvant therapy using Iparomlimab and Tuvonralimab in patients with locally advanced colorectal cancer that is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR). The intervention involves the use of these immunotherapy drugs, which may significantly impact the preoperative treatment approach for this specific cancer type. This is a Phase 2 trial that is currently active but not recruiting, indicating that it is in the process of evaluating its outcomes but is not accepting new participants at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, and RAS, which are essential for determining suitable candidates for this treatment.

NCT: NCT07160647 · Enrollment: 29 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

PM8002 (BNT327) in Combination With Chemotherapy in Patients With Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of PM8002 (BNT327) in combination with various chemotherapy regimens for patients with metastatic colorectal cancer. PM8002 is a bispecific antibody that targets PD-L1 and VEGF, which may enhance the effectiveness of chemotherapy in this patient population. Currently in Phase II and actively recruiting, this trial aims to enroll 100 participants to assess the objective response rate (ORR). Eligible patients may have specific biomarkers such as MSS, MSI-L, pMMR, or RAS.

NCT: NCT07133750 · Enrollment: 100 · Sites: Beijing, China, Changchun, China, Chengdu, China, Guangzhou, China, Hangzhou, China

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

A First in Human Study of ALX2004 With Advanced or Metastatic Selected Solid Tumors

NCT: NCT07085091 · Enrollment: 170 · Sites: Columbus, Fairfax, Farmington Hills, Grand Rapids, Houston

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 6

AK112 Plus FOLFIRI Versus Bevacizumab Plus FOLFIRI as Second-line Treatment of MSS/pMMR Metastatic Colorectal Cancer

This clinical trial is testing the combination of AK112 plus FOLFIRI against bevacizumab plus FOLFIRI as a second-line treatment for patients with MSS/pMMR metastatic colorectal cancer who have not tolerated oxaliplatin-containing first-line therapy or have experienced disease progression or recurrence within six months after oxaliplatin adjuvant therapy. The intervention involves the investigational drug AK112 alongside the standard chemotherapy regimen FOLFIRI, which is important for evaluating its safety and antitumor activity compared to an established treatment. This is a Phase II trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarker assessments such as MSI, MSI-H, MMR, DMMR, and NGS.

NCT: NCT07021950 · Enrollment: 130 ·

MEDIUM PHASE1 Stage IV KRAS TARGETED THERAPY • RECRUITING 6

A Phase 1 Study of TGI-5 as Monotherapy and in Combination With Nivolumab in Subjects With Locally Advanced/Metastatic Solid Tumors

This clinical trial is testing the drug TGI-5, both as a standalone treatment and in combination with Nivolumab, for patients with unresectable locally advanced or metastatic solid tumors. The study aims to evaluate the tolerability, safety, pharmacokinetics, pharmacodynamics, and preliminary antitumor activity of TGI-5. It is currently in Phase 1 and is actively recruiting participants, which means eligible patients may have the opportunity to access this investigational treatment. Eligibility criteria include specific biomarkers such as KRAS, BRAF, MSI, MSI-H, and RAS.

NCT: NCT07376707 · Enrollment: 194 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 6

Clinical Study on the First-line Treatment of Advanced Colorectal Cancer With Simultaneous Infusion of Levozolinate for Injection and 5-FU

NCT: NCT07391618 · Enrollment: 583 · Sites: Beijing, China

MEDIUM N/A Stage III MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 6

A Real-World Study of Neoadjuvant/Conversion Therapy for Colorectal Cancer With Chemotherapy And Anti-Angiogenic Targeted Agents

NCT: NCT06470178 · Enrollment: 70 · Sites: Wuhan, China

SIGNAL PHASE3 Stage IV MSI/MMR RESEARCH • RECRUITING 6

SBRT + PD-1 Monoclonal Antibody in Unresectable Colorectal Liver Metastases

This clinical trial is testing the efficacy and safety of combining stereotactic body radiation therapy (SBRT) with a PD-1 monoclonal antibody in patients with unresectable colorectal cancer liver metastases. The intervention involves the use of SBRT, a precise form of radiation therapy, alongside immunotherapy to potentially enhance treatment outcomes. Currently in Phase 3 and actively recruiting participants, this trial aims to provide high-level evidence for this combination therapy's effectiveness. Eligibility criteria include the presence of specific biomarkers, such as mismatch repair (MMR) status.

NCT: NCT06794086 · Enrollment: 24 · Sites: Guangzhou, China

MEDIUM PHASE2 All Stages KRAS LIQUID BIOPSY • RECRUITING 6

Valproic Acid to Potentiate Anti-EGFR Treatment Efficacy and Prevent/Revert Resistance in Colorectal Cancer

NCT: NCT06714357 · Enrollment: 130 · Sites: Avellino, Italy, Caserta, Italy, Frattamaggiore, Italy, Naples, Italy, Pozzuoli, Italy

MEDIUM N/A Stage III MSI/MMR TARGETED THERAPY • RECRUITING 6

Exploring the Treatment Duration of PD-1 Neoadjuvant Therapy in Stage II-III dMMR Rectal Cancer

NCT: NCT06613165 · Enrollment: 100 · Sites: Xi'an, China

MEDIUM PHASE1 Stage IV BRAF TARGETED THERAPY • RECRUITING 6

A First-in-Human Escalation and Expansion Study of Patients With Advanced Solid Tumors

This clinical trial is testing the investigational drug JR8603 in patients with advanced solid tumors who have not responded to prior treatments. The study aims to evaluate the safety and effectiveness of JR8603, which is significant for patients seeking alternative options after standard therapies have failed. Currently in Phase 1 and actively recruiting, this trial provides an opportunity for patients to access a potential treatment while contributing to research. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, HER2, and RAS.

NCT: NCT06783569 · Enrollment: 94 · Sites: Harbin, China, Shenyang, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

Phase 2 Study of SR-8541A in Combination With Botensilimab and Balstilimab in Subjects With Refractory Metastatic Microsatellite Stable Colorectal Cancer (MSS-CRC)

This clinical trial is testing the combination of SR-8541A, botensilimab, and balstilimab in patients with refractory metastatic microsatellite stable colorectal cancer (MSS-CRC). The intervention involves administering SR-8541A orally alongside the other two drugs intravenously, aiming to evaluate its safety, tolerability, efficacy, and determine the optimal dose for further study. This is a Phase 2 trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include specific biomarkers related to microsatellite instability and mismatch repair status.

NCT: NCT06589440 · Enrollment: 70 · Sites: Austin, Dallas, Houston, Morristown, Seattle

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

Immune Checkpoint Inhibitors for Organ Preservation in Non-metastatic dMMR/MSI-H Gastric or Colon Cancers

This clinical trial is testing the effectiveness of immune checkpoint inhibitors for organ preservation in patients with non-metastatic dMMR/MSI-H gastric or colon cancers. The intervention includes PD1/PDL1 antibody monotherapy and a combination of intensive immunotherapy with anti-VEGF treatment, both administered for up to 24 weeks, alongside the option of radical surgery. Currently in Phase 2 and actively recruiting, this trial aims to assess the organ preservation rate, which is a critical outcome for patients considering treatment options. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS, as well as NGS testing.

NCT: NCT06580574 · Enrollment: 34 · Sites: Beijing, China

SIGNAL PHASE2, PHASE3 All Stages RAS RESEARCH • RECRUITING 6

COLDFIRE-III Trial: Efficacy of Irreversible Electroporation and Stereotactic Body Radiotherapy for Perivascular and Peribiliary Colorectal Liver Metastases

The COLDFIRE-III trial is testing the efficacy of irreversible electroporation (IRE) and stereotactic body radiotherapy (SBRT) in patients with perivascular or peribiliary colorectal liver metastases that are not suitable for surgical resection or thermal ablation. The intervention involves comparing IRE, a procedure that uses electrical fields to induce cell death, with SBRT, a form of targeted radiation therapy, to determine which method provides better local control of the disease. This is a phase IIb/III trial currently recruiting participants, which means eligible patients may have the opportunity to receive one of the interventions while contributing to important clinical research. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT06185556 · Enrollment: 78 · Sites: Amsterdam, Netherlands

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 6

Chemopreventive Effect of Combination of Celecoxib and Metformin in Patients With Familial Adenomatous Polyposis

This clinical trial is testing the chemopreventive effect of a combination of celecoxib and metformin in patients with familial adenomatous polyposis (FAP). The intervention involves administering celecoxib alone and in combination with metformin to assess their impact on the number and size of polyps in the colon and duodenum. Currently in Phase 3 and actively recruiting participants, this trial aims to provide insights into effective strategies for delaying the progression of adenomas and cancers in FAP patients. Eligibility criteria may include specific biomarkers related to FAP, although detailed requirements are not provided.

NCT: NCT06545526 · Enrollment: 28 · Sites: Seoul, South Korea

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 6

Regorafenib in Combination With Multimodal Metronomic Chemotherapy for Chemo-resistant Metastatic Colorectal Cancers

NCT: NCT06425133 · Enrollment: 174 · Sites: Auxerre, France, Besançon, France, Chalon-sur-Saône, France, Colmar, France, Dijon, France

MEDIUM PHASE1 All Stages BRAF LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 6

ELVN-002 With Trastuzumab +/- Chemotherapy in HER2+ Solid Tumors, Colorectal and Breast Cancer

This clinical trial is testing the safety and tolerability of ELVN-002 in combination with trastuzumab, with and without chemotherapy, for patients with advanced-stage HER2-positive solid tumors, specifically colorectal and breast cancer. The intervention involves the drug ELVN-002 alongside trastuzumab and chemotherapy agents such as 5-Fluorouracil, Oxaliplatin, and Capecitabine, which is important for understanding potential treatment options for this patient population. This is a Phase 1 trial that is currently active but not recruiting, indicating that it is in the early stages of evaluating safety and dosage rather than efficacy. Eligibility criteria include specific biomarkers such as HER2, BRAF, and RAS, which may determine patient suitability for participation.

NCT: NCT06328738 · Enrollment: 275 · Sites: Fairfax, Plantation, St Louis

SIGNAL PHASE2, PHASE3 Stage IV BRAF RESEARCH • RECRUITING 6

Fruquintinib With or Without HAI-FOLFOX for Refractory Colorectal Cancer

NCT: NCT06441565 · Enrollment: 84 · Sites: Guangzhou, China

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 6

Reducing Inflammatory Syndrome in Surgery

The RISIS-CR trial is testing the effectiveness of alpha-ketoglutarate (AKG) in reducing postoperative inflammatory responses in patients undergoing colorectal cancer surgery. The intervention involves administering AKG as a dietary supplement to patients identified as high inflammatory responders, with the aim of minimizing complications associated with surgical inflammatory syndrome. This is a Phase 3 trial currently recruiting 80 participants, which means eligible patients may have the opportunity to receive a potentially beneficial treatment while contributing to important research outcomes. Eligibility criteria include the identification of patients based on their inflammatory responsiveness prior to surgery.

NCT: NCT06646809 · Enrollment: 80 · Sites: Singapore, Singapore

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

NCT: NCT06149481 · Enrollment: 60 · Sites: Bethesda

MEDIUM PHASE2 Stage III BRAF LIQUID BIOPSY • RECRUITING 6

CtDNA in Adjuvant Chemotherapy of Stage III Colon Cancer (REVISE Trial)

The REVISE trial is testing the effectiveness of circulating tumor DNA (ctDNA) as a biomarker to guide adjuvant chemotherapy decisions in patients with stage III colon cancer. The intervention involves two chemotherapy regimens, FOLFOXIRI and XELOX, which are standard treatments for this stage of cancer, and the trial aims to assess how ctDNA concentration changes in response to these therapies. This is a Phase 2 trial currently recruiting 60 participants, which means patients may have the opportunity to contribute to research that could refine treatment approaches for stage III colon cancer. Eligibility criteria include specific biomarker assessments such as BRAF, MSI, and mismatch repair status.

NCT: NCT06242418 · Enrollment: 60 · Sites: Chengde, China, Chengdu, China, Guiyang, China, Kunming, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 6

Liposomal Irinotecan and Leucovorin/5-fluorouracil Plus Bevacizumab in Metastatic Colorectal Cancer

NCT: NCT06184698 · Enrollment: 173 · Sites: Shijiazhuang, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

Study to Assess Adverse Events and Change in Disease Activity in Previously Treated Adult Participants Receiving Intravenous (IV) ABBV-400 With Unresectable Metastatic Colorectal Cancer in Combination With IV Fluorouracil, Folinic Acid, and Bevacizumab

NCT: NCT06107413 · Enrollment: 280 · Sites: Charleston, Charlotte, Chicago, Duarte, Durham

SIGNAL PHASE3 All Stages RESEARCH • NOT_YET_RECRUITING 6

Preoperative Imatinib Mesylate Combined With Rectal-sparing Surgery in Patients With c-KIT Gene-mutant Rectal GIST

This clinical trial is testing the safety and viability of preoperative imatinib mesylate combined with local resection in patients with c-KIT gene-mutant rectal gastrointestinal stromal tumors (GIST). The intervention involves administering imatinib mesylate prior to surgery to potentially preserve organ function and avoid more invasive procedures like abdominoperineal resection. This is a Phase 3 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include having a confirmed c-KIT gene mutation associated with rectal GIST.

NCT: NCT05970900 · Enrollment: 23 · Sites: Fuzhou, China

MEDIUM PHASE2 Stage IV BRAF LIQUID BIOPSY • RECRUITING 6

Cetuximab Plus Irinotecan in Patients With NeoRAS Wild-type Metastatic Colorectal Cancer In Third-line Therapy

This clinical trial is testing the efficacy and safety of cetuximab plus irinotecan in patients with NeoRAS wild-type metastatic colorectal cancer (mCRC) who are receiving third-line therapy. The intervention involves the combination of cetuximab, a monoclonal antibody, and irinotecan, a chemotherapy drug, which may provide a targeted approach for this specific patient population. Currently in phase II and actively recruiting, this trial aims to assess the objective response rate (ORR) to the treatment. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, CTDNA, and RAS status.

NCT: NCT05962502 · Enrollment: 54 · Sites: Guangzhou, China

SIGNAL PHASE2, PHASE3 All Stages RAS RESEARCH • ACTIVE_NOT_RECRUITING 6

REDEL Trial: Reduced Elective Nodal Dose for Anal Cancer Toxicity Mitigation

The REDEL trial is testing the efficacy of reduced elective nodal radiation in patients with anal cancer undergoing chemoradiation. The intervention involves administering a reduced dose of radiation (30.6 Gy) along with the drugs Capecitabine and Mitomycin C, aiming to mitigate treatment-related toxicity. This is a Phase 2 and Phase 3 trial that is currently active but not recruiting, indicating that patients are not being enrolled at this time. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT05902533 · Enrollment: 33 · Sites: Burlington, Cincinnati, Columbus

MEDIUM NA Stage III MSI/MMR TARGETED THERAPY • RECRUITING 6

MRD-guided Deferred Adjuvant Therapy in Resectable Early-stage Colon Cancer

NCT: NCT06204484 · Enrollment: 349 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 6

A Study to Explore the Third-line Treatment of Fruquintinib Combined With Serplulimab in Advanced Non-liver-limited Metastatic Colorectal Cancer: a Single-center, Phase 2 Study

NCT: NCT05954429 · Enrollment: 50 · Sites: Shanghai, China

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

A Phase 1-2 of ST316 With Selected Advanced Unresectable and Metastatic Solid Tumors

This clinical trial is testing the safety and efficacy of the drug ST316 in patients with selected advanced unresectable and metastatic solid tumors. The intervention involves administering ST316 intravenously, alongside established regimens such as FOLFIRI and bevacizumab, to evaluate its effects on tumors with specific abnormalities in the WNT/β-catenin signaling pathway. Currently, the trial is in Phase 1 and is active but not recruiting, indicating that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT05848739 · Enrollment: 130 · Sites: Birmingham, Boston, Denver, Durham, Grand Rapids

MEDIUM PHASE1, PHASE2 Stage IV TARGETED THERAPY • RECRUITING 6

Spevatamig (PT886) as Monotherapy or in Combination With Chemo and/or Targeted Therapy and/or ICI, for the Treatment of Patients With Advanced Gastrointestinal Cancers, Including Gastric, Gastroesophageal Junction, Pancreatic Ductal or Biliary Tract Carcinomas (the TWINPEAK Study)

NCT: NCT05482893 · Enrollment: 350 · Sites: Boston, Denver, Duarte, Durham, Fairfax

SIGNAL PHASE2, PHASE3 All Stages BRAF RESEARCH • RECRUITING 6

DETERMINE Trial Treatment Arm 05: Vemurafenib in Combination With Cobimetinib in Adult Patients With BRAF Positive Cancers.

The DETERMINE Trial Treatment Arm 05 is investigating the efficacy of the drug combination vemurafenib and cobimetinib in adult patients with BRAF positive cancers. Vemurafenib and cobimetinib are already approved for treating unresectable or metastatic melanoma, and this trial aims to assess their effectiveness in other cancer types with BRAF V600 mutations. Currently in Phase 2 and Phase 3, the trial is actively recruiting participants, which means eligible patients may have the opportunity to access these treatments during the study. Eligibility criteria include having a BRAF V600 mutation, confirmed through next-generation sequencing (NGS).

NCT: NCT05768178 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

A Study Investigating the Efficacy and Safety of Alcestobart (LBL-007) Plus Tislelizumab in Combination With Bevacizumab Plus Fluoropyrimidine Versus Bevacizumab Plus Fluoropyrimidine in Participants With Unresectable or Metastatic Colorectal Cancer

This clinical trial is investigating the efficacy and safety of alcestobart (LBL-007) combined with tislelizumab, alongside bevacizumab and fluoropyrimidine, compared to bevacizumab and fluoropyrimidine alone in participants with unresectable or metastatic colorectal cancer. The intervention involves multiple drugs, which may provide insights into their combined effects on treatment outcomes. Currently, the trial is in Phase 1b/2 and is active but not recruiting, indicating that it is ongoing and data is being collected, but no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT05609370 · Enrollment: 113 · Sites: Albuquerque, Anchorage, Billings, Cerritos, Covington

SIGNAL PHASE3 All Stages RESEARCH • RECRUITING 6

The Possible Protective Role of Ketotifen Against Oxaliplatin Induced Peripheral Neuropathy

NCT: NCT05624138 · Enrollment: 64 · Sites: Tanta, Egypt

MEDIUM NA Stage III BRAF LIQUID BIOPSY • RECRUITING 6

Circulating Tumor DNA Guided Adjuvant Chemotherapy for Colon Cancer

NCT: NCT05529615 · Enrollment: 2684 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 Stage IV RAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

Study of JK08 in Patients with Unresectable Locally Advanced or Metastatic Cancer

NCT: NCT05620134 · Enrollment: 263 · Sites: Barcelona, Spain, Brussels, Belgium, Edegem, Belgium, Ghent, Belgium, Madrid, Spain

MEDIUM PHASE1 Stage IV BRAF LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 6

A Study to Learn About the Study Medicine Called PF-07799933 in People With Advanced Solid Tumors With BRAF Alterations.

NCT: NCT05355701 · Enrollment: 103 · Sites: Aurora, Boston, Cleveland, Detroit, Fayetteville

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

A Study of AK112 With or Without AK117 in Metastatic Colorectal Cancer

This clinical trial is evaluating the efficacy, safety, and pharmacokinetics of AK112 with or without AK117 in patients with metastatic colorectal cancer who are not suitable for surgery. The intervention involves the use of AK112 and AK117, alongside standard treatments such as Oxaliplatin, Capecitabine, and Irinotecan, which are important for assessing

potential improvements in treatment outcomes. Currently, the trial is in Phase II and is active but not recruiting, indicating that it is ongoing and data is being collected, but no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and microsatellite status.

NCT: NCT05382442 · Enrollment: 254 · Sites: Fairfax, Los Angeles, Murrieta

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 6

A Phase II Study of Tislelizumab as Adjuvant Therapy for Patients With Stage II and Stage III Colon Cancer and dMMR/MSI.

This Phase II clinical trial is testing the efficacy of Tislelizumab as adjuvant therapy for patients with high-risk Stage II and Stage III colon cancer characterized by deficient mismatch repair (dMMR) or microsatellite instability high (MSI-H). Tislelizumab is an anti-PD-1 antibody that may enhance the immune response against cancer cells, which is particularly relevant given the sensitivity of dMMR colon cancers to immune checkpoint inhibition. The trial is currently not yet recruiting participants, which means that patients interested in this study will need to wait until enrollment begins. Eligibility criteria include the presence of specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and mismatch repair status.

NCT: NCT05231850 · Enrollment: 70 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

Testing Nivolumab and Ipilimumab With Short-Course Radiation in Locally Advanced Rectal Cancer

This phase II clinical trial is investigating the combination of nivolumab and ipilimumab with short-course radiation therapy in patients with locally advanced rectal cancer that has spread to nearby tissue or lymph nodes. The intervention involves the use of two immunotherapy drugs, nivolumab and ipilimumab, alongside radiation therapy, which may enhance the immune response against cancer cells while also targeting them directly with radiation. Currently, the trial is active but not recruiting new participants, indicating that it is ongoing with the existing enrolled patients. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT04751370 · Enrollment: 31 · Sites: Aberdeen, Altamonte Springs, Ames, Anchorage, Appleton

MEDIUM PHASE1 All Stages BRAF TARGETED THERAPY • RECRUITING 6

Study to Assess Adverse Events and Pharmacokinetics in Adult Participants With Non-Small Cell Lung Cancer, Head and Neck Squamous Cell Carcinoma and Other Solid Tumors, Receiving Intravenous Infusion of Azirkitug Alone or in Combination(s) With Budigalimab, Bevacizumab, or Telisotuzumab Adizutecan

NCT: NCT05005403 · Enrollment: 694 · Sites: Austin, Chicago, Duarte, Durham, Fairfax

MEDIUM PHASE1, PHASE2 All Stages KRAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

Study of LY3537982 in Cancer Patients With a Specific Genetic Mutation (KRAS G12C)

NCT: NCT04956640 · Enrollment: 540 · Sites: Baton Rouge, Birmingham, Boston, Chapel Hill, Charlotte

SIGNAL PHASE3 All Stages RESEARCH • ACTIVE_NOT_RECRUITING 6

Chemotherapy Followed by Pelvic Reirradiation Versus Chemotherapy Alone as Pre-operative Treatment for Locally Recurrent Rectal Cancer

The GRECCAR 15 trial is testing the efficacy of neoadjuvant chemotherapy followed by pelvic reirradiation compared to neoadjuvant chemotherapy alone in patients with locally recurrent rectal cancer who have previously undergone pelvic radiotherapy. The intervention involves administering six cycles of the chemotherapy regimen FOLFIRINOX, followed by radiochemotherapy and subsequent surgery, which is significant for addressing the increased risk of non-curative resection in this patient population. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the advanced stages of evaluation and may provide insights into treatment options for patients with this condition. Eligibility criteria include patients with a history of pelvic radiotherapy for primary rectal cancer and locally recurrent disease.

NCT: NCT03879109 · Enrollment: 58 · Sites: Avignon, France, Bordeaux, France, Grenoble, France, Lille, France, Lyon, France

SIGNAL PHASE3 Stage IV RESEARCH • ACTIVE_NOT_RECRUITING 6

QoL in mCRC Elderly Patients Receiving First-line Therapy Based on Simplified Geriatric Parameters.

This clinical trial is testing the effectiveness of two treatment regimens, OPTIMOX-bevacizumab and capecitabine plus bevacizumab, in elderly patients with metastatic colorectal cancer (mCRC). The intervention focuses on assessing health-related quality of life (HRQoL) at six months, which is crucial for understanding the impact of treatment on this patient population. Currently in Phase 3 and active but not recruiting, this trial aims to provide insights into optimal therapeutic strategies for elderly patients. Eligibility criteria include being an elderly patient diagnosed with mCRC, although specific biomarker requirements are not mentioned.

NCT: NCT03828227 · Enrollment: 49 · Sites: Abbeville, France, Amiens, France, Beauvais, France, Boulogne-sur-Mer, France, Cannes, France

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 6

Neoadjuvant Toripalimab With or Without Celecoxib in dMMR/MSI-H Colorectal Cancer

This clinical trial is investigating the effectiveness of neoadjuvant toripalimab, with or without celecoxib, in patients with mismatch repair-deficient (dMMR) or microsatellite instability-high (MSI-H) colorectal cancer. The intervention involves administering toripalimab, an anti-PD-1 immunotherapy, either alone or in combination with celecoxib for a total of 6 or 12 cycles, which is significant as these patients typically do not respond well to traditional chemotherapy. Currently in Phase 2 and actively recruiting, this trial aims to assess the pathological complete response (pCR) rates, which may inform treatment options for patients with this specific cancer subtype. Eligibility criteria include specific biomarker requirements such as dMMR, MSI-H, and RAS status.

NCT: NCT03926338 · Enrollment: 270 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 All Stages KRAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 6

Phase 1/2 Study of MRTX849 in Patients With Cancer Having a KRAS G12C Mutation KRYSTAL-1

The KRYSTAL-1 trial is evaluating the safety and tolerability of MRTX849 (adagrasib) in patients with advanced solid tumors that have a KRAS G12C mutation. The intervention includes MRTX849, along with other drugs such as Pembrolizumab, Cetuximab, and Afatinib, which are being tested to understand their effects on this specific mutation. This is a Phase 1/2 study that is currently active but not recruiting new participants, indicating that it is in the process of gathering data on safety and efficacy for those already enrolled. Eligibility for the trial requires patients to have a confirmed KRAS G12C mutation.

NCT: NCT03785249 · Enrollment: 731 · Sites: Albany, Allentown, Altamonte Springs, Anchorage, Arlington

MEDIUM N/A All Stages KRAS TARGETED THERAPY • RECRUITING 6

SYNERGY-AI: Artificial Intelligence Based Precision Oncology Clinical Trial Matching and Registry

NCT: NCT03452774 · Enrollment: 50000 · Sites: Albuquerque, Ann Arbor, Atlanta, Aurora, Baltimore

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 6

Neoadjuvant Immune Checkpoint Inhibition and Novel IO Combinations in Early-stage Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant immune checkpoint inhibition and novel immuno-oncology combinations in patients with early-stage colon cancer (stage 1-3 adenocarcinoma) who do not have distant metastases. The intervention involves administering a combination of drugs, including Nivolumab, Ipilimumab, Celecoxib, BMS-986253, and BMS-986016, to evaluate their safety and potential benefits before surgical resection of the tumor. This is a Phase 2 trial currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the understanding of its safety profile. Eligibility criteria include specific biomarker requirements related to MMR and DMMR status.

NCT: NCT03026140 · Enrollment: 353 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Haarlem, Netherlands, Hilversum, Netherlands, The Hague, Netherlands

MEDIUM NA All Stages BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

Liver Transplantation for Non-Resectable Colorectal Liver Metastases Without Local Treatment Options

The GTO-001 clinical trial is testing the effectiveness of liver transplantation in adults aged 18 to 65 with non-resectable colorectal liver metastases. The intervention involves liver transplantation, which may offer a significant survival advantage by completely removing the liver and its contained tumors, as opposed to traditional palliative chemotherapy. This trial is currently not yet recruiting participants, indicating that it is in the preparatory phase and not yet open for enrollment. Eligible participants must have stable or improved liver metastases that cannot be surgically removed or ablated, and may need to meet specific biomarker criteria including BRAF, MSI, MSI-H, MMR, and DMMR.

NCT: NCT07811453 · Enrollment: 260 · Sites: Berlin, Germany

MEDIUM PHASE1, PHASE2 Stage IV KRAS TARGETED THERAPY • NOT_YET_RECRUITING 5

A Study of DES-1357 in People With Deficient Mismatch Repair (dMMR)/Microsatellite Instability-High (MSI-H) Metastatic Colorectal Cancer (mCRC)

NCT: NCT07807800 · Enrollment: 88 ·

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

Efficacy and Safety of Tunlametinib Combination Therapy in the Treatment of Second-line and Above Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of Tunlametinib Combination Therapy for patients with second-line and above metastatic colorectal cancer. The intervention involves multiple drug arms, which is significant for understanding how different combinations may impact treatment outcomes. Currently, the trial is in Phase 2 and is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT07714447 · Enrollment: 91 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage II MSI/MMR LIQUID BIOPSY • NOT_YET_RECRUITING 5

A Study to Evaluate INCA33890 in the Treatment of Circulating Tumor DNA-Positive Stage II/III Microsatellite Stable Colorectal Cancer After Curative Resection and Chemotherapy

NCT: NCT07823322 · Enrollment: 40 ·

SIGNAL NA All Stages RESEARCH • NOT_YET_RECRUITING 5

Evaluation of Versatile Intelligent Staining Technology (VIST) and Spectral Flexible Imaging (SFI) for Detection and Optical Characterization of Colorectal Lesions: A Prospective Randomized Controlled Study

NCT: NCT07797361 · Enrollment: 990 · Sites: Porto Alegre, Brazil

MEDIUM N/A All Stages TARGETED THERAPY • NOT_YET_RECRUITING 5

Multimodal Liquid-Biopsy Omics Biomarkers of Treatment Response in Breast Cancer

NCT: NCT07880106 · Enrollment: 2000 · Sites: Shanwei, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

mFOLFOXIRI Plus Bevacizumab for Organ Preservation in Locally Advanced Rectal Cancer

NCT: NCT07817888 · Enrollment: 119 ·

SIGNAL NA All Stages RAS RESEARCH • NOT_YET_RECRUITING 5

Preoperative Nasal Intestinal Obstruction Catheter Mediated Terminal Ileum Fecal Diversion Versus Intraoperative Protective Ileostomy for Preventing Anastomotic Leakage After Mid Low Rectal Cancer Surgery: A Non Inferiority, Prospective, Multicenter Randomized Controlled Trial

NCT: NCT07821437 · Enrollment: 500 ·

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

A Study of MRG007 Combination Therapy for Advanced Colorectal Cancer

NCT: NCT07737600 · Enrollment: 316 · Sites: Beijing, China

MEDIUM NA All Stages TARGETED THERAPY • RECRUITING 5

Effect of Meal Timing During Cancer Treatment in Patients in Alaska: A Randomized Clinical Trial

This clinical trial is testing the effect of meal timing during cancer treatment in patients with rectal or breast cancer receiving neoadjuvant therapy at the Alaska Native Medical Center. The intervention includes time-restricted eating, health coaching, and biospecimen collection, aiming to improve therapeutic response and metabolic health. The trial is currently in the recruiting phase, which means eligible patients can still enroll to participate in the study. Eligibility criteria include having HER2 biomarkers and being part of the specified patient population at the ANMC.

NCT: NCT06802172 · Enrollment: 100 · Sites: Anchorage

MEDIUM N/A All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

Non-Operative Management and Following Immunotherapy for Colorectal Cancer and Other GI Cancers

This clinical trial is evaluating the safety and efficacy of Non-Operative Management (NOM) and Organ-Preserving Functional Surgery (OPFS) in patients with mismatch repair-deficient/microsatellite instability-high (dMMR/MSI-H) or POLE-mutated gastrointestinal cancers who have received neoadjuvant immunotherapy. The intervention involves a "Watch & Wait" strategy for patients achieving a clinical complete response or near-complete response, while others may undergo local excision or endoscopic resection, with radical operation serving as a control. The study is currently not yet recruiting and is designed to enroll 50 participants, which means patients will need to wait for recruitment to begin before they can participate. Eligibility criteria include specific biomarker requirements such as MSI, MSI-H, MMR, and DMMR status.

NCT: NCT07656740 · Enrollment: 50 · Sites: Beijing, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

Becotatug Vedotin Combined With Cetuximab in the Later-line Treatment of Metastatic RAS Wild-type Colorectal Cancer

This clinical trial is testing the combination of becotatug vedotin and cetuximab for patients with metastatic RAS wild-type colorectal cancer who have already received prior treatments. The intervention involves using these two drugs together as a salvage therapy, which is important for patients who have limited options left. This is a Phase 2 trial that is not yet recruiting, meaning that patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarker assessments such as BRAF, MSI, MMR, and RAS status.

NCT: NCT07490119 · Enrollment: 31 ·

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

FULIRI Plus Targeted Therapy for First-line Conversion Therapy of Colorectal Cancer Liver Metastases.

NCT: NCT07515963 · Enrollment: 24 · Sites: Wuhan, China

MEDIUM PHASE2 Stage IV MSI/MMR PHASE II TRIAL • RECRUITING 5

Neoadjuvant Radiotherapy Combined With NALIRIFOX and Adebrelimab in pMMR/MSS Locally Advanced Rectal Cancer: A Prospective, Randomized, Phase II Clinical Trial

NCT: NCT07622771 · Enrollment: 84 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

FOLFIRI and Bevacizumab With or Without Pelareorep for Second-Line Treatment of Metastatic RAS-Mutated, Microsatellite-Stable Colorectal Cancer

NCT: NCT07446322 · Enrollment: 60 · Sites: Atlanta, Beverly Hills, Canton, Chicago, Detroit

MEDIUM PHASE1 Stage IV MSI/MMR LIQUID BIOPSY • RECRUITING 5

GUCY2C Prime-Boost Vaccination for Advanced Colorectal and Small Bowel Adenocarcinomas

This clinical trial is testing a heterologous prime-boost vaccination strategy for patients with advanced colorectal and small bowel adenocarcinomas who have progressed on standard therapies. The intervention involves administering the Ad5.F35-hGUCY2C-PADRE vaccine followed by the Lm-GUCY2C vaccine to evaluate safety and immune response. This is a Phase 1 trial currently recruiting participants, which means patients may have the opportunity to access this investigational treatment while contributing to the understanding of its safety profile. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, CTDNA, and RAS status.

NCT: NCT07417488 · Enrollment: 18 · Sites: Philadelphia

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

LPM6690176 in Combination With Chemotherapy and Bevacizumab in Metastatic Colorectal Cancer Patients With RAS Mutation

NCT: NCT07391566 · Enrollment: 99 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 Stage III KRAS TARGETED THERAPY • RECRUITING 5

KRAS Neoantigen Nanovaccine as Adjuvant Therapy for Colorectal Cancer/Pancreatic Cancer

This clinical trial is testing the KRAS Neoantigen Nanovaccine as adjuvant therapy for patients with colorectal or pancreatic cancer who have KRAS mutations and are at high risk of recurrence. The intervention involves a nanovaccine constructed from the bacterial membranes of an engineered Lactococcus lactis strain, which expresses KRAS antigenic peptides, aiming to enhance the immune response against cancer cells. Currently in Phase 1 and Phase 2, the trial is recruiting participants, which means eligible patients may have the opportunity to receive this investigational therapy. Eligibility criteria include the presence of specific biomarkers such as KRAS, BRAF, MSI, MSI-H, MMR, DMMR, RAS, and MICROSATELLITE.

NCT: NCT07353645 · Enrollment: 49 · Sites: Nanjing, China

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

Everolimus in CDK12-Deficient Metastatic Colorectal Cancer (EVER-RECODE)

NCT: NCT07435584 · Enrollment: 38 ·

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

A Clinical Study on the Treatment of Metastatic Colorectal Cancer at the Second-line or Beyond.

NCT: NCT07395063 · Enrollment: 68 ·

MEDIUM N/A Stage III MSI/MMR TARGETED THERAPY • RECRUITING 5

Neoadjuvant Therapy With Tislelizumab for dMMR/MSI-H Stage II-III Colorectal Cancer

NCT: NCT07352280 · Enrollment: 30 · Sites: Beijing, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

A Phase II Study of Pirfenidone Plus PD-1 Inhibitor With or Without Hypofractionated Radiotherapy for Refractory pMMR/MSS Colorectal Cancer

This clinical trial is testing the combination of pirfenidone and a PD-1 inhibitor, with or without hypofractionated radiotherapy, for patients with advanced refractory pMMR/MSS colorectal cancer. The intervention aims to assess the safety, tolerability, and efficacy of these treatments, which may impact progression-free survival. Currently in Phase II and actively recruiting, this trial offers eligible patients the opportunity to participate in a study focused on potentially improving outcomes in a challenging cancer setting. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT07644286 · Enrollment: 48 · Sites: Wuhan, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 5

A Study to Evaluate the Safety and Efficacy of Two Dose Levels of ONO-4578 With Opdivo®, in Combination With mFOLFOX6 and Bevacizumab Versus Standard of Care in Participants With Non-MSI-H/dMMR, PD-L1 Positive Advanced Colorectal Cancer

NCT: NCT06948448 · Enrollment: 144 · Sites: Columbus, Jacksonville, Lone Tree, Los Angeles, Norfolk

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 5

The Safety and Efficacy of Cetuximab Beta Plus Fruquintinib With or Without Immune Checkpoint Inhibitorrs in First-line Treatment of RAS/BRAF Wild Type Unresectable Metastatic Colorectal Cancer

NCT: NCT07257653 · Enrollment: 70 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage II MSI/MMR TARGETED THERAPY • RECRUITING 5

Immunotherapy (Toripalimab) for Reducing Recurrence Risk After Surgery for Mismatch Repair Deficient Stage IIB, IIC, or III Colon Cancer

NCT: NCT07140679 · Enrollment: 40 · Sites: Atlanta, Johns Creek

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

A Study on the Efficacy and Safety of Irinotecan Liposome (II), 5-FU/LV in Combination With Bevacizumab for the Treatment of Advanced Colorectal Cancer

NCT: NCT07123441 · Enrollment: 300 · Sites: Shanghai, China

MEDIUM N/A Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Real-world Efficacy and Safety of Neoadjuvant Dostarlimab in Patients With dMMR/MSI-H Locally Advanced Rectal Cancer

This clinical trial is investigating the real-world efficacy and safety of neoadjuvant dostarlimab in patients with locally advanced rectal cancer characterized by dMMR/MSI-H. The intervention, dostarlimab, is a drug that targets specific biomarkers associated with this type of cancer, which is important for understanding its effectiveness in a real-world context. The trial is currently in the recruiting phase, indicating that eligible patients can still enroll to participate and contribute to the study's findings. Eligibility criteria include the presence of biomarkers such as MSI, MSI-H, MMR, and DMMR, as well as enrollment in the AIFA monitoring registry for dostarlimab.

NCT: NCT07269249 · Enrollment: 50 · Sites: Naples, Italy

MEDIUM PHASE1 All Stages KRAS TARGETED THERAPY • RECRUITING 5

A Study to Find a Suitable Dose of ASP5834 in Adults With Solid Tumors

NCT: NCT07094204 · Enrollment: 404 · Sites: Atlanta, Buffalo, Cleveland, Fairfax, Grand Rapids

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

QL1706 Plus Bevacizumab for Unresectable or Metastatic MSI-H/dMMR CRC

NCT: NCT07009145 · Enrollment: 22 · Sites: Jinan, China

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Clinical Trial of an Anti-cancer Drug, CA-4948 (Emavusertib), in Combination With Chemotherapy Treatment (FOLFOX Plus Bevacizumab) in Metastatic Colorectal Cancer

NCT: NCT06696768 · Enrollment: 24 · Sites: City of Saint Peters, Creve Coeur, Fairway, Gainesville, Kansas City

MEDIUM EARLY_PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

A Dose-Expansion Trial of Intravenous HNF4α srRNA for Unresectable or Metastatic Colorectal Cancer

NCT: NCT07050394 · Enrollment: 20 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Phase II Clinical Trial of Cadonilimab Combined With Anti-angiogenic Agents in Metastatic dMMR/MSI-H CRC

This Phase II clinical trial is evaluating the efficacy and safety of cadonilimab combined with anti-angiogenic agents in patients with recurrent or metastatic colorectal cancer characterized by dMMR/MSI-H. The intervention involves cadonilimab in combination with an anti-angiogenic agent, selected from bevacizumab, regorafenib, or fruquintinib, which may enhance treatment outcomes for this specific patient population. The trial is currently recruiting 40 participants, indicating that eligible patients may have the opportunity to receive this combination therapy while the study is ongoing. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS, as well as prior exposure to PD-1/PD-L1 antibody therapy.

NCT: NCT07003022 · Enrollment: 40 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Toripalimab Plus Celecoxib for dMMR/MSI-H Locally Advanced Colorectal Cancer

NCT: NCT06903858 · Enrollment: 105 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 5

Glumetinib Combined With Fruquintinib in the Treatment of MET Amplification or Protein Overexpression in Third-Line Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the combination of Glumetinib and Fruquintinib for patients with third-line unresectable metastatic colorectal cancer who have MET amplification or protein overexpression. The intervention involves the use of these two drugs to evaluate their efficacy and safety in this specific patient population. Currently in Phase 1 and Phase 2, the trial is recruiting participants, which means eligible patients may have the opportunity to access this treatment option. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and microsatellite status, as well as next-generation sequencing (NGS) results.

NCT: NCT06980532 · Enrollment: 42 · Sites: Wuhan, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Y-90, Capecitabine, and Atezolizumab for Oligometastatic CRC

The BrUOG-430 trial is evaluating the combination of yttrium-90 radioembolization, capecitabine, and atezolizumab for patients with unresectable colorectal cancer liver metastases who have previously received two or more lines of systemic therapy. This intervention aims to assess its preliminary efficacy based on the intrahepatic disease control rate at 12 weeks. Currently in Phase 2 and actively recruiting, this trial offers patients the opportunity to participate in a study that may provide insights into treatment effectiveness. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, RAS, and microsatellite status.

NCT: NCT06555133 · Enrollment: 18 · Sites: Providence

SIGNAL PHASE2 Stage III ctDNA LIQUID BIOPSY • RECRUITING 5

A Randomized Controlled Trial of Plasma ctDNA Methylation-Guided Adjuvant Chemotherapy Decision-Making in High-Risk Stage III (T4N+ or T1-3N2) Colorectal Cancer

NCT: NCT07562490 · Enrollment: 100 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

Short-Course Radiotherapy Combined with Intracavitary Brachytherapy Followed by Pucotenlimab, Bevacizumab, Oxaliplatin, and Trifluridine/Tipiracil (TAS-102) for Total Neoadjuvant Therapy of Microsatellite Stable (MSS) Locally Advanced Low Rectal Cancer

This clinical trial is testing a combination of short-course radiotherapy and several drugs, including Pucotenlimab, Bevacizumab, Oxaliplatin, and Trifluridine/Tipiracil, for patients with microsatellite stable (MSS) locally advanced low rectal cancer. The intervention aims to evaluate the organ preservation rate following total neoadjuvant therapy, which is significant for potentially improving treatment outcomes and quality of life. This is a Phase II trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MMR, and RAS status, which may determine suitability for the study.

NCT: NCT06872606 · Enrollment: 33 · Sites: Hangzhou, China

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

A Clinical Study to Evaluate the Safety and Tolerability of AWT020 in Patients With Advanced Malignancies

This clinical trial is evaluating the safety and tolerability of AWT020 in patients with advanced malignancies, including colorectal cancer. The intervention involves AWT020 used alone or in combination with other chemotherapy agents such as Taxol, Cisplatin, Carboplatin, Pemetrexed, Oxaliplatin, and Capecitabine, which is important for understanding its potential role in treatment regimens. Currently in Phase 1 and actively recruiting, this trial aims to assess the dose-limiting toxicity (DLT) rate, which is crucial for determining the safety profile of AWT020. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS status.

NCT: NCT06839105 · Enrollment: 214 · Sites: Changchun, China, Changsha, China, Fuzhou, China, Guangzhou, China, Hangzhou, China

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

Systemic Antitumor Treatment with or Without Pressurized Intraperitoneal Aerosol Chemotherapy for Colon Peritoneal Metastases (PIPOX02)

The PIPOX02 clinical trial is investigating the effectiveness of Pressurized Intraperitoneal Aerosol Chemotherapy (PIPAC) in combination with standard medical therapy for patients with advanced peritoneal metastases from colorectal cancer. The intervention involves administering systemic treatments alongside PIPAC, which may enhance progression-free survival (PFS) compared to standard treatment alone. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and microsatellite status.

NCT: NCT06681038 · Enrollment: 114 · Sites: Caen, France, Dijon, France, Lille, France, Limoges, France, Marseille, France

MEDIUM PHASE1 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 5

A Phase 1, First-in-human Study of OKN4395 and Pembrolizumab in Patients With Solid Tumors

This clinical trial is testing the safety and tolerability of the drug OKN4395, both as a monotherapy and in combination with pembrolizumab, in patients with solid tumors. The intervention involves administering OKN4395 and pembrolizumab to evaluate their effects on tumor activity and the incidence of dose-limiting toxicities (DLTs). This is a Phase 1 study currently recruiting participants, which means it is in the early stages of testing to determine safety and appropriate dosing. Eligibility criteria include specific biomarkers such as HER2 and microsatellite instability, as well as the use of next-generation sequencing (NGS).

NCT: NCT06789172 · Enrollment: 146 · Sites: Beverly Hills, Houston, Santa Monica

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

SCRT Combined With Chemotherapy and Iparomlimab and Tuvonralimab in MSS or pMMR Patients With Locally Advanced Rectal Cancer

This clinical trial is testing a novel neoadjuvant regimen for patients with locally advanced rectal cancer (LARC) classified as T3-4/N+ without distant metastasis. The intervention involves short-course radiotherapy followed by four cycles of CAPEOX combined with the drugs Iparomlimab and Tuvonralimab, aiming to improve organ preservation and treatment efficacy. Currently in Phase 2 and actively recruiting, this trial offers patients the opportunity to participate in research that may enhance treatment outcomes. Eligibility criteria include patients with microsatellite stable (MSS) or mismatch repair proficient (pMMR) tumors, as well as specific biomarker assessments related to microsatellite instability and mismatch repair.

NCT: NCT06864013 · Enrollment: 116 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Different Cycles of Preoperative Neoadjuvant Sintilimab in Mismatch-repair Deficient/microsatellite Instability-high, Locally Advanced Colorectal Cancer

This clinical trial is testing the efficacy of different cycles of preoperative neoadjuvant sintilimab in patients with mismatch-repair deficient/microsatellite instability-high, locally advanced colorectal cancer. The intervention involves the biological agent sintilimab, which is being evaluated for its potential to improve the pathological complete response rate (pCR) in this specific patient population. This is a Phase 2 trial that is currently recruiting participants, indicating that eligible patients may have the opportunity to receive this treatment as part of the study. Eligibility criteria include the presence of biomarkers such as MSI, MSI-H, MMR, DMMR, and microsatellite instability.

NCT: NCT06698757 · Enrollment: 96 · Sites: Zheng'zhou, China

MEDIUM EARLY_PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

A Clinical Trial Evaluating the Safety and Efficacy of Intravenous HNF4α SrRNA in Treating Metastatic CRC Patients

NCT: NCT06621173 · Enrollment: 9 · Sites: Shanghai, China

SIGNAL PHASE2 Stage III ctDNA LIQUID BIOPSY • RECRUITING 5

Randomized Trial of Plasma ctDNA Methylation-Guided Adjuvant Therapy in T4N0 and Low-Risk Stage III Colorectal Cancer

This clinical trial is testing the effectiveness of plasma ctDNA methylation-guided adjuvant therapy in patients with T4N0 and low-risk stage III colorectal cancer. The intervention involves the CAPEOX regimen, which is being evaluated for its impact on patient outcomes compared to a control group receiving a shorter duration of therapy. Currently in Phase 2 and actively recruiting, this trial aims to assess the primary outcome of patients' 2-year progression-free survival. Eligible participants will have specific criteria related to their cancer stage and may be monitored through blood samples for ctDNA analysis.

NCT: NCT07562503 · Enrollment: 340 · Sites: Shanghai, China

SIGNAL PHASE2 Stage III ctDNA LIQUID BIOPSY • RECRUITING 5

Randomized Study of Plasma ctDNA Methylation to Guide Adjuvant Chemotherapy Decisions in High-Risk T3N0 Colorectal Cancer

This clinical trial is testing the use of plasma ctDNA methylation to guide adjuvant chemotherapy decisions in patients with high-risk T3N0 stage II colorectal cancer. The intervention involves administering Capecitabine and FOLFOX chemotherapy regimens to evaluate their impact on ctDNA clearance rates and prognostic significance. Currently in Phase 2 and actively recruiting, this trial aims to improve patient outcomes by tailoring chemotherapy based on postoperative ctDNA status. Eligible participants must be ctDNA-positive one month after surgery.

NCT: NCT07562711 · Enrollment: 340 · Sites: Shanghai, China

MEDIUM PHASE1, PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 5

A Study of Oral 7HP349 (Alintegimod) in Combination With Ipilimumab Followed by Nivolumab Monotherapy

NCT: NCT06362369 · Enrollment: 126 · Sites: Aurora, Houston, Lake Mary, Lebanon, Providence

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

First-Line Treatment Induced With mFOLFOX6 and HLX04 Regimen, Following Combined With Serplulimab in MSS Initially Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the effectiveness and safety of the mFOLFOX6 and HLX04 regimen combined with Serplulimab as a first-line treatment for patients with MSS-type initially unresectable metastatic colorectal cancer. The intervention involves administering the mFOLFOX6 and HLX04 regimen followed by Serplulimab, which is significant for understanding potential treatment options for this patient population. This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include having untreated MSS-type initial unresectable metastatic colorectal cancer, with specific biomarkers such as MSI, MSI-H, MMR, and DMMR being relevant to the study.

NCT: NCT06491355 · Enrollment: 72 · Sites: Hanzhou, China

SIGNAL N/A Stage IV BRAF RESEARCH • RECRUITING 5

cfDNA 5mC/5hmC Biomarkers to Predict Chemotherapy Response in Metastatic Colorectal Cancer

The EpiCORE study is testing cfDNA-based epigenetic markers to predict chemotherapy response in patients with metastatic colorectal cancer (mCRC). The intervention involves cfDNA 5mC/5hmC sequencing and the EpiCORE assay, which aim to create a non-invasive biomarker panel to differentiate between responders and non-responders to first-line chemotherapy regimens (FOLFOX or FOLFIRI). This trial is currently in the recruiting phase, indicating that eligible patients can still enroll to participate. Eligibility criteria include the presence of specific biomarkers such as BRAF and RAS.

NCT: NCT07224815 · Enrollment: 600 · Sites: Duarte

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

Liposomal Irinotecan Plus Bevacizumab in Irinotecan-refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of liposomal irinotecan and bevacizumab in patients with irinotecan-refractory metastatic colorectal cancer. The intervention involves administering liposomal irinotecan, a formulation designed to enhance the delivery and effectiveness of the drug, alongside bevacizumab, which targets blood vessel growth in tumors. This study is in Phase 1 and Phase 2 and is currently not yet recruiting, meaning that patients will need to wait until recruitment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status.

NCT: NCT06434090 · Enrollment: 74 · Sites: Guangzhou, China

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

The Phoenix Trial: Phase II Trial of Cemiplimab for the Non-operative Management of Localized dMMR Colon Cancer

The Phoenix Trial is a Phase II clinical trial evaluating the efficacy of cemiplimab for the non-operative management of localized dMMR colon cancer. Cemiplimab is an immunotherapy drug that targets specific biomarkers associated with this

type of cancer, which may influence treatment outcomes. Currently, the trial is active but not recruiting new participants, meaning that it is ongoing with the existing enrolled population of 50 patients. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite instability.

NCT: NCT05961709 · Enrollment: 50 · Sites: Camden, Gilbert, Houston, Jacksonville, New Orleans

MEDIUM N/A All Stages MSI/MMR TARGETED THERAPY • RECRUITING 5

Immune Checkpoint Inhibitor Response in Solid Tumors Using a Live Tumor Diagnostic Platform

This clinical trial is focused on evaluating the accuracy of the Elephas Score in predicting clinical response to various immunotherapies and chemoimmunotherapy in patients with solid tumors, including colorectal cancer. The intervention involves the collection of biospecimens and tissue samples, which are essential for assessing the effectiveness of the Elephas diagnostic platform. The trial is currently in the recruiting phase, indicating that eligible patients can still enroll to participate in the study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, and microsatellite status.

NCT: NCT06349642 · Enrollment: 324 · Sites: Jacksonville, Phoenix, Rochester

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 5

Liposomal Irinotecan and Capecitabine Plus Bevacizumab as Second-line Therapy in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of liposomal irinotecan, capecitabine, and bevacizumab as a second-line therapy for patients with metastatic colorectal cancer. The combination of these drugs is being evaluated to determine its impact on progression-free survival. This is a Phase 2 trial that is currently not yet recruiting participants, which means patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status.

NCT: NCT06192680 · Enrollment: 63 ·

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 5

Platform Study of Immunotherapy Combinations in Colorectal Cancer Liver Metastases

NCT: NCT06300463 · Enrollment: 24 · Sites: New York

MEDIUM NA Stage IV IMMUNOTHERAPY • RECRUITING 5

A Randomized, Multicenter Phase II Basket Study of Hypofractionated Radiotherapy/Stereotactic Body Radiotherapy Followed by Immunotherapy-Based Systemic Therapy +/- L. Rhamnosus M9 for the First-Line Treatment of Advanced Digestive System Malignancies.

This clinical trial is testing the effectiveness of hypofractionated radiotherapy or stereotactic body radiotherapy followed by immunotherapy-based systemic therapy, with or without L. Rhamnosus M9, for patients with advanced digestive system malignancies. The intervention includes a combination of radiation therapy and various drugs, such as anti-PD-1 monoclonal antibodies and chemotherapy agents, aimed at enhancing immune response and treatment efficacy. Currently in the recruiting phase, this Phase II study seeks to enroll 120 participants, which means eligible patients can potentially access this integrated treatment approach. Eligibility criteria may include specific biomarkers related to the patient's cancer type and treatment history.

NCT: NCT06349044 · Enrollment: 120 · Sites: Hangzhou, China

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Dose Escalation and Expansion Clinical Trial of Irinotecan Liposome Combined With Oxaliplatin and 5-FU/LV Plus Bevacizumab as First-line Treatment of Metastatic Colorectal Cancer

This clinical trial is testing the combination of irinotecan liposome, oxaliplatin, 5-FU/LV, and bevacizumab as a first-line treatment for patients with advanced metastatic colorectal cancer. The intervention aims to determine the maximum tolerated dose and assess the safety and efficacy of this drug combination, which is significant for optimizing treatment regimens. Currently in Phase 1 and actively recruiting participants, this trial offers patients the opportunity to receive a potentially effective treatment while contributing to the understanding of dose-limiting toxicities. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06225622 · Enrollment: 78 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Efficacy and Safety of Chemotherapy With XELOX (Oxaliplatin + Capecitabine) and Bevacizumab in Combination With Adebrelimab in First-line Treatment of Microsatellite Stable (MSS) Initially Unresectable Metastatic Colorectal Cancer

NCT: NCT06282445 · Enrollment: 36 · Sites: Jinhua, China

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

A Study to Evaluate the Safety, Pharmacokinetics, and Anti-Tumor Activity of VVD-133214 as Monotherapy and in Combination in Participants With Advanced Solid Tumors

NCT: NCT06004245 · Enrollment: 280 · Sites: Ann Arbor, Atlanta, Chicago, Duarte, Durham

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Cadonilimab in Combination With Bevacizumab and FOLFOX Regimen for the First-Line Treatment of Advanced Unresectable MSS-Type, RAS-Mutated Metastatic Colorectal Cancer

This clinical trial is investigating the combination of cadonilimab, bevacizumab, and the FOLFOX regimen for the first-line treatment of advanced unresectable MSS-type, RAS-mutated metastatic colorectal cancer. The intervention involves a dual immunotherapy approach, which is significant as it aims to improve outcomes compared to previous chemotherapy or immunotherapy options. Currently in Phase 2 and actively recruiting participants, this trial seeks to evaluate the 1-year progression-free survival rate among enrolled patients. Eligibility criteria include the presence of specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS mutations.

NCT: NCT06218810 · Enrollment: 53 · Sites: Shanghai, China

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 5

The Efficacy and Safety of Short-course Radiation Combined With Adebrelimab and CAPEOX Neo-adjuvant Therapy for Organ-retention in Patients With MSS/pMMR Ultra Low Rectal Adenocarcinoma

This clinical trial is testing the efficacy and safety of short-course radiation combined with Adebrelimab and CAPEOX neoadjuvant therapy in patients with MSS/pMMR ultra low rectal adenocarcinoma. The intervention involves the use of Adebrelimab, Oxaliplatin, Capecitabine, and short-course radiation, which aims to improve organ retention rates in this patient population. Currently in Phase 2 and actively recruiting, this trial may provide patients with access to a novel treatment approach while contributing to the understanding of its effectiveness. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06266832 · Enrollment: 30 · Sites: Hangzhou, China

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 5

Clinical Study of Short-course Radiotherapy Followed by Fruquintinib Plus Sintilimab vs Bevacizumab Plus Capecitabine as First Line Treatment in Advanced mCRC

NCT: NCT06195670 · Enrollment: 220 · Sites: Hangzhou, China

MEDIUM PHASE2 All Stages BRAF LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 5

Clinical Trial of All-trans-retinoic Acid, Bevacizumab and Atezolizumab in Colorectal Cancer

This clinical trial is testing the combination of all-trans-retinoic acid (ATRA), bevacizumab, and atezolizumab for patients with advanced colorectal cancer. The intervention involves administering ATRA orally, while bevacizumab and atezolizumab are given intravenously, which may provide insights into their combined effects on treatment outcomes. Currently in Phase 2 and marked as active but not recruiting, this trial indicates that it is ongoing but not accepting new participants at this time. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, HER2, ctDNA, RAS, and microsatellite status.

NCT: NCT05999812 · Enrollment: 22 · Sites: Dallas

MEDIUM PHASE1, PHASE2 All Stages RAS IMMUNOTHERAPY • RECRUITING 5

Safety and Efficacy of NEO212 in Patients With Astrocytoma IDH-mutant, Glioblastoma IDH-wildtype or Brain Metastasis

NCT: NCT06047379 · Enrollment: 134 · Sites: Beverly Hills, Boston, Boynton Beach, Dallas, Glendale

MEDIUM PHASE4 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Survival Benefit of Compound Kushen Injection in the Treatment of Advanced Colorectal Cancer

This clinical trial is testing the effectiveness and safety of compound kushen injection in patients with advanced colorectal cancer. The intervention involves a first-line treatment scheme combined with compound kushen injection, which is being evaluated for its potential benefits in this patient population. Currently in Phase 4 and actively recruiting, this trial aims to provide insights into progression-free survival (PFS) for participants. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT05894694 · Enrollment: 320 · Sites: Beijing, China

MEDIUM PHASE1 All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

Study of HRO761 Alone or in Combination in Cancer Patients With Specific DNA Alterations Called Microsatellite Instability or Mismatch Repair Deficiency.

NCT: NCT05838768 · Enrollment: 123 · Sites: Boston, Houston, Los Angeles, New York, San Francisco

MEDIUM N/A All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

Observational Basket Trial to Collect Tissue to Develop and Train a Live Tumor Diagnostic Platform (ELEPHAS-02)

The ELEPHAS-02 trial is an observational study aimed at developing and training a live tumor diagnostic platform specifically for patients with colorectal cancer. The intervention involves various biopsy procedures, including core needle, forceps, and punch biopsies, which are essential for collecting tissue samples to assess tumor characteristics. This trial is currently active but not recruiting participants, meaning that while the study is ongoing, no new patients are being enrolled at this time. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and TMB, which are important for understanding the tumor's genetic profile.

NCT: NCT05520099 · Enrollment: 416 · Sites: Buffalo, Chapel Hill, Hackensack, Hagerstown, Little Rock

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 5

Second-line Treatment of Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of trifluridine/tipiracil in combination with irinotecan and bevacizumab for patients with metastatic colorectal cancer who have previously received oxaliplatin and fluoropyrimidine-based chemotherapy. The intervention involves the use of trifluridine/tipiracil, which is significant as it may provide an alternative treatment option for patients who have progressed after standard therapies. This is a Phase 2 trial currently recruiting 50 participants, which means eligible patients may have the opportunity to access this treatment while contributing to the research. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, DMMR, and RAS status.

NCT: NCT06242067 · Enrollment: 50 · Sites: Jinan, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

A Study of Dostarlimab in Untreated dMMR/MSI-H Locally Advanced Rectal Cancer

This clinical trial is investigating the efficacy of dostarlimab in patients with untreated locally advanced Mismatch-repair deficient (dMMR) or Microsatellite instability-high (MSI-H) rectal cancer. The intervention involves dostarlimab monotherapy, which is significant as it aims to provide an alternative treatment option that may allow patients to avoid chemotherapy, radiation, and surgery. This is a Phase 2 trial that is currently active but not recruiting, indicating that while

the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include having dMMR or MSI-H biomarkers, which are essential for determining suitability for this specific treatment approach.

NCT: NCT05723562 · Enrollment: 154 · Sites: Albuquerque, Dallas, Los Angeles, Nashville, New York

MEDIUM PHASE2 All Stages BRAF TARGETED THERAPY • RECRUITING 5

A Phase IIa Randomized, Double-Blinded Clinical Trial of Naproxen or Aspirin for Cancer Immune Interception in Lynch Syndrome

This clinical trial is testing the effects of naproxen and aspirin on individuals with Lynch Syndrome, a hereditary condition that increases the risk of colorectal cancer. The intervention involves administering either naproxen or aspirin to evaluate their impact on T cell abundance and other immune responses in the normal colon. Currently in Phase IIa and actively recruiting participants, this trial aims to provide insights that could inform future cancer prevention strategies for those at risk. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, RAS, and microsatellite status.

NCT: NCT05411718 · Enrollment: 40 · Sites: Houston

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

Combination Immunotherapy in Colorectal Cancer

This clinical trial is testing the combination of two investigational drugs, botensilimab and balstilimab, in patients with colorectal cancer prior to surgical tumor removal. The intervention involves administering these monoclonal antibodies, which target the immune system, to evaluate their effects on the cancer and any associated side effects. Currently in Phase 2 and marked as active but not recruiting, this trial is focused on understanding the pathological overall response rate in participants. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, and mismatch repair status, particularly for participants in Cohort C.

NCT: NCT05571293 · Enrollment: 26 · Sites: Brooklyn, Flushing, New York

MEDIUM PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 5

Vitamin E Combined With Fruquintinib and Tislelizumab in Microsatellite Stabilized Metastatic Colorectal Cancer Patients

This clinical trial is testing the efficacy of Vitamin E combined with Fruquintinib and Tislelizumab in patients with microsatellite stabilized metastatic colorectal cancer (mCRC) who have not responded to standard therapy. The intervention involves the use of these three drugs to evaluate their impact on survival time, safety, and tolerability, while also analyzing various biomarkers to understand treatment efficacy and resistance mechanisms. This is a Phase 2 trial that is currently recruiting participants, which means eligible patients may have the opportunity to access this treatment option while contributing to research. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT05771181 · Enrollment: 25 · Sites: Shanghai, China

MEDIUM PHASE1, PHASE2 Stage IV KRAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

Study of DECOY20 With or Without Tislelizumab in Patients With Advanced Solid Tumors

This clinical trial is evaluating the safety, tolerability, and clinical activity of Decoy20, both as a standalone treatment and in combination with tislelizumab, in patients with advanced solid tumors. The intervention involves Decoy20, a drug that may have implications for treating these types of cancers, particularly in the context of immune response modulation. Currently, the trial is in Phase 1/2 and is active but not recruiting, which means that while it is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as KRAS, BRAF, MSI, MSI-H, MMR, DMMR, and RAS, which may be relevant for patient selection.

NCT: NCT05651022 · Enrollment: 120 · Sites: Atlanta, Buffalo, Canton, Cleveland, Detroit

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 5

Immunotherapy in Patients With Early dMMR Rectal Cancer

This clinical trial is testing the efficacy and tolerability of nivolumab and ipilimumab in patients with early-stage (stage 1-3) mismatch repair-deficient (dMMR) rectal cancer. The intervention involves administering two cycles of combination immunotherapy, which is significant for its potential to induce a complete clinical response in this patient population. This is a phase II trial that is currently recruiting participants, indicating that patients may have the opportunity to receive this

treatment while the study is ongoing. Eligibility criteria include having tumors that are microsatellite instability-high (MSI-H) or dMMR.

NCT: NCT05732389 · Enrollment: 39 · Sites: Aalborg, Denmark, Aarhus, Denmark, Copenhagen, Denmark, Odense, Denmark, Roskilde, Denmark

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

A Study of a Selective T Cell Receptor (TCR) Targeting, Bifunctional Antibody-fusion Molecule STAR0602 in Participants With Advanced Solid Tumors

NCT: NCT05592626 · Enrollment: 366 · Sites: Bethesda, Boston, Celebration, Columbus, Detroit

MEDIUM PHASE2 All Stages KRAS TARGETED THERAPY • NOT_YET_RECRUITING 5

A Combination Therapy Including Anti-PD-1 Immunotherapy in MSS Rectal Cancer With Resectable Distal Metastasis

This clinical trial is testing a combination therapy for patients with microsatellite-stable (MSS) rectal cancer who have resectable distal metastasis. The intervention involves the use of tislelizumab, an anti-PD-1 immunotherapy, in conjunction with preoperative short-course radiotherapy and neoadjuvant chemotherapy, which may enhance treatment outcomes for this patient population. This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MMR, DMMR, HER2, RAS, and microsatellite status.

NCT: NCT05359393 · Enrollment: 52 · Sites: Shanghai, China

MEDIUM PHASE2 All Stages BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

A Study of Botensilimab and Balstilimab for the Treatment of Colorectal Cancer

NCT: NCT05608044 · Enrollment: 234 · Sites: Ann Arbor, Aurora, Austin, Boston, Cleveland

MEDIUM PHASE1 Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

A Study of Safety and Efficacy of KFA115 Alone and in Combo With Pembrolizumab in Patients With Select Advanced Cancers

This clinical trial is testing the safety and efficacy of KFA115 alone and in combination with pembrolizumab in patients with select advanced cancers, including colorectal cancer. The intervention involves administering KFA115, a drug being evaluated for its tolerability and potential effectiveness, both as a single agent and in combination with pembrolizumab, an established immunotherapy. Currently in Phase 1 and marked as active but not recruiting, this trial focuses on determining the maximum tolerated dose and assessing dose-limiting toxicities, which is crucial for understanding the drug's safety profile for future studies. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, and RAS.

NCT: NCT05544929 · Enrollment: 126 · Sites: Boston, Nashville, New York, Pittsburgh

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

METIMMOX-2: Metastatic pMMR/MSS Colorectal Cancer

NCT: NCT05504252 · Enrollment: 80 · Sites: Lørenskog, Norway, Oslo, Norway, Trondheim, Norway

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Safety and Efficacy of PD-1 ± mFOLFOX6 Neoadjuvant Therapy in Local Advanced sMPCC

This clinical trial is testing the safety and efficacy of PD-1 monoclonal antibody therapy, both as a combination with mFOLFOX6 and as a single agent, in patients with locally advanced synchronous multiple primary colorectal cancer (sMPCC). The intervention involves neoadjuvant therapy aimed at improving treatment outcomes for patients with different microsatellite instability statuses (MSI-H and MSS). Currently in Phase 2 and actively recruiting participants, this trial may provide insights into effective treatment options for a rare subset of colorectal cancer patients. Eligibility criteria include specific biomarker requirements such as MSI, MMR, and RAS status, which are essential for determining the appropriate treatment approach.

NCT: NCT06002789 · Enrollment: 17 · Sites: Guangzhou, China

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 5

Single cEll pRofiling PErsistaNce To ImmuNothErapy

NCT: NCT06680739 · Enrollment: 60 · Sites: Barcelona, Spain

MEDIUM NA Stage IV BRAF TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

RESOLUTE Trial Aims to Investigate the Value of Adding Local Ablative Treatment to Standard Systemic Treatment for Unresectable Oligometastatic Colorectal Cancer

The RESOLUTE trial is investigating the clinical benefit of adding local ablative therapy to standard first-line systemic treatment for patients with unresectable oligometastatic colorectal cancer. The intervention involves local ablative therapy, which may enhance treatment outcomes by targeting metastatic lesions. This trial is currently active but not recruiting, indicating that while it is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT05862051 · Enrollment: 6 · Sites: Albury, Australia, Bendigo, Australia, Box Hill, Australia, Epping, Australia, Fitzroy, Australia

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Study of Pembrolizumab Treatment After CYAD-101 With FOLFOX Preconditioning in Metastatic Colorectal Cancer

NCT: NCT04991948 · Enrollment: 34 · Sites: Jacksonville, Tampa

MEDIUM N/A All Stages ctDNA LIQUID BIOPSY • RECRUITING 5

ORACLE: Observation of ResiduAl Cancer With Liquid Biopsy Evaluation

The ORACLE trial is testing the ability of a novel ctDNA assay, Guardant Reveal, to detect recurrence in individuals who have been treated for early-stage solid tumors, specifically colorectal cancer. The intervention involves a liquid biopsy that analyzes circulating tumor DNA, which is important for monitoring potential cancer recurrence. This trial is currently in the recruiting phase, indicating that eligible patients can still enroll to participate in the study. Eligibility criteria include the requirement for biomarkers such as HER2 and the use of liquid biopsy for assessment.

NCT: NCT05059444 · Enrollment: 2020 · Sites: Appleton, Aurora, Birmingham, Boston, Broomall

MEDIUM NA Stage IV TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

Patient Reported Outcomes Study Using Electronic Monitoring System for Advanced or Metastatic Solid Cancer

This clinical trial is testing the effectiveness of electronic patient-reported outcomes (e-PRO) monitoring in patients with unresectable advanced cancers or metastatic/recurrent solid tumors receiving systemic drug therapy. The intervention involves the use of e-PRO monitoring to assess its impact on overall survival and health-related quality of life compared to usual care. The trial is currently active but not recruiting new participants, indicating that it is in the process of collecting data from enrolled patients. Eligibility criteria include patients with specific biomarkers, such as HER2, and a total enrollment of 500 participants is planned.

NCT: NCT05931445 · Enrollment: 500 · Sites: Kobe, Japan

MEDIUM N/A Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

Retrospective Study on the Use of Immunotherapy in Patients With MSI-H Metastatic Colorectal Cancer

NCT: NCT04612309 · Enrollment: 100 · Sites: Carpi, Italy

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 5

BOLD-100 in Combination With FOLFOX With or Without Bevacizumab for the Treatment of Advanced, Solid Tumours

NCT: NCT04421820 · Enrollment: 220 · Sites: Santa Monica, Tampa

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 5

Neo-adjuvant Pembrolizumab and Radiotherapy in Localised MSS Rectal Cancer

NCT: NCT04109755 · Enrollment: 25 · Sites: Geneva, Switzerland

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

Pd1 Antibody Sintilimab ± Chemoradiotherapy for Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy of the PD-1 antibody Sintilimab in combination with chemoradiotherapy for patients with locally advanced rectal cancer, specifically focusing on those with different MMR/MSI statuses. The intervention involves administering Sintilimab alongside oxaliplatin, capecitabine, and radiotherapy, which may influence treatment outcomes based on the tumor's biomarker profile. This is a Phase 2 trial that is currently active but not recruiting, indicating that patients are not being enrolled at this time but results may still be forthcoming. Eligibility criteria include specific biomarker requirements such as MMR, MSI, and RAS status, which will determine the treatment approach for participants.

NCT: NCT04304209 · Enrollment: 195 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 5

Pembrolizumab With Ataluren in Patients With Metastatic pMMR and dMMR Colorectal Carcinoma or Metastatic dMMR Endometrial Carcinoma: the ATAPEMBRO Study

The ATAPEMBRO study is a clinical trial evaluating the safety and efficacy of the combination of Ataluren and Pembrolizumab in patients with metastatic colorectal carcinoma with either proficient or deficient mismatch repair (pMMR and dMMR) and metastatic dMMR endometrial carcinoma. The intervention involves administering Ataluren alongside Pembrolizumab, which is significant for its potential to enhance treatment outcomes in these specific cancer types. This trial is currently in Phase 1 and Phase 2 and is actively recruiting participants, indicating that eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include the presence of specific biomarkers related to mismatch repair status, namely MMR and dMMR.

NCT: NCT04014530 · Enrollment: 47 · Sites: Amsterdam, Netherlands

MEDIUM PHASE1, PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 5

Study of IDE196 in Patients With Solid Tumors Harboring GNAQ/11 Mutations or PRKC Fusions

NCT: NCT03947385 · Enrollment: 336 · Sites: Cincinnati, Cleveland, Denver, Durham, Grand Rapids

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 5

KN035 for dMMR/MSI-H Advanced Solid Tumors

NCT: NCT03667170 · Enrollment: 188 · Sites: Beijing, China, Changchun, China, Changsha, China, Fuzhou, China, Ganzhou, China

MEDIUM PHASE2 Stage IV RAS IMMUNOTHERAPY • RECRUITING 5

Immunotherapy Using Tumor Infiltrating Lymphocytes for Patients With Metastatic Cancer

This clinical trial is testing an experimental immunotherapy using Tumor Infiltrating Lymphocytes (TIL) for patients with metastatic cancer, specifically targeting those with digestive tract, urothelial, breast, or ovarian/endometrial cancers. The intervention involves the administration of pembrolizumab, fludarabine, cyclophosphamide, aldesleukin, and young TIL, which are designed to enhance the body's immune response against tumors. Currently in Phase 2 and actively recruiting,

this trial aims to evaluate the response rate of the treatment, providing patients with an opportunity to participate in a study that may contribute to understanding the safety and efficacy of this approach. Eligible participants include adults aged 18-72 with upper or lower gastrointestinal cancers, and those with specific RAS biomarkers may also be considered.

NCT: NCT01174121 · Enrollment: 332 · Sites: Bethesda

MEDIUM PHASE2 Stage IV MSI/MMR PHASE II TRIAL • NOT_YET_RECRUITING 4

A Prospective, Biomarker-Stratified Phase 2 Study: β-Hydroxybutyrate as a Predictive Biomarker for Short-Course Radiotherapy Followed by Chemotherapy Plus PD-1 Inhibitor Versus Chemotherapy Alone in Locally Advanced Rectal Cancer

NCT: NCT07847229 · Enrollment: 144 · Sites: Wuhan, China

MEDIUM NA All Stages MSI/MMR LIQUID BIOPSY • NOT_YET_RECRUITING 4

ctDNA and TEP Levels in Colon Cancer

NCT: NCT07701408 · Enrollment: 80 · Sites: Ljubljana, Slovenia

SIGNAL NA Early Detection LIQUID BIOPSY • NOT_YET_RECRUITING 4

Blood Biomarkers Based Screening for HPV-driven OPC

NCT: NCT06528353 · Enrollment: 10000 ·

SIGNAL PHASE1, PHASE2 Stage III RESEARCH • NOT_YET_RECRUITING 4

Dietary Arginine Deprivation Combined With Chemoradiotherapy for Neoadjuvant Treatment of Rectal Cancer: A Randomized Controlled Trial

This clinical trial is testing the effectiveness of an arginine-free diet combined with chemoradiotherapy in patients with locally advanced rectal cancer. The intervention includes a liquid nutritional formula that deprives the body of arginine, alongside short-course radiotherapy and the CAPOX regimen with a PD-1 inhibitor, aiming to enhance treatment outcomes. This study is in the Phase 1 and Phase 2 stages and is currently not yet recruiting, meaning patients will need to wait for enrollment to participate. Eligibility criteria may include specific health conditions related to rectal cancer, but further details on biomarkers are not provided.

NCT: NCT07755124 · Enrollment: 140 ·

MEDIUM PHASE2 Stage IV KRAS LIQUID BIOPSY • RECRUITING 4

Ivosidenib Plus mFOLFOXIRI and Celecoxib for BRAF-targeted Therapy-refractory BRAF V600E-mutant Colorectal Cancer Patients

This clinical trial is testing the combination of ivonescimab, mFOLFOXIRI, and celecoxib in patients with BRAF V600E-mutant colorectal cancer who have previously undergone BRAF-targeted therapy. The intervention involves a bispecific antibody, ivonescimab, which targets PD-1 and VEGF, alongside a chemotherapy regimen and a COX-2 inhibitor, aiming to address resistance mechanisms and improve treatment outcomes. This is a Phase 2 trial currently recruiting 45 participants, which means patients may have the opportunity to access this investigational treatment while contributing to the understanding of its efficacy. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, and others related to tumor characteristics.

NCT: NCT07809893 · Enrollment: 45 · Sites: Guangzhou, China

MEDIUM PHASE2 All Stages KRAS TARGETED THERAPY • NOT_YET_RECRUITING 4

A Study of Trastuzumab Deruxtecan (T-DXd) and Cetuximab in People With Colorectal Cancer

This clinical trial is testing the effectiveness of the combination of trastuzumab deruxtecan (T-DXd) and cetuximab in individuals with metastatic and/or unresectable colorectal cancer that expresses low levels of HER2 and has progressed after standard treatment. The intervention involves the drug fam-trastuzumab deruxtecan-Nxki, which is being evaluated for its potential to improve treatment outcomes. This is a Phase 2 trial that is currently not yet recruiting participants,

meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include specific biomarker requirements such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, and HER2 status.

NCT: NCT07691489 · Enrollment: 27 · Sites: Basking Ridge, Commack, Harrison, Middletown, Montvale

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

SHR-A1904 in Advanced Colorectal Cancer: an Exploratory Study

NCT: NCT07700719 · Enrollment: 17 · Sites: Beijing, China

MEDIUM N/A All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Safety and Efficacy of NOM and OPFS Versus RO for dMMR/MSI-H or POLE-Mutated Gastrointestinal Cancers

NCT: NCT07642323 · Enrollment: 22 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Sintilimab Plus Gossypol Acetate in Advanced Colorectal Cancer

This clinical trial is testing the combination of sintilimab and gossypol acetate in patients with advanced colorectal cancer who have previously failed at least two lines of standard therapy. The intervention involves administering oral gossypol acetate daily, followed by intravenous sintilimab every three weeks, which may provide a new approach to treatment in this patient population. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before they can participate. Eligible participants must have confirmed advanced colorectal adenocarcinoma, measurable disease, and meet specific performance and organ function criteria.

NCT: NCT07656766 · Enrollment: 32 · Sites: Beijing, China

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

KORTUC (KRC-01) Plus Radiotherapy for Organ Preservation in Rectal Cancer

The KORTUC (KRC-01) clinical trial is testing the combination of KRC-01 injections with standard chemoradiotherapy in patients with locally advanced rectal cancer without metastases. KRC-01 is an investigational drug designed to enhance the effectiveness of radiotherapy by increasing oxygen levels within tumors, which may improve treatment outcomes. This is a Phase II trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, microsatellite status, and mismatch repair status.

NCT: NCT07622017 · Enrollment: 24 · Sites: Bordeaux, France, Villejuif, France

SIGNAL PHASE1 Stage IV KRAS PHASE I TRIAL • NOT_YET_RECRUITING 4

A Phase I Multicentre Randomized Double-Blind Parallel-Controlled Study of HLX05-N vs. ERBITUX® in Metastatic Colorectal Cancer

NCT: NCT07543471 · Enrollment: 387 ·

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

BAL/BOT/agenT-797 in pMMR CRC With Liver Metastases

This clinical trial is testing the safety and effectiveness of a combination treatment involving balstilimab, botensilimab, and agenT-797 in adults with previously treated metastatic colorectal cancer that is microsatellite stable (pMMR) and has spread to the liver. The intervention consists of three drugs administered in repeating 42-day treatment cycles, which is important for evaluating their combined impact on tumor response and side effects. This is a Phase 2 trial that is not yet recruiting participants, meaning that patients will need to wait until enrollment begins to participate. Eligibility criteria include having pMMR colorectal cancer and the presence of liver metastases, with specific biomarker assessments related to microsatellite stability and mismatch repair.

NCT: NCT07550088 · Enrollment: 17 · Sites: La Jolla

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

SHR-3821 in Advanced Colorectal Cancer: an Exploratory Study

This clinical trial is testing the combination of SHR-3821 and fruquintinib in patients with advanced colorectal cancer who have experienced failure of standard therapy. The intervention involves two drugs, SHR-3821 and fruquintinib, which are being evaluated for their safety and efficacy in this patient population. This is a Phase 2 trial that is currently not yet recruiting, meaning patients cannot enroll at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and NGS.

NCT: NCT07515820 · Enrollment: 12 ·

MEDIUM EARLY_PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

ASCEND-CRC: Profiling and Targeting Dynamic Tumor Resistance in Patients With Metastatic Colorectal Cancer

NCT: NCT07318389 · Enrollment: 100 · Sites: Houston

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 4

Zanidatamab Before Surgery for the Treatment of HER2 Positive Colon and Rectal Cancer in Patients Planned for Curative Intent Treatment

This phase II clinical trial is investigating the effectiveness of zanidatamab as a neoadjuvant treatment for patients with HER2 positive colon and rectal cancer who are scheduled for curative intent surgery. Zanidatamab is a monoclonal antibody designed to inhibit tumor cell growth and spread, which may enhance surgical outcomes. The trial is currently recruiting 38 participants, and its primary outcome is the rate of complete and major pathologic regression. Eligibility criteria include specific biomarkers such as MSI, MSI-H, HER2, RAS, and mismatch repair status.

NCT: NCT07405476 · Enrollment: 38 · Sites: Atlanta, Decatur

MEDIUM NA Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Clinical Efficacy of Transarterial Infusion Chemotherapy for Unresectable Colorectal Cancer

NCT: NCT07333053 · Enrollment: 50 · Sites: Xicheng, China

SIGNAL NA Stage IV ctDNA LIQUID BIOPSY • RECRUITING 4

Prospective National Cohort Evaluating Prognostic Factors for Efficacy of Fruquintinib Treatment in Patients With Metastatic Colorectal Cancer

NCT: NCT07124858 · Enrollment: 300 · Sites: Albi, France, Amiens, France, Annecy, France, Aurillac, France, Bayeux, France

MEDIUM PHASE1, PHASE2 All Stages BRAF TARGETED THERAPY • NOT_YET_RECRUITING 4

A Study of AK112 in Combination With VG2025 in Colorectal Cancer With Liver Metastases

This clinical trial is investigating the combination of AK112 and VG2025 for patients with advanced colorectal cancer that has metastasized to the liver. The intervention involves the use of the drug AK112 alongside the biological agent VG2025, which is significant for its potential to enhance treatment efficacy. Currently, the trial is in Phase 1 and Phase 2 and is not yet recruiting participants, meaning that patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT07357220 · Enrollment: 72 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 4

A Phase II Study of Sintilimab Combined With Ipilimumab N01, Cetuximab and Dabrafenib in Patients With Microsatellite-Stable, BRAF V600E-Mutated Metastatic Colorectal Cancer

NCT: NCT07506109 · Enrollment: 49 · Sites: Beijing, China, Chengdu, China, Tianjin, China, Wuhan, China

MEDIUM PHASE1 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

PIN in Combination With Sintilimab in Previously Treated pMMR/MSS CRC With Hepatic Metastases

This clinical trial is testing the combination of PIN and sintilimab in patients with advanced proficient mismatch repair/microsatellite stable (pMMR/MSS) colorectal cancer (CRC) who have hepatic metastases. The intervention involves the use of PIN alongside the anti-PD1 antibody sintilimab to evaluate its safety and efficacy in this patient population. Currently, the trial is in Phase 1 and is not yet recruiting, which means patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MMR, and RAS, which will be assessed to determine suitability for the study.

NCT: NCT07411781 · Enrollment: 25 · Sites: Beijing, China

MEDIUM PHASE1 Stage IV RAS TARGETED THERAPY • RECRUITING 4

A Study of IDE034 in Adult Participants With Locally Advanced/Metastatic Solid Tumors Types

NCT: NCT07503808 · Enrollment: 150 · Sites: Austin, Denver, Detroit, Fairfax, Fort Worth

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 4

Safety and Efficacy of Tegavivint in Patients With Metastatic Colorectal Carcinoma

This clinical trial is testing the safety and efficacy of tegavivint in patients with metastatic colorectal carcinoma (mCRC). The intervention involves administering tegavivint as a monotherapy and in combination with standard therapies, which is important for understanding its potential role in treatment. Currently in Phase 1 and Phase 2, the trial is recruiting participants, indicating that patients may have the opportunity to access this treatment option. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MMR, DMMR, RAS, microsatellite status, and mismatch repair status.

NCT: NCT07463599 · Enrollment: 126 · Sites: Scottsdale

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 4

A Single-arm, Single-center, Phase II Clinical Study of Aipalolitovorelizumab (QL1706) Combined With Bevacizumab and Standard Chemotherapy as First-line Treatment for MSS/pMMR Metastatic Colorectal Cancer With BRAF V600E Mutation

This clinical trial is testing the combination of Aipalolitovorelizumab (QL1706) with bevacizumab and standard chemotherapy as a first-line treatment for patients with MSS/pMMR metastatic colorectal cancer that has a BRAF V600E mutation. The intervention involves administering Aipalolitovorelizumab alongside various chemotherapy regimens, which is significant for potentially improving treatment outcomes in this specific patient population. This is a Phase II study that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include having the BRAF V600E mutation and meeting specific MMR status requirements.

NCT: NCT07229846 · Enrollment: 30 · Sites: Harbin, China

SIGNAL NA Early Detection RESEARCH • RECRUITING 4

Randomized Trial Comparing Fecal Testing (FIT) to Colonoscopy for Post-polypectomy Surveillance

This clinical trial is testing the effectiveness of fecal occult blood testing (FIT) compared to colonoscopy for post-polypectomy surveillance in patients who have had colorectal polyps removed. The intervention involves using fecal testing as a less burdensome alternative to colonoscopy, which is significant given the limitations in colonoscopy availability and the fact that many surveillance colonoscopies yield no findings. The trial is currently in the recruiting phase and aims to enroll 4,000 participants, which means eligible patients can still join the study. Eligibility criteria may include a history of polyp removal, but specific biomarker requirements are not mentioned.

NCT: NCT06983639 · Enrollment: 4000 · Sites: Arendal, Norway

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING 4

A Phase Ib/II Trial of HS-20110 Combination Therapies in Advanced Colorectal Cancer Patients.

This clinical trial is evaluating the safety and efficacy of HS-20110 combination therapies in patients with advanced colorectal cancer. The interventions include HS-20110 combined with Bevacizumab and various chemotherapy regimens, which is important for assessing potential treatment options for this patient population. The trial is currently in Phase Ib/II and is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as BRAF and MSI, which may be relevant for patient selection.

NCT: NCT07283367 · Enrollment: 502 ·

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Combination Immunotherapy for the Treatment of Chemotherapy-refractory Metastatic MSS CRC

This clinical trial is testing the combination of nadunolimab and toripalimab for patients with chemotherapy-refractory metastatic microsatellite stable (MSS) colorectal cancer. The intervention involves two immunotherapy drugs: nadunolimab, which targets IL-1RAP, and toripalimab, which targets PD-1, aiming to enhance the immune response against cancer cells. Currently in Phase 1b/2 and actively recruiting, this trial will assess safety and efficacy, providing patients with the opportunity to participate in a study focused on potentially improving treatment outcomes. Eligibility criteria include specific biomarker assessments related to microsatellite instability (MSI) and mismatch repair (MMR) status.

NCT: NCT07281716 · Enrollment: 24 · Sites: New York

SIGNAL PHASE2 Stage IV RAS RESEARCH • NOT_YET_RECRUITING 4

Trifluridine/Tipiracil Plus Fruquintinib vs. Trifluridine/Tipiracil Plus Bevacizumab in Refractory Metastatic Colorectal Cancer: A Randomized, Controlled, Open-Label, Non-Inferiority Trial

This clinical trial is testing the efficacy and safety of trifluridine/tipiracil combined with fruquintinib compared to trifluridine/tipiracil combined with bevacizumab in patients with refractory metastatic colorectal cancer. The intervention involves two drug combinations, which are being evaluated to determine if one is non-inferior to the other in terms of progression-free survival (PFS). This is a Phase 2 trial that is currently not yet recruiting participants, meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT07261709 · Enrollment: 236 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 Stage IV BRAF IMMUNOTHERAPY • RECRUITING 4

A FIH, Phase I/IIa, Trial Assessing Feasibility of Administrations of TIL-based Immunotherapy in Patients With Metastatic CRC and PC

This clinical trial is testing the feasibility of TIL-based immunotherapy in patients with metastatic colorectal cancer (CRC) and pancreatic cancer (PC). The intervention involves the administration of a novel autologous TIL product, CC-38, along with pembrolizumab, cyclophosphamide, interleukin-2, and uromitexan, which is important for evaluating safety and tolerability. Currently in Phase I/IIa and actively recruiting, this trial aims to assess the repeated administration of CC-38 in a small cohort of 12 participants. Eligibility criteria include the presence of the BRAF biomarker.

NCT: NCT07255664 · Enrollment: 12 · Sites: Frankfurt, Germany

MEDIUM PHASE2 All Stages MSI/MMR LIQUID BIOPSY • RECRUITING 4

A Study of Botensilimab and Balstilimab for Colorectal Cancer With ctDNA+ After Surgery and Chemotherapy

This clinical trial is testing the effectiveness of the combination of botensilimab and balstilimab for patients with microsatellite stable (MSS) colorectal cancer or colorectal liver metastases (CRLM) who have measurable residual disease (MRD) after surgery and chemotherapy. The intervention involves administering botensilimab and balstilimab, followed by balstilimab alone, to evaluate its impact on ctDNA clearance. This is a Phase 2 trial currently recruiting participants, which means eligible patients may have the opportunity to receive this investigational treatment while contributing to the research. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, ctDNA, RAS, and NGS.

NCT: NCT07227636 · Enrollment: 284 · Sites: Basking Ridge, Commack, Harrison, Middletown, Montvale

MEDIUM PHASE1 Stage IV IMMUNOTHERAPY • RECRUITING 4

A Study of STRO-004 in Adults With Refractory/Recurrent Metastatic Cancer

This clinical trial is testing the safety and preliminary anti-tumor activity of STRO-004 in adults with refractory or recurrent metastatic cancer. The intervention involves administering STRO-004, both as a monotherapy and in combination with pembrolizumab, to evaluate its effects on tumors expressing Tissue Factor (TF). Currently in Phase 1 and actively recruiting participants, this trial aims to assess dose-limiting toxicities and establish dosing for future studies. Eligibility criteria include adults with specific metastatic cancer types that express the relevant biomarkers.

NCT: NCT07227168 · Enrollment: 200 · Sites: Austin, Boston, Denver, Fairfax, New York

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Liposomal Irinotecan, Capecitabine and Enlonstobart With Short-Course Radiotherapy for Locally Advanced Rectal Cancer

This clinical trial is testing the combination of liposomal irinotecan, capecitabine, and enlonstobart with short-course radiotherapy for patients with locally advanced rectal cancer. The intervention aims to evaluate the effectiveness of this regimen as neoadjuvant therapy, which is important for potentially improving treatment outcomes before surgery. Currently, the trial is in Phase 2 and is actively recruiting participants, indicating that eligible patients may have the opportunity to receive this treatment approach. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, microsatellite status, and mismatch repair.

NCT: NCT07281157 · Enrollment: 30 · Sites: Tianjin, China

MEDIUM PHASE4 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 4

SIS-Reinforced vs. Conventional Anastomosis for Mid-to-Low Rectal Cancer: A Multicenter RCT on Anastomotic Leak

NCT: NCT07209787 · Enrollment: 966 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 4

A Study to Investigate the Safety and Preliminary Efficacy of GSK5460025 Alone or in Combination With Other Anti-cancer Agents in Participants With Solid Tumors

This clinical trial is investigating the safety and preliminary efficacy of GSK5460025, a drug intended for participants with solid tumors that exhibit specific genetic characteristics. The intervention involves administering GSK5460025 alone or in combination with other anti-cancer agents to assess its impact on tumor size, safety, and tolerability. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and microsatellite status.

NCT: NCT07213609 · Enrollment: 47 · Sites: Aurora, Canton, Detroit, Nashville

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

STREAM-2: Second-line Treatment With REgorafenib in Advanced RAS-Mutant Colorectal Cancer

NCT: NCT07213570 · Enrollment: 60 · Sites: Naples, Italy

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Ivonescimab Plus Chemotherapy in Patients With Initially Unresectable Colorectal Cancer Liver Metastases

NCT: NCT06848842 · Enrollment: 40 · Sites: Beijing, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 4

Amplitude-Modulated Radiofrequency Electromagnetic Fields (AM RF EMF) in Combination With Fruquintinib in Refractory Metastatic Colorectal Cancer

NCT: NCT07130903 · Enrollment: 102 · Sites: Bay City, Clarkston, Detroit, Flint, Lansing

MEDIUM PHASE2 Stage IV MSI/MMR PHASE II TRIAL • RECRUITING 4

Node-Sparing Hypofractionated Radiotherapy Plus Chemotherapy and PD-1 Inhibitor in pMMR/MSS High-Risk Locally Advanced Colon Cancer: A Prospective, Randomized, Phase II Trial

NCT: NCT07230639 · Enrollment: 52 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 Stage IV KRAS TARGETED THERAPY • NOT_YET_RECRUITING 4

Efficacy Study of Digoxin Combined With Serplulimab and Chemotherapy in First-Line Treatment of MSS Advanced Colorectal Cancer

NCT: NCT07025850 · Enrollment: 20 · Sites: Anyang, China, Shanghai, China

MEDIUM PHASE1 Stage IV KRAS IMMUNOTHERAPY • RECRUITING 4

AMG 410 Alone and in Combination With Other Agents in Participants With KRAS Altered Advanced or Metastatic Solid Tumors

This clinical trial is testing the safety and efficacy of AMG 410, both alone and in combination with pembrolizumab and panitumumab, in participants with advanced or metastatic solid tumors that have KRAS alterations. The intervention involves a dose-escalation study to identify the Maximum Tolerated Dose (MTD) and the Recommended Phase 2 Dose (RP2D) of AMG 410, which is important for determining the optimal dosing for future studies. This is a Phase 1 trial currently recruiting participants, which means that it is in the early stages of testing and aims to establish safety and dosing parameters. Eligibility criteria include having KRAS alterations, which is a specific biomarker requirement for participation.

NCT: NCT07094113 · Enrollment: 434 · Sites: Atlanta, Boston, Duarte, Durham, Fairfax

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 4

TQB2922 and TAS-102 Tablets for Injection With or Without Bevacizumab in Chemotherapy-failed RAS/BRAF Wild-type Advanced Colorectal Cancer

This clinical trial is evaluating the safety and efficacy of TQB2922 injection in combination with TAS-102 tablets, with or without Bevacizumab, for patients with RAS/BRAF wild-type advanced colorectal cancer that has not responded to previous chemotherapy. The intervention involves administering TQB2922 and TAS-102 to assess their effects on patients who have exhausted standard treatment options. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive these investigational treatments. Eligibility criteria include specific biomarker assessments such as BRAF, MSI, and RAS status.

NCT: NCT07044908 · Enrollment: 72 · Sites: Beijing, China, Changchun, China, Changsha, China, Chongqing, China, Guangzhou, China

MEDIUM PHASE1, PHASE2 All Stages KRAS IMMUNOTHERAPY • RECRUITING 4

A Phase 1/2 Study of VS-7375 in Patients With KRAS G12D-Mutated Solid Tumors

NCT: NCT07020221 · Enrollment: 616 · Sites: Ann Arbor, Baltimore, Boston, Charlottesville, Fairfax

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • RECRUITING 4

A Phase II Trial of Neoadjuvant PD-1 Vaccine PD1-Vaxx in Operable MSI High Colorectal Cancer

This clinical trial is testing the neoadjuvant PD-1 vaccine IMU-201 (PD1-Vaxx) in patients with operable microsatellite instability-high (MSI-H) colorectal cancer. The intervention involves administering three doses of the PD1-Vaxx prior to surgical resection, with the goal of achieving a Major Pathological Response (MPR) defined as less than 10% viable tumor cells. This is a Phase II trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include having operable MSI-H colorectal cancer, with specific biomarker requirements such as MSI, MMR, and NGS.

NCT: NCT06692959 · Enrollment: 44 · Sites: Adelaide, Australia, Guildford, United Kingdom, Manchester, United Kingdom, Perth, Australia

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Exploring Sintilimab + Bevacizumab + Decitabine for Advanced pMMR/MSS Colorectal Cancer (After 2+ Prior Therapies)

This clinical trial is testing the combination of sintilimab, bevacizumab, and decitabine for patients with advanced proficient mismatch repair/microsatellite stable (pMMR/MSS) colorectal cancer who have received two or more prior therapies. The intervention involves intravenous infusions of these three drugs, which may provide a new approach to managing this specific cancer type. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include having pMMR/MSS colorectal cancer and having undergone at least three prior lines of systemic therapy.

NCT: NCT07007767 · Enrollment: 32 ·

SIGNAL NA All Stages RESEARCH • RECRUITING 4

Intraoperative Rectal Lavage to Prevent Local Recurrence After Laparoscopic Mid-to-Low Rectal Cancer Resection: A Multicenter Randomized Trial

This clinical trial is testing the effectiveness of intraoperative rectal irrigation during laparoscopic radical resection for patients with low-to-mid rectal cancer. The intervention involves rectal irrigation, which aims to reduce postoperative local recurrence rates compared to no irrigation. The trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate in the study. Participants will undergo surgery and follow specific postoperative protocols, with scheduled follow-ups to monitor their recovery and outcomes.

NCT: NCT07512232 · Enrollment: 1598 · Sites: Guangzhou, China

SIGNAL NA Stage IV RESEARCH • RECRUITING 4

Randomized Controlled Trial of the Effects of Combined Resistance and AerobiC Exercise on Health-related Quality of Life in Patients Undergoing First-line Chemotherapy for Metastatic Colorectal Cancer (REACH)

The REACH trial is testing the effects of combined resistance and aerobic exercise on health-related quality of life in patients undergoing first-line chemotherapy for metastatic colorectal cancer. The intervention involves 18 weeks of structured exercise training, which aims to assess its impact on improving quality of life during treatment. This trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate. Eligibility criteria include being a patient with metastatic colorectal cancer who is starting first-line chemotherapy.

NCT: NCT06938971 · Enrollment: 150 · Sites: Copenhagen, Denmark, Herlev, Denmark, Hillerød, Denmark, Roskilde, Denmark

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Sintilimab Plus Bevacizumab and Chemotherapy in MSS/pMMR Colorectal With no Liver Metastasis

NCT: NCT06973343 · Enrollment: 20 · Sites: Hefei, China

MEDIUM PHASE1, PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING 4

A Clinical Study of Arfolitixorin in Patients With mCRC

This clinical trial is testing the drug arfolitixorin in patients with metastatic colorectal cancer (mCRC). The intervention involves administering arfolitixorin as a replacement for the standard treatment leucovorin, alongside other established therapies, to evaluate its potential benefits. Currently in Phase 1 and Phase 2, the trial is recruiting up to 90 participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status.

NCT: NCT06922383 · Enrollment: 90 · Sites: Berlin, Germany, Essen, Germany, München, Germany

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Chidamide in Combination With PD-1 Inhibitor, Bevacizumab, and XELOX for Metastatic Colorectal Cancer

NCT: NCT06858969 · Enrollment: 130 ·

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Papaverine in Combination With Radiation Therapy for the Treatment of Locally Advanced Rectal Cancer, DINOMITE Trial

NCT: NCT06834126 · Enrollment: 36 · Sites: Duarte

MEDIUM PHASE1 All Stages KRAS IMMUNOTHERAPY • RECRUITING 4

BBO-11818 in Adult Subjects With KRAS Mutant Cancer

NCT: NCT06917079 · Enrollment: 665 · Sites: Boston, Dallas, Houston, Los Angeles, Nashville

MEDIUM NA Stage IV MSI/MMR IMMUNOTHERAPY • NOT_YET_RECRUITING 4

The Study on the Efficacy and Safety of Lactobacillus Johnsonii in Combination with CapeOX and Pembrolizumab for the Treatment of MSS/pMMR Metastatic Colorectal Cancer.

This clinical trial is testing the efficacy and safety of Lactobacillus johnsonii in combination with CapeOX and Pembrolizumab for patients with MSS/pMMR metastatic colorectal cancer who have previously failed standard chemotherapy. The intervention involves a dietary supplement, Lactobacillus johnsonii, which is being evaluated for its potential benefits when used alongside established cancer treatments. The trial is currently not yet recruiting participants, indicating that it is in the preparatory phase and will begin enrollment in the future. Eligibility criteria include having MSS/pMMR metastatic colorectal cancer and prior treatment failure with standard chemotherapy.

NCT: NCT06823323 · Enrollment: 150 ·

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Short-course Radiotherapy Followed by CAPOX and Ivonescimab for Locally Advanced Rectal Cancer

NCT: NCT06760520 · Enrollment: 40 ·

MEDIUM PHASE2 All Stages KRAS TARGETED THERAPY • RECRUITING 4

A Phase 2 Study of Zanidatamab in Patients With HER2-expressing Tumors

This clinical trial is testing the efficacy and safety of zanidatamab in patients with HER2-expressing tumors who have previously received treatment. Zanidatamab is a drug that targets HER2, which is significant for patients with tumors that overexpress this receptor, as it may provide a targeted treatment option. The trial is currently in Phase 2 and is actively recruiting participants, indicating that patients may have the opportunity to receive this intervention while the study is ongoing. Eligibility criteria include the presence of specific biomarkers such as HER2, KRAS, NRAS, and BRAF.

NCT: NCT06695845 · Enrollment: 200 · Sites: Amarillo, Chicago Ridge, Dallas, Detroit, Fort Myers

MEDIUM PHASE4 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

A Trial to Evaluate the Safety and Activity of Fruquintinib in Minority Populations With Advanced, Previously Treated Colorectal Cancer

NCT: NCT06562543 · Enrollment: 78 · Sites: Atlanta, Baltimore, Baton Rouge, Birmingham, Boston

MEDIUM PHASE1, PHASE2 All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING

A Study to Investigate the Safety, Pharmacokinetics, and Preliminary Effectiveness of GSK4418959 Alone or in Combination With Other Anti-cancer Agents in Participants With Solid Tumors

4

NCT: NCT06710847 · Enrollment: 14 · Sites: Denver, Detroit, Los Angeles, New York, Philadelphia

SIGNAL PHASE1 Stage IV KRAS LIQUID BIOPSY • RECRUITING

4

A First-in-human Study of BGB-53038, a Pan-KRAS Inhibitor, Alone or in Combinations in Participants With Advanced or Metastatic Solid Tumors With KRAS Mutations or Amplification

This clinical trial is testing BGB-53038, a pan-KRAS inhibitor, in participants with advanced or metastatic solid tumors that have KRAS mutations or amplification. The intervention includes BGB-53038 alone and in combination with tislelizumab or cetuximab, which is significant for understanding its safety and potential effectiveness in treating these specific cancers. Currently in Phase 1 and actively recruiting, this trial aims to assess the safety, tolerability, and preliminary antitumor activity of the drug, which is crucial for determining the next steps in treatment options for eligible patients. Participants must have tumors with KRAS mutations or amplification, and the study includes a liquid biopsy component for biomarker assessment.

NCT: NCT06585488 · Enrollment: 514 · Sites: Baltimore, Houston, Kansas City, Los Angeles

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING

4

Botensilimab and Balstilimab Optimization in Colorectal Cancer

This clinical trial is testing the combination of botensilimab and balstilimab in patients with colorectal cancer. The intervention involves administering these drugs alongside oxaliplatin, leucovorin, and fluorouracil, which is significant for evaluating their effectiveness in managing the disease. This is a Phase 2 trial that is currently active but not recruiting, meaning that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT06268015 · Enrollment: 16 · Sites: Durham

SIGNAL PHASE2 Stage III ctDNA LIQUID BIOPSY • RECRUITING

4

Randomized Controlled Study of Plasma ctDNA Methylation-Guided Adjuvant Chemotherapy in Stage I/Low-Risk Stage II Colorectal Cancer

This clinical trial is testing the effectiveness of plasma ctDNA methylation-guided adjuvant chemotherapy in patients with stage I or low-risk stage II colorectal cancer. The intervention involves the use of the drug capecitabine, which is being evaluated for its impact on ctDNA clearance rates and the prognostic significance of these rates. Currently in Phase 2 and actively recruiting, this trial aims to identify patients at high risk of recurrence and implement intensified chemotherapy strategies to improve outcomes. Eligible participants must be ctDNA-positive one month after surgery.

NCT: NCT07562659 · Enrollment: 80 · Sites: Shanghai, China

SIGNAL NA Early Detection RESEARCH • RECRUITING

4

Total Underwater Colonoscopy (TUC) for Improved Colorectal Cancer Screening: A Randomized Controlled Trial

NCT: NCT06333392 · Enrollment: 1070 · Sites: Drammen, Norway, Gothenburg, Sweden, Grålum, Norway, Lørenskog, Norway, Oslo, Norway

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING

4

A Study of Dostarlimab in Participants With Untreated Locally Advanced Rectal Cancer in China

This clinical trial is evaluating the effect of dostarlimab in participants with untreated locally advanced rectal cancer in China, specifically those with Mismatch-repair deficient (dMMR) or Microsatellite instability-high (MSI-H) tumors. The intervention, dostarlimab, is a biological therapy that targets specific cancer markers, which is important for potentially improving treatment outcomes in this patient population. This is a Phase 2 trial that is currently active but not recruiting, indicating that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include the presence of specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06640049 · Enrollment: 23 · Sites: Chengdu, China, Chongqing, China, Guangzhou, China, Hangzhou, China, Jinan, China

SIGNAL NA All Stages RAS RESEARCH • NOT_YET_RECRUITING 4

A Multicenter, Open-Label, Non-Inferiority Randomized Controlled Trial of Postoperative VTE Prevention in Chinese Patients After Colorectal Cancer Surgery

NCT: NCT06583330 · Enrollment: 1448 ·

MEDIUM N/A All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

EC_ItaLynch: Mainstreaming the Diagnosis of Lynch Syndrome

NCT: NCT06501417 · Enrollment: 600 · Sites: Roma, Italy

MEDIUM PHASE2 All Stages BRAF TARGETED THERAPY • RECRUITING 4

Cetuximab, Irinotecan, Toripalimab in RAS/BRAF Wild-type Ultraselected Right-sided Colorectal Cancer Study

NCT: NCT06547203 · Enrollment: 34 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 All Stages HER2 TARGETED THERAPY • RECRUITING 4

Beamion BCGC-1: A Study to Find a Suitable Dose of Zongertinib Used Alone and in Combination With Other Treatments to Test Whether it Helps People With Different Types of HER2+ Cancer That Has Spread

This clinical trial is testing the drug zongertinib, both alone and in combination with other treatments, for adults aged 18 years and older with various types of HER2+ cancer that has metastasized and is not amenable to surgical removal. The intervention involves zongertinib, which inhibits HER2, a protein that promotes cancer cell growth, and aims to determine the most tolerable dose while assessing its effectiveness in shrinking tumors. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible individuals may have the opportunity to receive investigational treatment. Eligibility criteria include having tumors with HER2 aberrations and a history of unsuccessful prior treatments.

NCT: NCT06324357 · Enrollment: 768 · Sites: Ann Arbor, Boston, Cerritos, Fairfax, Hershey

SIGNAL N/A Stage IV BRAF RESEARCH • RECRUITING 4

Development of a cfDNA 5mC/5hmC-based Biomarker Panel to Predict Targeted Therapy Efficacy in mCRC

NCT: NCT07224841 · Enrollment: 500 · Sites: Duarte

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Chemotherapy Plus Bevacizumab and Anti-PD-1 Followed by Induction Therapy of Chemotherapy Plus Bevacizumab

This clinical trial is testing the combination of chemotherapy, bevacizumab, and an anti-PD-1 monoclonal antibody for patients with initial unresectable metastatic colorectal cancer (mCRC) of the MSS type. The intervention involves the mFOLFOX6 regimen along with bevacizumab and a PD-1 monoclonal antibody, which may enhance treatment efficacy and improve patient outcomes. This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06415851 · Enrollment: 36 · Sites: Hanzhou, China

MEDIUM PHASE1 Stage IV RAS IMMUNOTHERAPY • RECRUITING 4

A Study of Sofe-M in Participants With Selected Advanced Solid Tumors

NCT: NCT06400472 · Enrollment: 575 · Sites: Atlanta, Columbus, Falls Church, Grand Rapids, Houston

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Nanoliposomal Irinotecan, Oxaliplatin Plus Capecitabine As Conversion Therapy of Locally Advanced Colorectal Cancer

This clinical trial is testing the efficacy and safety of nanoliposomal irinotecan, oxaliplatin, and capecitabine as conversion therapy for patients with locally advanced colorectal cancer. The intervention involves a combination of these drugs, which is significant as it aims to improve the organ-sparing R0 resection rate in this patient population. Currently in Phase 2 and actively recruiting participants, this trial may provide insights into treatment options for patients with locally advanced disease. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06405139 · Enrollment: 36 · Sites: Beijing, China

SIGNAL NA Stage III ctDNA LIQUID BIOPSY • RECRUITING 4

Circulating Tumor DNA Methylation Guiding Postoperative Adjuvant Chemotherapy in Stage III Colorectal Cancer

This clinical trial is testing the effectiveness of circulating tumor DNA (ctDNA) methylation as a guide for postoperative adjuvant chemotherapy in patients with stage III colorectal cancer. The intervention involves using ctDNA methylation to determine treatment strategies, which may allow for more tailored chemotherapy approaches based on individual risk levels. Currently in the recruiting phase, this trial aims to enroll 990 participants and will assess recurrence-free survival at three years, providing insights into treatment efficacy for both high-risk and low-risk patients. Eligibility criteria include being diagnosed with stage III colorectal cancer and having specific tumor characteristics that classify patients into high-risk or low-risk cohorts.

NCT: NCT06167967 · Enrollment: 990 · Sites: Hanzhou, China

SIGNAL N/A All Stages ctDNA LIQUID BIOPSY • RECRUITING 4

Colon Cancer Liquid Biopsy Investigation of Methylation-based Biomarkers Evaluation Using Delta-HLD (CLIMBED)

The CLIMBED trial is investigating the effectiveness of Delta-HLD, a proprietary liquid biopsy technology, for detecting colorectal cancer in an average-risk population undergoing screening colonoscopy. This observational study aims to evaluate the sensitivity of this liquid biopsy method, which is significant for improving early detection of colorectal cancer. Currently in the recruiting phase, this study involves an enrollment of 3,200 participants, providing an opportunity for individuals to contribute to research that may enhance CRC screening practices. Eligibility criteria include being part of the average-risk population for colorectal cancer.

NCT: NCT07168876 · Enrollment: 3200 · Sites: Mendoza, Argentina

MEDIUM NA Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Study on the Safety and Efficacy of MR-Linac Technique in Patients With Unresectable Locally Advanced Colon Cancer

This clinical trial is testing the safety and efficacy of the MR-Linac technique in patients with unresectable locally advanced colon cancer. The intervention involves the use of MR-Linac for short course radiotherapy, followed by chemotherapy and surgical resection, which is significant for potentially improving treatment outcomes in this patient population. The study is currently in the recruiting phase and aims to enroll 20 participants, indicating that eligible patients may still join the trial. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06244537 · Enrollment: 20 · Sites: Chengdu, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

AK104 in Locally Advanced MSI-H/dMMR Gastric Carcinoma and Colorectal Cancer

This clinical trial is testing the drug AK104 in patients with locally advanced MSI-H/dMMR gastric carcinoma and colorectal cancer. AK104 is a PD-1/CTLA-4 bispecific antibody, and its evaluation focuses on safety, tolerability, and anti-tumor activities during the perioperative period. This is a Phase II trial that is currently active but not recruiting, meaning that while the study is ongoing, no new participants are being enrolled at this time. Eligible patients must have MSI, MSI-H, MMR, or dMMR biomarkers and will receive AK104 for three cycles before surgery and up to six months after surgery.

NCT: NCT04556253 · Enrollment: 47 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 Stage IV IMMUNOTHERAPY • RECRUITING 4

GM103 Intratumoral Injection in Patients With Locally Advanced, Unresectable, Refractory and/or Metastatic Solid Tumors

This clinical trial is testing the safety and efficacy of GM103, administered alone and in combination with pembrolizumab, in patients with locally advanced, unresectable, refractory, and/or metastatic solid tumors, including colorectal cancer. The intervention involves the intratumoral injection of GM103, which is being evaluated for its potential to improve treatment outcomes in this patient population. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include having specific types of solid tumors, and the study will assess the percentage of patients experiencing dose-limiting toxicities across different treatment cohorts.

NCT: NCT06265025 · Enrollment: 125 · Sites: Goyang-si, South Korea, Seoul, South Korea

MEDIUM PHASE1 All Stages BRAF TARGETED THERAPY • RECRUITING 4

FIH, Bispecific CD276xCD3 Antibody CC-3 in Patients With Colorectal Cancer

This clinical trial is testing the bispecific antibody CC-3 in patients with colorectal cancer who have experienced failure of at least three prior therapies. CC-3 targets CD276 on cancer cells and tumor vessels, which may enhance its anti-cancer effects while minimizing off-target T cell activation and associated side effects. Currently in Phase 1 and actively recruiting participants, this trial focuses on assessing the safety of CC-3 through the evaluation of adverse events. Eligible participants may have specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, or RAS.

NCT: NCT05999396 · Enrollment: 89 · Sites: Tübingen, Germany

MEDIUM N/A All Stages MSI/MMR LIQUID BIOPSY • RECRUITING 4

ctDNA in CRC Patients Undergoing Curative-intent Surgery for Liver Metastases

This clinical trial is investigating the prognostic value of circulating tumor DNA (ctDNA) in patients with colorectal cancer (CRC) who are undergoing curative-intent surgery for liver metastases. The intervention focuses on assessing ctDNA as a biomarker to evaluate disease response and the risk of recurrence or death following surgical resection. The study is currently in the recruiting phase, which means eligible patients can still enroll to participate and contribute to the research. Eligibility criteria include specific biomarkers such as microsatellite instability (MSI), mismatch repair (MMR), and others related to ctDNA.

NCT: NCT05787197 · Enrollment: 232 · Sites: Besançon, France, Bordeaux, France, Lille, France, Lyon, France, Paris, France

MEDIUM PHASE2 All Stages BRAF TARGETED THERAPY • RECRUITING 4

Efficacy and Safety of AK104 Combined With Chemotherapy and Cetuximab or Bevacizumab

NCT: NCT07253896 · Enrollment: 40 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Neoadjuvant Immunotherapy for T4 dMMR Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant immunotherapy using Camrelizumab in patients with T4 dMMR colon cancer. The intervention aims to improve the rate of R0 resection, which is crucial for achieving better surgical outcomes and preserving adjacent organs in this patient population. Currently in Phase 2 and marked as active but not recruiting, this trial indicates that patients may not be able to enroll at this time but can still benefit from the findings once available. Eligibility for the study requires patients to have dMMR status, which is a specific biomarker associated with their cancer.

NCT: NCT06215677 · Enrollment: 18 · Sites: Beijing, China

MEDIUM PHASE1, PHASE2 Stage IV KRAS TARGETED THERAPY • RECRUITING 4

Study of Elironrasib and Daraxonrasib as Monotherapies and Combination Therapy in Participants With Advanced KRAS G12C Mutant Solid Tumors

NCT: NCT06128551 · Enrollment: 534 · Sites: Aurora, Boston, Dallas, Detroit, Duarte

SIGNAL NA Stage IV ctDNA LIQUID BIOPSY • RECRUITING 4

Circulating Tumor DNA Methylation Guided Postoperative Follow-up Strategy for Non-metastatic Colorectal Cancer

NCT: NCT05904665 · Enrollment: 584 · Sites: Shanghai, China

MEDIUM PHASE2 Stage II MSI/MMR TARGETED THERAPY • NOT_YET_RECRUITING 4

Tislelizumab Combined With Chemotherapy (CAPOX) in the Perioperative Treatment of MSI-H/dMMR Stage II or III Colorectal Cancer

NCT: NCT05841134 · Enrollment: 25 ·

MEDIUM PHASE1 All Stages BRAF TARGETED THERAPY • RECRUITING 4

A Study of ASP1002 in Adults for Treatment of Solid Tumors

NCT: NCT05719558 · Enrollment: 798 · Sites: Cleveland, Dallas, Detroit, Edmonds, Fairfax

MEDIUM PHASE1, PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Cadonilimab for PD-1/PD-L1 Blockade-refractory, MSI-H/dMMR, Advanced Colorectal Cancer

This clinical trial is investigating the efficacy of cadonilimab in patients with advanced colorectal cancer who are refractory to PD-1/PD-L1 blockade and have microsatellite instability-high (MSI-H) or mismatch repair-deficient (dMMR) tumors. The intervention involves the administration of cadonilimab, which is significant as it targets specific biomarkers associated with this cancer type. Currently, the trial is in Phase 1 and Phase 2 and is active but not recruiting, indicating that it is ongoing but not accepting new participants at this time. Eligibility criteria include the presence of biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite instability.

NCT: NCT05426005 · Enrollment: 28 · Sites: Guangzhou, China

MEDIUM PHASE2 All Stages MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Short-course Radiotherapy or Long-course Chemoradiation Followed by mFOLFOXIRI Consolidation Chemotherapy for Organ Preservation in Low Rectal Cancer

NCT: NCT06417476 · Enrollment: 66 · Sites: Guangzhou, China

MEDIUM PHASE1 Stage IV IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 4

A Study to Evaluate the Safety, Pharmacokinetics, and Activity of Enzelkitug as a Single Agent and in Combination With Checkpoint Inhibitor in Participants With Locally Advanced or Metastatic Solid Tumors

This clinical trial is evaluating the safety, pharmacokinetics, and anti-tumor activity of enzelkitug, both as a single agent and in combination with checkpoint inhibitors atezolizumab and pembrolizumab, for adult participants with locally advanced or metastatic solid tumors, including colorectal cancer. The intervention involves administering enzelkitug alongside established immunotherapies to assess its potential effectiveness and safety profile. Currently in Phase 1 and marked as active but not recruiting, this trial is focused on determining dose-limiting toxicities among the 165 enrolled participants. Eligibility criteria include adults with various solid tumors, such as colorectal cancer, but specific biomarker requirements are not detailed in the provided information.

NCT: NCT05581004 · Enrollment: 165 · Sites: Atlanta, Aurora, Boston, Germantown, Nashville

SIGNAL NA All Stages RESEARCH • RECRUITING 4

GENOCARE: A Prospective, Randomized Clinical Trial of Genotype-Guided Dosing Versus Usual Care

NCT: NCT05391126 · Enrollment: 178 · Sites: Lexington

SIGNAL N/A Stage III ctDNA LIQUID BIOPSY • RECRUITING 4

ctDNA Methylation Application in Postoperative Relapse and Adjuvant Chemotherapy Efficacy Evaluation

NCT: NCT05536089 · Enrollment: 2000 · Sites: Hangzhou, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING 4

Nivolumab and Ipilimumab in Anti-PD1-Resistant dMMR/MSI mCRC

NCT: NCT05310643 · Enrollment: 30 · Sites: Besançon, France, Bordeaux, France, Lille, France, Limoges, France, Lyon, France

MEDIUM PHASE1 Stage IV IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 4

Study of NGM831 as Monotherapy and in Combination With Pembrolizumab or Pembrolizumab and NGM438 in Advanced or Metastatic Solid Tumors

NCT: NCT05215574 · Enrollment: 130 · Sites: Austin, Gilbert, Grand Rapids, Houston, Los Angeles

MEDIUM PHASE2 Stage III MSI/MMR LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 4

Circulating Tumor DNA Based Decision for Adjuvant Treatment in Colon Cancer Stage II-III

NCT: NCT07821333 · Enrollment: 45 · Sites: Barcelona, Spain, Córdoba, Spain, L'Hospitalet de Llobregat, Spain, Madrid, Spain, Valencia, Spain

MEDIUM PHASE1 All Stages MSI/MMR IMMUNOTHERAPY • RECRUITING 4

Study of Zanzalintinib in Combination With Immuno-Oncology Agents in Participants With Solid Tumors

This clinical trial is evaluating the safety and efficacy of zanzalintinib, both as a monotherapy and in combination with immuno-oncology agents, in participants with advanced solid tumors. The intervention involves zanzalintinib, nivolumab, and ipilimumab, which are being tested to assess their effects on tumor response and safety profile. This is a Phase 1b trial currently recruiting participants, indicating that it is in the early stages of testing for safety and tolerability before further evaluation in larger cohorts. Eligibility criteria include the presence of specific biomarkers, such as microsatellite instability, which may influence treatment response.

NCT: NCT05176483 · Enrollment: 1314 · Sites: Austin, Baltimore, Boston, Celebration, Charlottesville

MEDIUM PHASE2 Stage III MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Atezolizumab in Patients With MSI-h/MMR-D Stage II High Risk and Stage III Colorectal Cancer Ineligible for Oxaliplatin

This clinical trial is testing the efficacy of atezolizumab in patients with stage II high-risk and stage III colorectal cancer who are ineligible for oxaliplatin-based adjuvant chemotherapy. The intervention involves the use of atezolizumab, a drug that targets the immune system, alongside IMM-101, to potentially enhance disease-free survival in this specific patient population. Currently in Phase 2 and marked as active but not recruiting, this trial indicates that patients are not being enrolled at this time, but results may inform future treatment options. Eligibility criteria include biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and NGS.

NCT: NCT05118724 · Enrollment: 80 · Sites: Essen, Germany

MEDIUM PHASE1 Stage IV IMMUNOTHERAPY • RECRUITING 4

A Study Evaluating the Safety and Efficacy of Targeted Therapies in Subpopulations of Patients With Metastatic Colorectal Cancer (INTRINSIC)

The INTRINSIC trial is evaluating the safety and efficacy of targeted therapies and immunotherapy in patients with metastatic colorectal cancer (mCRC) whose tumors are biomarker positive. The intervention includes a combination of drugs such as Inavolisib, Bevacizumab, Cetuximab, Atezolizumab, and Tiragolumab, which are being tested to determine their effectiveness in this specific patient population. This is a Phase 1 study currently recruiting participants, which means it is in the early stages of testing to assess safety and dosage. Eligible participants will be enrolled based on specific biomarker assay results that define their treatment arm.

NCT: NCT04929223 · Enrollment: 542 · Sites: Arlington Heights, Aurora, Baton Rouge, Birmingham, Boston

MEDIUM PHASE1, PHASE2 All Stages IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 4

A Beta-only IL-2 ImmunoTherapY Study

NCT: NCT05086692 · Enrollment: 160 · Sites: Atlanta, Boca Raton, Detroit, Houston, San Diego

MEDIUM PHASE1, PHASE2 Stage IV IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 4

Study of NGM707 As Monotherapy and in Combination with Pembrolizumab in Advanced or Metastatic Solid Tumor Malignancies

This clinical trial is investigating the efficacy of NGM707, both as a monotherapy and in combination with pembrolizumab (KEYTRUDA®), in patients with advanced or metastatic solid tumor malignancies, including colorectal cancer. The intervention involves administering NGM707 alone and in combination with pembrolizumab to assess its safety and potential effectiveness. Currently, the trial is in Phase 1 and Phase 2 and is active but not recruiting, indicating that it is ongoing but not accepting new participants at this time. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

NCT: NCT04913337 · Enrollment: 179 · Sites: Albany, Baltimore, Blacksburg, Boston, Dallas

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Two Combination Treatment Regimens of Nivolumab and Ipilimumab in Patients With dMMR and / or MSI mCRC.

This clinical trial is testing two combination treatment regimens of nivolumab and ipilimumab in patients with metastatic colorectal cancer (mCRC) who have deficient mismatch repair (dMMR) or microsatellite instability (MSI). The interventions involve administering nivolumab at two different dosages (480 mg and 240 mg) alongside ipilimumab, aiming to find a regimen that offers high clinical activity with reduced toxicity. This is a Phase 2 trial that is currently active but not recruiting, indicating that patients are not being enrolled at this time, but results may still be forthcoming. Eligibility criteria include specific biomarkers such as MSI, MMR, dMMR, RAS, and microsatellite status.

NCT: NCT04730544 · Enrollment: 96 · Sites: Avignon, France, Besançon, France, Brest, France, Créteil, France, Dijon, France

MEDIUM PHASE1 All Stages HER2 TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

CAR-macrophages for the Treatment of HER2 Overexpressing Solid Tumors

This clinical trial is testing the safety and tolerability of CAR-macrophages in patients with HER2 overexpressing solid tumors. The intervention involves the use of CT-0508 and Pembrolizumab, which are biological agents aimed at targeting and treating these specific tumors. Currently in Phase 1 and marked as active but not recruiting, this trial is focused on evaluating adverse events in the enrolled participants. Eligibility criteria include the requirement for tumors to overexpress the HER2 biomarker.

NCT: NCT04660929 · Enrollment: 48 · Sites: Chapel Hill, Duarte, Houston, Nashville, Philadelphia

MEDIUM PHASE1 Stage IV RAS TARGETED THERAPY • RECRUITING 4

Study of CRX100 as Monotherapy and in Combination With Pembrolizumab in Patients With Advanced Solid Malignancies

NCT: NCT04282044 · Enrollment: 60 · Sites: La Jolla, Scottsdale, Stanford

MEDIUM NA All Stages MSI/MMR TARGETED THERAPY • RECRUITING 4

Watch and Wait in PD-1 Monoclonal Antibody Treated dMMR/MSI-H Distal Rectal Cancer

NCT: NCT04643041 · Enrollment: 47 · Sites: Guangzhou, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

PD-1 Inhibitors Combined With VEGF Inhibitors for Locally Advanced dMMR/MSI-H Colorectal Cancer

This clinical trial is testing the combination of PD-1 inhibitors and VEGF inhibitors in patients with locally advanced dMMR/MSI-H colorectal cancer. The intervention involves the use of Camrelizumab (a PD-1 inhibitor) and Apatinib (a VEGF inhibitor) to assess their effectiveness in achieving clinical or pathological complete response. This is a Phase II study that is currently active but not recruiting, meaning that while the trial is ongoing, no new participants are being enrolled at this time. Eligible patients must have locally advanced colorectal cancer with specific biomarkers, including dMMR, MSI-H, and RAS mutations.

NCT: NCT04715633 · Enrollment: 53 · Sites: Guangzhou, China

MEDIUM PHASE1, PHASE2 Stage IV IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 4

Safety, PK and Efficacy of ONC-392 in Monotherapy and in Combination of Anti-PD-1 in Advanced Solid Tumors and NSCLC

NCT: NCT04140526 · Enrollment: 733 · Sites: Ann Arbor, Atlanta, Atlantis, Aurora, Baltimore

MEDIUM PHASE2 All Stages HER2 TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Tucatinib Plus Trastuzumab and Oxaliplatin-based Chemotherapy or Pembrolizumab-containing Combinations for HER2+ Gastrointestinal Cancers

NCT: NCT04430738 · Enrollment: 40 · Sites: Albuquerque, Aurora, City of Saint Peters, Cleveland, Creve Coeur

SIGNAL NA All Stages LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 4

Circulating DNA to Improve Outcome of Oncology PatiEnt. A Randomized Study

This clinical trial is testing the effectiveness of a liquid biopsy intervention in improving outcomes for patients with colorectal cancer. The intervention involves using genetic analysis of circulating tumor DNA (ctDNA) to guide treatment decisions and monitor disease progression. Currently, the trial is in an active status but is not recruiting new participants, indicating that it is ongoing and data is being collected from enrolled patients. Eligibility criteria include being diagnosed with colorectal cancer and participating in a randomized study comparing two different follow-up approaches based on ctDNA analysis versus standard imaging.

NCT: NCT04258137 · Enrollment: 332 · Sites: Bayonne, France, Bordeaux, France, Brest, France, Pau, France

MEDIUM PHASE2 All Stages HER2 TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

A Phase 2 Study of T-DXd in Patients With Selected HER2 Expressing Tumors

NCT: NCT04482309 · Enrollment: 477 · Sites: Boston, Chicago, Duarte, Durham, Harrison

MEDIUM PHASE1, PHASE2 All Stages MSI/MMR TARGETED THERAPY • RECRUITING 4

Sintilimab Plus Hypofractionated Radiotherapy for MSI-H/dMMR Rectal Cancer

This clinical trial is testing the combination of Sintilimab and hypofractionated radiotherapy for patients with microsatellite instability-high (MSI-H) or DNA mismatch repair-deficient (dMMR) non-metastatic rectal cancer. The intervention involves the use of Sintilimab, a drug that may enhance the effects of radiotherapy, which is important for improving treatment

outcomes in this specific patient population. Currently, the trial is in Phase 1/Phase 2 and is actively recruiting participants, indicating that eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite status.

NCT: NCT04636008 · Enrollment: 20 · Sites: Chengdu, China

MEDIUM PHASE1 Stage IV KRAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

A Study to Evaluate the Safety, Pharmacokinetics, and Activity of GDC-6036 Alone or in Combination in Participants With Advanced or Metastatic Solid Tumors With a KRAS G12C Mutation

This clinical trial is evaluating the safety, pharmacokinetics, and preliminary activity of GDC-6036, both alone and in combination with other therapies, in participants with advanced or metastatic solid tumors that have a KRAS G12C mutation. The intervention includes GDC-6036 and additional drugs such as Atezolizumab, Cetuximab, Bevacizumab, and Erlotinib, which are being tested to understand their effects on this specific mutation. Currently, the trial is in Phase I and is active but not recruiting, indicating that it is ongoing but not accepting new participants at this time. Eligibility criteria include having a confirmed KRAS G12C mutation, which is essential for participation in the study.

NCT: NCT04449874 · Enrollment: 498 · Sites: Boston, Duarte, La Jolla, New Haven, Oklahoma City

SIGNAL NA Stage III ctDNA LIQUID BIOPSY • RECRUITING 4

Circulating Tumor DNA Based Adjuvant Chemotherapy in Stage II Colon Cancer Patients: the MEDOCC-CrEATE Trial

The MEDOCC-CrEATE trial is testing the effectiveness of circulating tumor DNA (ctDNA) analysis to guide adjuvant chemotherapy decisions in patients with stage II colon cancer. The intervention involves analyzing blood samples for ctDNA after surgery to determine whether patients should receive chemotherapy, which may impact treatment decisions for those not initially indicated for adjuvant therapy. This trial is currently in the recruiting phase, meaning eligible patients can still enroll and participate in the study. Participants must be part of the Prospective Dutch ColoRectal Cancer cohort and provide informed consent for biobanking and future studies.

NCT: NCT06434896 · Enrollment: 1320 · Sites: 's-Hertogenbosch, Netherlands, Alkmaar, Netherlands, Almelo, Netherlands, Almere Stad, Netherlands, Amersfoort, Netherlands

MEDIUM PHASE2 Stage IV RAS PHASE II TRIAL • ACTIVE_NOT_RECRUITING 4

Radiochemotherapy +/- Durvalumab for Locally-advanced Anal Carcinoma. A Multicenter, Randomized, Phase II Trial of the German Anal Cancer Study Group

NCT: NCT04230759 · Enrollment: 180 · Sites: Berlin, Germany, Darmstadt, Germany, Dresden, Germany, Essen, Germany, Frankfurt, Germany

SIGNAL NA Stage III ctDNA LIQUID BIOPSY • ACTIVE_NOT_RECRUITING 4

ctDNA-guided Surveillance for Stage III CRC, a Randomized Intervention Trial

NCT: NCT04084249 · Enrollment: 359 · Sites: Aalborg, Denmark, Aarhus, Denmark, Copenhagen, Denmark, Herlev, Denmark, Herning, Denmark

MEDIUM NA All Stages TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

An Integrated Genomic Approach to Assess Genetic Risk and Drug Sensitivity

NCT: NCT07555639 · Enrollment: 4000 · Sites: Roma, Italy

MEDIUM PHASE1 Stage IV RAS TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Naptumomab Estafenatox in Combination With Durvalumab in Subjects With Selected Advanced or Metastatic Solid Tumor, Including a Cohort Expansion in Esophageal Cancer.

NCT: NCT03983954 · Enrollment: 120 · Sites: Ahmedabad, India, Haifa, Israel, Hyderabad, India, Jhajjar, India, New Delhi, India

MEDIUM PHASE1 Stage IV MSI/MMR IMMUNOTHERAPY • RECRUITING 4

Testing an Immunotherapy Anti-cancer Drug, Nivolumab, for Advanced Cancers in Patients With Autoimmune Disorders, AIM-NIVO

NCT: NCT03816345 · Enrollment: 300 · Sites: Atlanta, Baltimore, Bethesda, Birmingham, Boston

MEDIUM PHASE2 Stage IV PHASE II TRIAL • ACTIVE_NOT_RECRUITING 4

Microwave Ablation Versus Stereotactic Body Radiotherapy for Colorectal Liver Metastases in Oligometastatic Disease: a Prospective, Randomised, Phase 2 Trial

The LAVA-CRLM trial is testing the effectiveness of microwave ablation (MWA) compared to stereotactic body radiotherapy (SBRT) in patients with colorectal liver metastases and oligometastatic disease. The intervention involves using MWA, a device-based treatment, and SBRT, a form of radiation therapy, to assess their impact on local control and safety. This is a Phase 2 trial that is currently active but not recruiting new participants, indicating that the study is ongoing and data is being collected from enrolled patients. Eligibility criteria for participation may include specific characteristics related to the patient's metastatic disease status.

NCT: NCT03654131 · Enrollment: 100 · Sites: Copenhagen, Denmark

MEDIUM PHASE1, PHASE2 Stage IV NTRK TARGETED THERAPY • RECRUITING 4

A Study of Repotrectinib (TPX-0005) in Patients With Advanced Solid Tumors Harboring ALK, ROS1, or NTRK1-3 Rearrangements

This clinical trial is investigating the efficacy and safety of repotrectinib (TPX-0005) in patients with advanced solid tumors that have ALK, ROS1, or NTRK1-3 gene rearrangements. The intervention involves oral administration of repotrectinib, which is significant for targeting specific genetic alterations in tumors. Currently, the trial is in Phase 1 and Phase 2 stages and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include the presence of specific gene rearrangements, as determined by next-generation sequencing (NGS) or other biomarker assessments.

NCT: NCT03093116 · Enrollment: 500 · Sites: Ann Arbor, Appleton, Athens, Aurora, Baltimore

MEDIUM PHASE2 All Stages RAS IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 4

Nivolumab and Ipilimumab in Treating Patients With Rare Tumors

This phase II clinical trial is investigating the efficacy of nivolumab and ipilimumab in treating patients with rare tumors. The intervention involves the use of these monoclonal antibodies, which may enhance the immune system's ability to combat cancer and inhibit tumor growth and spread. Currently, the trial is active but not recruiting new participants, indicating that it is ongoing with enrolled patients being monitored for outcomes. Eligibility criteria include specific tumor types, such as epithelial tumors of the nasal cavity and salivary glands, with a focus on RAS biomarkers.

NCT: NCT02834013 · Enrollment: 798 · Sites: Aberdeen, Adrian, Aitkin, Albuquerque, Allentown

MEDIUM PHASE2 All Stages NTRK TARGETED THERAPY • ACTIVE_NOT_RECRUITING 4

Basket Study of Entrectinib (RXDX-101) for the Treatment of Patients With Solid Tumors Harboring NTRK 1/2/3 (Trk A/B/C), ROS1, or ALK Gene Rearrangements (Fusions)

NCT: NCT02568267 · Enrollment: 534 · Sites: Athens, Atlanta, Aurora, Baltimore, Boston

MEDIUM PHASE2 Stage IV RAS IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING 4

Nivolumab With or Without Ipilimumab in Treating Patients With Refractory Metastatic Anal Canal Cancer

This phase II clinical trial is investigating the effectiveness of nivolumab, with or without ipilimumab, in patients with refractory metastatic anal canal cancer. The intervention involves the use of these monoclonal antibodies, which may enhance the immune system's ability to target and eliminate cancer cells. Currently, the trial is active but not recruiting new participants, indicating that it is ongoing with the existing enrolled population of 143 patients. Eligibility criteria include the presence of specific biomarkers, such as RAS, which may be relevant for patient selection.

NCT: NCT02314169 · Enrollment: 143 · Sites: Ann Arbor, Aurora, Bellevue, Buffalo, Chicago

SIGNAL NA All Stages RESEARCH ● RECRUITING 4

Association Between Health Care Provider (HCP)-Assessed ECOG Performance Status (PS) and Overall Survival, and Objectively Measure of Physical Activity (PA) Levels in Advance-cancer Patients"

This clinical trial is testing the association between health care provider-assessed ECOG performance status and objectively measured physical activity levels in patients with advanced cancer. The intervention includes a behavioral exercise program and multiple health telemonitoring components, which aim to enhance the understanding of how physical activity impacts performance status and overall survival. Currently in the recruiting phase, this study seeks to enroll 590 participants, which means eligible patients can join to contribute to this important research. Specific eligibility criteria have not been detailed, but participation may involve assessments related to physical activity and performance status.

NCT: NCT01365169 · Enrollment: 590 · Sites: Houston

Appendix A — Patient Guide to Active Trials

For patients, caregivers, and family members. Always discuss with your oncologist before considering a clinical trial.

What is a clinical trial?

Clinical trials test new treatments, combinations, or approaches to care. Participating can give you access to treatments not yet widely available. Trials are listed by the biomarker your tumor must have — because many newer treatments only work for specific mutations. Ask your oncologist which biomarkers your tumor has been tested for, then look for matching trials below.

MSI/MMR

Anti-PCSK9 Antibody Tafolecimab and Anti-PD-1 Antibody Sintilimab Combined With Neoadjuvant Chemoradiotherapy for pMMR/ MSS Locally Advanced Rectal Cancer : A Prospective, Multicenter, Randomized, Open-Label, Parallel-Controlled Trial

Phase: PHASE3 · Status: NOT_YET_RECRUITING · NCT07686796

KRAS

Study of Petosemtamab Plus Chemotherapy Versus Cetuximab Plus Chemotherapy in RAS and BRAF Wild-type, Unresectable or Metastatic, Left-sided Colorectal Cancer (LiGeR-CRC1)

Phase: PHASE3 · Status: RECRUITING · NCT07702032

BRAF

JSKN003 Versus Physician Choiced Treatment in Patients With HER2-positive and Advanced Colorectal Cancer Who Had Failed to Respond to Oxaliplatin, 5-Fu, and Irinotecan

Phase: PHASE3 · Status: NOT_YET_RECRUITING · NCT07384377

ctDNA

ctDNA-guided Selection of Adjuvant Chemotherapy Regimens for Elderly Colon Cancer Patients After Surgery: A Single-center, Randomized, Controlled Study

Phase: PHASE3 · Status: RECRUITING · NCT06609551

This clinical trial is testing the effectiveness of ctDNA-guided selection of adjuvant chemotherapy regimens in elderly patients with high-risk stage II and stage III colon cancer following surgery. Participants will receive either 6 months of 5-FU monotherapy or an intensive XELOX treatment regimen, which is significant for tailoring chemotherapy based on ctDNA detection. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into disease-free survival outcomes for eligible patients. Eligibility criteria include being an elderly patient with high-risk stage II or stage III colon cancer and undergoing ctDNA testing post-surgery.

NGS

A Phase III Trial to Evaluate the Efficacy, Immunogenicity and Safety of a Recombinant Human Papillomavirus 9-Valent (Types 6, 11, 16, 18, 31, 33, 45, 52, 58) Vaccine (E.Coli) in Chinese Males

Phase: PHASE3 · Status: ACTIVE_NOT_RECRUITING · NCT06866574

HER2

A Study of Tucatinib With Trastuzumab and mFOLFOX6 Versus Standard of Care Treatment in First-line HER2+ Metastatic Colorectal Cancer

Phase: PHASE3 · Status: RECRUITING · NCT05253651

RAS

EA2176: Phase 3 Clinical Trial of Carboplatin and Paclitaxel +/- Nivolumab in Metastatic Anal Cancer Patients

Phase: PHASE3 · Status: ACTIVE_NOT_RECRUITING · NCT04444921

The EA2176 trial is testing the effectiveness of adding nivolumab to a chemotherapy regimen of carboplatin and paclitaxel in patients with metastatic anal cancer. The intervention involves the use of nivolumab, an immunotherapy drug, alongside standard chemotherapy, which may enhance the immune response against cancer cells while also targeting tumor growth through traditional chemotherapy mechanisms. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients are not currently being enrolled. Eligibility for participation includes specific biomarker requirements related to RAS.

NTRK

A Study of Repotrectinib (TPX-0005) in Patients With Advanced Solid Tumors Harboring ALK, ROS1, or NTRK1-3 Rearrangements

Phase: PHASE1, PHASE2 · Status: RECRUITING · NCT03093116

This clinical trial is investigating the efficacy and safety of repotrectinib (TPX-0005) in patients with advanced solid tumors that have ALK, ROS1, or NTRK1-3 gene rearrangements. The intervention involves oral administration of repotrectinib, which is significant for targeting specific genetic alterations in tumors. Currently, the trial is in Phase 1 and Phase 2 stages and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include the presence of specific gene rearrangements, as determined by next-generation sequencing (NGS) or other biomarker assessments.

What this means for you

Ask your oncologist whether any active trials match your biomarker profile and disease stage. The most actively trialed biomarkers in colorectal cancer right now are **MSI/MMR, KRAS, BRAF, ctDNA, and NGS**. If your tumor has not been tested for these markers, ask your oncologist before your next treatment decision — biomarker results can determine which trials you qualify for.

Questions to ask your oncologist about trials

"Am I eligible for any active Phase III trials based on my biomarker profile?"
"Does my tumor need to be retested before I can qualify for a biomarker-directed trial?"
"What are the travel requirements and visit frequency for the trials I might qualify for?"

Source: ClinicalTrials.gov · 2026-11-09 07:30:00
This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Clinical Trial Reference Guide

What the phases mean

Phase I — First testing in humans. Focus on safety and dosing. Small groups.

Phase II — Testing effectiveness. Larger groups. Determines if Phase III is warranted.

Phase III — Randomized comparison against standard treatment. Required for FDA approval.

Key biomarkers that determine trial eligibility

RAS (KRAS/NRAS) — mutation status affects anti-EGFR eligibility and KRAS G12C trial eligibility

BRAF V600E — required for encorafenib-based trials (~8–10% of CRC)

MSI-H/dMMR — required for immunotherapy trials; ~5% of mCRC, ~15% of Stage II/III

HER2 — required for HER2-directed trials (~2–3% of CRC)

NTRK fusion — required for TRK inhibitor trials (<1% of CRC)

KRAS G12C — required for sotorasib/adagrasib trials (~3–4% of CRC)

TMB-H — tumor mutational burden; may qualify for immunotherapy

ctDNA — liquid biopsy; used in MRD monitoring trials

How to find trials you may qualify for

1. Get complete biomarker testing: RAS, BRAF, MSI/MMR, HER2, NTRK, TMB

2. Search ClinicalTrials.gov with your biomarker and stage

3. Ask your oncologist about NCI-designated cancer centers with trial access

4. Contact the Colorectal Cancer Alliance: ccalliance.org

Key Resources

ClinicalTrials.gov · NCI Cancer Information (cancer.gov) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org)

Colon & Colorectal Cancer — Clinical Trials Intelligence

About This Report

This report is generated automatically from publicly available medical databases including PubMed, ClinicalTrials.gov, and the FDA Drug@FDA database. Entries are classified using deterministic rule-based logic and scored by clinical significance. AI-generated summaries are grounded exclusively in the source data provided.

Limitations

This report does not represent a complete picture of all published research or approved treatments. Database coverage, feed availability, and relevance filtering affect what appears. New drug approvals and trial status changes may not be reflected immediately. Always verify current information at the original source before making clinical decisions.

Disclaimer

Trial data is sourced from ClinicalTrials.gov, a registry maintained by the U.S. National Library of Medicine. Entry summaries are AI-generated from publicly available trial records. Trial status and eligibility criteria may change — always verify at ClinicalTrials.gov before acting on this information. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

Copyright

© 2026 Randy Hinton. All rights reserved. This report may be shared freely for informational purposes. It may not be modified, republished, or presented as original work without attribution. ·
https://petroai1492.github.io/delta/timelines/coloncancer_clinicaltrials_timeline.pdf