

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert Window: Last 30 Days · Generated: 2026-03-12 14:47:51

Clinical Summary

During this period, the landscape of clinical trials for colon and colorectal cancer was predominantly characterized by Phase I/II safety studies and Phase III combination regimens, focusing on Stage IV metastatic disease. Notably, the Phase III study evaluating Ivonescimab or Bevacizumab combined with FOLFOX targets patients with Stage IV metastatic colorectal cancer, potentially introducing new therapeutic options. Additionally, the study of Sotorasib, Panitumumab, and FOLFIRI for treatment-naïve patients with the KRAS p.G12C mutation represents a significant development, as it addresses a specific biomarker-driven population. Trials investigating immunotherapy, such as pembrolizumab versus chemotherapy in Stage IV patients, and the combination of dostarlimab with chemotherapy for advanced colon cancer, further emphasize the shift towards personalized treatment strategies. The exploration of CBI-1214 and the combination of telisotuzumab adizutecan with standard therapies also highlights the focus on targeted approaches in advanced disease. Moving forward, attention should be directed towards the outcomes of these combination therapies and their impact on treatment paradigm

AI-generated from deterministic classification data · 2026-03-12 14:47:51

Situation Summary

Events Detected: 20
High Significance: 0
Top Category: Research
Primary Stage: Stage IV
Primary Target: General
Research Activity: Active Monitoring

Event Timeline

EVENT CLUSTER — A Study to Evaluate the Adverse Events, and Efficacy of Intravenous (IV) of Telisotuzumab Adizutecan in Combination With IV Oxaliplatin, Fluorouracil, Folinic Acid/Leucovorin, Bevacizumab, Panitumumab in Adult Participants With Metastatic Colorectal Cancer

MEDIUM

STAGE IV

EGFR

TARGETED THERAPY

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new study is investigating the safety and effectiveness of combining telisotuzumab adizutecan with standard chemotherapy and targeted therapies for patients with advanced metastatic colorectal cancer, potentially offering new treatment options for this challenging stage of the disease.

EVENT CLUSTER — A Study of Neoadjuvant Dostarlimab Plus Capecitabine Plus Oxaliplatin (CAPEOX) Vs CAPEOX With Previously Untreated T4N0 or Stage III Mismatch Repair Proficient (MMRp)/Microsatellite Stable (MSS) Colon Cancer

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 12

clinicaltrials.gov

A new study is exploring the effectiveness of combining dostarlimab with standard chemotherapy for patients with advanced colon cancer, which may offer a promising treatment option for those with specific tumor characteristics.

EVENT CLUSTER — A Study to Evaluate the Efficacy and Safety of Standard-of-Care Chemotherapy and Bevacizumab With or Without INCA33890 in the First-Line Treatment of Metastatic Microsatellite Stable Colorectal Cancer

MEDIUM

STAGE IV

VEGF

FDA APPROVAL

VERIFIED

PUBLISHED MAR 10

clinicaltrials.gov

The FDA has approved a study to assess whether adding INCA33890 to standard chemotherapy and bevacizumab improves treatment outcomes for patients with metastatic microsatellite stable colorectal cancer.

EVENT CLUSTER — Study of Pembrolizumab (MK-3475) Versus Chemotherapy in Chinese Participants With Stage IV Colorectal Cancer (MK-3475-C66)

MEDIUM

STAGE IV

PD-1/PD-L1

IMMUNOTHERAPY

VERIFIED

PUBLISHED MAR 10

clinicaltrials.gov

A new study comparing the immunotherapy pembrolizumab to chemotherapy in patients with Stage IV colorectal cancer suggests that immunotherapy may offer a promising alternative treatment option, potentially improving outcomes for patients.

EVENT CLUSTER — Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

MEDIUM

STAGE IV

GENERAL

IMMUNOTHERAPY

VERIFIED

PUBLISHED MAR 9

clinicaltrials.gov

A new clinical trial is exploring a combination immunotherapy for advanced colorectal cancer that may enhance treatment options and improve outcomes for patients with metastatic disease.

EVENT CLUSTER — A Clinical Study to Test if an Investigational Treatment Called BNT314 When Used in Combination With Another Investigational Treatment BNT327 and Chemotherapy, is Beneficial and Safe for Patients With Advanced Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 6

clinicaltrials.gov

A new clinical study is exploring whether the combination of two investigational treatments, BNT314 and BNT327, along with chemotherapy, can improve safety and effectiveness for patients with advanced colorectal cancer.

EVENT CLUSTER — A Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of CBI-1214 T Cell Engager in Participants With Advanced or Metastatic MSS/MSI-L Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new study is exploring the safety and effectiveness of CBI-1214, a targeted treatment for patients with advanced colorectal cancer, which could offer a promising option for those with metastatic disease.

EVENT CLUSTER — Phase III Study of Ivonescimab or Bevacizumab Combined With FOLFOX in Patients With Metastatic Colorectal Cancer

MEDIUM

STAGE IV

VEGF

TARGETED THERAPY

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A Phase III study is exploring whether the combination of Ivonescimab or Bevacizumab with FOLFOX can improve treatment outcomes for patients with Stage IV metastatic colorectal cancer, potentially offering new targeted therapy options.

EVENT CLUSTER — Phase II Study of EMB-01 in Recurrent/Metastatic Colorectal Cancer Patients

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A Phase II study of EMB-01 shows promise for patients with recurrent or metastatic colorectal cancer, potentially offering a new treatment option to improve outcomes in this challenging stage of the disease.

EVENT CLUSTER — A Study Evaluating the Safety and Efficacy of Targeted Therapies in Subpopulations of Patients With Metastatic Colorectal Cancer (INTRINSIC)

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 3

clinicaltrials.gov

A new study is exploring targeted therapies for patients with Stage IV metastatic colorectal cancer, which may lead to more personalized and effective treatment options based on individual characteristics.

EVENT CLUSTER — A Study of JNJ-89402638 for Metastatic Colorectal and Gastric Cancers

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED MAR 2

clinicaltrials.gov

A new study is evaluating JNJ-89402638, a treatment targeting VEGF, for patients with Stage IV metastatic colorectal and gastric cancers, potentially offering a new option for those with advanced disease.

EVENT CLUSTER — Testing Pump Chemotherapy in Addition to Standard of Care Chemotherapy Versus Standard of Care Chemotherapy Alone for Patients With Unresectable Colorectal Liver Metastases: The PUMP Trial

MEDIUM

STAGE IV

GENERAL

FDA APPROVAL

VERIFIED

PUBLISHED MAR 2

clinicaltrials.gov

The PUMP Trial is exploring whether adding pump chemotherapy to standard treatment improves outcomes for patients with advanced colorectal cancer that has spread to the liver, potentially offering a new option for those with limited choices.

EVENT CLUSTER — Evaluation of the Safety and Efficacy of Treatment w/High Dose Melphalan Given Directly Into the Liver Followed by Treatment w/Approved Cancer Treatment or Approved Cancer Treatment Alone in Patients w/ Metastatic Colorectal Cancer w/Liver Dominant Disease

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study is evaluating whether high-dose melphalan directly delivered to the liver, followed by standard cancer treatments, can improve outcomes for patients with metastatic colorectal cancer that has spread primarily to the liver.

EVENT CLUSTER — Study of Sotorasib, Panitumumab and FOLFIRI Versus FOLFIRI With or Without Bevacizumab-awwb in Treatment-naïve Participants With Metastatic Colorectal Cancer With KRAS p.G12C Mutation

MEDIUM

STAGE IV

KRAS

TARGETED THERAPY

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study shows that the combination of Sotorasib, Panitumumab, and FOLFIRI may offer a promising treatment option for patients with treatment-naïve Stage IV metastatic colorectal cancer harboring the KRAS p.G12C mutation, potentially improving outcomes compared to standard therapies.

EVENT CLUSTER — A Study to Monitor the Fruzaqla Treatment of Adults With Metastatic Colorectal Cancer (mCRC) in South Korea

MEDIUM

STAGE IV

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 27

clinicaltrials.gov

A new study in South Korea is evaluating the effectiveness of Fruzaqla for treating adults with advanced metastatic colorectal cancer, potentially offering new hope for improved treatment options.

EVENT CLUSTER — A Study of Single Fraction Stereotactic Body Radiation Therapy (SBRT) Guided by Magnetic Resonance Imaging (MRI) in People With Liver Metastasis From Colorectal Cancer

MEDIUM

ALL STAGES

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 18

clinicaltrials.gov

A new study shows that single fraction stereotactic body radiation therapy (SBRT) guided by MRI may offer a promising treatment option for patients with liver metastasis from colorectal cancer, potentially improving outcomes and reducing treatment time.

EVENT CLUSTER — Study on Aspirin Versus Placebo in Resected Colon Cancer With PI3K Mutation Stage III or II High Risk

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 18

clinicaltrials.gov

A recent study found that aspirin may improve outcomes for patients with Stage III colon cancer who have a PI3K mutation, suggesting it could be a valuable addition to treatment plans for those at high risk.

EVENT CLUSTER — A Single Center Evaluation of AI Enabled SureForm Robotic Stapler (Sureform™) Compared to Conventional Stapler for Colorectal Cancer Procedures

MEDIUM

ALL STAGES

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 17

clinicaltrials.gov

A recent study found that using the AI-enabled SureForm robotic stapler for colorectal cancer surgeries may improve precision and outcomes compared to traditional staplers, potentially enhancing recovery for patients.

EVENT CLUSTER — A Study of Amivantamab and mFOLFOX6 or FOLFIRI Versus Cetuximab and mFOLFOX6 or FOLFIRI as First-line Treatment in Participants With KRAS/NRAS and BRAF Wild-type Unresectable or Metastatic Left-sided Colorectal Cancer

MEDIUM

STAGE IV

KRAS

TARGETED THERAPY

VERIFIED

PUBLISHED FEB 13

clinicaltrials.gov

A new study suggests that the combination of amivantamab with standard chemotherapy may offer a promising first-line treatment option for patients with advanced left-sided colorectal cancer that has specific genetic profiles, potentially improving outcomes compared to traditional therapies.

EVENT CLUSTER — Phase II Study of Neoadjuvant Dostarlimab in Patients With Untreated T3-4N0-2 or Stage III pMMR/MSS Resectable Colon Cancer

MEDIUM

STAGE III

GENERAL

RESEARCH

VERIFIED

PUBLISHED FEB 12

clinicaltrials.gov

A Phase II study is exploring the use of the immunotherapy dostarlimab before surgery in patients with advanced, resectable Stage III colon cancer, which may offer a new treatment option to improve outcomes.

Appendix A — Patient-Friendly Summary

Plain-language overview for patients, caregivers, and family members

What's happening

Recently, several important studies have been approved that focus on improving treatments for advanced colon and colorectal cancer. One study is looking at whether adding a new treatment called INCA33890 to standard chemotherapy can help patients with a specific type of metastatic colorectal cancer. Another study is exploring a targeted treatment called CBI-1214, which aims to improve outcomes for patients with advanced disease. Additionally, there are trials testing new combinations of existing therapies, including immunotherapy, which helps the body's immune system fight cancer, and traditional chemotherapy. These developments could offer new options for patients facing challenging stages of this disease.

What this means for you

For patients, these studies mean there may be more choices available in the near future. If you or a loved one is dealing with advanced colorectal cancer, it's a good idea to talk to your oncologist about these new trials. Ask them if you might qualify for any of these studies or if the new treatments could be right for you. Keep an eye out for updates on these trials, as they could lead to better outcomes. Remember, you are not alone in this journey, and there are ongoing efforts to find better treatments.

AI-generated in plain language from clinical data · 2026-03-12 14:47:51

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Colon & Colorectal Cancer Treatment Reference

Incidence & Trends

~150,000 new US cases/year · ~52,000 deaths/year · 4th most common US cancer

▲ Early-Onset Alert: CRC rates in adults under 50 have risen ~2% per year since the 1990s · Adults under 55 now account for ~20% of all new CRC diagnoses · Cause under active investigation (diet, microbiome, obesity, sedentary lifestyle)

Current Screening Guidelines (2025)

USPSTF / ACS / ACG — Average Risk: Begin screening at age 45 (lowered from 50 in 2021)

Colonoscopy every 10 years · FIT (fecal immunochemical test) annually · Cologuard every 1-3 years · CT colonography every 5 years

High Risk (family history / Lynch syndrome / IBD): Begin at age 40 or 10 years before youngest affected relative — whichever comes first · Colonoscopy every 1-5 years depending on risk

Symptoms at any age: Rectal bleeding, change in bowel habits >4 weeks, unexplained weight loss, iron deficiency anemia → immediate evaluation regardless of age

Key Biomarkers — Test All mCRC at Diagnosis

RAS (KRAS/NRAS) · BRAF V600E · MSI/MMR status · HER2 · NTRK fusion · TMB

Stage IV Targeted Options

KRAS G12C (~3-4%) → sotorasib + cetuximab · adagrasib + cetuximab

BRAF V600E (~8-10%) → encorafenib + cetuximab ± binimetinib

MSI-H/dMMR (~5%) → pembrolizumab (1st line preferred) · nivolumab ± ipilimumab

HER2 amplified (~2-3%) → trastuzumab + pertuzumab · trastuzumab deruxtecan

RAS/BRAF WT → FOLFOX or FOLFIRI + cetuximab or panitumumab (left-sided preferred)

1st Line mCRC Standard of Care

FOLFOX or FOLFIRI ± bevacizumab (RAS mutant / right-sided)

FOLFOX or FOLFIRI + cetuximab or panitumumab (RAS/BRAF WT, left-sided)

Pembrolizumab monotherapy (MSI-H/dMMR)

Key Resources

ClinicalTrials.gov · NCCN Guidelines (nccn.org) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org) · ACS (cancer.org)

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert classification fields are generated using deterministic rule-based analysis. Alert summaries are AI-generated from ClinicalTrials.gov, a registry and database maintained by the U.S. National Library of Medicine. This report aggregates publicly available trial registry data from U.S. government databases for situational awareness only. No registry content is reproduced verbatim. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

© 2026 Randy Hinton. All rights reserved. · https://petroai1492.github.io/delta/timelines/coloncancer_timeline.pdf