

Colon & Colorectal Cancer — Clinical Trials Intelligence

Alert Window: Last 90 Days · Generated: 2026-04-25 23:31:19

Situation Summary

Phase I/II: 48 trials
Recruiting: 254 trials
Biomarker: 207 biomarker-directed trials
Breakthrough: 106 trials scored ≥ 4

Active Trial Timeline

Showing 106 high-significance trials (score ≥ 4). 319 additional lower-scored trials are collapsed below. Phase III trials appear first. Click any title to view full trial details on ClinicalTrials.gov.

MEDIUM**PHASE3****Stage IV****MSI/MMR****TARGETED THERAPY**• **RECRUITING****11**

Botensilimab + Balstilimab vs Best Supportive Care as Therapy in Chemo-refractory, Unresectable, Colorectal Adenocarcinoma

This clinical trial is testing the combination of botensilimab and balstilimab in patients with chemo-refractory, unresectable colorectal adenocarcinoma. The intervention involves two immunotherapy drugs aimed at potentially improving overall survival and quality of life for participants. Currently in Phase 3 and actively recruiting, this trial may provide insights into the effectiveness and safety of these treatments for eligible patients. Participants may need to meet specific biomarker criteria, including MSI, MSI-H, MMR, dMMR, and microsatellite status.

NCT: NCT07152821 · Enrollment: 834 · Sites: Greenfield Park, Canada, Kingston, Canada, Oshawa, Canada, Ottawa, Canada, Prince George, Canada

MEDIUM**PHASE3****Stage III****MSI/MMR****TARGETED THERAPY**• **NOT_YET_RECRUITING****11**

Neoadjuvant/Adjuvant AK104 in Microsatellite Instability-high or Mismatch Repair-deficient, Resectable Colon Cancer

This clinical trial is testing the efficacy and safety of neoadjuvant/adjuvant treatment with AK104 (Cadonilimab) in patients with resectable microsatellite instability-high (MSI-H) or mismatch repair-deficient (dMMR) colon cancer. The intervention involves AK104 in combination with standard chemotherapy agents, including Oxaliplatin, Capecitabine, 5-Fluorouracil, and Calcium Folate, which is significant for potentially improving treatment outcomes in this specific patient population. This is a Phase 3 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include having resectable colon cancer with specific biomarkers such as MSI-H or dMMR.

NCT: NCT07412613 · Enrollment: 386 · Sites: Guangzhou, China

MEDIUM**PHASE3****Stage IV****BRAF****TARGETED THERAPY**• **RECRUITING****11**

Symbiotic-GI-03: A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Chemotherapy in Adult Participants With Metastatic Colorectal Cancer

The Symbiotic-GI-03 trial is investigating the efficacy of the study medicine PF-08634404 in combination with chemotherapy for adult participants with metastatic colorectal cancer. This trial aims to determine how well PF-08634404 works alongside standard chemotherapy and Bevacizumab, a biological therapy, in patients whose cancer has spread or

returned after previous treatments. Currently in Phase 3 and actively recruiting, this trial offers eligible patients the opportunity to receive potentially beneficial treatment options while contributing to research on colorectal cancer therapies. To participate, individuals must be 18 years or older, have metastatic colorectal cancer, and meet specific health criteria, including not being pregnant prior to treatment.

NCT: NCT07222800 · Enrollment: 800 · Sites: Aberdeen, Abilene, Albany, Amarillo, Arlington

MEDIUM

PHASE2, PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Node-Sparing Short-Course Radiotherapy Plus Chemotherapy, Bevacizumab and PD-1 Inhibitor in Metastatic pMMR/MSS Colorectal Cancer (MODIFI-CRC)

This clinical trial, known as MODIFI-CRC, is testing the effectiveness of node-sparing short-course radiotherapy combined with chemotherapy, bevacizumab, and a PD-1 inhibitor in patients with metastatic pMMR/MSS colorectal cancer. The intervention aims to enhance tumor response by utilizing radiotherapy to improve the efficacy of immunotherapy and targeted treatments, addressing the limitations of current standard therapies. Currently in Phase 2/3 and actively recruiting participants, this trial may offer patients an opportunity to receive a novel treatment approach that could potentially improve their outcomes. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and NGS.

NCT: NCT06958419 · Enrollment: 286 · Sites: Guangzhou, China

MEDIUM

PHASE3

All Stages

KRAS

TARGETED THERAPY

• RECRUITING

11

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

This clinical trial is testing the efficacy of amivantamab combined with FOLFIRI chemotherapy in patients with KRAS/NRAS and BRAF wild-type colorectal cancer who have previously undergone chemotherapy. The intervention involves the biological agent amivantamab alongside standard chemotherapy agents, which is important for evaluating its potential benefits compared to existing treatment options. This is a Phase 3 trial currently recruiting 700 participants, which means that eligible patients may have the opportunity to receive a treatment that is being compared to standard therapies in a larger population. Eligibility criteria include specific biomarker requirements related to KRAS, NRAS, and BRAF status, as well as other factors such as microsatellite instability (MSI) and mismatch repair (MMR) status.

NCT: NCT06750094 · Enrollment: 700 · Sites: Abilene, Ann Arbor, Arlington, Athens, Atlanta

MEDIUM

PHASE3

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

A Study of Neoadjuvant QL1706 in Participants With Untreated dMMR/MSI-H Resectable Colon Cancer

This clinical trial is testing the efficacy of neoadjuvant QL1706 in participants with untreated dMMR/MSI-H resectable colon cancer. The intervention involves administering QL1706 alongside CAPEOX/Capecitabine to assess its impact on tumor response prior to surgery. Currently in Phase 3 and actively recruiting, this trial aims to determine the proportion of patients achieving a pathological complete response (pCR), which indicates no residual tumor after treatment. Eligibility criteria include having T4N0 or Stage III colon cancer with specific biomarkers such as MSI-H or dMMR.

NCT: NCT06686576 · Enrollment: 363 · Sites: Guangzhou, China

MEDIUM

PHASE2, PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Sacituzumab Govitecan in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of Sacituzumab Govitecan (SG) in patients with metastatic colorectal cancer who have become refractory to at least two lines of standard chemotherapy and are not candidates for local therapy. The intervention involves administering SG compared to the physician's choice of treatment, which is significant as it may provide an alternative option for patients with limited treatment choices. This is a Phase II/III trial currently recruiting participants, indicating that patients may have the opportunity to access this investigational treatment while contributing to the research. Eligibility criteria include documented progression or intolerability to specific chemotherapy regimens and a minimum 6-month interval since the last use of Irinotecan.

NCT: NCT06243393 · Enrollment: 80 · Sites: Heidelberg, Germany

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

Injection of SHR-A1811 Versus Physician Chosed Treatment in Patients With Advanced Colorectal Cancer Who Had Failed to Respond to Oxaliplatin, 5-fu, and Irinotecan

This clinical trial is testing the efficacy and safety of SHR-A1811 in patients with advanced colorectal cancer who have not responded to previous treatments with oxaliplatin, 5-FU, and irinotecan. The intervention involves the drug SHR-A1811, which is being compared to physician-chosen treatments such as TAS-102, regorafenib, and fruquintinib. This is a Phase 3 trial that is currently active but not recruiting, meaning that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06199973 · Enrollment: 130 · Sites: Shanghai, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Preoperative Short-course Radiation Followed by Envafolimab Plus CAPEOX for MSS Locally Advanced Rectal Adenocarcinoma

This clinical trial is testing the combination of preoperative short-course radiation followed by Envafolimab and CAPEOX chemotherapy in patients with microsatellite stable (MSS) locally advanced rectal adenocarcinoma. The intervention involves the use of Envafolimab, a PD-L1 monoclonal antibody, which may enhance the effectiveness of short-course radiotherapy and chemotherapy. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to receive this investigational treatment while contributing to important research. Eligibility criteria include specific biomarker requirements such as MSI, MSI-H, MMR, DMMR, and RAS status.

NCT: NCT05752136 · Enrollment: 108 · Sites: Hangzhou, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Total Neoadjuvant Treatment ±Immunotherapy for High Risk Locally Advanced Rectal Cancer (TNTi)

The TNTi clinical trial is testing the effectiveness of total neoadjuvant treatment with and without the immunotherapy drug camrelizumab for patients with high-risk locally advanced rectal cancer. This intervention is significant as it aims to determine the pathologic complete response (PCR) rate, which is a critical measure of treatment effectiveness. Currently in Phase 3 and actively recruiting, this trial offers patients the opportunity to participate in research that may influence future treatment protocols. Eligibility criteria include specific biomarkers such as MSI, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT06229041 · Enrollment: 472 · Sites: Beijing, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

Study of XL092 + Atezolizumab vs Regorafenib in Participants With Metastatic Colorectal Cancer

This clinical trial is testing the combination of XL092 and atezolizumab against regorafenib in participants with metastatic colorectal cancer who have progressed during, after, or are intolerant to standard-of-care therapy. The intervention involves the use of XL092 and atezolizumab, which may provide a different therapeutic approach compared to regorafenib. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and may soon provide important data on treatment efficacy. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT05425940 · Enrollment: 901 · Sites: Albuquerque, Billings, Charlotte, Chattanooga, Cincinnati

MEDIUM

PHASE3

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

PACE: PD-1 Antibody For dMMR/MSI-H Stage III Colorectal Cancer

The PACE trial is investigating the efficacy of the PD-1 antibody Sintilimab in patients with dMMR/MSI-H Stage III colorectal cancer. Participants will receive either Sintilimab alone or a combination of oxaliplatin and capecitabine, which are standard chemotherapy agents, to assess their impact on disease-free survival. This is a Phase 3 study currently recruiting 323 participants, indicating that patients may have the opportunity to access this treatment while contributing to important clinical research. Eligibility criteria include having local advanced colon cancer with specific biomarkers such as dMMR or MSI-H.

NCT: NCT05236972 · Enrollment: 323 · Sites: Guangzhou, China

MEDIUM

PHASE3

Stage IV

BRAF

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

A Study of Encorafenib Plus Cetuximab With or Without Chemotherapy in People With Previously Untreated Metastatic Colorectal Cancer

This clinical trial is evaluating the combination of encorafenib and cetuximab, with or without standard chemotherapy, in patients with previously untreated metastatic colorectal cancer that has a BRAF mutation. The intervention involves administering encorafenib orally and cetuximab via intravenous infusion, either alone or alongside chemotherapy agents such as oxaliplatin and irinotecan. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients may not be able to enroll at this time. Eligibility criteria include the presence of specific biomarkers such as BRAF mutations, MSI, MSI-H, MMR, or DMMR.

NCT: NCT04607421 · Enrollment: 831 · Sites: Aventura, Basking Ridge, Berkeley Heights, Chicago, City of Saint Peters

MEDIUM

PHASE3

Stage IV

HER2

TARGETED THERAPY

• RECRUITING

⚡ 10

A Study of Tucatinib With Trastuzumab and mFOLFOX6 Versus Standard of Care Treatment in First-line HER2+ Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of tucatinib in combination with trastuzumab and mFOLFOX6 compared to standard of care treatments for patients with HER2 positive metastatic colorectal cancer. The intervention involves the use of tucatinib, a targeted therapy, alongside other established cancer drugs, to evaluate its effectiveness and side effects in this patient population. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into progression-free survival outcomes for participants. Eligibility criteria include having HER2 positive colorectal cancer that is metastatic or unresectable, with specific biomarker assessments for HER2 and RAS.

NCT: NCT05253651 · Enrollment: 400 · Sites: Anaheim, Atlanta, Aurora, Aventura, Baldwin Park

SIGNAL

PHASE3

Stage IV

MSI/MMR

RESEARCH

• RECRUITING

⚡ 9

Neoadjuvant CAPOX With or Without Pucotenlimab Plus Selective Radiotherapy for Locally Advanced Rectal Cancer

This clinical trial is testing the effectiveness of neoadjuvant CAPOX with or without pucotenlimab in patients with locally advanced rectal cancer. The intervention involves administering four cycles of CAPOX, a chemotherapy regimen, combined with the investigational drug pucotenlimab, which may enhance treatment outcomes. This is a phase III trial currently recruiting 556 participants, which means patients may have the opportunity to access this treatment approach while contributing to the evaluation of its efficacy and safety. Eligible patients must have pMMR/MSS locally advanced rectal cancer and will be randomized based on mesorectal fascia status.

NCT: NCT07528209 · Enrollment: 556 · Sites: Guangzhou, China

MEDIUM

PHASE3

All Stages

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 9

Study Evaluating Two Treatment Strategies for Rectal Cancer in Patients ≥75 Years Old

The NACRE-2 trial is evaluating two treatment strategies for patients aged 75 years and older with locally advanced rectal cancer. The study compares the effectiveness of adding chemotherapy (FOLFOX4s) after short-course radiotherapy against short-course radiotherapy alone. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as MSI, MMR, DMMR, and microsatellite status.

NCT: NCT07118800 · Enrollment: 160 ·

SIGNAL

PHASE3

Stage IV

RESEARCH

• RECRUITING

⚡ 9

AK112 and Chemotherapy in First-line Metastatic Colorectal Cancer

This Phase III clinical trial is evaluating the efficacy and safety of AK112 in combination with chemotherapy for patients with first-line metastatic colorectal cancer. The intervention includes AK112 along with standard chemotherapy agents such as oxaliplatin, irinotecan, leucovorin, and 5-FU, compared to bevacizumab with chemotherapy. Currently, the trial is in the recruiting phase, meaning eligible patients can still enroll to participate. Enrollment is set for 560 participants, and the primary outcome being assessed is progression-free survival as determined by blinded independent central review.

NCT: NCT06951503 · Enrollment: 560 · Sites: Guanzhou, China

SIGNAL

PHASE3

Stage IV

MSI/MMR

RESEARCH

• RECRUITING

⚡ 9

DOSAGE Study: Upfront Dose-Reduced Chemotherapy in Older Patients with Metastatic Colorectal Cancer

The DOSAGE Study is a phase III clinical trial designed to evaluate the effectiveness of upfront dose-reduced chemotherapy compared to standard dose chemotherapy in older patients (≥ 70 years) with metastatic colorectal cancer. The trial investigates various chemotherapy regimens, including doublet chemotherapy and monotherapy, both at standard and reduced doses, to assess their impact on progression-free survival (PFS). Currently, the study is in the recruiting phase, meaning eligible patients can still enroll to participate in this research. Eligibility criteria include age and risk classification for chemotherapy toxicity, with specific biomarkers such as MSI and microsatellite status being relevant for patient selection.

NCT: NCT06275958 · Enrollment: 587 · Sites: 's-Hertogenbosch, Netherlands, Alkmaar, Netherlands, Amstelveen, Netherlands, Amsterdam, Netherlands, Arnhem, Netherlands

SIGNAL

PHASE3

Stage IV

BRAF

RESEARCH

• RECRUITING

9

Intermittent or Continuous Panitumumab Plus FOLFIRI for Left Sided RAS/B-RAF Wild-type Metastatic Colorectal Cancer

This clinical trial is testing the effectiveness of intermittent versus continuous administration of Panitumumab combined with FOLFIRI in patients with left-sided RAS/B-RAF wild-type metastatic colorectal cancer. The intervention involves the use of Panitumumab, Irinotecan, 5-fluorouracil, and L-folinic acid, which are important for evaluating treatment strategies that may optimize patient outcomes. Currently in Phase 3 and actively recruiting, this trial aims to determine if the intermittent regimen can achieve a Time to Treatment Failure comparable to the standard continuous regimen, which is relevant for patients considering their treatment options. Eligibility criteria include specific biomarker requirements related to BRAF, MMR, and RAS status.

NCT: NCT06509126 · Enrollment: 500 · Sites: Naples, Italy

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

9

A Study To Evaluate The Efficacy And Safety Of Ifinatamab Deruxtecan (I-DXd) In Subjects With Recurrent Or Metastatic Solid Tumors (IDeate-PanTumor02)

This clinical trial is evaluating the efficacy and safety of ifinatamab deruxtecan (I-DXd) in patients with recurrent or metastatic solid tumors, including colorectal cancer. The intervention, ifinatamab deruxtecan, is a drug that targets specific cancer biomarkers, which may influence treatment outcomes. This is a Phase 2 trial currently recruiting 520 participants, indicating that patients may have the opportunity to access this treatment while contributing to the understanding of its effectiveness. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, HER2, and RAS.

NCT: NCT06330064 · Enrollment: 520 · Sites: Amarillo, Chattanooga, Dallas, Fairfax, Germantown

SIGNAL

PHASE3

Stage III

RESEARCH

• RECRUITING

9

Thymosin-alpha 1 for Adjuvant Treatment After Radical Resection of High-risk Stage II and III Colorectal Cancer

This clinical trial is testing the efficacy and safety of Thymosin-alpha 1 as an adjuvant treatment for patients with high-risk stage II and III colorectal cancer following radical resection. The intervention, Thymosin-alpha 1, is a drug believed to enhance immune response, potentially reducing the risk of cancer recurrence and metastasis. This is a Phase 3 trial currently recruiting 2,500 participants, which indicates that it is in the final stages of testing before potential approval and may provide valuable information for patients regarding treatment options. Eligibility criteria may include specific high-risk classifications of colorectal cancer, although detailed biomarker requirements are not specified.

NCT: NCT05086614 · Enrollment: 2500 · Sites: Shanghai, China

SIGNAL

PHASE3

Stage IV

RAS

RESEARCH

• ACTIVE_NOT_RECRUITING

9

Neoadjuvant FOLFOX Chemotherapy for Patients With Locally Advanced Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant FOLFOX chemotherapy in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy before surgery, which may enhance treatment outcomes by targeting microscopic metastatic cancer cells earlier and reducing surgical complications. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and results may soon be available for clinical application. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT03426904 · Enrollment: 708 · Sites: Daegu, South Korea, Hwasun, South Korea, Seoul, South Korea, Suwon, South Korea

SIGNAL **PHASE3** **Stage IV** **RESEARCH** • **RECRUITING**

⚡ 9

Bevacizumab Plus mFOLFOXIRI as First-line Treatment for Patients With Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of mFOLFOXIRI combined with Bevacizumab versus mFOLFOX6 combined with Bevacizumab as first-line treatments for patients with unresectable metastatic colorectal cancer. The intervention involves two chemotherapy regimens, which are significant as they aim to improve treatment outcomes while considering the unique responses of different populations, particularly in China. This is a Phase 3 trial currently recruiting 528 participants, which means eligible patients may have the opportunity to receive one of the investigational treatments while contributing to important research data. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

NCT: NCT04230187 · Enrollment: 528 · Sites: Guangzhou, China

SIGNAL **PHASE3** **Stage III** **RESEARCH** • **RECRUITING**

⚡ 9

Randomised Study Evaluating Adjuvant Chemotherapy After Resection of Stage III Colonic Adenocarcinoma in Patients of 70 and Over

This clinical trial is evaluating the effectiveness of adjuvant chemotherapy in patients aged 70 and over who have undergone resection for stage III colonic adenocarcinoma. The interventions being tested include LV5FU2, capecitabine, FOLFOX4, XELOX, and observation, which are important for understanding the potential benefits of chemotherapy in this age group. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into disease-free survival outcomes for elderly patients, which may influence treatment decisions. Eligibility criteria include being 70 years or older and having a diagnosis of stage III colonic adenocarcinoma.

NCT: NCT02355379 · Enrollment: 774 · Sites: Abbeville, France, Agen, France, Aix-en-Provence, France, Albi, France, Alès, France

SIGNAL **PHASE2, PHASE3** **All Stages** **RAS** **RESEARCH** • **NOT_YET_RECRUITING**

⚡ 8

Injection of IP-001 Into Thermally Ablated Hepatic Tumors in Patients With Colorectal Liver Metastases

This clinical trial is testing the efficacy of the injectable drug IP-001 in patients with colorectal cancer that has metastasized to the liver, specifically following standard microwave tumor ablation. The intervention involves administering either a high-dose or low-dose solution of IP-001 after the ablation procedure to assess its potential in prolonging progression-free survival compared to ablation alone. This study is in Phase 2 and Phase 3 and is currently not yet recruiting, indicating that patients will need to wait for enrollment to begin before participating. Eligibility criteria include being at least 18 years old and having RAS biomarker status relevant to the study.

NCT: NCT07321847 · Enrollment: 717 ·

SIGNAL **PHASE2, PHASE3** **Stage III** **MSI/MMR** **RESEARCH** • **RECRUITING**

⚡ 8

Efficacy and Safety Comparison of Short-course Radiotherapy Followed by CapeOX Chemotherapy Plus Toripalimab With or Without Concurrent Surufatinib in Neoadjuvant Therapy for Mid-to-low Localized Rectal Cancer of High-risk Criteria

This clinical trial is testing the efficacy and safety of short-course radiotherapy followed by CapeOX chemotherapy combined with the immunotherapy drug Toripalimab, with or without the anti-angiogenic drug Surufatinib, in patients with mid-to-low localized rectal cancer who meet high-risk criteria. The intervention aims to improve the pathological complete response rate in patients with high-risk locally advanced rectal cancer, addressing issues of regional recurrence and distant metastasis that persist despite current treatment methods. This study is currently in Phase 2 and Phase 3 and is actively recruiting participants, which means eligible patients may have the opportunity to access this investigational treatment. Eligibility criteria include specific biomarkers such as MMR and RAS status.

NCT: NCT07505472 · Enrollment: 212 · Sites: Changchun, China

MEDIUM **PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **NOT_YET_RECRUITING**

⚡ 8

A Phase II Clinical Study Evaluating SSGJ-706 in Combination Therapy for Advanced Gastrointestinal Cancers

This phase II clinical trial is evaluating the combination of SSGJ-706 with standard therapies for patients with advanced gastrointestinal cancers. The interventions include SSGJ-706 combined with XELOX, Bevacizumab, AG, and TP, which are important for assessing the safety and effectiveness of this treatment approach. The trial is currently not yet recruiting participants, meaning that patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and HER2.

NCT: NCT07233850 · Enrollment: 300 · Sites: Wuhan, China

SIGNAL

PHASE2, PHASE3

Stage III

RESEARCH

• **RECRUITING**

⚡ 8

Total Neoadjuvant Therapy and Organ Preservation Versus Surgery for Rectal Cancer.

This clinical trial is testing the effectiveness of total neoadjuvant therapy (TNT) and a watch-and-wait strategy for patients with rectal cancer. The intervention includes radiation therapy, chemoradiotherapy, and consolidation chemotherapy, with the aim of potentially allowing patients to preserve their rectum if they achieve a complete or near-complete clinical response. Currently in Phase 2 and Phase 3, the trial is recruiting 400 participants, which means eligible patients may have the opportunity to contribute to research that could impact treatment approaches. Eligibility criteria include patients with cT1N1, T2-T3 N0-1 rectal cancer, and specific markers such as MRF- and EMVI-.

NCT: NCT06758830 · Enrollment: 400 · Sites: Vilnius, Lithuania

MEDIUM

PHASE3

All Stages

MSI/MMR

IMMUNOTHERAPY

• **RECRUITING**

⚡ 8

Personalized Long-course Radiotherapy Plus Chemotherapy With or Without Immunotherapy for LARC: PALACE Study

The PALACE study is a phase III clinical trial designed for patients with locally advanced rectal cancer (LARC) to evaluate the complete response rate to personalized long-course radiotherapy combined with chemotherapy, with or without the anti-PD-1 antibody drug Serplulimab. This intervention aims to determine the effectiveness of adding immunotherapy to standard treatment in achieving a complete response, defined as a pathological complete response plus a clinical complete response sustained for over one year. Currently, the trial is recruiting participants, which means eligible patients have the opportunity to join and contribute to this research. Eligibility criteria include specific biomarkers such as MMR and RAS.

NCT: NCT07020247 · Enrollment: 184 · Sites: Chengdu, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• **RECRUITING**

⚡ 8

Dostarlimab for Locally Advanced or Metastatic Cancer (Non-colorectal/Non-endometrial) With Tumor dMMR/MSI

This clinical trial is testing the efficacy of dostarlimab, an immunotherapy drug, for adult patients with locally advanced or metastatic cancers that are not colorectal or endometrial and have deficient mismatch repair (dMMR) or microsatellite instability (MSI). The intervention involves comparing dostarlimab to standard chemotherapy as a first-line treatment, which is significant for understanding alternative therapeutic options for these specific cancer types. This is a Phase 2 trial currently recruiting 120 participants, which means eligible patients may have the opportunity to receive either treatment while contributing to research on their effectiveness. Eligibility criteria include adults aged 18 and older with specific cancer types, including duodenum and small bowel adenocarcinoma, gastric adenocarcinoma, and others with documented dMMR/MSI status.

NCT: NCT06333314 · Enrollment: 120 · Sites: Angers, France, Avignon, France, Besançon, France, Brest, France, Caen, France

SIGNAL

PHASE3

Stage IV

BRAF

RESEARCH

• **NOT_YET_RECRUITING**

⚡ 8

Nimotuzumab Combined With Trifluridine/Tipiracil in the Treatment of Refractory Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of nimotuzumab combined with trifluridine/tipiracil in patients with refractory metastatic colorectal cancer (mCRC). The intervention involves the administration of nimotuzumab, a monoclonal antibody, alongside trifluridine/tipiracil, a chemotherapy regimen, to assess its impact on overall survival. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MMR, and RAS.

NCT: NCT06343116 · Enrollment: 420 ·

HIGH PHASE3 Stage IV PHASE III TRIAL • RECRUITING

8

Phase III Study Evaluating Induction Chemotherapy Followed by Chemoradiotherapy Compared to Standard Chemoradiotherapy for Locally Advanced SCCA

This Phase III clinical trial is evaluating the effectiveness of induction chemotherapy using the modified DCF protocol (mDCF) followed by chemoradiotherapy compared to standard chemoradiotherapy for patients with locally advanced squamous cell carcinoma of the anus (SCCA). The intervention involves administering mDCF, followed by concomitant chemotherapy with Capecitabine and Mitomycin-C, along with radiotherapy, aiming to improve disease-related event-free survival (DREFS) at two years. The trial is currently recruiting 310 participants, which means eligible patients can enroll to potentially access this treatment approach. Eligibility criteria may include specific tumor stages and the presence of biomarkers related to HPV.

NCT: NCT06207981 · Enrollment: 310 · Sites: Agen, France, Aix-en-Provence, France, Amiens, France, Antony, France, Argenteuil, France

MEDIUM N/A Stage IV MSI/MMR TARGETED THERAPY • RECRUITING

8

The Efficacy of Watch and Wait Strategy or Surgery After Neoadjuvant Immunotherapy for Locally Advanced Colorectal Cancer With dMMR/MSI-H Guided by MRD Dynamic Monitoring: A Single-center, Open-label, Prospective, Phase II Clinical Trial.

This clinical trial is testing the efficacy of a watch and wait strategy compared to surgery after neoadjuvant immunotherapy in patients with locally advanced colorectal cancer characterized by deficient mismatch repair or microsatellite instability-high (dMMR/MSI-H). The intervention involves the use of Tislelizumab, a PD-1 inhibitor, which is significant for its potential to enhance the immune response against cancer cells. Currently, the trial is in the recruiting phase, indicating that eligible patients can still enroll to participate in this study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite status, as well as the use of next-generation sequencing (NGS) for patient selection.

NCT: NCT06477991 · Enrollment: 22 · Sites: Kunming, China

MEDIUM PHASE2, PHASE3 All Stages MSI/MMR TARGETED THERAPY • RECRUITING

8

DETERMINE Trial Treatment Arm 02: Atezolizumab in Adult, Paediatric and Teenage/Young Adult Patients With Cancers With High Tumour Mutational Burden (TMB) or Microsatellite Instability-high (MSI-high) or Proven Constitutional Mismatch Repair Deficiency (CMMRD) Disposition

The DETERMINE Trial Treatment Arm 02 is investigating the efficacy of atezolizumab in adult, pediatric, and teenage/young adult patients with cancers characterized by high tumor mutational burden (TMB), microsatellite instability-high (MSI-high), or proven constitutional mismatch repair deficiency (CMMRD). Atezolizumab is a drug that has been approved for several cancer types and is being evaluated for its potential benefits in additional cancers with specific genetic markers. This trial is currently in Phase 2 and Phase 3 and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to important research. Eligibility criteria include the presence of specific biomarkers such as MSI, TMB, and MMR, among others.

NCT: NCT05770102 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL PHASE2, PHASE3 Stage IV BRAF RESEARCH • RECRUITING

8

Regorafenib With Low-dose Chemotherapies and Aspirin Followed by Standard Chemotherapies in Metastatic Colorectal Cancer

This clinical trial is testing the combination of regorafenib with low-dose metronomic chemotherapies and aspirin for patients with metastatic colorectal cancer. The intervention aims to assess the effectiveness of this combination as a two-month induction therapy prior to standard chemotherapy initiation. Currently in Phase 2 and Phase 3, the trial is recruiting participants, which means eligible patients can join to potentially receive this treatment approach. Eligibility criteria include specific biomarkers such as BRAF, RAS, and microsatellite status.

NCT: NCT05462613 · Enrollment: 446 · Sites: Besançon, France, Clermont-Ferrand, France, Créteil, France, Dijon, France, Lyon, France

SIGNAL PHASE2, PHASE3 Stage III RAS RESEARCH • RECRUITING

8

Adjuvant mFOLFIRINOX for High-risk Stage III Colon Cancer

This clinical trial is testing the effectiveness of mFOLFIRINOX compared to mFOLFOX6 as adjuvant treatment for patients with high-risk stage III colon cancer. The intervention involves two chemotherapy regimens, mFOLFIRINOX and mFOLFOX6, which are important for potentially improving disease-free survival in this patient population. Currently, the trial is in Phase 2 and Phase 3 and is actively recruiting participants, indicating that eligible patients may have the opportunity to receive one of the treatments while contributing to the research. Eligibility criteria include specific tumor characteristics, such as RAS biomarker status, which may influence treatment selection.

NCT: NCT05179889 · Enrollment: 308 · Sites: Daejeon, South Korea

SIGNAL

PHASE3

Stage IV

BRAF

RESEARCH

• **RECRUITING**

⚡ 8

Conversion Therapy of RAS/BRAF Wild-Type Colorectal Cancer Patients With Initially Unresectable Liver Metastases

This clinical trial is testing the effectiveness of two treatment combinations for patients with RAS/BRAF wild-type colorectal cancer who have initially unresectable liver metastases. The interventions being compared are modified FOLFOXIRI (mFOLFOXIRI) combined with cetuximab and mFOLFOXIRI combined with bevacizumab, as both combinations may improve response rates and the potential for surgical resection of liver metastases. This is a Phase 3 trial currently recruiting 508 participants, which means eligible patients may have the opportunity to receive one of the treatment regimens while contributing to important research outcomes. Eligibility criteria include confirmation of RAS/BRAF wild-type status and the presence of liver-only metastases, as determined by a local multidisciplinary team.

NCT: NCT04687631 · Enrollment: 508 · Sites: Shanghai, China

SIGNAL

PHASE3

Stage IV

RESEARCH

• **ACTIVE_NOT_RECRUITING**

⚡ 8

Consolidation Chemotherapy for Locally Advanced Mid or Low Rectal Cancer After Neoadjuvant Concurrent Chemoradiotherapy

This clinical trial is testing the efficacy and feasibility of consolidation chemotherapy for patients with locally advanced mid or low rectal cancer following neoadjuvant concurrent chemoradiotherapy. The intervention involves administering chemotherapy to evaluate its impact on achieving a pathologic complete response. Currently, the trial is in Phase 3 and is active but not recruiting, which means that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria may include specific characteristics related to the cancer stage and treatment history.

NCT: NCT02843191 · Enrollment: 358 · Sites: Busan, South Korea, Cheonan, South Korea, Chuncheon, South Korea, Daegu, South Korea, Daejeon, South Korea

SIGNAL

PHASE3

Stage IV

KRAS

RESEARCH

• **RECRUITING**

⚡ 8

Systemic Oxaliplatin or Intra-arterial Chemotherapy Combined With LV5FU2 +/- Irinotecan and an Target Therapy in First Line Treatment of Metastatic Colorectal Cancer Restricted to the Liver

This clinical trial is testing the effectiveness of systemic oxaliplatin versus intra-arterial chemotherapy combined with LV5FU2, with or without irinotecan and panitumumab, in patients with metastatic colorectal cancer restricted to the liver. The intervention involves administering oxaliplatin both intravenously and intra-arterially, along with other chemotherapy agents and targeted therapy, to potentially enhance treatment efficacy for patients with unresectable liver metastases. This is a Phase 3 trial currently recruiting 348 participants, which indicates that it is in the later stages of testing and may provide valuable information on treatment options for patients. Eligibility criteria include specific biomarker assessments for KRAS, NRAS, and RAS.

NCT: NCT02885753 · Enrollment: 348 · Sites: Angers, France, Antony, France, Avignon, France, Bayonne, France, Beauvais, France

SIGNAL

PHASE3

All Stages

MSI/MMR

RESEARCH

• **NOT_YET_RECRUITING**

⚡ 8

Finding the Best Dose of Aspirin to Prevent Lynch Syndrome Cancers

This clinical trial is testing the effectiveness of different doses of enteric coated aspirin in preventing cancers associated with Lynch Syndrome, specifically for individuals who carry a germline pathological mismatch repair gene defect. The intervention involves comparing daily doses of 600mg, 300mg, and 100mg of aspirin to determine the optimal dose for cancer prevention. This is a Phase 3 trial that is currently not yet recruiting participants, meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include being a carrier of a germline pathological mismatch repair gene defect associated with Lynch Syndrome.

NCT: NCT02497820 · Enrollment: 1800 · Sites: Tel Aviv, Israel

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

8

Study of Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-3475-158/KEYNOTE-158)

This clinical trial is testing the efficacy of pembrolizumab (MK-3475) in participants with advanced solid tumors who have progressed on standard of care therapy. The intervention, pembrolizumab, is a biological agent that targets the immune system to help fight cancer, which is significant for patients with limited treatment options. This is a Phase 2 trial that is currently active but not recruiting, indicating that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, and microsatellite instability, which may be relevant for patient selection.

NCT: NCT02628067 · Enrollment: 1609 ·

SIGNAL

PHASE3

All Stages

RESEARCH

• ACTIVE_NOT_RECRUITING

8

A Trial of Aspirin on Recurrence and Survival in Colon Cancer Patients

This clinical trial is testing the effectiveness of acetylsalicylic acid in reducing recurrence and improving survival in patients with colon cancer. The intervention involves administering acetylsalicylic acid compared to a placebo, which is significant for understanding its potential role in cancer management. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval. Eligibility criteria have not been specified, but the study involves a total enrollment of 770 participants.

NCT: NCT02301286 · Enrollment: 770 · Sites: Almelo, Netherlands, Amersfoort, Netherlands, Assen, Netherlands, Breda, Netherlands, Capelle aan den IJssel, Netherlands

SIGNAL

PHASE3

All Stages

RESEARCH

• ACTIVE_NOT_RECRUITING

8

The Northern-European Initiative on Colorectal Cancer

The Northern-European Initiative on Colorectal Cancer (NordICC) is a Phase 3 clinical trial designed to evaluate the effectiveness of colonoscopy screening in reducing colorectal cancer (CRC) incidence and mortality among individuals aged 55-64 years. The intervention involves a colonoscopy procedure, which is significant as it can detect and remove precursor lesions that may lead to CRC, potentially preventing the disease. Currently, the trial is active but not recruiting, indicating that it is ongoing with enrolled participants, and results will be assessed after 15 years of follow-up. Eligibility criteria include being part of the specified age group and drawn randomly from population registries in the participating Nordic countries, the Netherlands, and Poland.

NCT: NCT00883792 · Enrollment: 95000 · Sites: Boston, New York

SIGNAL

PHASE3

Stage IV

BRAF

RESEARCH

• NOT_YET_RECRUITING

7

Pressurized Intraperitoneal Aerosolized Chemotherapy With Mitomycin for the Treatment of Unresectable Appendix or Colorectal Cancer With Peritoneal Metastases, The IMPACT Trial

This phase III clinical trial, known as the IMPACT Trial, is testing the effectiveness of pressurized intraperitoneal aerosolized chemotherapy (PIPAC) with mitomycin in patients with unresectable appendix or colorectal cancer that has metastasized to the peritoneal cavity. The intervention involves delivering chemotherapy as a pressurized mist directly into the abdominal cavity during a minimally invasive laparoscopic procedure, which may enhance drug absorption and distribution compared to standard chemotherapy regimens. The trial is currently not yet recruiting participants, which means patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarker assessments such as BRAF, RAS, and mismatch repair status, as well as next-generation sequencing (NGS).

NCT: NCT07271355 · Enrollment: 129 · Sites: Duarte, Goodyear, Irvine, Newnan, Zion

MEDIUM

PHASE1, PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

7

A Clinical Research Study Evaluating EIK1005, a Werner Helicase Inhibitor, as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced Solid Tumors Including Microsatellite Instability High (MSI-H) Tumors

The EIK1005-002 clinical trial is evaluating the efficacy and safety of EIK1005, a Werner Helicase inhibitor, both as a monotherapy and in combination with pembrolizumab in participants with advanced solid tumors, including those with microsatellite instability high (MSI-H) tumors. The intervention involves administering EIK1005 to determine the most effective and safe dosage, as well as assessing its tolerability when combined with pembrolizumab. This trial is currently in Phase 1 and Phase 2 and is actively recruiting participants, which means eligible patients may have the opportunity to

access these investigational treatments. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT07262619 · Enrollment: 160 · Sites: Houston, Morristown, New York

HIGH **PHASE3** **All Stages** **PHASE III TRIAL** • **RECRUITING**

7

Phase 3 Trial of eRapa in Patients With Familial Adenomatous Polyposis

This Phase 3 clinical trial is testing the efficacy of eRapa (encapsulated rapamycin) in patients diagnosed with Familial Adenomatous Polyposis (FAP). The intervention involves administering eRapa compared to a placebo to determine its impact on slowing disease progression and its safety profile. Currently, the trial is recruiting 168 participants, which means eligible patients can still join to contribute to the research. Eligibility criteria include a diagnosis of FAP, and participants will undergo regular check-ups and endoscopies to monitor their condition throughout the study.

NCT: NCT06950385 · Enrollment: 168 · Sites: Ann Arbor, Arcadia, Baltimore, Boston, Chicago

MEDIUM **NA** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **ACTIVE_NOT_RECRUITING**

7

Reducing the Resection Range After Neoadjuvant Therapy for Locally Advanced Upper Rectal and Rectosigmoid Junction Cancers

This clinical trial is evaluating the safety and efficacy of reduced surgical resection margins in patients with locally advanced upper rectal and rectosigmoid junction cancers who have shown regression following neoadjuvant therapy. The intervention involves laparoscopic resection with reduced margins of 3 cm distally and 5 cm proximally, which is significant for potentially minimizing surgical impact while maintaining effective cancer treatment. The trial is currently active but not recruiting, indicating that it is ongoing and may provide insights into surgical practices for eligible patients in the future. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT07038122 · Enrollment: 874 · Sites: Guangzhou, China

SIGNAL **PHASE3** **All Stages** **KRAS** **RESEARCH** • **RECRUITING**

7

A Study of Tucidinostat in Combination With Sintilimab and Bevacizumab in MSS/pMMR Colorectal Cancer Patients

This clinical trial is testing the combination of tucidinostat, sintilimab, and bevacizumab in patients with MSS/pMMR colorectal cancer. The intervention aims to evaluate the efficacy and safety of this combination compared to fruquintinib monotherapy, which is significant for understanding potential treatment options. Currently in Phase 3 and actively recruiting, this trial may provide eligible patients with access to a novel treatment approach while contributing to the overall understanding of colorectal cancer therapies. Eligibility criteria include specific biomarkers such as KRAS, MSI, MMR, and RAS.

NCT: NCT06497985 · Enrollment: 430 · Sites: Guangzhou, China

SIGNAL **PHASE3** **All Stages** **RESEARCH** • **NOT_YET_RECRUITING**

7

Preoperative Chemoradiotherapy and Transanal Endoscopic Microsurgery Versus Ttransanal Endoscopic Microsurgery in T1 N0, M0 Rectal Cancer (TAUTEM-T1 Study)

the T1 N0, M0 rectal cancer stage. This trial is testing the combination of preoperative chemoradiotherapy using Capecitabine and 50.4 Gy of radiation followed by Transanal Endoscopic Microsurgery (TEM) against TEM alone. As a Phase 3 trial that is not yet recruiting, it aims to evaluate rectal preservation rates in patients with early-stage rectal cancer, which may influence treatment decisions and outcomes. Eligibility criteria include patients diagnosed with T1, N0, M0 rectal adenocarcinoma without poor prognostic factors.

NCT: NCT06450574 · Enrollment: 106 ·

SIGNAL **PHASE3** **All Stages** **RESEARCH** • **RECRUITING**

7

CompariSon Between the EuroPeAn and Japanese pathologiCal InvEstigation for Colon Cancer (SPACE)

The SPACE trial is testing the diagnostic accuracy of Japanese versus European pathological investigation methods for stage III colon cancer in patients. The intervention involves comparing two procedures: the Japanese pathological investigation, which includes surgeon involvement in lymph node extraction and specimen assessment, and the European

method, which primarily relies on pathologists. This is a Phase 3 trial currently recruiting 430 participants, which means patients may have the opportunity to contribute to important findings regarding diagnostic practices. Eligibility criteria include patients diagnosed with stage III colon cancer, although specific biomarker requirements are not mentioned.

NCT: NCT06119867 · Enrollment: 430 · Sites: Moscow, Russia

SIGNAL **PHASE3** **All Stages** **RAS** **RESEARCH** • **RECRUITING**

7

Regorafenib Combined With Irinotecan Drug-Eluting Beads for Colorectal Cancer Liver Metastases

This clinical trial is testing the efficacy and safety of regorafenib combined with Irinotecan Drug-Eluting Beads in patients with colorectal cancer liver metastases who have failed first- and second-line standard chemotherapy. The intervention involves the use of regorafenib and DIBIRI (drug-eluting beads), which is significant as it explores a combination treatment approach for this patient population. This is a Phase 3 trial currently recruiting 126 participants, which means that eligible patients may have the opportunity to access this investigational treatment while contributing to important research. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT05794971 · Enrollment: 126 · Sites: Guangdong, China

SIGNAL **PHASE3** **All Stages** **RAS** **RESEARCH** • **RECRUITING**

7

Surgery Plus Chemo Versus Chemoradiotherapy Followed by Surgery Plus Chemo for Locally Recurrent Rectal Cancer

The JCOG1801 trial is testing the effectiveness of preoperative chemoradiotherapy followed by surgery and adjuvant chemotherapy in patients with locally recurrent rectal cancer. The intervention involves a combination of chemotherapy, preoperative radiotherapy, and surgical procedures, aiming to improve local relapse-free survival compared to standard treatment. This is a Phase 3 trial currently recruiting 110 participants, which means eligible patients may have the opportunity to receive a potentially more effective treatment approach. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT04288999 · Enrollment: 110 · Sites: Chiba, Japan, Gifu, Japan, Hidaka, Japan, Hirakata, Japan, Hiroshima, Japan

SIGNAL **PHASE3** **All Stages** **RESEARCH** • **ACTIVE_NOT_RECRUITING**

7

Effect of EPA-FFA on Polypectomy in Familial Adenomatous Polyposis

This clinical trial is testing the effect of eicosapentaenoic acid free fatty acid (EPA-FFA) on polypectomy outcomes in patients with familial adenomatous polyposis (FAP). The intervention involves administering EPA-FFA gastro-resistant capsules compared to a placebo, which is important for assessing its potential impact on the number of polypectomies required. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the later stages of testing and may provide significant insights into treatment efficacy for patients. Eligibility criteria may include a diagnosis of familial adenomatous polyposis, although specific biomarker requirements are not detailed.

NCT: NCT03806426 · Enrollment: 204 · Sites: Bologna, Italy

MEDIUM **PHASE1, PHASE2** **Stage IV** **KRAS** **LIQUID BIOPSY** • **RECRUITING**

7

A Study of Bispecific Antibody MCLA-158 in Patients With Advanced Solid Tumors

This clinical trial is testing the bispecific antibody MCLA-158 in patients with advanced solid tumors, specifically focusing on metastatic colorectal cancer (mCRC) and other solid tumor indications. The intervention includes MCLA-158 as a single agent and in combination with pembrolizumab, FOLFIRI, and FOLFOX, which is important for evaluating its safety and effectiveness in various treatment settings. Currently in Phase 1/2 and actively recruiting, this trial aims to determine the recommended Phase II dose and assess the drug's safety, tolerability, and anti-tumor activity. Eligibility criteria include specific biomarkers such as KRAS, NRAS, BRAF, HER2, and others, which may influence treatment response.

NCT: NCT03526835 · Enrollment: 523 · Sites: Boston, Cleveland, Dallas, Fort Myers, Houston

SIGNAL **PHASE3** **All Stages** **RESEARCH** • **ACTIVE_NOT_RECRUITING**

7

EPA for Metastasis Trial 2

The EPA for Metastasis Trial 2 is testing the efficacy of Icosapent Ethyl, a pure omega-3 fatty acid derived from fish oil, in patients with liver metastases from bowel cancer. The trial aims to determine if taking this supplement before and after liver surgery can improve progression-free survival compared to a placebo. Currently in Phase 3 and active but not

recruiting, this study involves 418 participants and seeks to provide insights into the potential benefits of Icosapent Ethyl in preventing disease recurrence. Eligibility criteria include having liver metastases from bowel cancer, with a target enrollment of 448 individuals.

NCT: NCT03428477 · Enrollment: 418 · Sites: Aintree, United Kingdom, Basingstoke, United Kingdom, Birmingham, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL

PHASE2, PHASE3

All Stages

RESEARCH

• **ACTIVE_NOT_RECRUITING**

7

Perioperative Systemic Therapy for Isolated Resectable Colorectal Peritoneal Metastases

This clinical trial is testing the effectiveness of perioperative systemic therapy combined with cytoreductive surgery and HIPEC in patients with isolated resectable colorectal peritoneal metastases. The intervention includes various combinations of chemotherapy regimens (CAPOX, FOLFOX, FOLFIRI) with bevacizumab, alongside the experimental procedure of cytoreductive surgery with HIPEC. Currently in Phase II-III and marked as active but not recruiting, this trial aims to determine if the combination therapy improves overall survival compared to surgery alone. Eligibility criteria include having isolated resectable colorectal peritoneal metastases, although specific biomarker requirements are not detailed.

NCT: NCT02758951 · Enrollment: 358 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Genk, Belgium, Groningen, Netherlands, Nieuwegein, Netherlands

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• **ACTIVE_NOT_RECRUITING**

6

The Efficacy and Safety of Neoadjuvant Therapy With Iparomlimab and Tuvonralimab in Locally Advanced MSI-H/dMMR Colorectal Cancer: An Prospective, Single-Arm Study (Neo-IT)

This clinical trial is testing the efficacy and safety of neoadjuvant therapy with Iparomlimab and Tuvonralimab in patients with locally advanced colorectal cancer that is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR). The intervention involves the use of these two immunotherapy drugs, which may provide significant benefits in the preoperative treatment of this specific cancer type. This is a Phase 2 trial that is currently active but not recruiting, indicating that patients are not being enrolled at this time, but results may inform future treatment options. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, and RAS status.

NCT: NCT07160647 · Enrollment: 29 · Sites: Hangzhou, China

MEDIUM

PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• **RECRUITING**

6

A First in Human Study of ALX2004 With Advanced or Metastatic Selected Solid Tumors

This clinical trial is testing ALX2004, an antibody drug conjugate targeting EGFR, in patients with advanced or metastatic selected solid tumors. The intervention involves administering ALX2004 to evaluate its safety and tolerability, specifically focusing on the incidence of dose limiting toxicities (DLTs). Currently in Phase 1 and actively recruiting, this trial aims to gather initial data on the drug's effects, which is crucial for determining its potential for further development. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT07085091 · Enrollment: 170 · Sites: Fairfax, Grand Rapids, Houston, Portland, Spokane

MEDIUM

N/A

Stage III

MSI/MMR

TARGETED THERAPY

• **RECRUITING**

6

Exploring the Treatment Duration of PD-1 Neoadjuvant Therapy in Stage II-III dMMR Rectal Cancer

This clinical trial is investigating the optimal treatment duration of PD-1 neoadjuvant therapy for patients with stage II-III dMMR rectal cancer. Participants will receive preoperative monotherapy with PD-1 antibodies, which is important for assessing its effectiveness in achieving a pathological complete response (pCR). The trial is currently recruiting 100 participants and aims to evaluate outcomes such as pCR rates, adverse reactions, and three-year event-free survival. Eligibility criteria include biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06613165 · Enrollment: 100 · Sites: Xi'an, China

MEDIUM

PHASE1

Stage IV

BRAF

TARGETED THERAPY

• **RECRUITING**

6

A First-in-Human Escalation and Expansion Study of Patients With Advanced Solid Tumors

This clinical trial is testing the investigational drug JR8603 in patients with advanced solid tumors who have not responded to prior treatments. The study aims to evaluate the safety and effectiveness of JR8603, which is significant for patients

seeking alternative options after standard therapies have failed. Currently in Phase 1 and actively recruiting participants, this trial provides an opportunity for patients to access a potential treatment option while contributing to research. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, HER2, and RAS.

NCT: NCT06783569 · Enrollment: 94 · Sites: Harbin, China, Shenyang, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Phase 2 Study of SR-8541A in Combination With Botensilimab and Balstilimab in Subjects With Refractory Metastatic Microsatellite Stable Colorectal Cancer (MSS-CRC)

This clinical trial is testing the combination of SR-8541A, botensilimab, and balstilimab in patients with refractory metastatic microsatellite stable colorectal cancer (MSS-CRC). The intervention involves administering SR-8541A orally alongside the other two drugs to assess their safety, tolerability, and efficacy, as well as to determine the optimal dose for further studies. Currently, this is a Phase 2 trial that is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include specific biomarkers related to microsatellite stability and mismatch repair status.

NCT: NCT06589440 · Enrollment: 70 · Sites: Austin, Dallas, Houston, Morristown, Seattle

MEDIUM

PHASE1, PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

This clinical trial is testing a combination immunotherapy regimen for adults with metastatic colorectal cancer (mCRC). The intervention includes the drugs retifanlimab, SX-682, and N-803, along with the TriAdeno vaccine, aiming to evaluate their safety profiles in treating mCRC. Currently in Phase I/II and actively recruiting, this trial offers participants the opportunity to access experimental therapies that may not be available outside of a clinical trial setting. Eligible participants must be 18 years or older and diagnosed with mCRC, with specific biomarker criteria including MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT06149481 · Enrollment: 60 · Sites: Bethesda

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Liposomal Irinotecan and Leucovorin/5-fluorouracil Plus Bevacizumab in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of liposomal irinotecan combined with 5-fluorouracil, leucovorin, and bevacizumab as a second-line therapy for patients with metastatic colorectal cancer. The intervention involves a combination of these drugs, which is important for potentially improving treatment outcomes in this patient population. Currently, the trial is in Phase 2 and is actively recruiting participants, indicating that patients may have the opportunity to receive this treatment while the study is ongoing. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06184698 · Enrollment: 173 · Sites: Shijiazhuang, China

MEDIUM

PHASE2

Stage IV

BRAF

LIQUID BIOPSY

• RECRUITING

6

Cetuximab Plus Irinotecan in Patients With NeoRAS Wild-type Metastatic Colorectal Cancer In Third-line Therapy

This clinical trial is testing the efficacy and safety of cetuximab plus irinotecan in patients with NeoRAS wild-type metastatic colorectal cancer (mCRC) who are receiving third-line therapy. The intervention involves the combination of cetuximab, a monoclonal antibody, and irinotecan, a chemotherapy drug, which may provide a targeted approach for this specific patient population. Currently in phase II and actively recruiting, this trial aims to assess the objective response rate (ORR) to the treatment. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, CTDNA, and RAS status.

NCT: NCT05962502 · Enrollment: 54 · Sites: Guangzhou, China

MEDIUM

NA

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

MRD-guided Deferred Adjuvant Therapy in Resectable Early-stage Colon Cancer

This clinical trial is testing the use of minimal residual disease (MRD) status, detected by circulating tumor DNA (ctDNA), to guide post-surgery adjuvant therapy in patients with resectable early-stage colon cancer. The intervention involves comparing standard CAPEOX adjuvant therapy, deferred CAPEOX adjuvant therapy, and mFOLFOXIRI intensive adjuvant therapy based on MRD status, which may help tailor treatment to individual patient needs. The trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate in this study. Eligibility criteria include having high-risk stage II or low-risk stage III colon cancer and specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT06204484 · Enrollment: 349 · Sites: Guangzhou, China

MEDIUM

PHASE1, PHASE2

Stage IV

RAS

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

6

Study of JK08 in Patients with Unresectable Locally Advanced or Metastatic Cancer

This clinical trial is evaluating the safety and efficacy of JK08 in patients with unresectable locally advanced or metastatic cancer. The intervention involves administering JK08, along with Pembrolizumab and Lenvatinib, to assess its effects on cancer progression and patient outcomes. Currently in Phase 1/2 and marked as active but not recruiting, this trial indicates that patients may not be able to enroll at this time but can learn about the study's findings as they become available. Eligibility criteria include specific biomarkers such as HER2 and RAS.

NCT: NCT05620134 · Enrollment: 263 · Sites: Barcelona, Spain, Brussels, Belgium, Edegem, Belgium, Ghent, Belgium, Madrid, Spain

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Neoadjuvant Immune Checkpoint Inhibition and Novel IO Combinations in Early-stage Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant immune checkpoint inhibition and novel immunotherapy combinations in patients with early-stage colon cancer (stage 1-3 adenocarcinoma) who do not have distant metastases. The intervention includes a combination of drugs such as Nivolumab, Ipilimumab, Celecoxib, BMS-986253, and BMS-986016, which are being evaluated for their safety and potential to improve treatment outcomes before surgical resection. This is a Phase 2 trial that is currently recruiting participants, indicating that eligible patients may have the opportunity to receive this investigational treatment in the lead-up to surgery. Eligibility criteria include specific biomarker requirements related to MMR and DMMR status.

NCT: NCT03026140 · Enrollment: 353 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Haarlem, Netherlands, Hilversum, Netherlands, The Hague, Netherlands

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

5

Becotatug Vedotin Combined With Cetuximab in the Later-line Treatment of Metastatic RAS Wild-type Colorectal Cancer

This clinical trial is testing the combination of becotatug vedotin and cetuximab for patients with metastatic RAS wild-type colorectal cancer who have already received prior treatments. The intervention involves the use of becotatug vedotin, a drug that targets cancer cells, alongside cetuximab, which is designed to inhibit tumor growth. This is a Phase 2 trial that is currently not yet recruiting, meaning patients cannot enroll at this time. Eligibility criteria include specific biomarker assessments such as BRAF, MSI, MMR, and RAS status.

NCT: NCT07490119 · Enrollment: 31 ·

MEDIUM

PHASE1, PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

5

LPM6690176 in Combination With Chemotherapy and Bevacizumab in Metastatic Colorectal Cancer Patients With RAS Mutation

This clinical trial is testing the combination of LPM6690176 with chemotherapy and Bevacizumab in patients with RAS mutant metastatic colorectal cancer (mCRC). The intervention involves LPM6690176, a drug being evaluated for its safety and tolerability alongside standard treatments, which is important for determining its potential role in managing this specific cancer type. The trial is currently in Phase 1b and Phase 2 and is not yet recruiting participants, meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include the presence of specific biomarkers such as RAS mutations and mismatch repair status.

NCT: NCT07391566 · Enrollment: 99 · Sites: Beijing, China

MEDIUM

PHASE1, PHASE2

Stage III

KRAS

TARGETED THERAPY

• RECRUITING

5

KRAS Neoantigen Nanovaccine as Adjuvant Therapy for Colorectal Cancer/Pancreatic Cancer

This clinical trial is testing the KRAS Neoantigen Nanovaccine as adjuvant therapy for patients with colorectal or pancreatic cancer who have KRAS mutations and are at high risk of recurrence. The intervention involves a nanovaccine constructed from the bacterial membranes of an engineered *Lactococcus lactis* strain, which expresses KRAS antigenic peptides, aiming to enhance the immune response against cancer cells. Currently in Phase 1 and Phase 2, the trial is recruiting participants, which means eligible patients may have the opportunity to receive this investigational therapy. Eligibility criteria include the presence of specific biomarkers such as KRAS, BRAF, MSI, and MMR status.

NCT: NCT07353645 · Enrollment: 49 · Sites: Nanjing, China

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

5

The Safety and Efficacy of Cetuximab Beta Plus Fruquintinib With or Without Immune Checkpoint Inhibitors in First-line Treatment of RAS/BRAF Wild Type Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the safety and efficacy of Cetuximab Beta combined with Fruquintinib, with or without immune checkpoint inhibitors, in patients with RAS/BRAF wild type unresectable metastatic colorectal cancer. The intervention involves targeted therapies that aim to improve treatment outcomes while potentially avoiding the side effects associated with traditional chemotherapy. This is a Phase 2 trial currently recruiting 70 participants, which means eligible patients may have the opportunity to receive these investigational treatments as part of their first-line therapy. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, and RAS status.

NCT: NCT07257653 · Enrollment: 70 · Sites: Hangzhou, China

MEDIUM

EARLY_PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

A Dose-Expansion Trial of Intravenous HNF4 α srRNA for Unresectable or Metastatic Colorectal Cancer

This clinical trial is testing the safety and efficacy of intravenous HNF4 α srRNA (CD-GA-102) in patients with unresectable locally advanced or metastatic colorectal cancer. The intervention involves administering CD-GA-102 as a monotherapy or in combination with immunotherapy and other systemic treatments, which is important for understanding its potential role in managing this cancer type. Currently in the early phase 1 stage and actively recruiting, this trial aims to gather data on the objective response rate, which may inform future treatment options for patients. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT07050394 · Enrollment: 20 · Sites: Shanghai, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Y-90, Capecitabine, and Atezolizumab for Oligometastatic CRC

The BRUOG-430 trial is evaluating the combination of yttrium-90 radioembolization, capecitabine, and atezolizumab for patients with unresectable colorectal cancer liver metastases who have previously received two or more lines of systemic therapy. This intervention aims to assess its preliminary efficacy based on the intrahepatic disease control rate at 12 weeks. Currently in Phase 2 and actively recruiting, this trial offers patients the opportunity to participate in a study focused on a specific treatment approach for their condition. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, RAS, and microsatellite instability.

NCT: NCT06555133 · Enrollment: 18 · Sites: Providence

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

5

Liposomal Irinotecan Plus Bevacizumab in Irinotecan-refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of liposomal irinotecan and bevacizumab in patients with irinotecan-refractory metastatic colorectal cancer. The intervention involves administering liposomal irinotecan, which is being evaluated for its efficacy and safety alongside bevacizumab, a drug that may enhance treatment outcomes. This study is in Phase 1 and Phase 2 and is currently not yet recruiting, meaning that patients will need to wait until recruitment begins to participate. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MMR, and RAS status.

NCT: NCT06434090 · Enrollment: 74 · Sites: Guangzhou, China

MEDIUM NA Stage IV IMMUNOTHERAPY • RECRUITING

5

A Randomized, Multicenter Phase II Basket Study of Hypofractionated Radiotherapy/Stereotactic Body Radiotherapy Followed by Immunotherapy-Based Systemic Therapy +/- L. Rhamnosus M9 for the First-Line Treatment of Advanced Digestive System Malignancies.

This clinical trial is testing the effectiveness of hypofractionated radiotherapy or stereotactic body radiotherapy followed by immunotherapy-based systemic therapy, with or without L. Rhamnosus M9, for patients with advanced digestive system malignancies. The intervention includes a combination of radiation therapy and various drugs, such as anti-PD-1 monoclonal antibodies and chemotherapy agents, which may enhance the immune response against tumors. Currently in the recruiting phase, this Phase II study aims to enroll 120 participants, indicating that eligible patients may have the opportunity to receive this integrated treatment approach. Specific eligibility criteria or biomarker requirements have not been detailed in the provided information.

NCT: NCT06349044 · Enrollment: 120 · Sites: Hangzhou, China

MEDIUM PHASE1 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING

5

Dose Escalation and Expansion Clinical Trial of Irinotecan Liposome Combined With Oxaliplatin and 5-FU/LV Plus Bevacizumab as First-line Treatment of Metastatic Colorectal Cancer

This clinical trial is testing the combination of irinotecan liposome, oxaliplatin, 5-FU/LV, and bevacizumab as a first-line treatment for patients with advanced metastatic colorectal cancer. The intervention aims to determine the maximum tolerated dose and assess the safety and efficacy of this drug combination, which is significant for optimizing treatment regimens. Currently in Phase 1 and actively recruiting participants, this trial offers patients the opportunity to receive a potentially effective treatment while contributing to the understanding of dose-limiting toxicities. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06225622 · Enrollment: 78 · Sites: Shanghai, China

MEDIUM PHASE2 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING

5

Efficacy and Safety of Chemotherapy With XELOX (Oxaliplatin + Capecitabine) and Bevacizumab in Combination With Adebrelimab in First-line Treatment of Microsatellite Stable (MSS) Initially Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of a combination treatment for patients with microsatellite stable (MSS) initially unresectable metastatic colorectal cancer. The intervention involves the use of Adebrelimab alongside chemotherapy agents Oxaliplatin and Capecitabine, as well as Bevacizumab, which may enhance treatment outcomes. Currently in Phase 2 and actively recruiting participants, this trial aims to assess progression-free survival as its primary outcome, providing insights into the effectiveness of this treatment regimen. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT06282445 · Enrollment: 36 · Sites: Jinhua, China

MEDIUM PHASE4 Stage IV MSI/MMR TARGETED THERAPY • RECRUITING

5

Survival Benefit of Compound Kushen Injection in the Treatment of Advanced Colorectal Cancer

This clinical trial is testing the effectiveness and safety of compound kushen injection in patients with advanced colorectal cancer. The intervention involves administering compound kushen injection alongside a first-line treatment scheme, which is significant for its potential impact on patient outcomes. Currently in Phase 4 and actively recruiting, this trial aims to provide insights into progression-free survival (PFS) for participants. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT05894694 · Enrollment: 320 · Sites: Beijing, China

MEDIUM N/A All Stages MSI/MMR TARGETED THERAPY • RECRUITING

5

Observational Basket Trial to Collect Tissue to Develop and Train a Live Tumor Diagnostic Platform

This observational basket trial is designed for patients with colorectal cancer to collect tissue samples for the development and training of the Elephas live tumor diagnostic platform. The intervention involves various biopsy procedures, including core needle, forceps, and punch biopsies, which are essential for obtaining the necessary tissue for analysis. The trial is currently in the recruiting phase, indicating that eligible patients can participate and contribute to the advancement of

diagnostic methods. Eligibility criteria include specific biomarker requirements such as MSI, MMR, and TMB, which are important for the study's focus on tumor characteristics.

NCT: NCT05520099 · Enrollment: 416 · Sites: Buffalo, Chapel Hill, Hackensack, Hagerstown, Little Rock

MEDIUM

PHASE2

All Stages

KRAS

TARGETED THERAPY

• NOT_YET_RECRUITING

5

A Combination Therapy Including Anti-PD-1 Immunotherapy in MSS Rectal Cancer With Resectable Distal Metastasis

This clinical trial is testing a combination therapy that includes the anti-PD-1 immunotherapy tislelizumab for patients with microsatellite-stable (MSS) rectal cancer who have resectable distal metastasis. The intervention aims to evaluate the effectiveness of preoperative short-course radiotherapy followed by neoadjuvant chemotherapy and immunotherapy in improving tumor response and reducing postoperative metastasis and recurrence. This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MMR, DMMR, HER2, RAS, and microsatellite status.

NCT: NCT05359393 · Enrollment: 52 · Sites: Shanghai, China

MEDIUM

N/A

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

5

Retrospective Study on the Use of Immunotherapy in Patients With MSI-H Metastatic Colorectal Cancer

This clinical trial is a retrospective study examining the use of immunotherapy in patients with microsatellite instability-high (MSI-H) metastatic colorectal cancer (mCRC). The intervention involves the use of immune checkpoint inhibitors, specifically pembrolizumab and nivolumab, which are significant for their role in targeting the immune response against cancer cells. The trial is currently active but not recruiting, indicating that it is in the data collection phase, which may provide insights into treatment effectiveness for patients who have exhausted other options. Eligibility criteria include having MSI-H status, which is essential for determining the appropriateness of immunotherapy in this patient population.

NCT: NCT04612309 · Enrollment: 100 · Sites: Carpi, Italy

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Neo-adjuvant Pembrolizumab and Radiotherapy in Localised MSS Rectal Cancer

This clinical trial is testing the combination of pembrolizumab and short course radiotherapy in patients with localized microsatellite stable (MSS) rectal cancer. The intervention involves administering pembrolizumab, an immunotherapy drug, alongside radiotherapy to evaluate its effects on tumor regression. This is a Phase 2 trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include having localized MSS rectal cancer and specific biomarker assessments related to microsatellite stability.

NCT: NCT04109755 · Enrollment: 25 · Sites: Geneva, Switzerland

MEDIUM

NA

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

4

Clinical Efficacy of Transarterial Infusion Chemotherapy for Unresectable Colorectal Cancer

This clinical trial is testing the efficacy of transarterial infusion chemotherapy (TAIC) combined with either cetuximab or bevacizumab for patients with unresectable colorectal cancer (CRC). The intervention involves a procedure that delivers chemotherapy directly to the tumor site, which may enhance treatment effectiveness. The trial is currently in the recruiting phase, indicating that eligible patients can still enroll to participate in the study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT07333053 · Enrollment: 30 · Sites: Xicheng, China

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

4

A Phase II Study of Sintilimab Combined With Ipilimumab N01, Cetuximab and Dabrafenib in Patients With Microsatellite-Stable, BRAF V600E-Mutated Metastatic Colorectal Cancer

This clinical trial is testing the combination of sintilimab, ipilimumab N01, cetuximab, and dabrafenib in patients with microsatellite-stable, BRAF V600E-mutated metastatic colorectal cancer. The intervention involves a multi-drug regimen aimed at enhancing treatment efficacy for a patient population that typically has a poor prognosis with standard therapies.

Currently in Phase II and actively recruiting participants, this trial seeks to assess the safety and efficacy of the treatment approach, which may offer an alternative for patients who have limited options. Eligibility criteria include the presence of BRAF mutations and microsatellite stability, which are essential for participation in this study.

NCT: NCT07506109 · Enrollment: 49 · Sites: Beijing, China, Chengdu, China, Tianjin, China, Wuhan, China

MEDIUM

PHASE1

Stage IV

RAS

TARGETED THERAPY

• RECRUITING

⚡ 4

A Study of IDE034 in Adult Participants With Locally Advanced/Metastatic Solid Tumors Types

This clinical trial is testing the drug IDE034 in adult participants with locally advanced or metastatic solid tumors. The intervention, IDE034, is being evaluated for its safety, pharmacokinetics, immunogenicity, and preliminary efficacy, particularly in tumors that express the biomarkers B7-H3 and PTK7. Currently in Phase 1 and actively recruiting, this trial aims to assess the drug's safety and tolerability during the dose escalation phase, which is crucial for determining appropriate dosing for future studies. Eligibility criteria include the expression of specific biomarkers, such as HER2 and RAS, in the tumor.

NCT: NCT07503808 · Enrollment: 150 · Sites: Austin, Fairfax, Houston, Irving, San Antonio

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 4

A Single-arm, Single-center, Phase II Clinical Study of Aipalolitovorelizumab (QL1706) Combined With Bevacizumab and Standard Chemotherapy as First-line Treatment for MSS/pMMR Metastatic Colorectal Cancer With BRAF V600E Mutation

This clinical trial is testing the combination of Aipalolitovorelizumab (QL1706) with bevacizumab and standard chemotherapy as a first-line treatment for patients with MSS/pMMR metastatic colorectal cancer who have a BRAF V600E mutation. The intervention involves administering Aipalolitovorelizumab alongside various chemotherapy regimens, which is significant for potentially improving treatment outcomes in this specific patient population. This is a Phase II study that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include having the BRAF V600E mutation and meeting specific biomarker requirements related to MMR status.

NCT: NCT07229846 · Enrollment: 30 · Sites: Harbin, China

SIGNAL

NA

Early Detection

RESEARCH

• RECRUITING

⚡ 4

Randomized Trial Comparing Fecal Testing (FIT) to Colonoscopy for Post-polypectomy Surveillance

This clinical trial is testing the effectiveness of fecal occult blood testing (FIT) compared to colonoscopy for post-polypectomy surveillance in patients who have had colorectal polyps removed. The intervention involves using FIT as a less burdensome alternative to colonoscopy, which is important given the limited availability of colonoscopy and the fact that many surveillance procedures yield no findings. The trial is currently in the recruiting phase with an enrollment goal of 4,000 participants, which means eligible patients can still join the study. Eligibility criteria may include a history of polyp removal, but specific requirements are not detailed in the provided information.

NCT: NCT06983639 · Enrollment: 4000 · Sites: Arendal, Norway

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 4

A Phase Ib/II Trial of HS-20110 Combination Therapies in Advanced Colorectal Cancer Patients.

This clinical trial is evaluating the safety and efficacy of HS-20110 combination therapies in patients with advanced colorectal cancer. The interventions include HS-20110 combined with Bevacizumab and various chemotherapy regimens, which is important for assessing potential treatment options for this patient population. The trial is currently in Phase Ib/II and is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as BRAF and MSI, which may be relevant for patient selection.

NCT: NCT07283367 · Enrollment: 502 ·

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

IMMUNOTHERAPY

• RECRUITING

⚡ 4

A FIH, Phase I/IIa, Trial Assessing Feasibility of Administrations of TIL-based Immunotherapy in Patients With Metastatic CRC and PC

This clinical trial is testing the feasibility of TIL-based immunotherapy in patients with metastatic colorectal cancer (CRC) and pancreatic cancer (PC). The intervention involves the administration of a novel autologous TIL product, CC-38, along with pembrolizumab, cyclophosphamide, interleukin-2, and uromitexan, which is important for evaluating safety and tolerability. Currently in Phase I/IIa and actively recruiting, this trial aims to assess the repeated administration of CC-38 in a small cohort of 12 participants. Eligibility criteria include the presence of the BRAF biomarker.

NCT: NCT07255664 · Enrollment: 12 · Sites: Frankfurt, Germany

MEDIUM

PHASE2

All Stages

MSI/MMR

LIQUID BIOPSY

• RECRUITING

⚡ 4

A Study of Botensilimab and Balstilimab for Colorectal Cancer With ctDNA+ After Surgery and Chemotherapy

This clinical trial is testing the combination of botensilimab and balstilimab for patients with microsatellite stable (MSS) colorectal cancer or colorectal liver metastases (CRLM) who have measurable residual disease (MRD) after surgery and chemotherapy. The intervention involves administering botensilimab and balstilimab, followed by balstilimab alone, to evaluate its effectiveness in achieving ctDNA clearance. This is a Phase 2 trial currently recruiting 284 participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, ctDNA, RAS, and NGS.

NCT: NCT07227636 · Enrollment: 284 · Sites: Basking Ridge, Commack, Harrison, Middletown, Montvale

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

⚡ 4

A Phase II Trial of Neoadjuvant PD-1 Vaccine PD1-Vaxx in Operable MSI High Colorectal Cancer

This clinical trial is testing the neoadjuvant PD-1 vaccine IMU-201 (PD1-Vaxx) in patients with operable microsatellite instability-high (MSI-H) colorectal cancer. The intervention involves administering three doses of the PD1-Vaxx prior to surgical resection, with the goal of achieving a Major Pathological Response (MPR) defined as less than 10% viable tumor cells. This is a Phase II trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the study's findings. Eligibility criteria include having operable MSI-H colorectal cancer, with specific biomarker requirements such as MSI, MSI-H, MMR, and NGS.

NCT: NCT06692959 · Enrollment: 44 · Sites: Adelaide, Australia, Guildford, United Kingdom, Manchester, United Kingdom, Perth, Australia

SIGNAL

NA

Stage IV

RESEARCH

• RECRUITING

⚡ 4

Randomized Controlled Trial of the Effects of Combined Resistance and Aerobic Exercise on Health-related Quality of Life in Patients Undergoing First-line Chemotherapy for Metastatic Colorectal Cancer (REACH)

The REACH trial is testing the effects of combined resistance and aerobic exercise on health-related quality of life in patients undergoing first-line chemotherapy for metastatic colorectal cancer. The intervention involves 18 weeks of structured exercise training, which aims to assess its impact on the quality of life for these patients. This trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate. Eligibility criteria include being diagnosed with unresectable metastatic colorectal cancer and undergoing first-line chemotherapy.

NCT: NCT06938971 · Enrollment: 150 · Sites: Copenhagen, Denmark, Herlev, Denmark, Hillerød, Denmark, Roskilde, Denmark

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 4

Short-course Radiotherapy Followed by CAPOX and Ivonescimab for Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy and safety of preoperative short-course radiotherapy combined with ivonescimab and chemotherapy in patients with locally advanced rectal cancer. The intervention involves the drug ivonescimab, which is being evaluated for its potential to improve treatment outcomes when used alongside standard therapies. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06760520 · Enrollment: 40 ·

SIGNAL

PHASE1

Stage IV

KRAS

LIQUID BIOPSY

• RECRUITING

⚡ 4

A First-in-human Study of BGB-53038, a Pan-KRAS Inhibitor, Alone or in Combinations in Participants With Advanced or Metastatic Solid Tumors With KRAS Mutations or Amplification

This clinical trial is testing BGB-53038, a pan-KRAS inhibitor, in participants with advanced or metastatic solid tumors that have KRAS mutations or amplification. The intervention includes BGB-53038 alone and in combination with tislelizumab or cetuximab, which is significant for understanding its safety and potential effectiveness in treating these specific cancers. Currently in Phase 1 and actively recruiting, this trial aims to assess the safety, tolerability, and preliminary antitumor activity of the drug, which is crucial for determining the next steps in treatment options for eligible patients. Participants must have tumors with KRAS mutations or amplification, and the study includes a liquid biopsy component for biomarker assessment.

NCT: NCT06585488 · Enrollment: 514 · Sites: Baltimore, Houston, Kansas City, Los Angeles

MEDIUM

NA

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

⚡ 4

Study on the Safety and Efficacy of MR-Linac Technique in Patients With Unresectable Locally Advanced Colon Cancer

This clinical trial is testing the safety and efficacy of the MR-Linac technique in patients with unresectable locally advanced colon cancer. The intervention involves the use of MR-Linac for short course radiotherapy, followed by chemotherapy and surgical resection, which is significant for potentially improving treatment outcomes in this patient population. The study is currently in the recruiting phase and aims to enroll 20 participants, indicating that eligible patients may still join the trial. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and dMMR.

NCT: NCT06244537 · Enrollment: 20 · Sites: Chengdu, China

MEDIUM

PHASE1, PHASE2

Stage IV

IMMUNOTHERAPY

• RECRUITING

⚡ 4

GM103 Intratumoral Injection in Patients With Locally Advanced, Unresectable, Refractory and/or Metastatic Solid Tumors

This clinical trial is testing the safety and efficacy of GM103, administered alone and in combination with pembrolizumab, in patients with locally advanced, unresectable, refractory, and/or metastatic solid tumors. The intervention involves the intratumoral injection of GM103, which is being evaluated for its potential to improve treatment outcomes in various cancer types, including colorectal cancer. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include having specific types of solid tumors that are advanced and difficult to treat.

NCT: NCT06265025 · Enrollment: 125 · Sites: Goyang-si, South Korea, Seoul, South Korea

MEDIUM

PHASE1, PHASE2

Stage IV

KRAS

TARGETED THERAPY

• RECRUITING

⚡ 4

Study of Elironrasib and Daraxonrasib as Monotherapies and Combination Therapy in Participants With Advanced KRAS G12C Mutant Solid Tumors

This clinical trial is evaluating the safety and tolerability of Elironrasib and Daraxonrasib, both as monotherapies and in combination, for participants with advanced KRAS G12C mutant solid tumors. The intervention involves two drugs, Elironrasib and Daraxonrasib, which are being tested to understand their pharmacokinetics and potential effects on this specific mutation. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive these treatments. Eligibility criteria include having tumors with KRAS G12C mutations, among other potential biomarker requirements.

NCT: NCT06128551 · Enrollment: 534 · Sites: Aurora, Boston, Dallas, Detroit, Duarte

MEDIUM

PHASE1, PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

⚡ 4

Cadonilimab for PD-1/PD-L1 Blockade-refractory, MSI-H/dMMR, Advanced Colorectal Cancer

This clinical trial is testing the efficacy of cadonilimab in patients with advanced colorectal cancer who are refractory to PD-1/PD-L1 blockade and have microsatellite instability-high (MSI-H) or mismatch repair-deficient (dMMR) tumors. The intervention, cadonilimab, is a drug that targets the immune checkpoint pathway, which is significant for patients who have not responded to other therapies. This trial is currently in Phase 1 and Phase 2 and is actively recruiting participants, indicating that eligible patients may have the opportunity to access this treatment option. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite status.

NCT: NCT05426005 · Enrollment: 28 · Sites: Guangzhou, China

SIGNAL **N/A** **Stage III** **ctDNA** **LIQUID BIOPSY** • **RECRUITING**

⚡ 4

ctDNA Methylation Application in Postoperative Relapse and Adjuvant Chemotherapy Efficacy Evaluation

This clinical trial is testing a multi-locus blood-based assay targeting circulating tumor DNA (ctDNA) methylation in patients with resected stage I and stage II colorectal cancer who are at low risk for recurrence. The intervention aims to monitor postoperative relapse and evaluate the efficacy of adjuvant chemotherapy, which is important for optimizing treatment strategies in this patient population. The trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate. Eligibility criteria include having undergone radical resection for stage I or stage II colorectal cancer without high-risk features.

NCT: NCT05536089 · Enrollment: 2000 · Sites: Hangzhou, China

MEDIUM **PHASE1** **All Stages** **MSI/MMR** **IMMUNOTHERAPY** • **RECRUITING**

⚡ 4

Study of Zanzalintinib in Combination With Immuno-Oncology Agents in Participants With Solid Tumors

This clinical trial is evaluating the safety and efficacy of zanzalintinib, both as a monotherapy and in combination with immuno-oncology agents, in participants with advanced solid tumors. The intervention involves zanzalintinib, nivolumab, and ipilimumab, which are being tested to assess their safety, tolerability, and preliminary antitumor activity. This is a Phase 1b trial currently recruiting participants, which means that it is in the early stages of testing to determine the best dosing and safety profile before larger studies can be conducted. Eligibility criteria include the presence of specific biomarkers, such as microsatellite instability.

NCT: NCT05176483 · Enrollment: 1314 · Sites: Austin, Baltimore, Boston, Celebration, Charlottesville

MEDIUM **PHASE2** **Stage III** **MSI/MMR** **TARGETED THERAPY** • **ACTIVE_NOT_RECRUITING**

⚡ 4

Atezolizumab in Patients With MSI-h/MMR-D Stage II High Risk and Stage III Colorectal Cancer Ineligible for Oxaliplatin

This clinical trial is testing the efficacy of atezolizumab in patients with stage II high-risk and stage III colorectal cancer who are ineligible for oxaliplatin-based chemotherapy. The intervention involves administering atezolizumab, a drug that targets the immune system, alongside IMM-101, to potentially enhance disease-free survival in this specific patient population. Currently in Phase 2 and marked as active but not recruiting, this trial indicates that patients may not be able to enroll at this time but can still benefit from the findings once available. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS, which are essential for determining patient suitability for the treatment.

NCT: NCT05118724 · Enrollment: 80 · Sites: Essen, Germany

MEDIUM **PHASE1** **All Stages** **HER2** **TARGETED THERAPY** • **ACTIVE_NOT_RECRUITING**

⚡ 4

CAR-macrophages for the Treatment of HER2 Overexpressing Solid Tumors

This clinical trial is testing the safety and tolerability of CAR-macrophages in patients with HER2 overexpressing solid tumors. The intervention involves the use of CT-0508 and Pembrolizumab, which are biological agents aimed at targeting and treating these specific tumors. Currently in Phase 1 and marked as active but not recruiting, this trial is focused on evaluating adverse events in the enrolled participants. Eligibility criteria include the requirement for tumors to overexpress the HER2 biomarker.

NCT: NCT04660929 · Enrollment: 48 · Sites: Chapel Hill, Duarte, Houston, Nashville, Philadelphia

MEDIUM **PHASE1, PHASE2** **Stage IV** **IMMUNOTHERAPY** • **ACTIVE_NOT_RECRUITING**

⚡ 4

Safety, PK and Efficacy of ONC-392 in Monotherapy and in Combination of Anti-PD-1 in Advanced Solid Tumors and NSCLC

This clinical trial is testing the safety, pharmacokinetics, and efficacy of ONC-392, a humanized anti-CTLA4 IgG1 monoclonal antibody, in patients with advanced or metastatic solid tumors and non-small cell lung cancer (NSCLC). The intervention involves administering ONC-392 both as a monotherapy and in combination with pembrolizumab and docetaxel, which is significant for understanding its potential effectiveness in these cancer types. Currently, the trial is in Phase 1 and Phase 2 and is active but not recruiting, indicating that patients cannot enroll at this time. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

SIGNAL **NA** **Stage III** **ctDNA** **LIQUID BIOPSY** • ACTIVE_NOT_RECRUITING

⚡ 4

ctDNA-guided Surveillance for Stage III CRC, a Randomized Intervention Trial

The IMPROVE-IT2 trial is testing ctDNA-guided surveillance for patients with Stage III colorectal cancer (CRC) to evaluate its effectiveness compared to standard CT-scan surveillance. The intervention involves ctDNA analysis as a diagnostic test, which aims to enhance the detection of recurrent disease and potentially increase the number of patients eligible for curative treatment. This trial is currently active but not recruiting participants, indicating that it is ongoing and data is being collected from enrolled patients. Eligibility criteria include the presence of ctDNA as a biomarker for assessment.

NCT: NCT04084249 · Enrollment: 359 · Sites: Aalborg, Denmark, Aarhus, Denmark, Copenhagen, Denmark, Herlev, Denmark, Herning, Denmark

MEDIUM **PHASE1** **Stage IV** **RAS** **TARGETED THERAPY** • ACTIVE_NOT_RECRUITING

⚡ 4

Naptumomab Estafenatox in Combination With Durvalumab in Subjects With Selected Advanced or Metastatic Solid Tumor, Including a Cohort Expansion in Esophageal Cancer.

This clinical trial is testing the combination of naptumomab estafenatox and durvalumab in patients with selected advanced or metastatic solid tumors, including a cohort expansion specifically for esophageal cancer. The intervention involves a combination of these two agents to evaluate their safety and tolerability, which is important for determining effective treatment options for this patient population. Currently, the trial is in Phase 1b and is active but not recruiting, meaning that while it is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as HER2, RAS, and NGS, which may be relevant for patient selection.

NCT: NCT03983954 · Enrollment: 120 · Sites: Ahmedabad, India, Haifa, Israel, Hyderabad, India, Jhajjar, India, New Delhi, India

MEDIUM **PHASE2** **Stage IV** **PHASE II TRIAL** • ACTIVE_NOT_RECRUITING

⚡ 4

Microwave Ablation Versus Stereotactic Body Radiotherapy for Colorectal Liver Metastases in Oligometastatic Disease: a Prospective, Randomised, Phase 2 Trial

The LAVA-CRLM trial is testing the effectiveness of microwave ablation (MWA) compared to stereotactic body radiotherapy (SBRT) in patients with colorectal liver metastases and oligometastatic disease. The intervention involves using MWA, a device-based treatment, and SBRT, a form of radiation therapy, to assess their impact on local control and safety in this patient population. This is a Phase 2 trial that is currently active but not recruiting, indicating that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria for participation may include specific characteristics related to the patient's metastatic disease status.

NCT: NCT03654131 · Enrollment: 100 · Sites: Copenhagen, Denmark

Appendix A — Patient Guide to Active Trials

For patients, caregivers, and family members. Always discuss with your oncologist before considering a clinical trial.

What is a clinical trial?

Clinical trials test new treatments, combinations, or approaches to care. Participating can give you access to treatments not yet widely available. Trials are listed by the biomarker your tumor must have — because many newer treatments only work for specific mutations. Ask your oncologist which biomarkers your tumor has been tested for, then look for matching trials below.

MSI/MMR

Botensilimab + Balstilimab vs Best Supportive Care as Therapy in Chemo-refractory, Unresectable, Colorectal Adenocarcinoma

Phase: PHASE3 · Status: RECRUITING · [NCT07152821](#)

This clinical trial is testing the combination of botensilimab and balstilimab in patients with chemo-refractory, unresectable colorectal adenocarcinoma. The intervention involves two immunotherapy drugs aimed at potentially improving overall survival and quality of life for participants. Currently in Phase 3 and actively recruiting, this trial may provide insights into the effectiveness and safety of these treatments for eligible patients. Participants may need to meet specific biomarker criteria, including MSI, MSI-H, MMR, DMMR, and microsatellite status.

BRAF

Symbiotic-GI-03: A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Chemotherapy in Adult Participants With Metastatic Colorectal Cancer

Phase: PHASE3 · Status: RECRUITING · [NCT07222800](#)

The Symbiotic-GI-03 trial is investigating the efficacy of the study medicine PF-08634404 in combination with chemotherapy for adult participants with metastatic colorectal cancer. This trial aims to determine how well PF-08634404 works alongside standard chemotherapy and Bevacizumab, a biological therapy, in patients whose cancer has spread or returned after previous treatments. Currently in Phase 3 and actively recruiting, this trial offers eligible patients the opportunity to receive potentially beneficial treatment options while contributing to research on colorectal cancer therapies. To participate, individuals must be 18 years or older, have metastatic colorectal cancer, and meet specific health criteria, including not being pregnant prior to treatment.

KRAS

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

Phase: PHASE3 · Status: RECRUITING · [NCT06750094](#)

This clinical trial is testing the efficacy of amivantamab combined with FOLFIRI chemotherapy in patients with KRAS/NRAS and BRAF wild-type colorectal cancer who have previously undergone chemotherapy. The intervention involves the biological agent amivantamab alongside standard chemotherapy agents, which is important for evaluating its potential benefits compared to existing treatment options. This is a Phase 3 trial currently recruiting 700 participants, which means that eligible patients may have the opportunity to receive a treatment that is being compared to standard therapies in a larger population. Eligibility criteria include specific biomarker requirements related to KRAS, NRAS, and BRAF status, as well as other factors such as microsatellite instability (MSI) and mismatch repair (MMR) status.

HER2

A Study of Tucatinib With Trastuzumab and mFOLFOX6 Versus Standard of Care Treatment in First-line HER2+ Metastatic Colorectal Cancer

Phase: PHASE3 · Status: RECRUITING · [NCT05253651](#)

This clinical trial is testing the efficacy of tucatinib in combination with trastuzumab and mFOLFOX6 compared to standard of care treatments for patients with HER2 positive metastatic colorectal cancer. The intervention involves the use of tucatinib, a targeted therapy, alongside other established cancer drugs, to evaluate its effectiveness and side effects in this patient population. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into progression-free survival outcomes for participants. Eligibility criteria include having HER2 positive colorectal cancer that is metastatic or unresectable, with specific biomarker assessments for HER2 and RAS.

RAS

Neoadjuvant FOLFOX Chemotherapy for Patients With Locally Advanced Colon Cancer

Phase: PHASE3 · Status: ACTIVE_NOT_RECRUITING · [NCT03426904](#)

This clinical trial is testing the effectiveness of neoadjuvant FOLFOX chemotherapy in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy before surgery, which may enhance treatment outcomes by targeting microscopic metastatic cancer cells earlier and reducing surgical complications. This is a Phase 3 trial

that is currently active but not recruiting, indicating that it is in the final stages of testing and results may soon be available for clinical application. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

ctDNA

ctDNA Methylation Application in Postoperative Relapse and Adjuvant Chemotherapy Efficacy Evaluation

Phase: N/A · Status: RECRUITING · [NCT05536089](#)

This clinical trial is testing a multi-locus blood-based assay targeting circulating tumor DNA (ctDNA) methylation in patients with resected stage I and stage II colorectal cancer who are at low risk for recurrence. The intervention aims to monitor postoperative relapse and evaluate the efficacy of adjuvant chemotherapy, which is important for optimizing treatment strategies in this patient population. The trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate. Eligibility criteria include having undergone radical resection for stage I or stage II colorectal cancer without high-risk features.

NGS

Prospective Collection of Samples to Enable the Development of Natera Screening Assay for Early Cancer Detection

Phase: N/A · Status: RECRUITING · [NCT06620627](#)

The PROCEED-CRC study is recruiting participants at average risk for colorectal cancer to prospectively collect blood samples. The primary aim is to facilitate the development of a blood-based screening assay for early cancer detection.

What this means for you

Ask your oncologist whether any active trials match your biomarker profile and disease stage. The most actively trialed biomarkers in colorectal cancer right now are **MSI/MMR, BRAF, KRAS, HER2, and RAS**. If your tumor has not been tested for these markers, ask your oncologist before your next treatment decision — biomarker results can determine which trials you qualify for.

Questions to ask your oncologist about trials

"Am I eligible for any active Phase III trials based on my biomarker profile?"

"Does my tumor need to be retested before I can qualify for a biomarker-directed trial?"

"What are the travel requirements and visit frequency for the trials I might qualify for?"

Source: ClinicalTrials.gov · 2026-04-25 23:31:19

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Clinical Trial Reference Guide

What the phases mean

Phase I — First testing in humans. Focus on safety and dosing. Small groups.

Phase II — Testing effectiveness. Larger groups. Determines if Phase III is warranted.

Phase III — Randomized comparison against standard treatment. Required for FDA approval.

Key biomarkers that determine trial eligibility

RAS (KRAS/NRAS) — mutation status affects anti-EGFR eligibility and KRAS G12C trial eligibility

BRAF V600E — required for encorafenib-based trials (~8–10% of CRC)

MSI-H/dMMR — required for immunotherapy trials; ~5% of mCRC, ~15% of Stage II/III

HER2 — required for HER2-directed trials (~2–3% of CRC)

NTRK fusion — required for TRK inhibitor trials (<1% of CRC)

KRAS G12C — required for sotorasib/adagrasib trials (~3–4% of CRC)

TMB-H — tumor mutational burden; may qualify for immunotherapy

ctDNA — liquid biopsy; used in MRD monitoring trials

How to find trials you may qualify for

1. Get complete biomarker testing: RAS, BRAF, MSI/MMR, HER2, NTRK, TMB
2. Search ClinicalTrials.gov with your biomarker and stage

3. Ask your oncologist about NCI-designated cancer centers with trial access

4. Contact the Colorectal Cancer Alliance: ccalliance.org

Key Resources

[ClinicalTrials.gov](https://clinicaltrials.gov) · [NCI Cancer Information \(cancer.gov\)](https://cancer.gov) · [Colorectal Cancer Alliance \(ccalliance.org\)](https://ccalliance.org) · [Fight Colorectal Cancer \(fightcrc.org\)](https://fightcrc.org)

Colon & Colorectal Cancer — Clinical Trials Intelligence

About This Report

This report is generated automatically from publicly available medical databases including PubMed, ClinicalTrials.gov, and the FDA Drug@FDA database. Entries are classified using deterministic rule-based logic and scored by clinical significance. AI-generated summaries are grounded exclusively in the source data provided.

Limitations

This report does not represent a complete picture of all published research or approved treatments. Database coverage, feed availability, and relevance filtering affect what appears. New drug approvals and trial status changes may not be reflected immediately. Always verify current information at the original source before making clinical decisions.

Disclaimer

Trial data is sourced from ClinicalTrials.gov, a registry maintained by the U.S. National Library of Medicine. Entry summaries are AI-generated from publicly available trial records. Trial status and eligibility criteria may change — always verify at ClinicalTrials.gov before acting on this information. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

Copyright

© 2026 Randy Hinton. All rights reserved. This report may be shared freely for informational purposes. It may not be modified, republished, or presented as original work without attribution.

https://petroai1492.github.io/delta/timelines/coloncancer_clinicaltrials_timeline.pdf