

Optimization

Pieter Vanderschueren

Year 2024-2025

Contents

Mathematical Preliminaries	2
1 Vectors	3
2 Independence, Subspaces, Basis and Dimension	3
3 Orthogonality and Orthogonal Complements	3
4 Matrices	3
5 Sequences	3
6 Differential Calculus	3
Fundamental Concepts	4
7 Fundamental Concepts of Optimization	5
8 Types of Optimization Problems	6
9 Convex Optimization	9
10 The Lagrangian Function and Duality	14
Unconstrained Optimization Algorithms	18
11 Optimality Conditions	19
12 Estimation and Fitting Problems	21
13 Newton Type Optimization	22
14 Globalisation Strategies	22
15 Calculating Derivatives	22
Constrained Optimization Algorithms	23
16 Optimality Conditions for Constrained Optimization	24
17 Equality Constrained Optimization Algorithms	24
18 Inequality Constrained Optimization Algorithms	24
19 Optimal Control Problems	24

Mathematical Preliminaries

- 1 Vectors
- 2 Independence, Subspaces, Basis and Dimension
- 3 Orthogonality and Orthogonal Complements
- 4 Matrices
- 5 Sequences
- 6 Differential Calculus

Fundamental Concepts

7 Fundamental Concepts of Optimization

Theorem 7.1: Optimization problem in standard form

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && f(x) \\ & \text{subject to} && g(x) = 0, \\ & && h(x) \geq 0. \end{aligned}$$

Theorem 7.2: Level Set

The set

$$\{ x \in \mathbb{R}^n \mid f(x) = c \}$$

is the level set of f for the value c , i.e. the set of all points that map to the same value c .

Theorem 7.3: Feasible set

The set

$$\{ x \in \mathbb{R}^n \mid g(x) = 0 \wedge h(x) \geq 0 \}$$

is the feasible set Ω , i.e. the set of all points that satisfy the constraints.

Theorem 7.4: Global minimizer

The point x^* is a global minimizer if and only if

$$x^* \in \Omega, \forall x \in \Omega : f(x^*) \leq f(x).$$

Theorem 7.5: Strict global minimizer

The point x^* is a strict global minimizer if and only if

$$x^* \in \Omega, \forall x \in \Omega \setminus \{x^*\} : f(x^*) < f(x).$$

Theorem 7.6: Strict local minimizer

The point x^* is a strict local minimizer if and only if $x^* \in \Omega$ and there exists a neighborhood \mathcal{N} of x^* such that

$$\forall x \in (\Omega \cap \mathcal{N}) \setminus \{x^*\} : f(x^*) < f(x).$$

Theorem 7.7: Weierstrass

If $\Omega \in \mathbb{R}^n$ is compact, i.e. limited and closed, and $f : \Omega \rightarrow \mathbb{R}$ is continuous, then there exists a global minimizer (a solution) of the optimization problem

$$\begin{array}{ll} \text{minimize} & f(x) \\ & x \in \mathbb{R}^n \\ \text{subject to} & x \in \Omega. \end{array}$$

Proof 7.1: Weierstrass

Regard the graph of f , $G = \{(x, f(x)) \in \mathbb{R}^n \times \mathbb{R} \mid x \in \Omega\}$. G is a compact set, and so is the projection of G onto its last coordinate, the set $G = \{s \in \mathbb{R} \mid \exists x \text{ such that } (x, s) \in G\}$, which is a compact interval $[f_{\min}, f_{\max}] \subset \mathbb{R}$. By construction, there must be at least one x^* such that $(x^*, f(x^*)) \in G$. □

8 Types of Optimization Problems

Theorem 8.1: Convex set

A set $\Omega \subset \mathbb{R}^n$ is convex if

$$\forall x, y \in \Omega, \forall \lambda \in [0, 1] : x + \lambda(y - x) \in \Omega.$$

or if “all connecting lines lie inside the set”.

Theorem 8.2: Convex function

A function $f : \Omega \rightarrow \mathbb{R}$ is convex if

$$\forall x, y \in \Omega, \forall \lambda \in [0, 1] : f(x + \lambda(y - x)) \leq f(x) + \lambda(f(y) - f(x)).$$

or if “all secants (i.e. a line segment between two points on the graph) are above graph”. This definition is equivalent to saying that the Epigraph of f , i.e. the set $\{(x, s) \in \mathbb{R}^n \times \mathbb{R} \mid x \in \Omega, s \geq f(x)\}$, is a convex set.

Note: a concave function is the same but then with \geq instead of \leq .

Property 8.1: Convex function

If $f : D \rightarrow \mathbb{R}$ and $\Omega_f = \{(x, y) \mid x \in D, y \geq f(x)\}$ then the following holds:

$$f \text{ is convex} \Leftrightarrow \Omega_f \text{ is convex.}$$

Theorem 8.3: Convex optimization problem

An optimization problem with convex feasible set Ω and convex objective function $f : \Omega \rightarrow \mathbb{R}$ is called a convex optimization problem.

Property 8.2: Globality of local minima of convex optimization problem

For a convex optimization problem, a local minimum is also a global one.

Proof 8.1: Globality of local minima of convex optimization problem

Regard a local minimum x^* of the convex optimization problem


$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && f(x) \\ & \text{subject to} && x \in \Omega. \end{aligned}$$

We will show that for any point $y \in \Omega$ holds $f(y) \geq f(x^*)$, i.e. x^* is a global minimum.

First we choose, using local optimality, a neighborhood \mathcal{N} of x^* such that for all $\tilde{x} \in \Omega \cap \mathcal{N}$ holds $f(\tilde{x}) \geq f(x^*)$. Second, we regard the connecting line between x^* and y . This line is completely contained in Ω due to the convexity of Ω . Now we choose a point \tilde{x} on this line such that it is in the neighborhood, but not equal to x^* , i.e. $\tilde{x} = x^* + \lambda(y - x^*)$ for some $\lambda \in (0, 1)$, and $\tilde{x} \in \Omega \cap \mathcal{N}$. Due to local optimality, we have $f(\tilde{x}) \geq f(x^*)$, and due to convexity we have

$$f(\tilde{x}) = f(x^* + \lambda(y - x^*)) \leq f(x^*) + \lambda(f(y) - f(x^*)).$$

It follows that $\lambda(f(y) - f(x^*)) \geq 0$, and since $\lambda \in (0, 1)$, we have $f(y) \geq f(x^*)$, as desired. □



Theorem 8.4: Unconstrained optimization problem

An optimization problem with no constraints, i.e. $g(x) = 0$ and $h(x) \geq 0$ are empty, is called an unconstrained optimization problem.

Example 8.1: Nonlinear Programming (NLP)

Nonlinear Programming Problems are problems of the following form:

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && f(x) \\ & \text{subject to} && g(x) = 0, \\ & && h(x) \geq 0, \end{aligned}$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $g : \mathbb{R}^n \rightarrow \mathbb{R}^p$, and $h : \mathbb{R}^n \rightarrow \mathbb{R}^q$, are assumed to be continuously differentiable at least once.

Example 8.2: Linear Programming (LP)

When the functions f, g, h are affine (i.e., they can be expressed in the form $f(x) = a^T x + b$ for some vector a and scalar b) in the general formulation (see Theorem 8.1), the general NLP gets something easier to solve, namely a Linear Program (LP), which can be written as follows:

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && c^T x \\ & \text{subject to} && Ax - b = 0, \\ & && Cx - d \geq 0, \end{aligned}$$

where $c \in \mathbb{R}^n$, $A \in \mathbb{R}^{p \times n}$, $b \in \mathbb{R}^p$, $C \in \mathbb{R}^{q \times n}$, and $d \in \mathbb{R}^q$.

Example 8.3: Quadratic Programming (QP)

When the functions g, h are affine (as for an LP in Theorem 8.2), and the objective function f is a linear-quadratic function, the NLP becomes a Quadratic Program (QP), which can be written as follows:

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && c^T x + \frac{1}{2} x^T B x \\ & \text{subject to} && Ax - b = 0, \\ & && Cx - d \geq 0, \end{aligned}$$

where, in addition to the LP parameters, $B \in \mathbb{R}^{n \times n}$ is the Hessian matrix. Specifically, for the objective function $f(x) = c^T x + \frac{1}{2} x^T B x$, the Hessian is given by:

$$\nabla^2 f(x) = B.$$

Theorem 8.5: Convex QP

If the Hessian matrix B is positive semi-definite (i.e. if $\forall z \in \mathbb{R}^n \setminus \{0\} : z^T B z \geq 0$) we call the QP a convex QP. Convex QPs are tremendously easier to solve globally than “non-convex QPs” (i.e., where the Hessian B is not positive semi-definite), which might have different local minima (i.e. have a non-convex solution set).

Theorem 8.6: Strictly convex QP


If the Hessian matrix B is positive definite (i.e. if $\forall z \in \mathbb{R}^n : z^T B z > 0$) we call the QP a strictly convex QP. Strictly convex QPs are a subset of convex QPs (see Theorem 8.5).

Example 8.4: Mixed-Integer Programming (MIP)

A mixed-integer programming problem or mixed-integer program (MIP) is a problem with both real and integer decision variables. A MIP can be formulated as follows:

$$\begin{aligned} & \underset{\substack{x \in \mathbb{R}^n \\ z \in \mathbb{Z}^m}}{\text{minimize}} && f(x, z) \\ & \text{subject to} && g(x, z) = 0, \\ & && h(x, z) \geq 0, \end{aligned}$$

where $x \in \mathbb{R}^n$ are the continuous variables, and $z \in \mathbb{Z}^m$ are the integer variables.




9 Convex Optimization

Theorem 9.1: Convexity for C^1 functions

Assume that $f : \Omega \rightarrow \mathbb{R}$ is continuously differentiable and Ω is convex. Then holds that f is convex if and only if

$$\forall x, y \in \Omega : f(y) \geq f(x) + \nabla f(x)^T (y - x)$$

i.e. tangents lie below the graph.



Proof 9.1: Convexity for C^1 functions

“ \Rightarrow ”: Due to convexity of f holds for given $x, y \in \Omega$ and for any $\lambda \in [0, 1]$ that

$$f(x + \lambda(y - x)) - f(x) \leq \lambda(f(y) - f(x)).$$

and therefore that

$$\nabla f(x)^T (y - x) = \lim_{\lambda \rightarrow 0} \frac{f(x + \lambda(y - x)) - f(x)}{\lambda} \leq \frac{f(y) - f(x)}{y - x}.$$

“ \Leftarrow ”: To prove that $z = x + \lambda(y - x) = (1 - \lambda)x + \lambda y$ holds that $f(z) \leq (1 - \lambda)f(x) + \lambda f(y)$, we can use the equation from Theorem 9.1 twice to get

$$f(x) \geq f(z) + \nabla f(z)^T (x - z) \quad \text{and} \quad f(y) \geq f(z) + \nabla f(z)^T (y - z),$$

which yield, when weighted with $(1 - \lambda)$ and λ respectively, that

$$(1 - \lambda)f(x) + \lambda f(y) \geq f(z) + \underbrace{\nabla f(z)^T [(1 - \lambda)(x - z) + \lambda(y - z)]}_{=0}$$

□

Theorem 9.2: Generalized Inequality for Symmetric Matrices

We write for a symmetric matrix $B = B^T$, $B \in \mathbb{R}^{n \times n}$ that “ $B \succeq 0$ ” if and only if B is positive semi-definite, i.e.,

$$\forall z \in \mathbb{R}^n : z^T B z \geq 0,$$

or, equivalently, if all (real) eigenvalues of B are non-negative.

Theorem 9.3: $O(\cdot)$

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, we write

$$f(x) = O(g(x))$$

if and only if there exists a constant $C > 0$ and a neighborhood \mathcal{N} of 0 such that

$$\forall x \in \mathcal{N} : \|f(x)\| \leq Cg(x),$$

i.e. “ f shrinks as fast as g ”.

Theorem 9.4: $o(\cdot)$

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, we write

$$f(x) = o(g(x))$$

if and only if there exists a neighborhood \mathcal{N} of 0 and a function $c : \mathcal{N} \rightarrow \mathbb{R}$ with $\lim_{x \rightarrow 0} c(x) = 0$ such that

$$\forall x \in \mathcal{N} : \|f(x)\| \leq c(x)g(x),$$

i.e. “ f shrinks faster than g ”.

Theorem 9.5: Convexity for C^2 functions

Assume that $f : \Omega \rightarrow \mathbb{R}$ is twice continuously differentiable and Ω is convex and open. Then holds that f is convex if and only if

$$\forall x \in \Omega : \nabla^2 f(x) \succeq 0,$$

i.e. the Hessian of f is positive semi-definite.

Proof 9.2: Convexity for C^2 functions

To prove that the Theorem 9.1 implies the Theorem 9.5, we can use a second order Taylor expansion of y at x in an arbitrary direction p :

$$f(x + tp) = f(x) + t\nabla f(x)^T p + \frac{t^2}{2} p^T \nabla^2 f(x) p + o(t^2 \|p\|).$$

From this we obtain that

$$p^T \nabla^2 f(x) p = \lim_{t \rightarrow 0} \frac{2}{t^2} \underbrace{\left(f(x + tp) - f(x) - t \nabla f(x)^T p \right)}_{(9.1): \geq 0} \geq 0.$$

Conversely, to prove the other direction, we use the Taylor rest term formula with some arbitrary $\theta \in (0, 1)$:

$$f(y) = f(x) + \nabla f(x)^T (y - x) + \underbrace{\frac{1}{2} t^2 (y - x)^T \nabla^2 f(x + \theta(y - x)) (y - x)}_{(9.5): \geq 0}.$$

□

Property 9.1: Convexity perserving operations on convex functions

The following operations preserve the convexity of a function:

1. Non-negative weighted sum: Suppose that $\forall i \in [1, m] : f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are convex functions and $\forall i \in [1, m] : \lambda_i \geq 0$. Then the function

$$f(x) = \sum_{i=1}^m \lambda_i f_i(x)$$

is convex.

2. Affine input transformation: If $f : \Omega \rightarrow \mathbb{R}$ is convex, then also

$$A \in \mathbb{R}^{n \times m} : \tilde{f}(x) = f(Ax + b)$$

is convex on the domain $\tilde{\Omega} = \{x \mid Ax + b \in \Omega\}$.

3. Concatenation with monotone convex function: If $f : \Omega \rightarrow \mathbb{R}$ is convex and $g : \mathbb{R} \rightarrow \mathbb{R}$ is convex and monotonely increasing, then the composition $g \circ f$ is convex.
4. Pointwise supremum: The supremum over a set of convex functions $f_i(x)$, $i \in I$, where I can be an infinite set, i.e.,

$$f(x) = \sup_{i \in I} f_i(x)$$

is convex.

5. Composition: Let $h : \mathbb{R}^m \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $g = (g_1, \dots, g_m)$. Then $f(x) = (h \circ g)(x) = h(g(x))$ is convex if any of the followings holds:

- (a) h is convex and non-decreasing in each argument and $\forall i \in [1, m] : g_i$ is convex.
- (b) h is convex and non-decreasing in each argument and $\forall i \in [1, m] : g_i$ is concave, i.e. $-g_i$ is convex.

Proof 9.3: Convexity perserving operations on convex functions

1. TODO (use the definition of convexity).
2. TODO (use the definition of convexity).

3. Recall that g is a convex and monotonely increasing function, then:

$$\nabla^2(g \circ f) = \underbrace{g''(f(x))}_{\succeq 0} \underbrace{\nabla f(x) \nabla f(x)^T}_{\succeq 0} + \underbrace{g'(f(x))}_{\succeq 0} \underbrace{\nabla^2 f(x)}_{\succeq 0} \succeq 0,$$

i.e. $g \circ f$ is convex, since the Hessian is positive semi-definite.

4. Epigraph of f is the intersection of the epigraphs of f_i , which are convex.

5. Recall that g_i is convex and that h is convex and non-decreasing in each argument. Then:

$$\begin{aligned} f(\lambda x + (1 - \lambda)y) &= h(g(\lambda x + (1 - \lambda)y)) \\ &\leq h(\lambda g(x) + (1 - \lambda)g(y)) \end{aligned} \tag{1}$$

$$\begin{aligned} &\leq \lambda h(g(x)) + (1 - \lambda)h(g(y)) \tag{2} \\ &= \lambda f(x) + (1 - \lambda)f(y), \end{aligned}$$


where we used the convexity of g_i and the fact that h is non-decreasing to obtain inequality (1), and the convexity of h to obtain inequality (2). \square

Theorem 9.6: Convexity of Sublevel subsets

If $f : \Omega \rightarrow \mathbb{R}$ is a convex function, then all its level sets

$$\text{lev}_{\leq \gamma} f = \{ x \in \Omega \mid f(x) \leq \gamma \}$$

are convex.



Proof 9.4: Convexity of Sublevel subsets

If $f(x) \leq c$ and $f(y) \leq c$, then for any $\lambda \in [0, 1]$ holds also

$$f((1 - \lambda)x + \lambda y) \leq (1 - \lambda)f(x) + \lambda f(y) \leq \underbrace{(1 - \lambda)c + \lambda c}_{=c}.$$

\square

Property 9.2: Convexity perserving operations on convex sets

The following operations preserve the convexity of a set:

1. The intersection of finitely or infinitely many convex sets is convex.
2. Affine image: if Ω is convex, then for $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ also the set $A\Omega + b = \{y \in \mathbb{R}^m \mid \exists x \in \Omega : y = Ax + b\}$ is convex.
3. Affine pre-image: if Ω is convex, then for $A \in \mathbb{R}^{n \times m}$, $b \in \mathbb{R}^n$ also the set $\{z \in \mathbb{R}^m \mid Az + b \in \Omega\}$ is convex.

Example 9.1: Convex Feasible set

If $\forall i \in [1, m] : f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are convex functions, then the set

$$\Omega = \{x \in \mathbb{R}^n \mid f_i(x) \leq 0, i \in [1, m]\}$$

is a convex set, because it is the intersection of sublevel sets Ω_i of convex functions f_i , i.e.

$$\begin{aligned}\Omega &= \bigcap_{i=1}^m \Omega_i \\ &= \bigcap_{i=1}^m \{x \in \mathbb{R}^n \mid f_i(x) \leq 0\}.\end{aligned}$$

Theorem 9.7: Optimality condition for convex problems

Regard a convex optimization problem with continuously differentiable objective function f . A point $x^* \in \Omega$ is a global optimizer if and only if

$$\forall y \in \Omega : \nabla f(x^*)^T (y - x^*) \geq 0.$$

Proof 9.5: Optimality condition for convex problems

“ \Rightarrow ”: Assume for the sake of contradiction that $\exists y \in \Omega : \nabla f(x^*)^T (y - x^*) < 0$, then we could regard a Taylor expansion

$$f(x^* + \lambda(y - x^*)) = f(x^*) + \underbrace{\lambda \nabla f(x^*)^T (y - x^*)}_{<0} + \underbrace{o(\lambda)}_{\rightarrow 0}.$$

This implies that for sufficiently small positive λ , we have

$$f(x^* + \lambda(y - x^*)) < f(x^*),$$

which contradicts the optimality of x^* .

“ \Leftarrow ”: Due to the C^1 characterization of convexity of f in Theorem 9.1, we have for any feasible $y \in \Omega$:

$$f(y) \geq f(x^*) + \underbrace{\nabla f(x^*)^T (y - x^*)}_{\geq 0} \geq f(x^*),$$

which implies that x^* is a global optimizer. □

Theorem 9.8: Sufficient Condition for Convex NLP

For a nonlinear optimization problem (NLP) in standard form

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && f(x) \\ & \text{subject to} && g_i(x) = 0, i \in [1, m], \\ & && h_i(x) \geq 0, j \in [1, p], \end{aligned}$$

the following conditions are necessary for convexity:

- the objective function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ must be convex,
- the constraint set

$$X = \{ x \in \mathbb{R}^n \mid g(x) = 0, h(x) \geq 0 \}$$

must be convex. Since we know that the intersection of convex sets is convex, we can write X as the intersection of the sets G and H :

$$\begin{aligned} X &= \{x \in \mathbb{R}^n \mid g(x) = 0, h(x) \geq 0\} \\ &= \{x \in \mathbb{R}^n \mid g(x) = 0\} \cap \{x \in \mathbb{R}^n \mid h(x) \geq 0\} \\ &= \left(\bigcap_{i=1}^m \{x \in \mathbb{R}^n \mid g_i(x) = 0\} \right) \cap \left(\bigcap_{i=1}^m \{x \in \mathbb{R}^n \mid h_i(x) \geq 0\} \right) \\ &= G \cap H \end{aligned}$$

Now we must consider the requirements for the sets G and H to be convex:

- Suppose $-H_i = \{x \in \mathbb{R}^n \mid h_i(x) \leq 0\}$, the zero sublevel set of the function h_i . If h_i is convex, the set $-H_i$ is convex, as seen in Theorem 9.6. Now, consider the case where h_i is concave. Since h_i being concave means that $-h_i$ is convex, the set $H_i = \{x \in \mathbb{R}^n \mid h_i(x) \geq 0\}$ is a sublevel set of the convex function $-h_i$, and is therefore convex. Thus, the set H_i is convex when h_i is concave.
- On the other hand, G is the level set of g_i . Therefore it is certainly a convex set whenever g_i is a affine function

$$\forall i \in [i, m] : g_i(x) = a_i^T x + b_i.$$

TODO: examples

10 The Lagrangian Function and Duality

Theorem 10.1: Primal Optimization Problem

We will denote the globally optimal value of the objective function subject to the constraints as the primal optimal value p^* , i.e.,

$$p^* = \left(\min_{x \in \mathbb{R}^n} f(x) \quad \text{s.t.} \quad g(x) = 0 \wedge h(x) \geq 0 \right).$$

and we will denote this optimization problem as the primal optimization problem.

Theorem 10.2: Lagrangian Function and Lagrange Multipliers

We define the Lagrangian function to be

$$\begin{aligned}\mathcal{L}(x, \lambda, \mu) &= f(x) + \sum_{i=1}^l \lambda_i g_i(x) + \sum_{i=1}^m \mu_i h_i(x) \\ &= f(x) + \lambda^T g(x) + \mu^T h(x).\end{aligned}$$

Here, we have introduced Lagrange multipliers or dual variables $\lambda \in \mathbb{R}^l$ and strictly positive $\mu \in \mathbb{R}^m$.

Lemma 10.1: Lower Bound Property of Lagrangian

If \tilde{x} is a feasible point of (10.1) and $\mu \geq 0$, then

$$\mathcal{L}(\tilde{x}, \lambda, \mu) \leq f(\tilde{x}).$$

Proof 10.1: Lower Bound Property of Lagrangian

Since \tilde{x} is feasible, we have $g(\tilde{x}) = 0$ and $h(\tilde{x}) \geq 0$. Therefore, with $\mu \geq 0$, we have:

$$\mathcal{L}(\tilde{x}, \lambda, \mu) = f(\tilde{x}) + \lambda^T \underbrace{g(\tilde{x})}_{=0} + \underbrace{\mu^T}_{\geq 0} \underbrace{h(\tilde{x})}_{\geq 0} \leq f(\tilde{x}).$$

□

Theorem 10.3: Lagrange Dual Function

The Lagrange dual function is defined as the unconstrained infimum of the Lagrangian function over x , for fixed multipliers λ and μ :

$$q(\lambda, \mu) = \inf_{x \in \mathbb{R}^n} \mathcal{L}(x, \lambda, \mu).$$

Lemma 10.2: Lower Bound property of Lagrange Dual

If $\mu \geq 0$, then the dual function $q(\lambda, \mu)$ is a lower bound on the primal optimal value p^* , i.e.,

$$q(\lambda, \mu) \leq p^*.$$

Proof 10.2: Lower Bound property of Lagrange Dual

Since the Lagrange function is bounded from below by Lemma 10.1, we have

$$q(\lambda, \mu) = \inf_{x \in \mathbb{R}^n} \mathcal{L}(x, \lambda, \mu) \leq f(\tilde{x}) \quad \text{for any feasible } \tilde{x}.$$

Naturally, this inequality holds in particular for the global minimizer x^* , which yields:

$$q(\lambda, \mu) \leq f(x^*) = p^*.$$


□

Theorem 10.4: Concavity of Lagrange Dual

The function $q : \mathbb{R}^p \times \mathbb{R}^q \rightarrow \mathbb{R}$ is concave, even if the original NLP was not convex.

Proof 10.3: Concavity of Lagrange Dual

We will show that $-q$ is convex. The Lagrangian \mathcal{L} is an affine function in the multipliers λ and μ , which in particular implies that $-\mathcal{L}$ is convex in λ and μ . Thus, the function $-q(\lambda, \mu) = \sup_x -\mathcal{L}(x, \lambda, \mu)$ is the supremum of convex functions in λ and μ that are indexed by x , and therefore convex.



□

Theorem 10.5: Dual Problem

The dual problem is defined as the convex maximization problem, i.e.,

$$d^* = \left(\max_{\lambda \in \mathbb{R}^p, \mu \in \mathbb{R}^q} q(\lambda, \mu) \quad \text{s.t.} \quad \mu \geq 0 \right).$$

Theorem 10.6: Weak Duality

Consider a primal-dual pair. Then, the following inequality holds:

$$d^* \leq p^*.$$

Theorem 10.7: Strong Duality

If the primal optimization problem is convex and the Slater condition (see Theorem 10.8) holds, then strong duality holds, i.e.,

$$d^* = p^*.$$

Theorem 10.8: Slater Condition

If there exist one feasible point \tilde{x} such that all non-linear inequalities are strictly satisfied of a primal convex optimization problem hold, then the Slater condition is satisfied. More explicitly, for a convex problem we must have affine equality constraints, $g(x) = Ax + b$, and the inequality constraint functions can be either affine or concave functions, thus without loss of generality assume that the first $q_1 \leq q$ inequalities are affine and the remaining ones concave. Then the Slater condition holds if and only if there exists an \tilde{x} such that

$$\begin{aligned} A\tilde{x} + b &= 0, \\ h_i(\tilde{x}) &\geq 0, \quad \text{for } i = 1, \dots, q_1, \\ g_i(\tilde{x}) &> 0, \quad \text{for } i = q_1 + 1, \dots, q. \end{aligned}$$

Unconstrained Optimization Algorithms

11 Optimality Conditions

Theorem 11.1: Unconstrained optimization Problems

We define the unconstrained optimization problem as

$$\min_{x \in D} f(x),$$

where we regard objective function $f : D \rightarrow \mathbb{R}$ that are defined on some open domain $D \subseteq \mathbb{R}^n$.

Theorem 11.2: Stationary Point

A point \tilde{x} is called a stationary point of f if and only if

$$\nabla f(\tilde{x}) = 0.$$

Theorem 11.3: Descent Direction

A vector $p \in \mathbb{R}^n$ is called a descent direction at x if

$$\nabla f(x)^T p < 0.$$

Theorem 11.4: First Order Necessary Conditions (FONC)

If $x^* \in D$ is a local minimizer of $f : D \rightarrow \mathbb{R}$ and $f \in C^1$ then

$$\nabla f(x^*) = 0.$$

Proof 11.1: First Order Necessary Conditions (FONC)

Let us assume for contradiction that $\nabla f(x^*) \neq 0$. Then $p = -\nabla f(x^*)$ would be a descent direction in which the objective could be improved, as follows:

As D is open and $f \in C^1$, we could find a $t > 0$ that is small enough so that for all $\tau \in [0, t]$ holds $x^* + \tau p \in D$ and $\nabla f(x^* + \tau p)^T p < 0$. By Taylor's theorem, we would have for some $\theta \in (0, 1)$ that

$$f(x^* + tp) = f(x^*) + \underbrace{t \nabla f(x^* + \theta tp)^T p}_{< 0} < f(x^*).$$

□

Theorem 11.5: Second Order Necessary Conditions (SONC)

If $x^* \in D$ is a local minimizer of $f : D \rightarrow \mathbb{R}$ and $f \in C^2$ then

$$\nabla^2 f(x^*) \succeq 0.$$

Proof 11.2: Second Order Necessary Conditions (SONC)

Assume, for the sake of contradiction, that $\nabla^2 f(x^*) \prec 0$. This implies the existence of a vector $p \in \mathbb{R}^n$ such that $p^T \nabla^2 f(x^*) p < 0$. In this case, the objective function could be improved in the direction of p . By choosing a sufficiently small $t > 0$, we can ensure that for all $\tau \in [0, t]$, the following holds:

$$p^T \nabla^2 f(x^* + \tau p) p < 0.$$

Applying Taylor's theorem, we would have for some $\theta \in (0, 1)$ that

$$f(x^* + tp) = f(x^*) + \underbrace{t \nabla f(x^*)^T p}_{=0} + \frac{t^2}{2} \underbrace{p^T \nabla^2 f(x^* + \theta p) p}_{<0} < f(x^*),$$

which leads to a contradiction. □

Theorem 11.6: Convex First Order Sufficient Conditions (cFOSC)

Assume that $f : D \rightarrow \mathbb{R}$ is convex and $f \in C^1$. If $x^* \in D$ is a stationary point of f , then x^* is a global minimizer of f .

Theorem 11.7: Second Order Sufficient Conditions (SOSC)

Assume that $f : D \rightarrow \mathbb{R}$ and $f \in C^2$. If $x^* \in D$ is a stationary point of f and $\nabla^2 f(x^*) \succ 0$, then x^* is a strict local minimizer of f .

Proof 11.3: Second Order Sufficient Conditions (SOSC)

We can choose a sufficiently small closed ball B around x^* so that for all $x \in B$ holds $\nabla^2 f(x) \succ 0$. Restricted to this ball, we have a convex problem, so that Theorem 11.6 together with stationarity of x^* implies that x^* is a global minimizer of f . To prove that it is strict, we look for any $x \in B \setminus x^*$ at the Taylor expansion, which yields with some $\theta \in (0, 1)$:

$$f(x) = f(x^*) + \underbrace{\nabla f(x^*)^T (x - x^*)}_{=0} + \frac{1}{2} \underbrace{(x - x^*)^T \nabla^2 f(x^* + \theta(x - x^*)) (x - x^*)}_{>0} > f(x^*).$$

□

Theorem 11.8: Stability of Parametric Solutions

Assume that $f : D \times \mathbb{R}^m \rightarrow \mathbb{R}$, and regard the minimization of $f(\cdot, \tilde{a})$ for a given fixed value of $\tilde{a} \in \mathbb{R}^m$. If $\tilde{x} \in D$ satisfies the SOSC (see Theorem 11.7), then there is a neighborhood $\mathcal{N} \subset \mathbb{R}^m$ around \tilde{a} such that the parametric minimizer function $x^*(a)$ is well-defined for all $a \in \mathcal{N}$ (i.e. there is a unique minimizer $x^*(a)$ for each $a \in \mathcal{N}$), is differentiable in \mathcal{N} , and $x^*(\tilde{a}) = \tilde{x}$. Its derivative at \tilde{a} is given by

$$\frac{\partial(x^*(\tilde{a}))}{\partial a} = -(\nabla_x^2 f(\tilde{x}, \tilde{a}))^{-1} \frac{\partial(\nabla_x f(\tilde{x}, \tilde{a}))}{\partial a}.$$

Moreover, each such $x^*(a)$ with $a \in \mathcal{N}$ satisfies again the SOSC and is thus a strict local minimizer.

Proof 11.4: Stability of Parametric Solutions

The existence of the differentiable map $x^* : \mathcal{N} \rightarrow D$ follows from the implicit function theorem applied to the stationarity condition $\nabla_x f(x^*(a), a) = 0$. The derivative of $x^*(a)$ at \tilde{a} is given by

$$\frac{d(\nabla_x f(x^*(a), a))}{da} = \underbrace{\frac{\partial(\nabla_x f(x^*(a), a))}{\partial x}}_{=\nabla_x^2 f} \frac{\partial x^*(a)}{\partial a} + \frac{\partial(\nabla_x f(x^*(a), a))}{\partial a} = 0$$

The fact that all points $x^*(a)$ satisfy the SOSC follows from the continuity of the second derivative. \square

12 Estimation and Fitting Problems

Theorem 12.1: Estimation and Fitting Problems

Estimation and fitting problems are optimization problems with a special objective, namely a least squares objective. We define the estimation problem as

$$\min_{x \in \mathbb{R}^n} \frac{1}{2} \|\eta - M(x)\|_2^2,$$

where $\eta \in \mathbb{R}^m$ is the measurement vector, $M : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is the model function, and $x \in \mathbb{R}^n$ is the parameter vector. Many models in estimation and fitting problems are linear functions of x . If M is linear, $M(x) = Jx$, then $f(x) = \frac{1}{2} \|\eta - Jx\|_2^2$ which is a convex function, as $\nabla^2 f(x) = J^T J \succeq 0$. Therefore local minimizers are found by:

$$\begin{aligned} \nabla f(x) = 0 &\Leftrightarrow J^T Jx^* - J^T \eta = 0 \\ &\Leftrightarrow x^* = \underbrace{(J^T J)^{-1} J^T}_{J^+} \eta \end{aligned}$$

Theorem 12.2: Pseudo-inverse

J^+ is called the pseudo-inverse and is a generalization of the inverse matrix. If $J^T J \succ 0$, J^+ is given

by

$$J^+ = (J^T J)^{-1} J^T.$$

So far, $(J^T J)^{-1}$ is only defined if $J^T J \succ 0$. This holds if and only if $\text{rank}(J) = n$, i.e. if the columns of J are linearly independent.

Theorem 12.3: Moore Penrose Pseudo Inverse

Assume $J \in \mathbb{R}^{m \times n}$ and that the singular value decomposition (SVD) of J is given by $J = U \Sigma V^T$. Then, the Moore-Penrose pseudo-inverse J^+ is given by

$$J^+ = V S^+ U^T,$$

where for

$$S = \left[\begin{array}{ccccccc} \sigma_1 & & & & & & \\ & \sigma_2 & & & & & \\ & & \ddots & & & & \\ & & & \sigma_r & & & \\ & & & & 0 & & \\ & & & & & \ddots & \\ & & & & & & 0 \\ \hline 0 & \dots & \dots & 0 & \dots & \dots & 0 \end{array} \right] \quad \text{holds} \quad S^+ = \left[\begin{array}{ccccccc} \sigma_1^{-1} & & & & & & 0 \\ & \sigma_2^{-1} & & & & & \vdots \\ & & \ddots & & & & \vdots \\ & & & \sigma_r^{-1} & & & 0 \\ & & & & 0 & & \vdots \\ & & & & & \ddots & \vdots \\ & & & & & & 0 \end{array} \right]$$

Lemma 12.1

or $\epsilon \rightarrow 0$

13 Newton Type Optimization

14 Globalisation Strategies

15 Calculating Derivatives

Constrained Optimization Algorithms

- 16 Optimality Conditions for Constrained Optimization
- 17 Equality Constrained Optimization Algorithms
- 18 Inequality Constrained Optimization Algorithms
- 19 Optimal Control Problems