

T-Axi (declarative)

Wojciech Kołowski
Mateusz Pyzik

TODO

- Kinding.
- Polymorphism.
- Higher-order quantification.
- Totality checker.
- Computation.
- Inductives and records.
- Algorithmic system.
- Distinguish partial terms.

Quantities

Quantities:

$r ::= 0 \mid 1 \mid ? \mid + \mid *$

- 0 means a resource has been used up and is no longer available.
- 1 means a resource must be used exactly once.
- ? (pronounced “few”) means a resource must be used at most once.
- + (pronounced “many”) means a resource must be used at least once.
- * (pronounced “any”) means no restrictions on usage.

* is the default quantity, so when there's nothing to indicate quantity, it means it's *.

Subusage ordering

$r_1 \sqsubseteq r_2$ means that a resource with quantity r_1 may be used when quantity r_2 is expected. This ordering is called sub-usaging. The definition below is just the skeleton, the full ordering is the reflexive-transitive closure of it.

$$* \sqsubseteq ?$$

$$* \sqsubseteq +$$

$$+ \sqsubseteq 1$$

$$? \sqsubseteq 1$$

$$? \sqsubseteq 0$$

We will denote the greatest lower bound in this order with $r_1 \sqcap r_2$ and the least upper bound (if it exists) with $r_1 \sqcup r_2$.

Addition of quantities

When we have two quantities of the same resource, we can sum the quantities.

$$0 + r = r$$

$$r + 0 = r$$

$$? + ? = *$$

$$? + * = *$$

$$* + ? = *$$

$$* + * = *$$

$$_ + _ = +$$

Multiplication of quantities

When we have a quantity r_1 of resource A that contains quantity r_2 of resource B , then we in fact have quantity $r_1 \cdot r_2$ of resource B .

$$0 \cdot r = 0$$

$$r \cdot 0 = 0$$

$$1 \cdot r = r$$

$$r \cdot 1 = r$$

$$?.? = ?$$

$$+ \cdot + = +$$

$$_ \cdot _ = *$$

The algebra of quantities

Quantities \mathcal{Q} form a positive ordered commutative semiring with no zero divisors, i.e.:

- $(\mathcal{Q}, +, 0)$ is a commutative monoid.
- $(\mathcal{Q}, \cdot, 1)$ is a commutative monoid.
- 0 annihilates multiplication.
- Multiplication distributes over addition.
- Addition and multiplication preserve the subusage ordering in both arguments.
- If $r_1 + r_2 = 0$, then $r_1 = 0$ and $r_2 = 0$.
- If $r_1 \cdot r_2 = 0$, then $r_1 = 0$ or $r_2 = 0$.

Types

Types:

$A, B ::=$

$r A \rightarrow B \mid !_r A \mid A \otimes B \mid A \oplus B \mid \text{Unit} \mid \text{Empty} \mid$
 $\forall a : \text{Type}_r. A \mid \forall \{a : \text{Type}_r\}. A$

Terms

Terms:

$e ::=$

$x \mid \lambda_r x : A. e \mid e_1 e_2 \mid$
 $\text{box}_r e \mid \text{let box } x = e_1 \text{ in } e_2$
 $(e_1, e_2) \mid \text{let } (x, y) = e_1 \text{ in } e_2 \mid$
 $\text{inl}_A e \mid \text{inr}_A e \mid \text{case } e \text{ of } (e_1, e_2) \mid$
 $\text{unit} \mid \text{let}_A \text{unit} = e_1 \text{ in } e_2 \mid$
 $\text{Empty-elim}_A e \mid$
 $\text{let}_r x : A = e_1 \text{ in } e_2 \mid$
 $\text{let noncomputable } x : A = e_1 \text{ in } e_2 \mid$
 $\text{choose}_A p \mid \text{choose-witness } x \text{ } h \text{ for } p \text{ in } e$

$\text{choose}_A p$ and $\text{choose-witness } x \text{ } h \text{ for } p \text{ in } e$ are noncomputable terms, whereas all others are computable.

Propositions

Propositions:

$P, Q ::=$

$\top \mid \perp \mid P \Rightarrow Q \mid P \wedge Q \mid P \vee Q \mid$

$\forall x : A. P \mid \exists x : A. P \mid$

$e_1 =_A e_2$

Notations:

$\neg P$ stands for $P \Rightarrow \perp$

$P \Leftrightarrow Q$ stands for $(P \Rightarrow Q) \wedge (Q \Rightarrow P)$

Proofterms

Proofterms (P, Q are propositions, e are terms, h are variables):

$p, q ::=$

$h \mid$ **assumption** \mid **trivial** \mid **absurd** p
assume $h : P$ **in** $q \mid$ **apply** $p_1 p_2 \mid$
both $p_1 p_2 \mid$ **and-left** $p \mid$ **and-right** $p \mid$
or-left $p \mid$ **or-right** $p \mid$ **cases** $p_1 p_2 p_3 \mid$
lemma $h : P$ **by** p **in** $q \mid$ **proving** P **by** $p \mid$
suffices P **by** q **in** $p \mid$
pick-any $x : A$ **in** $e \mid$ **instantiate** p **with** $e \mid$
witness e **such that** $p \mid$ **pick-witness** $x h$ **for** p_1 **in** $p_2 \mid$
refl $e \mid$ **rewrite** p_1 **in** $p_2 \mid$ **funext** $x : A$ **in** p
by-contradiction $h : \neg P$ **in** $q \mid$
choose-spec $p \mid$ **choose-witness** $x h$ **for** p **in** q

Contexts

Contexts:

$$\Gamma ::= \cdot \mid \Gamma, r x : A \mid \Gamma, r x : A := e \mid \Gamma, h : P \mid \Gamma, a : \text{Type}_r$$

Judgements

Well-formed context judgement: $\Gamma \text{ ctx}_i$, where i is either c or nc .

Well-formed type judgement: $\Gamma \vdash A \text{ type}$

Type conversion judgement: $\Gamma \vdash A \equiv B \text{ type}$

Typing judgement: $\Gamma \vdash_i e : A$, where i is either c or nc

Conversion judgement: $\Gamma \vdash e_1 \equiv e_2 : A$

Well-formed proposition judgement: $\Gamma \vdash P \text{ prop}$

Proposition conversion judgement: $\Gamma \vdash P \equiv Q \text{ prop}$

Proof judgement: $\Gamma \vdash p : P$

Sanity checks

We'll set up the system so that:

- If $\Gamma \vdash e_1 \equiv e_2 : A$, then $\Gamma \vdash_{\text{nc}} e_1 : A$ and $\Gamma \vdash_{\text{nc}} e_2 : A$.
- If $\Gamma \vdash_i e : A$, then $|\Gamma| \vdash A \text{ type}$ and $\Gamma \text{ ctx}_i$.
- If $\Gamma \vdash A \equiv B \text{ type}$, then $\Gamma \vdash A \text{ type}$ and $\Gamma \vdash B \text{ type}$.
- If $\Gamma \vdash A \text{ type}$, then $\Gamma \text{ ctx}_{\text{nc}}$.
- If $\Gamma \vdash p : P$, then $\Gamma \vdash P \text{ prop}$.
- If $\Gamma \vdash P \equiv Q \text{ prop}$, then $\Gamma \vdash P \text{ prop}$ and $\Gamma \vdash Q \text{ prop}$.
- If $\Gamma \vdash P \text{ prop}$, then $\Gamma \text{ ctx}_{\text{nc}}$.
- If $\Gamma \text{ ctx}_{\text{nc}}$, then $\Gamma \text{ ctx}_c$.

Operations on contexts

Operations on contexts:

$\Gamma_1 \sqsubseteq \Gamma_2$ – context subusaging

$\Gamma_1 + \Gamma_2$ – context addition

$r \Gamma$ – context scaling

$|\Gamma|$ – cartesianization

$|\Gamma|_x$ – spotlight x

Context subusaging

$$\frac{}{\cdot \sqsubseteq \cdot}$$

$$\frac{\Gamma_1 \sqsubseteq \Gamma_2 \quad r_1 \sqsubseteq r_2}{\Gamma_1, r_1 x : A \sqsubseteq \Gamma_2, r_2 x : A}$$

$$\frac{\Gamma_1 \sqsubseteq \Gamma_2 \quad r_1 \sqsubseteq r_2}{\Gamma_1, r_1 x : A := e \sqsubseteq \Gamma_2, r_2 x : A := e}$$

$$\frac{\Gamma_1 \sqsubseteq \Gamma_2}{\Gamma_1, h : P \sqsubseteq \Gamma_2, h : P}$$

Context addition

$$\cdot + \cdot = \cdot$$

$$(\Gamma_1, r_1 x : A) + (\Gamma_2, r_2 x : A) = (\Gamma_1 + \Gamma_2), (r_1 + r_2) x : A$$

$$(\Gamma_1, r_1 x : A := e) + (\Gamma_2, r_2 x : A := e) = (\Gamma_1 + \Gamma_2), (r_1 + r_2) x : A := e$$

$$(\Gamma_1, p : P) + (\Gamma_2, h : P) = (\Gamma_1 + \Gamma_2), h : P$$

Context scaling

$$s \cdot = \cdot$$

$$s(\Gamma, rx : A) = s\Gamma, (s \cdot r)x : A$$

$$s(\Gamma, rx : A := e) = s\Gamma, (s \cdot r)x : A := e$$

$$s(\Gamma, h : P) = s\Gamma, h : P$$

Spotlight

$|\cdot|_x = \text{undefined}$

$|\Gamma, r x : A|_x = 0 \Gamma, 1 x : A$

$|\Gamma, r y : A|_x = |\Gamma|_x, 0 y : A$

$|\Gamma, r x : A := e|_x = 0 \Gamma, 1 x : A := e$

$|\Gamma, r y : A := e|_x = |\Gamma|_x, 0 y : A := e$

$|\Gamma, h : P|_x = |\Gamma|_x, h : P$

Cartesianization

Cartesianization turns a context into a context with the same shape but unlimited resources.

$$\begin{aligned} |\cdot| &= \cdot \\ |\Gamma, r x : A| &= |\Gamma|, x : A \\ |\Gamma, r x : A := e| &= |\Gamma|, x : A := e \\ |\Gamma, h : P| &= |\Gamma|, h : P \end{aligned}$$

Well-formed contexts

$$\frac{}{\cdot \text{ctx}_c}$$

$$\frac{\Gamma \text{ ctx}_c \quad |\Gamma| \vdash A \text{ type} \quad x \notin \Gamma}{\Gamma, r x : A \text{ ctx}_c}$$

$$\frac{\Gamma \text{ ctx}_c \quad |\Gamma| \vdash_{\text{nc}} e : A \quad x \notin \Gamma}{\Gamma, r x : A := e \text{ ctx}_c}$$

$$\frac{\Gamma \text{ ctx}_c \quad |\Gamma| \vdash P \text{ prop} \quad h \notin \Gamma}{\Gamma, h : P \text{ ctx}_c}$$

$$\frac{\Gamma \text{ ctx}_c \quad \Gamma = |\Gamma|}{\Gamma \text{ ctx}_{\text{nc}}}$$

Well-formed types

$$\frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \text{Unit type}}$$

$$\frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \text{Empty type}}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash r A \rightarrow B \text{ type}}$$

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash !_r A \text{ type}}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A \otimes B \text{ type}}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A \oplus B \text{ type}}$$

The inherent quantity of a type

$$\begin{aligned}\text{qty}(\text{Unit}) &= * \\ \text{qty}(\text{Empty}) &= * \\ \text{qty}(!_0 A) &= * \\ \text{qty}(!_r A) &= r \cdot \text{qty}(A) \\ \text{qty}(A \otimes B) &= \text{qty}(A) \sqcup \text{qty}(B) \\ \text{qty}(A \oplus B) &= \text{qty}(A) \sqcup \text{qty}(B) \\ \text{qty}(r A \rightarrow B) &= 1\end{aligned}$$

Total quantity added to context

$$r_A^{\text{nc}} = *$$
$$r_A^{\text{c}} = r \cdot \text{qty}(A)$$

Using variables

$$\frac{\Gamma \text{ ctx}_i \quad (x : A) \in \Gamma \quad \Gamma \sqsubseteq |\Gamma|_x}{\Gamma \vdash_i x : A}$$

Functions

$$\frac{\Gamma, r_A^i x : A \vdash_i e : B}{\Gamma \vdash_i \lambda_r x : A. e : r A \rightarrow B}$$

$$\frac{\Gamma \sqsubseteq \Gamma_1 + r \Gamma_2 \quad \Gamma_1 \vdash_i e_1 : r A \rightarrow B \quad \Gamma_2 \vdash_i e_2 : A}{\Gamma \vdash_i e_1 e_2 : B}$$

Box

$$\frac{\Gamma \sqsubseteq r \Gamma' \quad \Gamma' \vdash_i e : A}{\Gamma \vdash_i \text{box}_r e : !_r A}$$

$$\frac{\Gamma \sqsubseteq \Gamma_1 + \Gamma_2 \quad \Gamma_1 \vdash_i e_1 : !_r A \quad \Gamma_2, r_A^i x : A \vdash_i e_2 : B}{\Gamma \vdash_i \text{let box } x = e_1 \text{ in } e_2 : B}$$

Empty

$$\frac{|\Gamma| \vdash A \text{ type} \quad \Gamma \vdash_i e : \text{Empty}}{\Gamma \vdash_i \text{Empty-elim}_A e : A}$$

Unit

$$\frac{\Gamma \text{ ctx}; \quad \Gamma \sqsubseteq 0 \Gamma}{\Gamma \vdash_i \text{unit} : \text{Unit}}$$

$$\frac{\Gamma \sqsubseteq \Gamma_1 + \Gamma_2 \quad \Gamma_1 \vdash_i e_1 : \text{Unit} \quad \Gamma_2 \vdash_i e_2 : A}{\Gamma \vdash_i \text{let}_A \text{unit} = e_1 \text{ in } e_2 : A}$$

Products

$$\frac{\Gamma \sqsubseteq \Gamma_1 + \Gamma_2 \quad \Gamma_1 \vdash_i a : A \quad \Gamma_2 \vdash_i b : B}{\Gamma \vdash_i (a, b) : A \otimes B}$$

$$\frac{\Gamma \sqsubseteq \Gamma_1 + \Gamma_2 \quad \Gamma_1 \vdash_i e_1 : A \otimes B \quad \Gamma_2, 1_A^i x : A, 1_B^i y : B \vdash_i e_2 : C}{\Gamma \vdash_i \text{let } (x, y) = e_1 \text{ in } e_2 : C}$$

Sums

$$\frac{\Gamma \vdash_i e : A \quad |\Gamma| \vdash B \text{ type}}{\Gamma \vdash_i \text{inl}_B e : A \oplus B}$$

$$\frac{|\Gamma| \vdash A \text{ type} \quad \Gamma \vdash_i e : B}{\Gamma \vdash_i \text{inr}_A e : A \oplus B}$$

$$\frac{\Gamma \sqsubseteq \Gamma_1 + \Gamma_2 \quad \Gamma_1 \vdash_i e : A \oplus B \quad \Gamma_2 \vdash_i f : 1 A \rightarrow C \quad \Gamma_2 \vdash_i g : 1 B \rightarrow C}{\Gamma \vdash_i \text{case } e \text{ of } (f, g) : C}$$

Q: Do we want first-order representation of the branches?
Probably yes.

Let

$$\frac{\Gamma \sqsubseteq r \Gamma_1 + \Gamma_2 \quad \Gamma_1 \vdash_i e_1 : A \quad \Gamma_2, r_A^i x : A := e_1 \vdash_i e_2 : B}{\Gamma \vdash_i \text{let}_r x : A = e_1 \text{ in } e_2 : B}$$

Well-formed propositions

$$\frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \top \text{ prop}} \quad \frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \perp \text{ prop}}$$

$$\frac{\Gamma \vdash P \text{ prop} \quad \Gamma \vdash Q \text{ prop}}{\Gamma \vdash P \Rightarrow Q \text{ prop}}$$

$$\frac{\Gamma \vdash P \text{ prop} \quad \Gamma \vdash Q \text{ prop}}{\Gamma \vdash P \wedge Q \text{ prop}} \quad \frac{\Gamma \vdash P \text{ prop} \quad \Gamma \vdash Q \text{ prop}}{\Gamma \vdash P \vee Q \text{ prop}}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma, x : A \vdash P \text{ prop}}{\Gamma \vdash \forall x : A. P \text{ prop}}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma, x : A \vdash P \text{ prop}}{\Gamma \vdash \exists x : A. P \text{ prop}}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash_{\text{nc}} e_1 : A \quad \Gamma \vdash_{\text{nc}} e_2 : A}{\Gamma \vdash e_1 =_A e_2 \text{ prop}}$$

Substitution

The notation is $P[x := e]$ for substitution in propositions.

Assumptions and implication

$$\frac{\Gamma \text{ ctx}_{\text{nc}} \quad (h : P) \in \Gamma}{\Gamma \vdash h : P} \quad \frac{\Gamma \text{ ctx}_{\text{nc}} \quad (h : P) \in \Gamma}{\Gamma \vdash \mathbf{assumption} : P}$$

$$\frac{\Gamma, h : P \vdash q : Q}{\Gamma \vdash \mathbf{assume} \ h : P \ \mathbf{in} \ q : P \Rightarrow Q}$$

$$\frac{\Gamma \vdash q : P \Rightarrow Q \quad \Gamma \vdash p : P}{\Gamma \vdash \mathbf{apply} \ q \ p : Q}$$

Propositional logic

$$\frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \mathbf{trivial} : \top} \quad \frac{\Gamma \vdash Q \text{ prop} \quad \Gamma \vdash p : \perp}{\Gamma \vdash \mathbf{absurd} \ p : Q}$$

$$\frac{\Gamma \vdash p : P \quad \Gamma \vdash q : Q}{\Gamma \vdash \mathbf{both} \ p \ q : P \wedge Q}$$

$$\frac{\Gamma \vdash pq : P \wedge Q}{\Gamma \vdash \mathbf{and-left} \ pq : P} \quad \frac{\Gamma \vdash pq : P \wedge Q}{\Gamma \vdash \mathbf{and-right} \ pq : Q}$$

$$\frac{\Gamma \vdash p : P \quad \Gamma \vdash Q \text{ prop}}{\Gamma \vdash \mathbf{or-left} \ p : P \vee Q} \quad \frac{\Gamma \vdash P \text{ prop} \quad \Gamma \vdash q : Q}{\Gamma \vdash \mathbf{or-right} \ q : P \vee Q}$$

$$\frac{\Gamma \vdash pq : P \vee Q \quad \Gamma \vdash r_1 : P \Rightarrow R \quad \Gamma \vdash r_2 : Q \Rightarrow R}{\Gamma \vdash \mathbf{cases} \ pq \ r_1 \ r_2 : R}$$

Utilities

$$\frac{\Gamma \vdash p : P \quad \Gamma, h : P \vdash q : Q}{\Gamma \vdash \mathbf{lemma} \ h : P \ \mathbf{by} \ p \ \mathbf{in} \ q : Q}$$

$$\frac{\Gamma \vdash p : P}{\Gamma \vdash \mathbf{proving} \ P \ \mathbf{by} \ p : P}$$

$$\frac{\Gamma \vdash pq : P \Rightarrow Q \quad \Gamma \vdash p : P}{\Gamma \vdash \mathbf{suffices} \ P \ \mathbf{by} \ pq \ \mathbf{in} \ p : Q}$$

Quantifiers

$$\frac{\Gamma, x : A \vdash p : P}{\Gamma \vdash \text{pick-any } x : A \text{ in } p : \forall x : A. P}$$

$$\frac{\Gamma \vdash p : \forall x : A. P \quad \Gamma \vdash_{\text{nc}} e : A}{\Gamma \vdash \text{instantiate } p \text{ with } e : P[x := e]}$$

$$\frac{\Gamma \vdash_{\text{nc}} e : A \quad \Gamma, x : A \vdash P \text{ prop} \quad \Gamma \vdash p : P[x := e]}{\Gamma \vdash \text{witness } e \text{ such that } p : \exists x : A. P}$$

$$\frac{\Gamma \vdash p : \exists x : A. P \quad \Gamma \vdash Q \text{ prop} \quad \Gamma, x : A, h : P \vdash q : Q}{\Gamma \vdash \text{pick-witness } x \text{ } h \text{ for } p \text{ in } q : Q}$$

Equality

$$\frac{\Gamma \vdash_{\text{nc}} e : A}{\Gamma \vdash \mathbf{refl} \ e : e =_A e}$$

$$\frac{\Gamma \vdash q : e_1 =_A e_2 \quad \Gamma, x : A \vdash P \text{ prop} \quad \Gamma \vdash p : P[x := e_2]}{\Gamma \vdash \mathbf{rewrite} \ q \text{ in } p : P[x := e_1]}$$

$$\frac{\Gamma, x : A \vdash p : f \ x =_B g \ x}{\Gamma \vdash \mathbf{funext} \ x : A \text{ in } p : f =_{rA \rightarrow B} g}$$

Classical Logic

$$\frac{\Gamma, h : \neg P \vdash q : \perp}{\Gamma \vdash \text{by-contradiction } h : \neg P \text{ in } q : P}$$

$$\frac{|\Gamma| \vdash p : \exists x : A. P}{\Gamma \vdash_{\text{nc}} \text{choose } p : A}$$

$$\frac{\Gamma \vdash p : \exists x : A. P}{\Gamma \vdash \text{choose-spec } p : P [x := \text{choose } p]}$$

$$\frac{\Gamma \vdash p : \exists x : A. P \quad \Gamma \vdash Q \text{ prop} \quad \Gamma, x : A := \text{choose } p, h : P \vdash q : Q}{\Gamma \vdash \text{choose-witness } x \text{ for } p \text{ in } q : Q}$$

$$\frac{|\Gamma| \vdash p : \exists x : A. P \quad |\Gamma| \vdash B \text{ type} \quad \Gamma, x : A := \text{choose } p, h : P \vdash_{\text{nc}} e : B}{\Gamma \vdash_{\text{nc}} \text{choose-witness } x \text{ for } p \text{ in } e : B}$$

Conversion rules

$$\frac{\Gamma \vdash_i e : A \quad |\Gamma| \vdash A \equiv B \text{ type}}{\Gamma \vdash_i e : B}$$

$$\frac{\Gamma \vdash e_1 \equiv e_2 : A \quad \Gamma \vdash A \equiv B \text{ type}}{\Gamma \vdash e_1 \equiv e_2 : B}$$

$$\frac{\Gamma \vdash p : P \quad \Gamma \vdash P \equiv Q \text{ prop}}{\Gamma \vdash p : Q}$$

Type conversion

$$\frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \text{Unit} \equiv \text{Unit type}}$$

$$\frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \text{Empty} \equiv \text{Empty type}}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \text{ type} \quad \Gamma \vdash B_1 \equiv B_2 \text{ type}}{\Gamma \vdash r A_1 \rightarrow B_1 \equiv r A_2 \rightarrow B_2 \text{ type}}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \text{ type}}{\Gamma \vdash !_r A_1 \equiv !_r A_2 \text{ type}}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \text{ type} \quad \Gamma \vdash B_1 \equiv B_2 \text{ type}}{\Gamma \vdash A_1 \otimes B_1 \equiv A_2 \otimes B_2 \text{ type}}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \text{ type} \quad \Gamma \vdash B_1 \equiv B_2 \text{ type}}{\Gamma \vdash A_1 \oplus B_1 \equiv A_2 \oplus B_2 \text{ type}}$$

Type conversion – properties

The rules above are the complete definition of type conversion. For now the context Γ is not used, but soon it will. We don't need to take any closures – type conversion already is an equivalence relation.

- $\Gamma \vdash A \equiv A$ type
- If $\Gamma \vdash A \equiv B$ type then $\Gamma \vdash B \equiv A$ type
- If $\Gamma \vdash A \equiv B$ type and $\Gamma \vdash B \equiv C$ type then $\Gamma \vdash A \equiv C$ type

Proposition conversion

$$\frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \top \equiv \top \text{ prop}} \quad \frac{\Gamma \text{ ctx}_{\text{nc}}}{\Gamma \vdash \perp \equiv \perp \text{ prop}}$$

$$\frac{\Gamma \vdash P_1 \equiv P_2 \text{ prop} \quad \Gamma \vdash Q_1 \equiv Q_2 \text{ prop}}{\Gamma \vdash P_1 \Rightarrow Q_1 \equiv P_2 \Rightarrow Q_2 \text{ prop}}$$

$$\frac{\Gamma \vdash P_1 \equiv P_2 \text{ prop} \quad \Gamma \vdash Q_1 \equiv Q_2 \text{ prop}}{\Gamma \vdash P_1 \wedge Q_1 \equiv P_2 \wedge Q_2 \text{ prop}}$$

$$\frac{\Gamma \vdash P_1 \equiv P_2 \text{ prop} \quad \Gamma \vdash Q_1 \equiv Q_2 \text{ prop}}{\Gamma \vdash P_1 \vee Q_1 \equiv P_2 \vee Q_2 \text{ prop}}$$

Proposition conversion

$$\frac{\Gamma \vdash A_1 \equiv A_2 \text{ type} \quad \Gamma, x : A_1 \vdash P_1 \equiv P_2 \text{ prop}}{\Gamma \vdash \forall x : A_1. P_1 \equiv \forall x : A_2. P_2 \text{ prop}}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \text{ type} \quad \Gamma, x : A_1 \vdash P_1 \equiv P_2 \text{ prop}}{\Gamma \vdash \exists x : A_1. P_1 \equiv \exists x : A_2. P_2 \text{ prop}}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \text{ type} \quad \Gamma \vdash e_1 \equiv e_2 : A_1 \quad \Gamma \vdash e'_1 \equiv e'_2 : A_1}{\Gamma \vdash e_1 =_{A_1} e'_1 \equiv e_2 =_{A_2} e'_2 \text{ prop}}$$

Note that we don't worry about α -conversion and assume the variable is the same on both sides.

Term conversion – Empty

$$\frac{\Gamma \vdash_{\text{nc}} e_1 : \text{Empty} \quad \Gamma \vdash_{\text{nc}} e_2 : \text{Empty}}{\Gamma \vdash e_1 \equiv e_2 : \text{Empty}}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash e_1 \equiv e_2 : \text{Empty}}{\Gamma \vdash \text{Empty-elim } e_1 \equiv \text{Empty-elim } e_2 : A}$$

Term conversion – Unit

$$\frac{\Gamma \vdash_{\text{nc}} a : A}{\Gamma \vdash \text{let unit} = \text{unit in } a \equiv a : A}$$

$$\frac{\Gamma \vdash_{\text{nc}} u_1 : \text{Unit} \quad \Gamma \vdash_{\text{nc}} u_2 : \text{Unit}}{\Gamma \vdash u_1 \equiv u_2 : \text{Unit}}$$

$$\frac{\Gamma \vdash u_1 \equiv u_2 : \text{Unit} \quad \Gamma \vdash a_1 \equiv a_2 : A}{\Gamma \vdash \text{let unit} = u_1 \text{ in } a_1 \equiv \text{let unit} = u_2 \text{ in } a_2 : A}$$

Term conversion – functions

$$\frac{\Gamma, x : A \vdash_{\text{nc}} b : B \quad \Gamma \vdash_{\text{nc}} a : A}{\Gamma \vdash (\lambda_r x : A. b) a \equiv b[x := a] : B}$$

$$\frac{\Gamma, x : A \vdash f x \equiv g x : B}{\Gamma \vdash f \equiv g : r A \rightarrow B}$$

$$\frac{\Gamma \vdash f_1 \equiv f_2 : r A \rightarrow B \quad \Gamma \vdash a_1 \equiv a_2 : A}{\Gamma \vdash f_1 a_1 \equiv f_2 a_2 : B}$$

Term conversion – box

$$\frac{\Gamma \vdash_{\text{nc}} a : A \quad \Gamma, x : A \vdash_{\text{nc}} b : B}{\Gamma \vdash \text{let box } x = \text{box}_r a \text{ in } b \equiv b[x := a] : B}$$

$$\frac{\Gamma \vdash a_1 \equiv a_2 : A}{\Gamma \vdash \text{box } a_1 \equiv \text{box } a_2 : !_r A}$$

$$\frac{\Gamma \vdash a_1 \equiv a_2 : !_r A \quad \Gamma, x : A_1 \vdash b_1 \equiv b_2 : B}{\Gamma \vdash \text{let box } x = a_1 \text{ in } b_1 \equiv \text{let box } x = a_2 \text{ in } b_2 : B}$$

Term conversion – product

$$\frac{\Gamma \vdash_{\text{nc}} a : A \quad \Gamma \vdash_{\text{nc}} b : B \quad \Gamma, x : A, y : B \vdash_{\text{nc}} c : C}{\Gamma \vdash \text{let } (x, y) = (a, b) \text{ in } c \equiv c[x := a][y := b] : C}$$

$$\frac{\begin{array}{l} \Gamma \vdash \text{let } (x, y) = e_1 \text{ in } x \equiv \text{let } (x, y) = e_2 \text{ in } x : A \\ \Gamma \vdash \text{let } (x, y) = e_1 \text{ in } y \equiv \text{let } (x, y) = e_2 \text{ in } y : B \end{array}}{\Gamma \vdash e_1 \equiv e_2 : A \otimes B}$$

$$\frac{\Gamma \vdash a_1 \equiv a_2 : A \quad \Gamma \vdash b_1 \equiv b_2 : B}{\Gamma \vdash (a_1, b_1) \equiv (a_2, b_2) : A \otimes B}$$

$$\frac{\Gamma \vdash e_1 \equiv e_2 : A \otimes B \quad \Gamma, x : A, y : B \vdash c_1 \equiv c_2 : C}{\Gamma \vdash \text{let } (x, y) = e_1 \text{ in } c_1 \equiv \text{let } (x, y) = e_2 \text{ in } c_2 : C}$$

Term conversion – sum

$$\frac{\Gamma \vdash_{\text{nc}} a : A \quad \Gamma \vdash_{\text{nc}} f : 1 A \rightarrow C \quad \Gamma \vdash_{\text{nc}} g : 1 B \rightarrow C}{\Gamma \vdash \text{case inl } a \text{ of } (f, g) \equiv f \ a : C}$$

$$\frac{\Gamma \vdash_{\text{nc}} b : B \quad \Gamma \vdash_{\text{nc}} f : 1 A \rightarrow C \quad \Gamma \vdash_{\text{nc}} g : 1 B \rightarrow C}{\Gamma \vdash \text{case inr } b \text{ of } (f, g) \equiv g \ b : C}$$

$$\frac{\Gamma \vdash a_1 \equiv a_2 : A \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash \text{inl } a_1 \equiv \text{inl } a_2 : A \oplus B} \quad \frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash b_1 \equiv b_2 : B}{\Gamma \vdash \text{inr } b_1 \equiv \text{inr } b_2 : A \oplus B}$$

$$\frac{\Gamma \vdash e_1 \equiv e_2 : A \oplus B \quad \begin{array}{l} \Gamma \vdash f_1 \equiv f_2 : 1 A \rightarrow C \\ \Gamma \vdash g_1 \equiv g_2 : 1 B \rightarrow C \end{array}}{\Gamma \vdash \text{case } e_1 \text{ of } (f_1, g_1) \equiv \text{case } e_2 \text{ of } (f_2, g_2) : C}$$

Term conversion – let

$$\frac{\Gamma \vdash a_1 \equiv a_2 : A \quad \Gamma, x : A \vdash b_1 \equiv b_2 : B}{\Gamma \vdash \text{let } x = a_1 \text{ in } b_1 \equiv \text{let } x = a_2 \text{ in } b_2 : B}$$

Universes

$$U ::= \text{Type}_1 \mid \text{Type}_? \mid \text{Type}_+ \mid \text{Type}_*$$

Environments

Global environments:

$\Sigma ::=$

$$\emptyset \mid \Sigma, h : P := p \mid \Sigma, x : A := e \mid$$
$$\Sigma, \text{partial } x : A := e \mid \Sigma, \text{totality } x \ p$$

Well-formed environments

$$\overline{\emptyset \text{ env}}$$

$$\frac{\Sigma \text{ env} \quad h \notin \Sigma \quad \Sigma \mid \cdot \vdash p : P}{\Sigma, h : P := p \text{ env}}$$

$$\frac{\Sigma \text{ env} \quad x \notin \Sigma \quad \Sigma \mid \cdot \vdash_c e : A}{\Sigma, x : A := e \text{ env}}$$

$$\frac{\Sigma \text{ env} \quad x \notin \Sigma \quad \Sigma \mid \cdot \vdash_c e : A \quad e \text{ fails syntactic check}}{\Sigma, \text{partial } x : A := e \text{ env}}$$

$$\frac{\Sigma \text{ env} \quad \Sigma = \Sigma_1, \text{partial } x : A := e, \Sigma_2 \quad \Sigma \mid \cdot \vdash p : \exists r : A. e = ? = r}{\Sigma, \text{totality } x \text{ } p \text{ env}}$$