

# T-Axi (algorithmic)

Wojciech Kołowski  
Mateusz Pyzik

# Syntax

The syntax in the algorithmic system is the same, but we will have far fewer annotations in terms

# Judgements

Kind inference judgement:  $\Gamma \vdash A \Rightarrow \text{Type}_r \dashv \Gamma'$

Type conversion judgements:  $\Gamma \vdash A \equiv B \Rightarrow \text{Type}_r \dashv \Gamma'$  (all types),

$\Gamma \vdash A \hat{=} B \Rightarrow \text{Type}_r \dashv \Gamma'$  (types in whnf)

Type checking judgement:  $\Gamma \vdash_i e \Leftarrow A \dashv \Gamma'$

Type inference judgement:  $\Gamma \vdash_i e \Rightarrow A \dashv \Gamma'$

Term conversion judgements:  $\Gamma \vdash e_1 \equiv e_2 \Leftarrow A \dashv \Gamma'$  (all terms),

$\Gamma \vdash e_1 \equiv e_2 \Rightarrow A \dashv \Gamma'$  (all terms),  $\Gamma \vdash e_1 \hat{=} e_2 \Leftarrow A \dashv \Gamma'$  (whnfs),

$\Gamma \vdash e_1 \hat{=} e_2 \Rightarrow A \dashv \Gamma'$  (whnfs)

Proposition checking judgement:  $\Gamma \vdash P \Leftarrow \text{prop} \dashv \Gamma'$

Proposition conversion judgements:  $\Gamma \vdash P \equiv Q \dashv \Gamma'$  (all

propositions),  $\Gamma \vdash P \hat{=} Q \dashv \Gamma'$  (whnfs),  $\Gamma \vdash P \sim Q \dashv \Gamma'$  (whnfs)

# Subtraction of quantities

$r_1 - r_2$  is the least  $r'$  such that  $r_1 \sqsubseteq r' + r_2$ .

$r_1 - r_2$	0	1	?	+	*
0	0				
1	1	0			
?	?	0	0		
+	+	*	+	*	+
*	*	*	*	*	*

# Subtraction order on quantities

$r_1 \leq_{\text{sub}} r_2$  holds when  $r_2 - r_1$  is defined.

Explicitly:  $0 \leq_{\text{sub}} 1 \leq_{\text{sub}} ? \leq_{\text{sub}} + \leq_{\text{sub}} * \leq_{\text{sub}} +$

# Decrementation order on quantities

$r_1 \leq_{\text{dec}} r_2$  holds when  $r_2 - 1 = r_1$ .

$$\overline{* \leq_{\text{dec}} +}$$

$$\overline{0 \leq_{\text{dec}} 1}$$

$$\overline{0 \leq_{\text{dec}} ?}$$

# Arithmetic order on quantities

The arithmetic order on quantities is  $0 \leq 1 \leq ? \leq + \leq *$ . The idea is to compare the quantities by how “big” they are.

# Division with remainder

$a/b = (q, r)$  when  $a = b \cdot q + r$ , with  $q$  as large as possible and  $r$  being as small as possible according to the arithmetic order. Note that  $a/b = q$  means that  $r = 0$ .

$r_1/r_2$	0	1	?	+	*
0	*	0	0	0	0
1	$(*, 1)$	1	$(0, 1)$	$(0, 1)$	$(0, 1)$
?	$(*, ?)$	?	?	$(0, ?)$	$(0, ?)$
+	$(*, +)$	+	$(*, 1)$	+	$(*, 1)$
*	$(*, *)$	*	*	*	*



# Decrement variable in context

•  $-x = \mathbf{undefined}$

$$(\Gamma, rx : A) - x = \Gamma, (r-1)x : A$$

$$(\Gamma, ry : A) - x = \Gamma - x, ry : A$$

$$(\Gamma, rx : A := e) - x = \Gamma, (r-1)x : A := e$$

$$(\Gamma, ry : A := e) - x = \Gamma - x, ry : A := e$$

$$(\Gamma, h : P) - x = \Gamma - x, h : P$$

$$(\Gamma, a : \text{Type}_r) - x = \Gamma - x, a : \text{Type}_r$$

# Context division with remainder

$$\overline{\cdot/r} = \cdot$$

$$\frac{\Gamma/q = (\Gamma_1, \Gamma_2) \quad r/q = (r_1, r_2)}{(\Gamma, r x : A)/q = ((\Gamma_1, r_1 x : A), (\Gamma_2, r_2 x : A))}$$

$$\frac{\Gamma/q = (\Gamma_1, \Gamma_2) \quad r/q = (r_1, r_2)}{(\Gamma, r x : A := e)/q = ((\Gamma_1, r_1 x : A := e), (\Gamma_2, r_2 x : A := e))}$$

$$\frac{\Gamma/q = (\Gamma_1, \Gamma_2)}{(\Gamma, h : P)/q = ((\Gamma_1, h : P), (\Gamma_2, h : P))}$$

$$\frac{\Gamma/q = (\Gamma_1, \Gamma_2)}{(\Gamma, a : \text{Type}_r)/q = ((\Gamma_1, a : \text{Type}_r), (\Gamma_2, a : \text{Type}_r))}$$

# Kind inference

$$\overline{\Gamma \vdash \text{Unit} \Rightarrow \text{Type} \dashv \Gamma} \quad \overline{\Gamma \vdash \text{Empty} \Rightarrow \text{Type} \dashv \Gamma}$$

$$\frac{\Gamma \vdash A \Rightarrow \text{Type}_{s_1} \dashv \Gamma_1 \quad \Gamma_1 \vdash B \Rightarrow \text{Type}_{s_2} \dashv \Gamma_2}{\Gamma \vdash r A \rightarrow B \Rightarrow \text{Type}_1 \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash A \Rightarrow \text{Type}_s \dashv \Gamma'}{\Gamma \vdash !_0 A \Rightarrow \text{Type} \dashv \Gamma'} \quad \frac{\Gamma \vdash A \Rightarrow \text{Type}_s \dashv \Gamma' \quad r \neq 0}{\Gamma \vdash !_r A \Rightarrow \text{Type}_{r.s} \dashv \Gamma'}$$

$$\frac{\Gamma \vdash A \Rightarrow \text{Type}_{s_1} \dashv \Gamma_1 \quad \Gamma_1 \vdash B \Rightarrow \text{Type}_{s_2} \dashv \Gamma_2}{\Gamma \vdash A \otimes B \Rightarrow \text{Type}_{s_1 \sqcup s_2} \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash A \Rightarrow \text{Type}_{s_1} \dashv \Gamma_1 \quad \Gamma_1 \vdash B \Rightarrow \text{Type}_{s_2} \dashv \Gamma_2}{\Gamma \vdash A \oplus B \Rightarrow \text{Type}_{s_1 \sqcup s_2} \dashv \Gamma_2}$$

# Kind inference

$$\frac{(a : \text{Type}_r) \in \Gamma}{\Gamma \vdash a \Rightarrow \text{Type}_r \dashv \Gamma}$$

$$\frac{\Gamma, a : \text{Type}_r \vdash B \Rightarrow \text{Type}_s \dashv \Gamma', a : \text{Type}_r}{\Gamma \vdash \forall a : \text{Type}_r. B \Rightarrow \text{Type}_s \dashv \Gamma'}$$

$$\frac{\Gamma, a : \text{Type}_r \vdash B \Rightarrow \text{Type}_s \dashv \Gamma', a : \text{Type}_r}{\Gamma \vdash \forall \{a : \text{Type}_r\}. B \Rightarrow \text{Type}_s \dashv \Gamma'}$$

# Kind checking

$$\frac{\Gamma \vdash A \Rightarrow \text{Type}_s \dashv \Gamma' \quad r = s}{\Gamma \vdash A \Leftarrow \text{Type}_r \dashv \Gamma'}$$

# Context clean-up

$\Gamma, rx : A \vdash_i e \Leftarrow A \dashv \Gamma', 0x : A$  is a shorthand for  
 $\Gamma, rx : A \vdash_i e \Leftarrow A \dashv \Gamma', r'x : A$  with the additional condition  
 $r' \sqsubseteq 0$  when  $i = c$

$\Gamma, rx : A \vdash_i e \Rightarrow A \dashv \Gamma', 0x : A$  is a shorthand for  
 $\Gamma, rx : A \vdash_i e \Rightarrow A \dashv \Gamma', r'x : A$  with the additional condition  
 $r' \sqsubseteq 0$  when  $i = c$

# Subsumption and annotations

$$\frac{\Gamma \vdash_i e \Rightarrow A \dashv \Gamma_1 \quad \Gamma_1 \vdash A \equiv B \Rightarrow \text{Type}_r \dashv \Gamma_2}{\Gamma \vdash_i e \Leftarrow B \dashv \Gamma_2} \text{SUBSUMPTION}$$

$$\frac{\Gamma \vdash A \Rightarrow \text{Type}_r \dashv \Gamma_1 \quad \Gamma_1 \vdash_i e \Leftarrow A \dashv \Gamma_2}{\Gamma \vdash_i (e : A) \Rightarrow A \dashv \Gamma_2} \text{ANNOT}$$

# Using variables

$$\frac{\Gamma(x) = A}{\Gamma \vdash_c x \Rightarrow A \dashv \Gamma - x}$$

$$\frac{\Gamma(x) = A}{\Gamma \vdash_{nc} x \Rightarrow A \dashv \Gamma}$$



# Functions

$$\frac{\Gamma, r_A^i x : A \vdash_i e \Leftarrow B \dashv \Gamma', 0 x : A}{\Gamma \vdash_i \lambda x. e \Leftarrow r A \rightarrow B \dashv \Gamma'}$$

$$\frac{\Gamma \vdash_i f \Rightarrow r A \rightarrow B \dashv \Gamma_1 \quad \Gamma_1 / r = (\Gamma_2, \Gamma_3) \quad \Gamma_2 \vdash_i a \Leftarrow A \dashv \Gamma_4}{\Gamma \vdash_i f a \Rightarrow B \dashv \Gamma_3 + r \Gamma_4}$$

## Box

$$\frac{\Gamma/r = (\Gamma_1, \Gamma_2) \quad \Gamma_1 \vdash_i e \Leftarrow A \dashv \Gamma_3}{\Gamma \vdash_i \text{box } e \Leftarrow !_r A \dashv \Gamma_2 + r \Gamma_3}$$

$$\frac{\Gamma \vdash_i e_1 \Rightarrow !_r A \dashv \Gamma_1 \quad \Gamma_1, r_A^i x : A \vdash_i e_2 \Rightarrow B \dashv \Gamma_2, 0 x : A}{\Gamma \vdash_i \text{let box } x = e_1 \text{ in } e_2 \Rightarrow B \dashv \Gamma_2}$$

# Empty

$$\frac{\Gamma \vdash_i e \Leftarrow \text{Empty} \dashv \Gamma'}{\Gamma \vdash_i \text{Empty-elim } e \Leftarrow A \dashv \Gamma'}$$

# Unit

$$\overline{\Gamma \vdash_i \text{unit} \Leftarrow \text{Unit} \dashv \Gamma}$$

$$\frac{\Gamma \vdash_i e_1 \Rightarrow \text{Unit} \dashv \Gamma_1 \quad \Gamma_1 \vdash_i e_2 \Rightarrow A \dashv \Gamma_2}{\Gamma \vdash_i \text{let unit} = e_1 \text{ in } e_2 \Rightarrow A \dashv \Gamma_2}$$

# Products

$$\frac{\Gamma \vdash_i e_1 \Leftarrow A \dashv \Gamma_1 \quad \Gamma_1 \vdash_i e_2 \Leftarrow B \dashv \Gamma_2}{\Gamma \vdash_i (e_1, e_2) \Leftarrow A \otimes B \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash_i e_1 \Rightarrow A \otimes B \dashv \Gamma_1 \quad \Gamma_1, 1_A^i x : A, 1_B^i y : B \vdash_i e_2 \Rightarrow C \dashv \Gamma_2, 0x : A, 0y}{\Gamma \vdash_i \text{let } (x, y) = e_1 \text{ in } e_2 \Rightarrow C \dashv \Gamma_2}$$

## Sums

$$\frac{\Gamma \vdash_i e \Leftarrow A \dashv \Gamma'}{\Gamma \vdash_i \text{inl } e \Leftarrow A \oplus B \dashv \Gamma'} \quad \frac{\Gamma \vdash_i e \Leftarrow B \dashv \Gamma'}{\Gamma \vdash_i \text{inr } e \Leftarrow A \oplus B \dashv \Gamma'}$$

$$\frac{\Gamma \vdash_i e \Rightarrow A \oplus B \dashv \Gamma_1 \quad \begin{array}{l} \Gamma_1 \vdash_i f \Leftarrow 1A \rightarrow C \dashv \Gamma_2 \\ \Gamma_1 \vdash_i g \Leftarrow 1B \rightarrow C \dashv \Gamma_3 \end{array}}{\Gamma \vdash_i \text{case } e \text{ of } (f, g) \Leftarrow C \dashv \Gamma_2 \sqcup \Gamma_3}$$

$$\frac{\Gamma \vdash_i e \Rightarrow A \oplus B \dashv \Gamma_1 \quad \begin{array}{l} \Gamma_1 \vdash_i f \Rightarrow 1A \rightarrow C \dashv \Gamma_2 \\ \Gamma_1 \vdash_i g \Rightarrow 1B \rightarrow C \dashv \Gamma_3 \end{array}}{\Gamma \vdash_i \text{case } e \text{ of } (f, g) \Rightarrow C \dashv \Gamma_2 \sqcup \Gamma_3}$$

## Let

$$\frac{\Gamma/r = (\Gamma_1, \Gamma_2) \quad \Gamma_1 \vdash_i e_1 \Rightarrow A \dashv \Gamma_3 \quad (\Gamma_2 + r \Gamma_3), r_A^i x : A \vdash_i e_2 \Rightarrow B \dashv \Gamma_4}{\Gamma \vdash_i \text{let}_r x = e_1 \text{ in } e_2 \Rightarrow B \dashv \Gamma_4}$$

# Polymorphism

$$\frac{\Gamma, a : \text{Type}_r \vdash_i e \Leftarrow B \dashv \Gamma', a : \text{Type}_r}{\Gamma \vdash_i \Lambda a. e \Leftarrow \forall a : \text{Type}_r. B \dashv \Gamma'}$$

$$\frac{\Gamma \vdash_i e \Rightarrow \forall a : \text{Type}_r. B \dashv \Gamma_1 \quad \Gamma_1 \vdash A \Leftarrow \text{Type}_r \dashv \Gamma_2}{\Gamma \vdash_i e A \Rightarrow B[a := A] \dashv \Gamma_2}$$



# Polymorphism (implicit arguments)

$$\frac{\Gamma, a : \text{Type}_r \vdash_i e \Leftarrow B \dashv \Gamma', a : \text{Type}_r}{\Gamma \vdash_i \Lambda \{a\}. e \Leftarrow \forall \{a : \text{Type}_r\}. B \dashv \Gamma'}$$

$$\frac{\Gamma \vdash_i e \Rightarrow \forall \{a : \text{Type}_r\}. B \dashv \Gamma_1 \quad \Gamma_1 \vdash A \Leftarrow \text{Type}_r \dashv \Gamma_2}{\Gamma \vdash_i e @A \Rightarrow B[a := A] \dashv \Gamma_2}$$

# Subsumption and annotations

$$\frac{\Gamma \vdash p \Rightarrow P \dashv \Gamma_1 \quad \Gamma_1 \vdash P \equiv Q \dashv \Gamma_2}{\Gamma \vdash p \Leftarrow Q \dashv \Gamma_2} \text{SUBSUMPTION}$$

$$\frac{\Gamma \vdash P \Leftarrow \text{prop} \dashv \Gamma_1 \quad \Gamma_1 \vdash p \Leftarrow P \dashv \Gamma_2}{\Gamma \vdash \text{proving } P \text{ by } p \Rightarrow P \dashv \Gamma_2} \text{ANNOT}$$

# Assumptions and implication

$$\frac{\Gamma(h) = P}{\Gamma \vdash h \Rightarrow P \dashv \Gamma} \quad \frac{(h : P) \in \Gamma}{\Gamma \vdash \mathbf{assumption} \Leftarrow P \dashv \Gamma}$$

$$\frac{\Gamma, h : P \vdash q \Leftarrow Q \dashv \Gamma', h : P}{\Gamma \vdash \mathbf{assume} \ h \ \mathbf{in} \ q \Leftarrow P \Rightarrow Q \dashv \Gamma'}$$

$$\frac{\Gamma \vdash q \Rightarrow P \Rightarrow Q \dashv \Gamma_1 \quad \Gamma_1 \vdash p \Leftarrow P \dashv \Gamma_2}{\Gamma \vdash \mathbf{apply} \ q \ p \Rightarrow Q \dashv \Gamma_2}$$

# Propositional logic

$$\frac{}{\Gamma \vdash \mathbf{trivial} \Leftarrow \top \dashv \Gamma} \quad \frac{\Gamma \vdash p \Leftarrow \perp \dashv \Gamma'}{\Gamma \vdash \mathbf{absurd} \ p \Leftarrow Q \dashv \Gamma'}$$

$$\frac{\Gamma \vdash p \Leftarrow P \dashv \Gamma_1 \quad \Gamma_1 \vdash q \Leftarrow Q \dashv \Gamma_2}{\Gamma \vdash \mathbf{both} \ p \ q \Leftarrow P \wedge Q \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash pq \Rightarrow P \wedge Q \dashv \Gamma'}{\Gamma \vdash \mathbf{and-left} \ pq \Rightarrow P \dashv \Gamma'}$$

$$\frac{\Gamma \vdash pq \Rightarrow P \wedge Q \dashv \Gamma'}{\Gamma \vdash \mathbf{and-right} \ pq \Rightarrow Q \dashv \Gamma'}$$

$$\frac{\Gamma \vdash p \Leftarrow P \dashv \Gamma'}{\Gamma \vdash \mathbf{or-left} \ p \Leftarrow P \vee Q \dashv \Gamma'}$$

$$\frac{\Gamma \vdash q \Leftarrow Q \dashv \Gamma'}{\Gamma \vdash \mathbf{or-right} \ q \Leftarrow P \vee Q \dashv \Gamma'}$$

$$\frac{\Gamma \vdash pq \Rightarrow P \vee Q \dashv \Gamma_1 \quad \Gamma_1 \vdash r_1 \Leftarrow P \Rightarrow R \dashv \Gamma_2 \quad \Gamma_1 \vdash r_2 \Leftarrow Q \Rightarrow R \dashv \Gamma_3}{\Gamma \vdash \mathbf{cases} \ pq \ r_1 \ r_2 \Leftarrow R \dashv \Gamma_2 \sqcup \Gamma_3}$$

# Positive conjunction

$$\frac{\Gamma \vdash pq \Rightarrow P \wedge Q \dashv \vdash \Gamma_1 \quad \Gamma_1, h_1 : P, h_2 : Q \vdash r \Leftarrow R \dashv \vdash \Gamma_2, h_1 : P, h_2 : Q}{\Gamma \vdash \mathbf{destruct} \text{ } pq \text{ as } h_1 \ h_2 \text{ in } r \Leftarrow R \dashv \vdash \Gamma_2}$$

To make the system more checking, it makes sense to turn conjunction positive and get rid of the projections.

# Utilities

$$\frac{\Gamma \vdash P \Leftarrow \text{prop} \dashv \Gamma_1 \quad \Gamma_1 \vdash p \Leftarrow P \dashv \Gamma_2 \quad \Gamma_2, h : P \vdash q \Leftarrow Q \dashv \Gamma_3, h : P}{\Gamma \vdash \text{lemma } h : P \text{ by } p \text{ in } q \Leftarrow Q \dashv \Gamma_3}$$

$$\frac{\Gamma \vdash P \Leftarrow \text{prop} \dashv \Gamma_1 \quad \Gamma \vdash q \Leftarrow P \Rightarrow Q \dashv \Gamma_2 \quad \Gamma_2 \vdash p \Leftarrow P \dashv \Gamma_3}{\Gamma \vdash \text{suffices } P \text{ by } q \text{ in } p \Leftarrow Q \dashv \Gamma_2}$$

# Quantifiers

$$\frac{\Gamma \vdash p \Leftarrow P[x := y] \dashv \Gamma'}{\Gamma \vdash \text{pick-any } y \text{ in } p \Leftarrow \forall x : A. P \dashv \Gamma'}$$

$$\frac{\Gamma \vdash p \Rightarrow \forall x : A. P \dashv \Gamma_1 \quad \Gamma_1 \vdash_{\text{nc}} e \Leftarrow A \dashv \Gamma_2}{\Gamma \vdash \text{instantiate } p \text{ with } e \Rightarrow P[x := e] \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash_{\text{nc}} e \Leftarrow A \dashv \Gamma_1 \quad \Gamma_1 \vdash p \Leftarrow P[x := e] \dashv \Gamma_2}{\Gamma \vdash \text{witness } e \text{ such that } p \Leftarrow \exists x : A. P \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash p \Rightarrow \exists x : A. P \dashv \Gamma_1 \quad \Gamma_1, y : A, h : P[x := y] \vdash q \Leftarrow Q \dashv \Gamma_2}{\Gamma \vdash \text{pick-witness } y \text{ h for } p \text{ in } q \Leftarrow Q \dashv \Gamma_2}$$

# Equality

$$\frac{\Gamma \vdash e_1 \equiv e_2 \Leftarrow A \vdash \Gamma'}{\Gamma \vdash \mathbf{refl} \Leftarrow e_1 =_A e_2 \vdash \Gamma'}$$

$$\frac{\Gamma \vdash q \Rightarrow e_1 =_A e_2 \vdash \Gamma_1 \quad \begin{array}{l} \Gamma_1, x : A \vdash P \Leftarrow \text{prop} \vdash \Gamma_2, x : A \\ \Gamma_2 \vdash p \Leftarrow P[x := e_2] \vdash \Gamma_3 \end{array}}{\Gamma \vdash \mathbf{rewrite } q \text{ at } x.P \text{ in } p \Rightarrow P[x := e_1] \vdash \Gamma_3}$$

$$\frac{\Gamma, x : A \vdash p \Leftarrow f =_B g \vdash \Gamma', x : A}{\Gamma \vdash \mathbf{funext } x \text{ in } p \Leftarrow f =_{rA \rightarrow B} g \vdash \Gamma'}$$



# Classical logic

$$\frac{\Gamma, h : \neg P \vdash q \Leftarrow \perp \dashv \Gamma', h : \neg P}{\Gamma \vdash \text{by-contradiction } h \text{ in } q \Leftarrow P \dashv \Gamma'}$$

$$\frac{\Gamma \vdash P \Leftarrow \text{prop} \dashv \Gamma_1 \quad \Gamma_1, h : \neg P \vdash q \Leftarrow \perp \dashv \Gamma_2, h : \neg P}{\Gamma \vdash \text{by-contradiction } h : \neg P \text{ in } q \Rightarrow P \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash p \Rightarrow \exists x : A. P \dashv \Gamma'}{\Gamma \vdash_{\text{nc}} \text{choose } p \Rightarrow A \dashv \Gamma'}$$

$$\frac{\Gamma \vdash p \Rightarrow \exists x : A. P \dashv \Gamma'}{\Gamma \vdash \text{choose-spec } p \Rightarrow P[x := \text{choose } p] \dashv \Gamma'}$$

# Classical logic

$$\Gamma \vdash p \Rightarrow \exists x : A. P \dashv \Gamma_1$$

$$\Gamma, y : A := \text{choose } p, h : P[x := y] \vdash q \Leftarrow Q \dashv \Gamma_2, y : A := \text{choose } p, h$$

---


$$\Gamma \vdash \text{choose-witness } y \text{ for } p \text{ in } q \Leftarrow Q \dashv \Gamma_2$$

$$\Gamma \vdash p \Rightarrow \exists x : A. P \dashv \Gamma_1$$

$$\Gamma, y : A := \text{choose } p, h : P[x := y] \vdash_{\text{nc}} e \Rightarrow B \dashv \Gamma_2, y : A := \text{choose } p, h$$

---


$$\Gamma \vdash_{\text{nc}} \text{choose-witness } y \text{ for } p \text{ in } e \Rightarrow B \dashv \Gamma_2$$

# Type conversion

$$\frac{\Gamma \vdash A \equiv B \Rightarrow \text{Type}_s \vdash \Gamma' \quad r = s}{\Gamma \vdash A \equiv B \Leftarrow \text{Type}_r \vdash \Gamma'}$$

$$\frac{\Gamma \vdash A \triangleq B \Rightarrow \text{Type}_r \vdash \Gamma'}{\Gamma \vdash A \equiv B \Rightarrow \text{Type}_r \vdash \Gamma'}$$

# Type conversion

$$\overline{\Gamma \vdash \text{Unit} \hat{=} \text{Unit} \Rightarrow \text{Type} \vdash \Gamma}$$

$$\overline{\Gamma \vdash \text{Empty} \hat{=} \text{Empty} \Rightarrow \text{Type} \vdash \Gamma}$$

$$\frac{r_1 = r_2 \quad \Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_{s_A} \vdash \Gamma_1 \quad \Gamma_1 \vdash B_1 \equiv B_2 \Rightarrow \text{Type}_{s_B} \vdash \Gamma_2}{\Gamma \vdash r_1 A_1 \rightarrow B_1 \hat{=} r_2 A_2 \rightarrow B_2 \Rightarrow \text{Type}_1 \vdash \Gamma_2}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_s \vdash \Gamma'}{\Gamma \vdash !_0 A_1 \hat{=} !_0 A_2 \Rightarrow \text{Type} \vdash \Gamma'}$$

$$\frac{r_1 = r_2 \quad r_1 \neq 0 \quad \Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_s \vdash \Gamma'}{\Gamma \vdash !_r A_1 \hat{=} !_r A_2 \Rightarrow \text{Type}_{r_1.s} \vdash \Gamma'}$$

# Type conversion

$$\frac{\Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_{s_A} \dashv \Gamma_1 \quad \Gamma_1 \vdash B_1 \equiv B_2 \Rightarrow \text{Type}_{s_B} \dashv \Gamma_2}{\Gamma \vdash A_1 \otimes B_1 \hat{=} A_2 \otimes B_2 \Rightarrow \text{Type}_{s_A \sqcup s_B} \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_{s_A} \dashv \Gamma_1 \quad \Gamma_1 \vdash B_1 \equiv B_2 \Rightarrow \text{Type}_{s_B} \dashv \Gamma_2}{\Gamma \vdash A_1 \oplus B_1 \hat{=} A_2 \oplus B_2 \Rightarrow \text{Type}_{s_A \sqcup s_B} \dashv \Gamma_2}$$

# Type conversion

$$\frac{(a : \text{Type}_r) \in \Gamma}{\Gamma \vdash a \triangleq a \Rightarrow \text{Type}_r \dashv \Gamma}$$

$$\frac{r_1 = r_2 \quad \Gamma, a_1 : \text{Type}_{r_1} \vdash B_1 \equiv B_2 [a_2 := a_1] \Rightarrow \text{Type}_s \dashv \Gamma', a_1 : \text{Type}_{r_1}}{\Gamma \vdash \forall a_1 : \text{Type}_{r_1}. B_1 \triangleq \forall a_2 : \text{Type}_{r_2}. B_2 \Rightarrow \text{Type}_s \dashv \Gamma'}$$

$$\frac{r_1 = r_2 \quad \Gamma, a_1 : \text{Type}_{r_1} \vdash B_1 \equiv B_2 [a_2 := a_1] \Rightarrow \text{Type}_s \dashv \Gamma', a_1 : \text{Type}_{r_1}}{\Gamma \vdash \forall \{a_1 : \text{Type}_{r_1}\}. B_1 \triangleq \forall \{a_2 : \text{Type}_{r_2}\}. B_2 \Rightarrow \text{Type}_s \dashv \Gamma'}$$

# Proposition conversion

$$\frac{\Gamma \vdash P_1 \triangleq P_2 \dashv \Gamma'}{\Gamma \vdash P_1 \equiv P_2 \dashv \Gamma'}$$

# Proposition conversion

$$\overline{\Gamma \vdash \top \triangleq \top \dashv \Gamma} \quad \overline{\Gamma \vdash \perp \triangleq \perp \dashv \Gamma}$$

$$\frac{\Gamma \vdash P_1 \equiv P_2 \dashv \Gamma_1 \quad \Gamma_1 \vdash Q_1 \equiv Q_2 \dashv \Gamma_2}{\Gamma \vdash P_1 \Rightarrow Q_1 \triangleq P_2 \Rightarrow Q_2 \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash P_1 \equiv P_2 \dashv \Gamma_1 \quad \Gamma_1 \vdash Q_1 \equiv Q_2 \dashv \Gamma_2}{\Gamma \vdash P_1 \wedge Q_1 \triangleq P_2 \wedge Q_2 \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash P_1 \equiv P_2 \dashv \Gamma_1 \quad \Gamma_1 \vdash Q_1 \equiv Q_2 \dashv \Gamma_2}{\Gamma \vdash P_1 \vee Q_1 \triangleq P_2 \vee Q_2 \dashv \Gamma_2}$$



# Proposition conversion

$$\frac{\Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_r \dashv \Gamma_1 \quad \Gamma_1, x_1 : A_1 \vdash P_1 \equiv P_2[x_2 := x_1] \dashv \Gamma_2, x_1 : A_1}{\Gamma \vdash \forall x_1 : A_1. P_1 \hat{=} \forall x_2 : A_2. P_2 \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_r \dashv \Gamma_1 \quad \Gamma_1, x_1 : A_1 \vdash P_1 \equiv P_2[x_2 := x_1] \dashv \Gamma_2, x_1 : A_1}{\Gamma \vdash \exists x_1 : A_1. P_1 \hat{=} \exists x_2 : A_2. P_2 \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \Rightarrow \text{Type}_r \dashv \Gamma_1 \quad \begin{array}{l} \Gamma_1 \vdash e_1 \equiv e_2 \leftarrow A_1 \dashv \Gamma_2 \\ \Gamma_2 \vdash e'_1 \equiv e'_2 \leftarrow A_1 \dashv \Gamma_3 \end{array}}{\Gamma \vdash e_1 =_{A_1} e'_1 \hat{=} e_2 =_{A_2} e'_2 \dashv \Gamma_3}$$

# Whnfs and neutral terms

Whnfs:

$\hat{e} ::=$

$$n \mid \lambda_r x. e \mid \text{box}_r e \mid (e_1, e_2) \mid \text{inl } e \mid \text{inr } e \mid \text{unit} \\ \Lambda a : \text{Type}_r. e \mid \Lambda \{a : \text{Type}_r\}. e$$

Neutral forms:

$n ::=$

$$x \mid n e \mid n A \mid n @ A \mid \\ \text{let unit} = n \text{ in } e \mid \text{let box } x = n \text{ in } e \mid \\ \text{let } (x, y) = n \text{ in } e \mid \text{case } n \text{ of } (e_1, e_2) \mid \\ \text{choose } p \mid \text{choose-witness } x \text{ } h \text{ for } p \text{ in } n$$

Note that `Empty-elim e` is neither whnf nor neutral.

# Computation

$e_1 \longrightarrow e'_1$  – single step

$e_1 \longrightarrow^* e'_1$  – multi step

# Term conversion (all terms)

$$\overline{\Gamma \vdash e_1 \equiv e_2 \Leftarrow \text{Unit} \vdash \Gamma}$$

$$\overline{\Gamma \vdash e_1 \equiv e_2 \Leftarrow \text{Empty} \vdash \Gamma}$$

$$\overline{\Gamma \vdash \text{Empty-elim } e_1 \equiv \text{Empty-elim } e_2 \Leftarrow A \vdash \Gamma'}$$

$$\frac{A \neq \text{Unit} \quad A \neq \text{Empty} \quad e_1 \longrightarrow^* e'_1 \quad e_2 \longrightarrow^* e'_2 \quad \Gamma \vdash e'_1 \hat{=} e'_2 \Leftarrow A \vdash \Gamma'}{\Gamma \vdash e_1 \equiv e_2 \Leftarrow A \vdash \Gamma'}$$

# Term conversion (whnfs)

$$\frac{\Gamma, x : A \vdash f \ x \equiv g \ x \Leftarrow B \dashv \Gamma', x : A}{\Gamma \vdash f \hat{=} g \Leftarrow r A \rightarrow B \dashv \Gamma'}$$

$$\frac{\Gamma \vdash \text{let box } x = e_1 \text{ in } x \equiv \text{let box } x = e_2 \text{ in } x \Leftarrow A \dashv \Gamma'}{\Gamma \vdash e_1 \hat{=} e_2 \Leftarrow !_r A \dashv \Gamma'}$$

$$\frac{\begin{array}{l} \Gamma \vdash \text{let } (x, y) = e_1 \text{ in } x \equiv \text{let } (x, y) = e_2 \text{ in } x \Leftarrow A \dashv \Gamma_1 \\ \Gamma_1 \vdash \text{let } (x, y) = e_1 \text{ in } y \equiv \text{let } (x, y) = e_2 \text{ in } y \Leftarrow B \dashv \Gamma_2 \end{array}}{\Gamma \vdash e_1 \hat{=} e_2 \Leftarrow A \otimes B \dashv \Gamma_2}$$

# Term conversion (whnfs)

$$\frac{\Gamma \vdash e_1 \equiv e_2 \Leftarrow A \vdash \Gamma'}{\Gamma \vdash \text{inl } e_1 \hat{=} \text{inl } e_2 \Leftarrow A \oplus B \vdash \Gamma'}$$

$$\frac{\Gamma \vdash e_1 \equiv e_2 \Leftarrow B \vdash \Gamma'}{\Gamma \vdash \text{inr } e_1 \hat{=} \text{inr } e_2 \Leftarrow A \oplus B \vdash \Gamma'}$$

$$\frac{\Gamma, a : \text{Type}_r \vdash f \ a \equiv g \ a \Leftarrow B \vdash \Gamma'}{\Gamma \vdash f \hat{=} g \Leftarrow \forall a : \text{Type}_r. B \vdash \Gamma'}$$

$$\frac{\Gamma, a : \text{Type}_r \vdash f \ @a \equiv g \ @a \Leftarrow B \vdash \Gamma'}{\Gamma \vdash f \hat{=} g \Leftarrow \forall \{a : \text{Type}_r\}. B \vdash \Gamma'}$$

# Term conversion (neutral)

$$\frac{\Gamma(x) = A}{\Gamma \vdash x \equiv x \Rightarrow A \dashv \Gamma}$$

$$\frac{\Gamma \vdash n_1 \hat{=} n_2 \Rightarrow r A \rightarrow B \dashv \Gamma_1 \quad \Gamma_1 \vdash e_1 \equiv e_2 \Leftarrow A \dashv \Gamma_2}{\Gamma \vdash n_1 e_1 \equiv n_2 e_2 \Rightarrow B \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash n_1 \hat{=} n_2 \Rightarrow \forall a : \text{Type}_r. B \dashv \Gamma_1 \quad \Gamma_1 \vdash A_1 \equiv A_2 \Leftarrow \text{Type}_r \dashv \Gamma_2}{\Gamma \vdash n_1 A_1 \equiv n_2 A_2 \Rightarrow B[a := A] \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash n_1 \hat{=} n_2 \Rightarrow \forall \{a : \text{Type}_r\}. B \dashv \Gamma_1 \quad \Gamma_1 \vdash A_1 \equiv A_2 \Leftarrow \text{Type}_r \dashv \Gamma_2}{\Gamma \vdash n_1 @A_1 \equiv n_2 @A_2 \Rightarrow B[a := A] \dashv \Gamma_2}$$

# Term conversion (neutral)

$$\frac{\Gamma \vdash n_1 \hat{=} n_2 \Rightarrow \text{Unit} \dashv \Gamma_1 \quad \Gamma_1 \vdash e_1 \equiv e_2 \Rightarrow A \dashv \Gamma_2}{\Gamma \vdash \text{let unit} = n_1 \text{ in } e_1 \equiv \text{let unit} = n_2 \text{ in } e_2 \Rightarrow A \dashv}$$

$$\frac{\Gamma \vdash n_1 \hat{=} n_2 \Rightarrow !_r A \dashv \Gamma_1 \quad \Gamma_1, x : A \vdash e_1 \equiv e_2 \Rightarrow B \dashv \Gamma_2, x : A}{\Gamma \vdash \text{let box } x = n_1 \text{ in } e_1 \equiv \text{let box } x = n_2 \text{ in } e_2 \Rightarrow B \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash n_1 \hat{=} n_2 \Rightarrow A \otimes B \dashv \Gamma_1 \quad \Gamma_1, x : A, y : B \vdash e_1 \equiv e_2 \Rightarrow C \dashv \Gamma_2, x : A, y : B}{\Gamma \vdash \text{let } (x, y) = n_1 \text{ in } e_1 \equiv \text{let } (x, y) = n_2 \text{ in } e_2 \Rightarrow C \dashv \Gamma_2}$$

$$\frac{\Gamma \vdash e_1 \equiv e_2 \Rightarrow A \oplus B \dashv \Gamma_1 \quad \begin{array}{l} \Gamma_1 \vdash f_1 \equiv f_2 \Rightarrow 1 A \rightarrow C \dashv \Gamma_2 \\ \Gamma_1 \vdash g_1 \equiv g_2 \Rightarrow 1 B \rightarrow C \dashv \Gamma_3 \end{array}}{\Gamma \vdash \text{case } e_1 \text{ of } (f_1, g_1) \equiv \text{case } e_2 \text{ of } (f_2, g_2) \Rightarrow C \dashv \Gamma_2 \sqcup \Gamma_3}$$



# Term conversion (choice)

$$\frac{\begin{array}{l} \Gamma \vdash p_1 \Rightarrow \exists x : A_1. P_1 \dashv \Gamma_1 \\ \Gamma_1 \vdash p_2 \Rightarrow \exists x : A_2. P_2 \dashv \Gamma_2 \end{array} \quad \Gamma_2 \vdash \exists x : A_1. P_1 \equiv \exists x : A_1. P_2 \dashv \Gamma_3}{\Gamma \vdash \text{choose } p_1 \equiv \text{choose } p_2 \Rightarrow A_1 \dashv \Gamma_3}$$