

### Exercise 1: Gaussian Processes - Prediction

Let  $\mathcal{X} = \mathbb{R}$  and assume the following statistical model

$$y = f(x) + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2),$$

where  $f(x) \in \mathcal{GP}(0, k(x, x'))$ . Suppose the covariance function of the GP is

$$k(x, x') = \mathbb{1}_{[|x-x'| < 1]} \cdot (1 - |x - x'|)$$

and we have seen the training data:

$i$	$\mathbf{x}^{(i)}$	$y^{(i)}$
1	1.6	3.0
2	2.8	3.3
3	0.5	2.0
4	3.9	2.7

As a test input we observe  $x_* = 1.2$ . Recall that the predictive distribution for  $f(x_*)$  is

$$f(x_*) \mid \mathbf{X}, \mathbf{y}, x_* \sim \mathcal{N}(m_{\text{post}}, k_{\text{post}}).$$

with

$$\begin{aligned} m_{\text{post}} &= \mathbf{K}_*^T (\mathbf{K} + \sigma^2 \cdot \mathbf{I})^{-1} \mathbf{y} \\ k_{\text{post}} &= K_{**} - \mathbf{K}_*^T (\mathbf{K} + \sigma^2 \cdot \mathbf{I})^{-1} \mathbf{K}_*, \end{aligned}$$

Here,  $\mathbf{K} = (k(\mathbf{x}^{(i)}, \mathbf{x}^{(j)}))_{i,j}$ ,  $\mathbf{K}_* = (k(x_*, \mathbf{x}^{(1)}), \dots, k(x_*, \mathbf{x}^{(n)}))^T$  and  $K_{**} = k(x_*, x_*)$ .

(a) Compute the predictive mean  $m_{\text{post}}$ .

(b) Compute the predictive variance  $k_{\text{post}}$ .

(c) Repeat the calculations from (a) and (b) by using as the test input  $x_* = \mathbf{x}^{(i)}$  for each  $i = 1, 2, 3, 4$ , respectively.

(d) Based on your calculations so far, try to sketch the posterior Gaussian process.

(e) If the nugget  $\sigma^2$  would be zero, how would the posterior Gaussian process (roughly) look like?

(f) Quiz time: Log in to Particify (<https://partici.fi/63221686>) and try to answer the questions for **Week 13**.