

# Kilohertz electron paramagnetic resonance spectroscopy of single nitrogen centers at zero magnetic field



**Fei Kong<sup>1,2,3\*</sup>, Pengju Zhao<sup>1,2,3\*</sup>, Pei Yu<sup>1,2,3\*</sup>, Zhuoyang Qin<sup>1,2,3</sup>, Zhehua Huang<sup>1,2,3</sup>,  
Zhecheng Wang<sup>1,2,3</sup>, Mengqi Wang<sup>1,2,3</sup>, Fazhan Shi<sup>1,2,3†</sup>, Jiangfeng Du<sup>1,2,3†</sup>**

Hefei National Laboratory for Physical Sciences at the Microscale and Department of Modern Physics, University of Science and Technology of China, Hefei 230026, China.  
CAS Key Laboratory of Microscale Magnetic Resonance, University of Science and Technology of China, Hefei 230026, China.

Synergetic Innovation Center of Quantum Information and Quantum Physics,  
University of Science and Technology of China, Hefei 230026, China.



# NV center–based detection

NV center  
(sensor)



$S = 1/2, I = 1/2$  system

P1 center  
(target)



$$H_t = \mathbf{S} \cdot \mathbf{A} \cdot \mathbf{I}$$

$$|T_{+1}\rangle = |\uparrow\uparrow\rangle,$$

$$|S_0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle),$$

$$|T_0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle),$$

$$|T_{-1}\rangle = |\downarrow\downarrow\rangle,$$


$$H_{\text{dd}} \approx \frac{\mu_0 \gamma_{\text{NV}} \gamma_{\text{tar}} \hbar}{4\pi r^3} (\cos \theta_e - 3 \cos \theta_r \cos \theta_{r'}) S_z^{\text{NV}} S_{zz}^{\text{T}}$$

$$= C(\theta_e, r, \theta_r, \theta_{r'}) S_z^{\text{NV}} S_{zz}^{\text{T}},$$

# NV center–based zero-field EPR spectrometer

$$H_0 = A_{\perp}(S_x I_x + S_y I_y) + A_{\parallel} S_z I_z$$

$$\delta H = \sum_{j=x,y,z} \delta_j S_j$$




$$\omega_{S_0} = -\frac{A_{\perp}}{2} - \frac{A_{\parallel}}{4},$$

$$\omega_{T_0} = \frac{A_{\perp}}{2} - \frac{A_{\parallel}}{4},$$

$$\omega_{T_{\pm 1}} = \frac{A_{\parallel}}{4}$$

$$\begin{aligned} \delta \omega_{S_0} &\approx -\frac{\delta_x^2 + \delta_y^2}{2(A_{\parallel} + A_{\perp})} - \frac{\delta_z^2}{4A_{\perp}}, \\ \delta \omega_{T_0} &\approx -\frac{\delta_x^2 + \delta_y^2}{2(A_{\parallel} - A_{\perp})} + \frac{\delta_z^2}{4A_{\perp}}, \\ \delta \omega_{T_{\pm 1}} &\approx \pm \frac{\delta_z}{2} \end{aligned}$$




$$A_{\perp} = 114 \text{ MHz}$$

$$A_{\parallel} = 160 \text{ MHz}$$

Energy levels of the target spin

# modified correlation detection protocol for zero-field EPR spectroscopy



**A**



$$S_{\text{corr}} = \frac{1}{2} [1 + \langle \cos 2\varphi_1 \cos 2\varphi_2 \rangle]$$

$$|0\rangle \longrightarrow (|1\rangle + |-1\rangle)/\sqrt{2} \longrightarrow (e^{i\varphi_1} |1\rangle + e^{-i\varphi_1} |-1\rangle)/\sqrt{2}$$


$$\cos \varphi_1 |\psi_+\rangle + i \sin \varphi_1 |\psi_-\rangle$$




Rabi oscillations for the  $ST_{\pm 1}$  and  $ST_0$  transitions.  
 $f = 30 \text{ MHz}$ ,  $T_{1p} \sim 150 \text{ us}$

spin-locking  
relaxation time

# High-resolution EPR spectroscopy of single P1 centers




FWHM:  
 $230 \pm 20$  kHz and  $260 \pm 20$  kHz for  $ST_{\pm 1}$   
 $11.6 \pm 0.6$  kHz and  $8.6 \pm 0.4$  kHz for  $ST_0$



Magnetic-field dependence  
 of the line shape  
 $ST_{\pm 1}$  and  $ST_0$

# Coherence properties of the NV center




Magnetic-field dependence of the line shape

$$H_I = \Omega_1 \cos \left[ Dt + \frac{2\Omega_2}{\Omega_1} \sin \Omega_1 t \right] S_x$$

$$H_{II} = e^{i\frac{\Omega_1 t}{2} S_x} H_I e^{-i\frac{\Omega_1 t}{2} S_x} - \frac{\Omega_1}{2} S_x$$

$$= -\frac{\Omega_2^2}{2} (S_z^2 - S_y^2)$$

# Schematics of the compensation process



Schematics of the compensation process