

Chapter 5. The Vector Space \mathbb{R}^n

- 1 Subspace and Spanning
- 2 Independence and Dimension
- 3 Orthogonality
- 4 Rank of a Matrix

5.1 Subspace and Spanning

$$\mathbb{R}^n = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, x_i \in \mathbb{R} \right\}$$

We consider \mathbb{R}^n as a set of all matrices of size $n \times 1$ that has two operations: addition and scalar multiplication. Denote

$$\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, \mathbf{e}_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

is the **standard basis** of \mathbb{R}^n . For any vector $\mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T$ can be expressed via the standard basis: $\mathbf{x} = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \cdots + x_n \mathbf{e}_n$.

Definition

A set U of vectors in \mathbb{R}^n is called a **subspace** of \mathbb{R}^n if it satisfies the following properties:

- (i) If $\mathbf{x}, \mathbf{y} \in U$ then $\mathbf{x} + \mathbf{y} \in U$.
- (ii) If $\mathbf{x} \in U$ then $a\mathbf{x} \in U$, for any $a \in \mathbb{R}$.

Definition

A set U of vectors in \mathbb{R}^n is called a **subspace** of \mathbb{R}^n if it satisfies the following properties:

- (i) If $\mathbf{x}, \mathbf{y} \in U$ then $\mathbf{x} + \mathbf{y} \in U$.
- (ii) If $\mathbf{x} \in U$ then $a\mathbf{x} \in U$, for any $a \in \mathbb{R}$.

Definition

A set U of vectors in \mathbb{R}^n is called a **subspace** of \mathbb{R}^n if it satisfies the following properties:

- (i) If $\mathbf{x}, \mathbf{y} \in U$ then $\mathbf{x} + \mathbf{y} \in U$.
- (ii) If $\mathbf{x} \in U$ then $a\mathbf{x} \in U$, for any $a \in \mathbb{R}$.

Note.

- If U is a subspace then the **zero vector** $\mathbf{0} = \begin{bmatrix} 0 & 0 & \cdots & 0 \end{bmatrix}^T \in U$.

Definition

A set U of vectors in \mathbb{R}^n is called a **subspace** of \mathbb{R}^n if it satisfies the following properties:

- (i) If $\mathbf{x}, \mathbf{y} \in U$ then $\mathbf{x} + \mathbf{y} \in U$.
- (ii) If $\mathbf{x} \in U$ then $a\mathbf{x} \in U$, for any $a \in \mathbb{R}$.

Note.

- If U is a subspace then the **zero vector** $\mathbf{0} = \begin{bmatrix} 0 & 0 & \cdots & 0 \end{bmatrix}^T \in U$.
- In \mathbb{R}^2 , all subspaces of \mathbb{R}^2 is either zero vector, or lines pass the origin, or \mathbb{R}^2 .

Definition

A set U of vectors in \mathbb{R}^n is called a **subspace** of \mathbb{R}^n if it satisfies the following properties:

- (i) If $\mathbf{x}, \mathbf{y} \in U$ then $\mathbf{x} + \mathbf{y} \in U$.
- (ii) If $\mathbf{x} \in U$ then $a\mathbf{x} \in U$, for any $a \in \mathbb{R}$.

Note.

- If U is a subspace then the **zero vector** $\mathbf{0} = \begin{bmatrix} 0 & 0 & \cdots & 0 \end{bmatrix}^T \in U$.
- In \mathbb{R}^2 , all subspaces of \mathbb{R}^2 is either zero vector, or lines pass the origin, or \mathbb{R}^2 .
- In \mathbb{R}^3 , all subspaces of \mathbb{R}^3 is either zero vector, or lines pass the origin, or planes containing the origin, or \mathbb{R}^3 .

Definition

A set U of vectors in \mathbb{R}^n is called a **subspace** of \mathbb{R}^n if it satisfies the following properties:

- (i) If $\mathbf{x}, \mathbf{y} \in U$ then $\mathbf{x} + \mathbf{y} \in U$.
- (ii) If $\mathbf{x} \in U$ then $a\mathbf{x} \in U$, for any $a \in \mathbb{R}$.

Note.

- If U is a subspace then the **zero vector** $\mathbf{0} = \begin{bmatrix} 0 & 0 & \cdots & 0 \end{bmatrix}^T \in U$.
- In \mathbb{R}^2 , all subspaces of \mathbb{R}^2 is either zero vector, or lines pass the origin, or \mathbb{R}^2 .
- In \mathbb{R}^3 , all subspaces of \mathbb{R}^3 is either zero vector, or lines pass the origin, or planes containing the origin, or \mathbb{R}^3 .
- ...

Definition

A set U of vectors in \mathbb{R}^n is called a **subspace** of \mathbb{R}^n if it satisfies the following properties:

- (i) If $\mathbf{x}, \mathbf{y} \in U$ then $\mathbf{x} + \mathbf{y} \in U$.
- (ii) If $\mathbf{x} \in U$ then $a\mathbf{x} \in U$, for any $a \in \mathbb{R}$.

Note.

- If U is a subspace then the **zero vector** $\mathbf{0} = \begin{bmatrix} 0 & 0 & \cdots & 0 \end{bmatrix}^T \in U$.
- In \mathbb{R}^2 , all subspaces of \mathbb{R}^2 is either zero vector, or lines pass the origin, or \mathbb{R}^2 .
- In \mathbb{R}^3 , all subspaces of \mathbb{R}^3 is either zero vector, or lines pass the origin, or planes containing the origin, or \mathbb{R}^3 .
- ...

Example

Which of the following are subspaces of \mathbb{R}^3 ?

(i) $U = \{(x, y, z) | z = 2x + 3y + 2\}$

(ii) $V = \{(x, y, z) | x^2 + y^2 = 2z^2\}$

Answer:

Example

Which of the following are subspaces of \mathbb{R}^3 ?

(i) $U = \{(x, y, z) | z = 2x + 3y + 2\}$

(ii) $V = \{(x, y, z) | x^2 + y^2 = 2z^2\}$

Answer:

- (i) Since $\mathbf{0} = \begin{bmatrix} 0 & 0 & 0 \end{bmatrix}^T$ is not in U . Hence U is not a subspace.
(U is a plane which does not pass through the origin in \mathbb{R}^3)

Example

Which of the following are subspaces of \mathbb{R}^3 ?

- (i) $U = \{(x, y, z) | z = 2x + 3y + 2\}$
- (ii) $V = \{(x, y, z) | x^2 + y^2 = 2z^2\}$

Answer:

- (i) Since $\mathbf{0} = \begin{bmatrix} 0 & 0 & 0 \end{bmatrix}^T$ is not in U . Hence U is not a subspace.
(U is a plane which does not pass through the origin in \mathbb{R}^3)
- (ii) V is not closed under addition (although V is closed under scalar multiplication as well as $\mathbf{0} \in V$). Indeed, we can see that

$$\mathbf{v}_1 = \begin{bmatrix} 2 & 2 & 2 \end{bmatrix}^T, \mathbf{v}_2 = \begin{bmatrix} 2 & 2 & -2 \end{bmatrix}^T \in V$$

but $\mathbf{v}_1 + \mathbf{v}_2 = \begin{bmatrix} 4 & 4 & 0 \end{bmatrix}^T \notin V$. Hence V is not a subspace.
(V is a cone in the space \mathbb{R}^3)

Example

Which of the following are subspaces of \mathbb{R}^3 ?

- (i) $U = \{(x, y, z) | z = 2x + 3y + 2\}$
- (ii) $V = \{(x, y, z) | x^2 + y^2 = 2z^2\}$

Answer:

- (i) Since $\mathbf{0} = \begin{bmatrix} 0 & 0 & 0 \end{bmatrix}^T$ is not in U . Hence U is not a subspace.
(U is a plane which does not pass through the origin in \mathbb{R}^3)
- (ii) V is not closed under addition (although V is closed under scalar multiplication as well as $\mathbf{0} \in V$). Indeed, we can see that

$$\mathbf{v}_1 = \begin{bmatrix} 2 & 2 & 2 \end{bmatrix}^T, \mathbf{v}_2 = \begin{bmatrix} 2 & 2 & -2 \end{bmatrix}^T \in V$$

but $\mathbf{v}_1 + \mathbf{v}_2 = \begin{bmatrix} 4 & 4 & 0 \end{bmatrix}^T \notin V$. Hence V is not a subspace.
(V is a cone in the space \mathbb{R}^3)

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

You can check by definition. For any $\mathbf{u} = (x_1, y_1)$, $\mathbf{v} = (x_2, y_2)$ in U_1 and $a \in \mathbb{R}$, then we have $2x_1 + 5y_1 = 0$ and $2x_2 + 5y_2 = 0$. Hence

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

You can check by definition. For any $\mathbf{u} = (x_1, y_1)$, $\mathbf{v} = (x_2, y_2)$ in U_1 and $a \in \mathbb{R}$, then we have $2x_1 + 5y_1 = 0$ and $2x_2 + 5y_2 = 0$. Hence

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \in U_1$$

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

You can check by definition. For any $\mathbf{u} = (x_1, y_1)$, $\mathbf{v} = (x_2, y_2)$ in U_1 and $a \in \mathbb{R}$, then we have $2x_1 + 5y_1 = 0$ and $2x_2 + 5y_2 = 0$. Hence

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \in U_1$$

since $2(x_1 + x_2) + 5(y_1 + y_2) = (2x_1 + 5y_1) + (2x_2 + 5y_2) = 0$. And

$$a\mathbf{u} = (ax_1, ay_1) \in U_1$$

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

You can check by definition. For any $\mathbf{u} = (x_1, y_1)$, $\mathbf{v} = (x_2, y_2)$ in U_1 and $a \in \mathbb{R}$, then we have $2x_1 + 5y_1 = 0$ and $2x_2 + 5y_2 = 0$. Hence

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \in U_1$$

since $2(x_1 + x_2) + 5(y_1 + y_2) = (2x_1 + 5y_1) + (2x_2 + 5y_2) = 0$. And

$$a\mathbf{u} = (ax_1, ay_1) \in U_1$$

since $2ax_1 + 5ay_1 = a(2x_1 + 5y_1) = 0$.

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

You can check by definition. For any $\mathbf{u} = (x_1, y_1)$, $\mathbf{v} = (x_2, y_2)$ in U_1 and $a \in \mathbb{R}$, then we have $2x_1 + 5y_1 = 0$ and $2x_2 + 5y_2 = 0$. Hence

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \in U_1$$

since $2(x_1 + x_2) + 5(y_1 + y_2) = (2x_1 + 5y_1) + (2x_2 + 5y_2) = 0$. And

$$a\mathbf{u} = (ax_1, ay_1) \in U_1$$

since $2ax_1 + 5ay_1 = a(2x_1 + 5y_1) = 0$.

(ii) U_2 is the y -axis (a line pass through the origin). Hence it is a subspace of \mathbb{R}^2 .

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

You can check by definition. For any $\mathbf{u} = (x_1, y_1)$, $\mathbf{v} = (x_2, y_2)$ in U_1 and $a \in \mathbb{R}$, then we have $2x_1 + 5y_1 = 0$ and $2x_2 + 5y_2 = 0$. Hence

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \in U_1$$

since $2(x_1 + x_2) + 5(y_1 + y_2) = (2x_1 + 5y_1) + (2x_2 + 5y_2) = 0$. And

$$a\mathbf{u} = (ax_1, ay_1) \in U_1$$

since $2ax_1 + 5ay_1 = a(2x_1 + 5y_1) = 0$.

(ii) U_2 is the y -axis (a line pass through the origin). Hence it is a subspace of \mathbb{R}^2 .

Example

Which of the following are subspaces of \mathbb{R}^2 ?

(i) $U_1 = \{(x, y) : 2x + 5y = 0\}$

(ii) $U_2 = \{(x, y) : x = 0\}$

(i) U_1 is the line through the origin with normal vector $\begin{bmatrix} 2 & 5 \end{bmatrix}^T$ in the plane \mathbb{R}^2 . Hence U_1 is a subspace of \mathbb{R}^2 .

You can check by definition. For any $\mathbf{u} = (x_1, y_1)$, $\mathbf{v} = (x_2, y_2)$ in U_1 and $a \in \mathbb{R}$, then we have $2x_1 + 5y_1 = 0$ and $2x_2 + 5y_2 = 0$. Hence

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \in U_1$$

since $2(x_1 + x_2) + 5(y_1 + y_2) = (2x_1 + 5y_1) + (2x_2 + 5y_2) = 0$. And

$$a\mathbf{u} = (ax_1, ay_1) \in U_1$$

since $2ax_1 + 5ay_1 = a(2x_1 + 5y_1) = 0$.

(ii) U_2 is the y -axis (a line pass through the origin). Hence it is a subspace of \mathbb{R}^2 .

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- Null space of A :

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- Null space of A :

$$\text{null}(A) = \left\{ \mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0} \right\} \subseteq \mathbb{R}^n$$

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- Null space of A :

$$\text{null}(A) = \left\{ \mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0} \right\} \subseteq \mathbb{R}^n$$

the set of all solutions of homogeneous system $A\mathbf{x} = \mathbf{0}$.

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- **Null space** of A :

$$\text{null}(A) = \left\{ \mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0} \right\} \subseteq \mathbb{R}^n$$

the set of all solutions of homogeneous system $A\mathbf{x} = \mathbf{0}$.

- **Image space** of A :

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- **Null space** of A :

$$\text{null}(A) = \left\{ \mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0} \right\} \subseteq \mathbb{R}^n$$

the set of all solutions of homogeneous system $A\mathbf{x} = \mathbf{0}$.

- **Image space** of A :

$$\text{im}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} \subseteq \mathbb{R}^m$$

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- **Null space** of A :

$$\text{null}(A) = \left\{ \mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0} \right\} \subseteq \mathbb{R}^n$$

the set of all solutions of homogeneous system $A\mathbf{x} = \mathbf{0}$.

- **Image space** of A :

$$\text{im}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} \subseteq \mathbb{R}^m$$

is the set of all $\mathbf{b} \in \mathbb{R}^m$ such that the system $A\mathbf{x} = \mathbf{b}$ has at least one solution.

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- **Null space** of A :

$$\text{null}(A) = \left\{ \mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0} \right\} \subseteq \mathbb{R}^n$$

the set of all solutions of homogeneous system $A\mathbf{x} = \mathbf{0}$.

- **Image space** of A :

$$\text{im}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} \subseteq \mathbb{R}^m$$

is the set of all $\mathbf{b} \in \mathbb{R}^m$ such that the system $A\mathbf{x} = \mathbf{b}$ has at least one solution.

Let A be an $m \times n$ matrix.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \\ \mathbf{x} & \mapsto & A\mathbf{x} \end{array}$$

Definition

- **Null space** of A :

$$\text{null}(A) = \left\{ \mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0} \right\} \subseteq \mathbb{R}^n$$

the set of all solutions of homogeneous system $A\mathbf{x} = \mathbf{0}$.

- **Image space** of A :

$$\text{im}(A) = \{ A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n \} \subseteq \mathbb{R}^m$$

is the set of all $\mathbf{b} \in \mathbb{R}^m$ such that the system $A\mathbf{x} = \mathbf{b}$ has at least one solution.

Example

Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \\ 0 & 2 \end{bmatrix}$. We have

$$\text{null}(A) = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\} = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid \begin{cases} x - y = 0 \\ 2x + 3y = 0 \\ 2y = 0 \end{cases} \right\} = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$$

$$\text{im}(A) = \left\{ A \begin{bmatrix} x \\ y \end{bmatrix} \mid \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2 \right\} = \left\{ \begin{bmatrix} x - y \\ 2x + 3y \\ 2y \end{bmatrix} \mid x, y \in \mathbb{R} \right\}$$

Example

Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \\ 0 & 2 \end{bmatrix}$. We have

$$\text{null}(A) = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\} = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid \begin{cases} x - y = 0 \\ 2x + 3y = 0 \\ 2y = 0 \end{cases} \right\} = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$$

$$\text{im}(A) = \left\{ A \begin{bmatrix} x \\ y \end{bmatrix} \mid \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2 \right\} = \left\{ \begin{bmatrix} x - y \\ 2x + 3y \\ 2y \end{bmatrix} \mid x, y \in \mathbb{R} \right\}$$

Example

Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \\ 0 & 2 \end{bmatrix}$. We have

$$\text{null}(A) = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\} = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid \begin{cases} x - y = 0 \\ 2x + 3y = 0 \\ 2y = 0 \end{cases} \right\} = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$$

$$\text{im}(A) = \left\{ A \begin{bmatrix} x \\ y \end{bmatrix} \mid \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2 \right\} = \left\{ \begin{bmatrix} x - y \\ 2x + 3y \\ 2y \end{bmatrix} \mid x, y \in \mathbb{R} \right\}$$

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .
- $\text{im}(A)$ is a subspace of \mathbb{R}^m .

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .
- $\text{im}(A)$ is a subspace of \mathbb{R}^m .

Example

The set

$$U = \left\{ \begin{bmatrix} x & y & z \end{bmatrix}^T \in \mathbb{R}^3 \mid 2x + y - z = 0 \text{ and } 3x - y + 5z = 0 \right\}$$

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .
- $\text{im}(A)$ is a subspace of \mathbb{R}^m .

Example

The set

$$U = \left\{ \begin{bmatrix} x & y & z \end{bmatrix}^T \in \mathbb{R}^3 \mid 2x + y - z = 0 \text{ and } 3x - y + 5z = 0 \right\}$$

is a subspace of \mathbb{R}^3 . In fact, $U = \text{null}(A)$, for $A = \begin{bmatrix} 2 & 1 & -1 \\ 3 & -1 & 5 \end{bmatrix}$.

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .
- $\text{im}(A)$ is a subspace of \mathbb{R}^m .

Example

The set

$$U = \left\{ \begin{bmatrix} x & y & z \end{bmatrix}^T \in \mathbb{R}^3 \mid 2x + y - z = 0 \text{ and } 3x - y + 5z = 0 \right\}$$

is a subspace of \mathbb{R}^3 . In fact, $U = \text{null}(A)$, for $A = \begin{bmatrix} 2 & 1 & -1 \\ 3 & -1 & 5 \end{bmatrix}$.

The set

$$V = \left\{ \begin{bmatrix} 2x - y + z + t & y - 3z - t \end{bmatrix}^T \mid x, y, z, t \text{ are real numbers} \right\}$$

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .
- $\text{im}(A)$ is a subspace of \mathbb{R}^m .

Example

The set

$$U = \left\{ \begin{bmatrix} x & y & z \end{bmatrix}^T \in \mathbb{R}^3 \mid 2x + y - z = 0 \text{ and } 3x - y + 5z = 0 \right\}$$

is a subspace of \mathbb{R}^3 . In fact, $U = \text{null}(A)$, for $A = \begin{bmatrix} 2 & 1 & -1 \\ 3 & -1 & 5 \end{bmatrix}$.

The set

$$V = \left\{ \begin{bmatrix} 2x - y + z + t & y - 3z - t \end{bmatrix}^T \mid x, y, z, t \text{ are real numbers} \right\} \text{ is a}$$

subspace of \mathbb{R}^2 .

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .
- $\text{im}(A)$ is a subspace of \mathbb{R}^m .

Example

The set

$$U = \left\{ \begin{bmatrix} x & y & z \end{bmatrix}^T \in \mathbb{R}^3 \mid 2x + y - z = 0 \text{ and } 3x - y + 5z = 0 \right\}$$

is a subspace of \mathbb{R}^3 . In fact, $U = \text{null}(A)$, for $A = \begin{bmatrix} 2 & 1 & -1 \\ 3 & -1 & 5 \end{bmatrix}$.

The set

$V = \left\{ \begin{bmatrix} 2x - y + z + t & y - 3z - t \end{bmatrix}^T \mid x, y, z, t \text{ are real numbers} \right\}$ is a subspace of \mathbb{R}^2 . In fact, $V = \text{im}(B)$ with $B = \begin{bmatrix} 2 & -1 & 1 & 1 \\ 0 & 1 & -3 & -1 \end{bmatrix}$.

Theorem

For any matrix A of size $m \times n$,

- $\text{null}(A)$ is a subspace of \mathbb{R}^n .
- $\text{im}(A)$ is a subspace of \mathbb{R}^m .

Example

The set

$$U = \left\{ \begin{bmatrix} x & y & z \end{bmatrix}^T \in \mathbb{R}^3 \mid 2x + y - z = 0 \text{ and } 3x - y + 5z = 0 \right\}$$

is a subspace of \mathbb{R}^3 . In fact, $U = \text{null}(A)$, for $A = \begin{bmatrix} 2 & 1 & -1 \\ 3 & -1 & 5 \end{bmatrix}$.

The set

$V = \left\{ \begin{bmatrix} 2x - y + z + t & y - 3z - t \end{bmatrix}^T \mid x, y, z, t \text{ are real numbers} \right\}$ is a subspace of \mathbb{R}^2 . In fact, $V = \text{im}(B)$ with $B = \begin{bmatrix} 2 & -1 & 1 & 1 \\ 0 & 1 & -3 & -1 \end{bmatrix}$.

Spanning sets

Given vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ in \mathbb{R}^n .

Definition

- A vector of the form

$$t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \cdots + t_k\mathbf{x}_k$$

where the $t_i \in \mathbb{R}$ are **scalars** is called a **linear combination** of the \mathbf{x}_i , and t_i is called the **coefficient** of \mathbf{x}_i in the linear combination.

Spanning sets

Given vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ in \mathbb{R}^n .

Definition

- A vector of the form

$$t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \cdots + t_k\mathbf{x}_k$$

where the $t_i \in \mathbb{R}$ are **scalars** is called a **linear combination** of the \mathbf{x}_i , and t_i is called the **coefficient** of \mathbf{x}_i in the linear combination. • The set of all such linear combinations is called the **span** of the \mathbf{x}_i and is denoted

$$\text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\} = \left\{ t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \cdots + t_k\mathbf{x}_k \mid t_i \in \mathbb{R} \right\}$$

If $V = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$, we say that V is **spanned by** the vectors $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$.

Spanning sets

Given vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ in \mathbb{R}^n .

Definition

- A vector of the form

$$t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \cdots + t_k\mathbf{x}_k$$

where the $t_i \in \mathbb{R}$ are **scalars** is called a **linear combination** of the \mathbf{x}_i , and t_i is called the **coefficient** of \mathbf{x}_i in the linear combination. • The set of all such linear combinations is called the **span** of the \mathbf{x}_i and is denoted

$$\text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\} = \left\{ t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \cdots + t_k\mathbf{x}_k \mid t_i \in \mathbb{R} \right\}$$

If $V = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$, we say that V is **spanned by** the vectors $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$.

Example

Let $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \in \mathbb{R}^2$. Then

$$\text{span}\{\mathbf{x}\} = \{t\mathbf{x} \mid t \in \mathbb{R}\} = \left\{ \begin{bmatrix} t \\ 2t \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

Example

Let $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \in \mathbb{R}^2$. Then

$$\text{span}\{\mathbf{x}\} = \{t\mathbf{x} \mid t \in \mathbb{R}\} = \left\{ \begin{bmatrix} t \\ 2t \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

is the line through the origin with direction vector $\begin{bmatrix} 1 & 2 \end{bmatrix}^T$ in the plane \mathbb{R}^2 .

Example

Let $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \in \mathbb{R}^2$. Then

$$\text{span}\{\mathbf{x}\} = \{t\mathbf{x} \mid t \in \mathbb{R}\} = \left\{ \begin{bmatrix} t \\ 2t \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

is the line through the origin with direction vector $\begin{bmatrix} 1 & 2 \end{bmatrix}^T$ in the plane \mathbb{R}^2 .

Example

Given two vectors $\mathbf{u} = \begin{bmatrix} -1 & 0 & 1 \end{bmatrix}^T$ and $\mathbf{v} = \begin{bmatrix} 2 & 1 & 3 \end{bmatrix}^T$ in \mathbb{R}^3 .

Example

Let $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \in \mathbb{R}^2$. Then

$$\text{span}\{\mathbf{x}\} = \{t\mathbf{x} \mid t \in \mathbb{R}\} = \left\{ \begin{bmatrix} t \\ 2t \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

is the line through the origin with direction vector $\begin{bmatrix} 1 & 2 \end{bmatrix}^T$ in the plane \mathbb{R}^2 .

Example

Given two vectors $\mathbf{u} = \begin{bmatrix} -1 & 0 & 1 \end{bmatrix}^T$ and $\mathbf{v} = \begin{bmatrix} 2 & 1 & 3 \end{bmatrix}^T$ in \mathbb{R}^3 . Then

$$\text{span}\{\mathbf{u}, \mathbf{v}\} = \{a\mathbf{u} + b\mathbf{v} \mid a, b \in \mathbb{R}\} = \left\{ \begin{bmatrix} -a + 2b \\ b \\ a + 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

Example

Let $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \in \mathbb{R}^2$. Then

$$\text{span}\{\mathbf{x}\} = \{t\mathbf{x} \mid t \in \mathbb{R}\} = \left\{ \begin{bmatrix} t \\ 2t \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

is the line through the origin with direction vector $\begin{bmatrix} 1 & 2 \end{bmatrix}^T$ in the plane \mathbb{R}^2 .

Example

Given two vectors $\mathbf{u} = \begin{bmatrix} -1 & 0 & 1 \end{bmatrix}^T$ and $\mathbf{v} = \begin{bmatrix} 2 & 1 & 3 \end{bmatrix}^T$ in \mathbb{R}^3 . Then

$$\text{span}\{\mathbf{u}, \mathbf{v}\} = \{a\mathbf{u} + b\mathbf{v} \mid a, b \in \mathbb{R}\} = \left\{ \begin{bmatrix} -a + 2b \\ b \\ a + 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

is a plane through the origin with a normal vector $\mathbf{u} \times \mathbf{v} = \begin{bmatrix} -1 & 5 & -1 \end{bmatrix}^T$.

Example

Let $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \in \mathbb{R}^2$. Then

$$\text{span}\{\mathbf{x}\} = \{t\mathbf{x} \mid t \in \mathbb{R}\} = \left\{ \begin{bmatrix} t \\ 2t \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

is the line through the origin with direction vector $\begin{bmatrix} 1 & 2 \end{bmatrix}^T$ in the plane \mathbb{R}^2 .

Example

Given two vectors $\mathbf{u} = \begin{bmatrix} -1 & 0 & 1 \end{bmatrix}^T$ and $\mathbf{v} = \begin{bmatrix} 2 & 1 & 3 \end{bmatrix}^T$ in \mathbb{R}^3 . Then

$$\text{span}\{\mathbf{u}, \mathbf{v}\} = \{a\mathbf{u} + b\mathbf{v} \mid a, b \in \mathbb{R}\} = \left\{ \begin{bmatrix} -a + 2b \\ b \\ a + 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

is a plane through the origin with a normal vector $\mathbf{u} \times \mathbf{v} = \begin{bmatrix} -1 & 5 & -1 \end{bmatrix}^T$.

Example

Find all values of a so that the vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$.

Answer:

Example

Find all values of a so that the vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$.

Answer: The vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$ if and only if the linear system

$$[5 \ 3 \ a]^T = s[3 \ 2 \ 0]^T + t[1 \ 0 \ 3]^T$$

has at least one solution s, t .

Example

Find all values of a so that the vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$.

Answer: The vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$ if and only if the linear system

$$[5 \ 3 \ a]^T = s[3 \ 2 \ 0]^T + t[1 \ 0 \ 3]^T$$

has at least one solution s, t . The above system is

$$5 = 3s + t, \quad 3 = 2s, \quad a = 3t.$$

Example

Find all values of a so that the vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$.

Answer: The vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$ if and only if the linear system

$$[5 \ 3 \ a]^T = s[3 \ 2 \ 0]^T + t[1 \ 0 \ 3]^T$$

has at least one solution s, t . The above system is

$$5 = 3s + t, \quad 3 = 2s, \quad a = 3t.$$

We find $s = 3/2, t = 1/2$ and $a = 3/2$.

Example

Find all values of a so that the vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$.

Answer: The vector $[5 \ 3 \ a]^T$ is in $\text{span}\{[3 \ 2 \ 0]^T, [1 \ 0 \ 3]^T\}$ if and only if the linear system

$$[5 \ 3 \ a]^T = s[3 \ 2 \ 0]^T + t[1 \ 0 \ 3]^T$$

has at least one solution s, t . The above system is

$$5 = 3s + t, \quad 3 = 2s, \quad a = 3t.$$

We find $s = 3/2, t = 1/2$ and $a = 3/2$.

Example

$$\begin{aligned} U &= \left\{ \begin{bmatrix} x + y - 3z & 2x - 3y + z & -x + 2y + 4z \end{bmatrix}^T \mid x, y, z \text{ are real} \right\} \\ &= \text{im} \left(\begin{bmatrix} 1 & 1 & -3 \\ 2 & -3 & 1 \\ -1 & 2 & 4 \end{bmatrix} \right) \\ &= \left\{ x \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} + y \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix} + z \begin{bmatrix} -3 \\ 1 \\ 4 \end{bmatrix} \mid x, y, z \text{ are real} \right\} \\ &= \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix}, \begin{bmatrix} -3 \\ 1 \\ 4 \end{bmatrix} \right\} \end{aligned}$$

Example

Let $\mathbf{x} = (2, -1, 1, 1)$ and $\mathbf{y} = (1, 1, 3, 4)$ in \mathbb{R}^4 . Determine whether $\mathbf{p} = (0, -2, 4, 5)$ or $\mathbf{q} = (7, -5, 1, 0)$ are in $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$.

Answer:

Example

Let $\mathbf{x} = (2, -1, 1, 1)$ and $\mathbf{y} = (1, 1, 3, 4)$ in \mathbb{R}^4 . Determine whether $\mathbf{p} = (0, -2, 4, 5)$ or $\mathbf{q} = (7, -5, 1, 0)$ are in $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$.

Answer:

- The vector \mathbf{p} is in U if and only if $\mathbf{p} = s\mathbf{x} + t\mathbf{y}$ for scalars s and t .
Equating components gives equations

$$2s + t = 0, \quad -s + t = -2, \quad s + 3t = 4, \quad s + 4t = 5$$

This system has no solution, hence $\mathbf{p} \notin U$.

Example

Let $\mathbf{x} = (2, -1, 1, 1)$ and $\mathbf{y} = (1, 1, 3, 4)$ in \mathbb{R}^4 . Determine whether $\mathbf{p} = (0, -2, 4, 5)$ or $\mathbf{q} = (7, -5, 1, 0)$ are in $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$.

Answer:

- The vector \mathbf{p} is in U if and only if $\mathbf{p} = s\mathbf{x} + t\mathbf{y}$ for scalars s and t .
Equating components gives equations

$$2s + t = 0, \quad -s + t = -2, \quad s + 3t = 4, \quad s + 4t = 5$$

This system has no solution, hence $\mathbf{p} \notin U$.

- The vector \mathbf{q} is in U if and only if $\mathbf{q} = a\mathbf{x} + b\mathbf{y}$ for scalars a and b .
This gives equations

$$2a + b = 7, \quad -a + b = -5, \quad a + 3b = 1, \quad a + 4b = 0$$

This system has solution $a = 4, b = -1$, hence $\mathbf{q} \in U$.

Example

Let $\mathbf{x} = (2, -1, 1, 1)$ and $\mathbf{y} = (1, 1, 3, 4)$ in \mathbb{R}^4 . Determine whether $\mathbf{p} = (0, -2, 4, 5)$ or $\mathbf{q} = (7, -5, 1, 0)$ are in $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$.

Answer:

- The vector \mathbf{p} is in U if and only if $\mathbf{p} = s\mathbf{x} + t\mathbf{y}$ for scalars s and t .
Equating components gives equations

$$2s + t = 0, \quad -s + t = -2, \quad s + 3t = 4, \quad s + 4t = 5$$

This system has no solution, hence $\mathbf{p} \notin U$.

- The vector \mathbf{q} is in U if and only if $\mathbf{q} = a\mathbf{x} + b\mathbf{y}$ for scalars a and b .
This gives equations

$$2a + b = 7, \quad -a + b = -5, \quad a + 3b = 1, \quad a + 4b = 0$$

This system has solution $a = 4, b = -1$, hence $\mathbf{q} \in U$.

Theorem

Let $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n . Then:

1. U is a subspace of \mathbb{R}^n containing each \mathbf{x}_i .

Theorem

Let $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n . Then:

1. U is a subspace of \mathbb{R}^n containing each \mathbf{x}_i .
2. If W is a subspace of \mathbb{R}^n and each $\mathbf{x}_i \in W$, then $U \subseteq W$.

Example

If $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$ in \mathbb{R}^n , show that $U = \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Theorem

Let $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n . Then:

1. U is a subspace of \mathbb{R}^n containing each \mathbf{x}_i .
2. If W is a subspace of \mathbb{R}^n and each $\mathbf{x}_i \in W$, then $U \subseteq W$.

Example

If $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$ in \mathbb{R}^n , show that $U = \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Answer:

Theorem

Let $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n . Then:

1. U is a subspace of \mathbb{R}^n containing each \mathbf{x}_i .
2. If W is a subspace of \mathbb{R}^n and each $\mathbf{x}_i \in W$, then $U \subseteq W$.

Example

If $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$ in \mathbb{R}^n , show that $U = \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Answer: Since $\mathbf{x} + \mathbf{y}$ and $\mathbf{x} - \mathbf{y}$ are in $\text{span}\{\mathbf{x}, \mathbf{y}\}$, thus $\text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\} \subseteq U$.

Theorem

Let $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n . Then:

1. U is a subspace of \mathbb{R}^n containing each \mathbf{x}_i .
2. If W is a subspace of \mathbb{R}^n and each $\mathbf{x}_i \in W$, then $U \subseteq W$.

Example

If $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$ in \mathbb{R}^n , show that $U = \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Answer: Since $\mathbf{x} + \mathbf{y}$ and $\mathbf{x} - \mathbf{y}$ are in $\text{span}\{\mathbf{x}, \mathbf{y}\}$, thus $\text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\} \subseteq U$.

Moreover, we have

$$\begin{aligned}\mathbf{x} &= \frac{1}{2}(\mathbf{x} + \mathbf{y}) + \frac{1}{2}(\mathbf{x} - \mathbf{y}) \\ \mathbf{y} &= \frac{1}{2}(\mathbf{x} + \mathbf{y}) - \frac{1}{2}(\mathbf{x} - \mathbf{y}).\end{aligned}$$

Theorem

Let $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n . Then:

1. U is a subspace of \mathbb{R}^n containing each \mathbf{x}_i .
2. If W is a subspace of \mathbb{R}^n and each $\mathbf{x}_i \in W$, then $U \subseteq W$.

Example

If $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$ in \mathbb{R}^n , show that $U = \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Answer: Since $\mathbf{x} + \mathbf{y}$ and $\mathbf{x} - \mathbf{y}$ are in $\text{span}\{\mathbf{x}, \mathbf{y}\}$, thus $\text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\} \subseteq U$.

Moreover, we have

$$\begin{aligned}\mathbf{x} &= \frac{1}{2}(\mathbf{x} + \mathbf{y}) + \frac{1}{2}(\mathbf{x} - \mathbf{y}) \\ \mathbf{y} &= \frac{1}{2}(\mathbf{x} + \mathbf{y}) - \frac{1}{2}(\mathbf{x} - \mathbf{y}).\end{aligned}$$

These show that $\mathbf{x}, \mathbf{y} \in \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$. Hence $U \subseteq \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Theorem

Let $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n . Then:

1. U is a subspace of \mathbb{R}^n containing each \mathbf{x}_i .
2. If W is a subspace of \mathbb{R}^n and each $\mathbf{x}_i \in W$, then $U \subseteq W$.

Example

If $U = \text{span}\{\mathbf{x}, \mathbf{y}\}$ in \mathbb{R}^n , show that $U = \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Answer: Since $\mathbf{x} + \mathbf{y}$ and $\mathbf{x} - \mathbf{y}$ are in $\text{span}\{\mathbf{x}, \mathbf{y}\}$, thus $\text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\} \subseteq U$.

Moreover, we have

$$\begin{aligned}\mathbf{x} &= \frac{1}{2}(\mathbf{x} + \mathbf{y}) + \frac{1}{2}(\mathbf{x} - \mathbf{y}) \\ \mathbf{y} &= \frac{1}{2}(\mathbf{x} + \mathbf{y}) - \frac{1}{2}(\mathbf{x} - \mathbf{y}).\end{aligned}$$

These show that $\mathbf{x}, \mathbf{y} \in \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$. Hence $U \subseteq \text{span}\{\mathbf{x} + \mathbf{y}, \mathbf{x} - \mathbf{y}\}$.

Example

It is easy to see that $\mathbb{R}^n = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. For example

$\mathbb{R}^2 = \text{span}\left\{\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$. Indeed, for any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ can be written as $\mathbf{u} = x\mathbf{e}_1 + y\mathbf{e}_2$.

Example

Show that $\mathbb{R}^2 = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}\right\}$.

Example

It is easy to see that $\mathbb{R}^n = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. For example

$\mathbb{R}^2 = \text{span}\left\{\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$. Indeed, for any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ can be written as $\mathbf{u} = x\mathbf{e}_1 + y\mathbf{e}_2$.

Example

Show that $\mathbb{R}^2 = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}\right\}$.

Answer:

Example

It is easy to see that $\mathbb{R}^n = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. For example

$\mathbb{R}^2 = \text{span}\left\{\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$. Indeed, for any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ can be written as $\mathbf{u} = x\mathbf{e}_1 + y\mathbf{e}_2$.

Example

Show that $\mathbb{R}^2 = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}\right\}$.

Answer: Any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix}$ in \mathbb{R}^2 can be written as

$$\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} = s \begin{bmatrix} 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \frac{2x+y}{3} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \frac{-x+y}{3} \begin{bmatrix} -1 \\ 2 \end{bmatrix},$$

Example

It is easy to see that $\mathbb{R}^n = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. For example

$\mathbb{R}^2 = \text{span}\left\{\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$. Indeed, for any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ can be written as $\mathbf{u} = x\mathbf{e}_1 + y\mathbf{e}_2$.

Example

Show that $\mathbb{R}^2 = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}\right\}$.

Answer: Any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix}$ in \mathbb{R}^2 can be written as

$$\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} = s \begin{bmatrix} 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \frac{2x+y}{3} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \frac{-x+y}{3} \begin{bmatrix} -1 \\ 2 \end{bmatrix},$$

this shows that $\mathbf{u} \in \text{span}\left\{\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}\right\}$.

Example

It is easy to see that $\mathbb{R}^n = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. For example

$\mathbb{R}^2 = \text{span}\left\{\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$. Indeed, for any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ can be written as $\mathbf{u} = x\mathbf{e}_1 + y\mathbf{e}_2$.

Example

Show that $\mathbb{R}^2 = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}\right\}$.

Answer: Any vector $\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix}$ in \mathbb{R}^2 can be written as

$$\mathbf{u} = \begin{bmatrix} x \\ y \end{bmatrix} = s \begin{bmatrix} 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \frac{2x+y}{3} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \frac{-x+y}{3} \begin{bmatrix} -1 \\ 2 \end{bmatrix},$$

this shows that $\mathbf{u} \in \text{span}\left\{\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}\right\}$.

5.2 Independence and Dimension

Definition

A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n is **linearly independent** if the following holds:

"If $t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \dots + t_k\mathbf{x}_k = \mathbf{0}$ then $t_1 = t_2 = \dots = t_k = 0$ ".

5.2 Independence and Dimension

Definition

A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n is **linearly independent** if the following holds:

"If $t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \dots + t_k\mathbf{x}_k = \mathbf{0}$ then $t_1 = t_2 = \dots = t_k = 0$ ".

Otherwise, we say $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is **linearly dependent**.

5.2 Independence and Dimension

Definition

A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n is **linearly independent** if the following holds:

"If $t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \dots + t_k\mathbf{x}_k = \mathbf{0}$ then $t_1 = t_2 = \dots = t_k = 0$ ".

Otherwise, we say $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is **linearly dependent**.

Example

The set $\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ is linear independent

5.2 Independence and Dimension

Definition

A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n is **linearly independent** if the following holds:

"If $t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \dots + t_k\mathbf{x}_k = \mathbf{0}$ then $t_1 = t_2 = \dots = t_k = 0$ ".

Otherwise, we say $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is **linearly dependent**.

Example

The set $\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ is linear independent since the homogeneous system

$$s \begin{bmatrix} 2 \\ 1 \end{bmatrix} + t \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow \begin{cases} 2s - t = 0 \\ s + t = 0 \end{cases}$$

5.2 Independence and Dimension

Definition

A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n is **linearly independent** if the following holds:

"If $t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \dots + t_k\mathbf{x}_k = \mathbf{0}$ then $t_1 = t_2 = \dots = t_k = 0$ ".

Otherwise, we say $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is **linearly dependent**.

Example

The set $\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ is linear independent since the homogeneous system

$$s \begin{bmatrix} 2 \\ 1 \end{bmatrix} + t \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow \begin{cases} 2s - t = 0 \\ s + t = 0 \end{cases}$$

has only trivial solution $s = t = 0$.

5.2 Independence and Dimension

Definition

A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ in \mathbb{R}^n is **linearly independent** if the following holds:

"If $t_1\mathbf{x}_1 + t_2\mathbf{x}_2 + \dots + t_k\mathbf{x}_k = \mathbf{0}$ then $t_1 = t_2 = \dots = t_k = 0$ ".

Otherwise, we say $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is **linearly dependent**.

Example

The set $\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ is linear independent since the homogeneous system

$$s \begin{bmatrix} 2 \\ 1 \end{bmatrix} + t \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow \begin{cases} 2s - t = 0 \\ s + t = 0 \end{cases}$$

has only trivial solution $s = t = 0$.

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 2 \\ -4 \\ 1 \end{bmatrix} \right\}$ is linearly dependent.

Answer:

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 2 \\ -4 \\ 1 \end{bmatrix} \right\}$ is linearly dependent.

Answer:

1st way: We can see that $\mathbf{w} = 2\mathbf{u} - \mathbf{v}$ or

$$2\mathbf{u} - \mathbf{v} - \mathbf{w} = \mathbf{0}.$$

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 2 \\ -4 \\ 1 \end{bmatrix} \right\}$ is linearly dependent.

Answer:

1st way: We can see that $\mathbf{w} = 2\mathbf{u} - \mathbf{v}$ or

$$2\mathbf{u} - \mathbf{v} - \mathbf{w} = \mathbf{0}.$$

Hence $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent.

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 2 \\ -4 \\ 1 \end{bmatrix} \right\}$ is linearly dependent.

Answer:

1st way: We can see that $\mathbf{w} = 2\mathbf{u} - \mathbf{v}$ or

$$2\mathbf{u} - \mathbf{v} - \mathbf{w} = \mathbf{0}.$$

Hence $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent.

2nd way: The system $t_1\mathbf{u} + t_2\mathbf{v} + t_3\mathbf{w} = \mathbf{0}$

$$t_1 + 2t_3 = 0, \quad -t_1 + 2t_2 - 4t_3 = 0, \quad 2t_1 + 3t_2 + t_3 = 0$$

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 2 \\ -4 \\ 1 \end{bmatrix} \right\}$ is linearly dependent.

Answer:

1st way: We can see that $\mathbf{w} = 2\mathbf{u} - \mathbf{v}$ or

$$2\mathbf{u} - \mathbf{v} - \mathbf{w} = \mathbf{0}.$$

Hence $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent.

2nd way: The system $t_1\mathbf{u} + t_2\mathbf{v} + t_3\mathbf{w} = \mathbf{0}$

$$t_1 + 2t_3 = 0, \quad -t_1 + 2t_2 - 4t_3 = 0, \quad 2t_1 + 3t_2 + t_3 = 0$$

has nontrivial solutions since $\text{rank} \begin{bmatrix} 1 & 0 & 2 \\ -1 & 2 & -4 \\ 2 & 3 & 1 \end{bmatrix} = 2$.

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 2 \\ -4 \\ 1 \end{bmatrix} \right\}$ is linearly dependent.

Answer:

1st way: We can see that $\mathbf{w} = 2\mathbf{u} - \mathbf{v}$ or

$$2\mathbf{u} - \mathbf{v} - \mathbf{w} = \mathbf{0}.$$

Hence $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent.

2nd way: The system $t_1\mathbf{u} + t_2\mathbf{v} + t_3\mathbf{w} = \mathbf{0}$

$$t_1 + 2t_3 = 0, \quad -t_1 + 2t_2 - 4t_3 = 0, \quad 2t_1 + 3t_2 + t_3 = 0$$

has nontrivial solutions since $\text{rank} \begin{bmatrix} 1 & 0 & 2 \\ -1 & 2 & -4 \\ 2 & 3 & 1 \end{bmatrix} = 2$. Hence $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is

linearly dependent.

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 2 \\ -4 \\ 1 \end{bmatrix} \right\}$ is linearly dependent.

Answer:

1st way: We can see that $\mathbf{w} = 2\mathbf{u} - \mathbf{v}$ or

$$2\mathbf{u} - \mathbf{v} - \mathbf{w} = \mathbf{0}.$$

Hence $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent.

2nd way: The system $t_1\mathbf{u} + t_2\mathbf{v} + t_3\mathbf{w} = \mathbf{0}$

$$t_1 + 2t_3 = 0, \quad -t_1 + 2t_2 - 4t_3 = 0, \quad 2t_1 + 3t_2 + t_3 = 0$$

has nontrivial solutions since $\text{rank} \begin{bmatrix} 1 & 0 & 2 \\ -1 & 2 & -4 \\ 2 & 3 & 1 \end{bmatrix} = 2$. Hence $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent.

Theorem

- $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is linearly independent if and only if

$$\text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = k .$$

- $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is linearly dependent if and only if

$$\text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m < k .$$

Theorem

- $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is linearly independent if and only if

$$\text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = k .$$

- $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is linearly dependent if and only if

$$\text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m < k .$$

And we only can take at most m linearly independent vectors from k vectors $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$.

Theorem

- $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is linearly independent if and only if

$$\text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = k .$$

- $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is linearly dependent if and only if

$$\text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m < k .$$

And we only can take at most m linearly independent vectors from k vectors $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$.

Example

For what values of a is the set $\{[1, 1, 1], [1, 0, 2], [1, a, 1]\}$ linearly dependent?

Answer:

Example

For what values of a is the set $\{[1, 1, 1], [1, 0, 2], [1, a, 1]\}$ linearly dependent?

Answer: The set $\{[1, 1, 1], [1, 0, 2], [1, a, 1]\}$ is linearly dependent if and only if

$$\begin{aligned} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & a \\ 1 & 2 & 1 \end{bmatrix} &\stackrel{-R_1+R_2}{=} \stackrel{-R_1+R_3}{=} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 0 & -1 & a-1 \\ 0 & 1 & 0 \end{bmatrix} \\ &\stackrel{R_2 \leftrightarrow R_3}{=} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & -1 & a-1 \end{bmatrix} \stackrel{R_2+R_3}{=} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & a-1 \end{bmatrix} < 3. \end{aligned}$$

Last inequality holds if $a - 1 = 0$ or $a = 1$.

Example

For what values of a is the set $\{[1, 1, 1], [1, 0, 2], [1, a, 1]\}$ linearly dependent?

Answer: The set $\{[1, 1, 1], [1, 0, 2], [1, a, 1]\}$ is linearly dependent if and only if

$$\begin{aligned} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & a \\ 1 & 2 & 1 \end{bmatrix} &\stackrel{-R_1+R_2}{\stackrel{-R_1+R_3}{=}} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 0 & -1 & a-1 \\ 0 & 1 & 0 \end{bmatrix} \\ &\stackrel{R_2 \leftrightarrow R_3}{=} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & -1 & a-1 \end{bmatrix} \stackrel{R_2+R_3}{=} \text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & a-1 \end{bmatrix} < 3. \end{aligned}$$

Last inequality holds if $a - 1 = 0$ or $a = 1$.

Dimension

Definition

If U is a subspace of \mathbb{R}^n , a set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ of vectors in U is called a **basis** of U if it satisfies the following two conditions:

1. $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is linearly independent.

Dimension

Definition

If U is a subspace of \mathbb{R}^n , a set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ of vectors in U is called a **basis** of U if it satisfies the following two conditions:

1. $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is linearly independent.
2. $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$.

Dimension

Definition

If U is a subspace of \mathbb{R}^n , a set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ of vectors in U is called a **basis** of U if it satisfies the following two conditions:

1. $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is linearly independent.
2. $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$.

Theorem

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ and $\{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k\}$ are basis of U then $m = k$.

Dimension

Definition

If U is a subspace of \mathbb{R}^n , a set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ of vectors in U is called a **basis** of U if it satisfies the following two conditions:

1. $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is linearly independent.
2. $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$.

Theorem

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ and $\{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k\}$ are basis of U then $m = k$.

Definition

If U is a subspace of \mathbb{R}^n and $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is a basis of U , then the number m is called the **dimension** of U , denoted $\dim(U) = m$.

Dimension

Definition

If U is a subspace of \mathbb{R}^n , a set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ of vectors in U is called a **basis** of U if it satisfies the following two conditions:

1. $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is linearly independent.
2. $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$.

Theorem

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ and $\{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k\}$ are basis of U then $m = k$.

Definition

If U is a subspace of \mathbb{R}^n and $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is a basis of U , then the number m is called the **dimension** of U , denoted $\dim(U) = m$.

Dimension

Definition

If U is a subspace of \mathbb{R}^n , a set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ of vectors in U is called a **basis** of U if it satisfies the following two conditions:

1. $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is linearly independent.
2. $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$.

Theorem

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ and $\{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k\}$ are basis of U then $m = k$.

Definition

If U is a subspace of \mathbb{R}^n and $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ is a basis of U , then the number m is called the **dimension** of U , denoted $\dim(U) = m$.

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -1 \\ 3 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^2 .

Answer:

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -1 \\ 3 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^2 .

Answer: Since $\text{rank} \begin{bmatrix} \mathbf{u} & \mathbf{v} \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = 2$, the set $\{\mathbf{u}, \mathbf{v}\}$ is linearly independent.

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -1 \\ 3 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^2 .

Answer: Since $\text{rank} \begin{bmatrix} \mathbf{u} & \mathbf{v} \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = 2$, the set $\{\mathbf{u}, \mathbf{v}\}$ is linearly independent.

Any vector $\begin{bmatrix} x & y \end{bmatrix}^T$ is a linear combination of \mathbf{u}, \mathbf{v} since

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -1 \\ 3 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^2 .

Answer: Since $\text{rank} \begin{bmatrix} \mathbf{u} & \mathbf{v} \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = 2$, the set $\{\mathbf{u}, \mathbf{v}\}$ is linearly independent.

Any vector $\begin{bmatrix} x & y \end{bmatrix}^T$ is a linear combination of \mathbf{u}, \mathbf{v} since

$$\begin{bmatrix} x & y \end{bmatrix}^T = \frac{3x + y}{5} \mathbf{u} + \frac{y - 2x}{5} \mathbf{v}.$$

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -1 \\ 3 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^2 .

Answer: Since $\text{rank} \begin{bmatrix} \mathbf{u} & \mathbf{v} \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = 2$, the set $\{\mathbf{u}, \mathbf{v}\}$ is linearly independent.

Any vector $\begin{bmatrix} x & y \end{bmatrix}^T$ is a linear combination of \mathbf{u}, \mathbf{v} since

$$\begin{bmatrix} x & y \end{bmatrix}^T = \frac{3x + y}{5} \mathbf{u} + \frac{y - 2x}{5} \mathbf{v}.$$

Thus, $\mathbb{R}^2 = \text{span}\{\mathbf{u}, \mathbf{v}\}$.

We have showed that $\{\mathbf{u}, \mathbf{v}\}$ is a basis of \mathbb{R}^2 .

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -1 \\ 3 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^2 .

Answer: Since $\text{rank} \begin{bmatrix} \mathbf{u} & \mathbf{v} \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = 2$, the set $\{\mathbf{u}, \mathbf{v}\}$ is linearly independent.

Any vector $\begin{bmatrix} x & y \end{bmatrix}^T$ is a linear combination of \mathbf{u}, \mathbf{v} since

$$\begin{bmatrix} x & y \end{bmatrix}^T = \frac{3x + y}{5} \mathbf{u} + \frac{y - 2x}{5} \mathbf{v}.$$

Thus, $\mathbb{R}^2 = \text{span}\{\mathbf{u}, \mathbf{v}\}$.

We have showed that $\{\mathbf{u}, \mathbf{v}\}$ is a basis of \mathbb{R}^2 .

Example

We can see that $\dim(\mathbb{R}^n) = n$ and $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is a basis of \mathbb{R}^n .

Example

Show that $\left\{ \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -1 \\ 3 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^2 .

Answer: Since $\text{rank} \begin{bmatrix} \mathbf{u} & \mathbf{v} \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = 2$, the set $\{\mathbf{u}, \mathbf{v}\}$ is linearly independent.

Any vector $\begin{bmatrix} x & y \end{bmatrix}^T$ is a linear combination of \mathbf{u}, \mathbf{v} since

$$\begin{bmatrix} x & y \end{bmatrix}^T = \frac{3x + y}{5} \mathbf{u} + \frac{y - 2x}{5} \mathbf{v}.$$

Thus, $\mathbb{R}^2 = \text{span}\{\mathbf{u}, \mathbf{v}\}$.

We have showed that $\{\mathbf{u}, \mathbf{v}\}$ is a basis of \mathbb{R}^2 .

Example

We can see that $\dim(\mathbb{R}^n) = n$ and $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is a basis of \mathbb{R}^n .

Theorem

Suppose that $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$. Then

$$\dim(U) = \text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m \leq k$$

And we can choose m vectors among k vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ to get a basis of U .

Example

Find the dimension of the subspace $U = \text{span}\{[1, 2], [2, -1], [0, 3], [1, 4]\}$.

Theorem

Suppose that $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$. Then

$$\dim(U) = \text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m \leq k$$

And we can choose m vectors among k vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ to get a basis of U .

Example

Find the dimension of the subspace $U = \text{span}\{[1, 2], [2, -1], [0, 3], [1, 4]\}$.

Answer:

Theorem

Suppose that $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$. Then

$$\dim(U) = \text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m \leq k$$

And we can choose m vectors among k vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ to get a basis of U .

Example

Find the dimension of the subspace $U = \text{span}\{[1, 2], [2, -1], [0, 3], [1, 4]\}$.

Answer: We have

$$\text{rank} \begin{bmatrix} 1 & 2 & 0 & 1 \\ 2 & -1 & 3 & 4 \end{bmatrix} \stackrel{-2R_1+R_2}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & -5 & 3 & 2 \end{bmatrix} = 2.$$

Theorem

Suppose that $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$. Then

$$\dim(U) = \text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m \leq k$$

And we can choose m vectors among k vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ to get a basis of U .

Example

Find the dimension of the subspace $U = \text{span}\{[1, 2], [2, -1], [0, 3], [1, 4]\}$.

Answer: We have

$$\text{rank} \begin{bmatrix} 1 & 2 & 0 & 1 \\ 2 & -1 & 3 & 4 \end{bmatrix} \stackrel{-2R_1+R_2}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & -5 & 3 & 2 \end{bmatrix} = 2.$$

Hence $\dim(U) = 2$. We can choose basis $\{[1, 2], [2, -1]\}$ for U . In fact, $U = \mathbb{R}^2$.

Theorem

Suppose that $U = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$. Then

$$\dim(U) = \text{rank} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} = m \leq k$$

And we can choose m vectors among k vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ to get a basis of U .

Example

Find the dimension of the subspace $U = \text{span}\{[1, 2], [2, -1], [0, 3], [1, 4]\}$.

Answer: We have

$$\text{rank} \begin{bmatrix} 1 & 2 & 0 & 1 \\ 2 & -1 & 3 & 4 \end{bmatrix} \stackrel{-2R_1+R_2}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & -5 & 3 & 2 \end{bmatrix} = 2.$$

Hence $\dim(U) = 2$. We can choose basis $\{[1, 2], [2, -1]\}$ for U . In fact, $U = \mathbb{R}^2$.

Example

Find the dimension of the subspace

$$U = \{[a + c, b + c, a + 2c + b] \mid a, b, c \text{ in } \mathbb{R}\}.$$

Answer:

Example

Find the dimension of the subspace

$$U = \{[a + c, b + c, a + 2c + b] \mid a, b, c \text{ in } \mathbb{R}\}.$$

Answer: We can see that

$$\begin{bmatrix} a + c \\ b + c \\ a + 2c + b \end{bmatrix} = a \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + b \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

Example

Find the dimension of the subspace

$$U = \{[a + c, b + c, a + 2c + b] \mid a, b, c \text{ in } \mathbb{R}\}.$$

Answer: We can see that

$$\begin{bmatrix} a + c \\ b + c \\ a + 2c + b \end{bmatrix} = a \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + b \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

$$\text{Hence } U = \text{span} \left\{ \begin{bmatrix} 1 & 0 & 1 \end{bmatrix}^T, \begin{bmatrix} 0 & 1 & 1 \end{bmatrix}^T, \begin{bmatrix} 1 & 1 & 2 \end{bmatrix}^T \right\}.$$

Example

Find the dimension of the subspace

$$U = \{[a + c, b + c, a + 2c + b] | a, b, c \text{ in } \mathbb{R}\}.$$

Answer: We can see that

$$\begin{bmatrix} a + c \\ b + c \\ a + 2c + b \end{bmatrix} = a \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + b \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

$$\text{Hence } U = \text{span} \left\{ \begin{bmatrix} 1 & 0 & 1 \end{bmatrix}^T, \begin{bmatrix} 0 & 1 & 1 \end{bmatrix}^T, \begin{bmatrix} 1 & 1 & 2 \end{bmatrix}^T \right\}.$$

Moreover,

$$\text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix} \stackrel{-R_1+R_3}{=} \text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \stackrel{-R_2+R_3}{=} \text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} = 2.$$

Example

Find the dimension of the subspace

$$U = \{[a + c, b + c, a + 2c + b] | a, b, c \text{ in } \mathbb{R}\}.$$

Answer: We can see that

$$\begin{bmatrix} a + c \\ b + c \\ a + 2c + b \end{bmatrix} = a \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + b \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

Hence $U = \text{span} \left\{ \begin{bmatrix} 1 & 0 & 1 \end{bmatrix}^T, \begin{bmatrix} 0 & 1 & 1 \end{bmatrix}^T, \begin{bmatrix} 1 & 1 & 2 \end{bmatrix}^T \right\}.$

Moreover,

$$\text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix} \stackrel{-R_1+R_3}{=} \text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \stackrel{-R_2+R_3}{=} \text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} = 2.$$

Hence $\dim(U) = 2$

Example

Find the dimension of the subspace

$$U = \{[a + c, b + c, a + 2c + b] | a, b, c \text{ in } \mathbb{R}\}.$$

Answer: We can see that

$$\begin{bmatrix} a + c \\ b + c \\ a + 2c + b \end{bmatrix} = a \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + b \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

Hence $U = \text{span} \left\{ \begin{bmatrix} 1 & 0 & 1 \end{bmatrix}^T, \begin{bmatrix} 0 & 1 & 1 \end{bmatrix}^T, \begin{bmatrix} 1 & 1 & 2 \end{bmatrix}^T \right\}.$

Moreover,

$$\text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix} \stackrel{-R_1+R_3}{=} \text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \stackrel{-R_2+R_3}{=} \text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} = 2.$$

Hence $\dim(U) = 2$

Example

Find the dimension of the subspace

$$V = \{[a + 2b + 2d, c + d, 2a + 4b + 2d, -c + d] \mid a, b, c, d \text{ in } \mathbb{R}\}.$$

Answer:

Example

Find the dimension of the subspace

$$V = \{[a + 2b + 2d, c + d, 2a + 4b + 2d, -c + d] \mid a, b, c, d \text{ in } \mathbb{R}\}.$$

Answer: We can see that V is

$$\text{span} \left\{ \begin{bmatrix} 1 & 0 & 2 & 0 \end{bmatrix}^T, \begin{bmatrix} 2 & 0 & 4 & 0 \end{bmatrix}^T, \begin{bmatrix} 0 & 1 & 0 & -1 \end{bmatrix}^T, \begin{bmatrix} 2 & 1 & 2 & 1 \end{bmatrix}^T \right\}.$$

Example

Find the dimension of the subspace

$$V = \{[a + 2b + 2d, c + d, 2a + 4b + 2d, -c + d] \mid a, b, c, d \text{ in } \mathbb{R}\}.$$

Answer: We can see that V is

$$\text{span} \left\{ \begin{bmatrix} 1 & 0 & 2 & 0 \end{bmatrix}^T, \begin{bmatrix} 2 & 0 & 4 & 0 \end{bmatrix}^T, \begin{bmatrix} 0 & 1 & 0 & -1 \end{bmatrix}^T, \begin{bmatrix} 2 & 1 & 2 & 1 \end{bmatrix}^T \right\}.$$

Moreover,

$$\begin{aligned} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 2 & 4 & 0 & 2 \\ 0 & 0 & -1 & 1 \end{bmatrix} &\stackrel{-2R_1+R_3}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix} \\ &\stackrel{R_2+R_4}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \stackrel{R_3 \leftrightarrow R_4}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} = 3. \Rightarrow \dim(V) = 3. \end{aligned}$$

Example

Find the dimension of the subspace

$$V = \{[a + 2b + 2d, c + d, 2a + 4b + 2d, -c + d] \mid a, b, c, d \text{ in } \mathbb{R}\}.$$

Answer: We can see that V is

$$\text{span} \left\{ \begin{bmatrix} 1 & 0 & 2 & 0 \end{bmatrix}^T, \begin{bmatrix} 2 & 0 & 4 & 0 \end{bmatrix}^T, \begin{bmatrix} 0 & 1 & 0 & -1 \end{bmatrix}^T, \begin{bmatrix} 2 & 1 & 2 & 1 \end{bmatrix}^T \right\}.$$

Moreover,

$$\begin{aligned} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 2 & 4 & 0 & 2 \\ 0 & 0 & -1 & 1 \end{bmatrix} &\stackrel{-2R_1+R_3}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix} \\ &\stackrel{R_2+R_4}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \stackrel{R_3 \leftrightarrow R_4}{=} \text{rank} \begin{bmatrix} 1 & 2 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} = 3. \Rightarrow \dim(V) = 3. \end{aligned}$$

Theorem

Any set consist of n linearly independent vectors in \mathbb{R}^n is a basis of \mathbb{R}^n .

Example

Which of the following sets are bases of \mathbb{R}^3 ?

- (i) $\{(1, 2, 1), (-1, 0, 3)\}$
- (ii) $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$

Theorem

Any set consist of n linearly independent vectors in \mathbb{R}^n is a basis of \mathbb{R}^n .

Example

Which of the following sets are bases of \mathbb{R}^3 ?

(i) $\{(1, 2, 1), (-1, 0, 3)\}$

(ii) $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$

Answer:

Theorem

Any set consist of n linearly independent vectors in \mathbb{R}^n is a basis of \mathbb{R}^n .

Example

Which of the following sets are bases of \mathbb{R}^3 ?

- (i) $\{(1, 2, 1), (-1, 0, 3)\}$
- (ii) $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$

Answer:

- (i) No, since any basis of \mathbb{R}^3 must have exactly three vectors.

Theorem

Any set consist of n linearly independent vectors in \mathbb{R}^n is a basis of \mathbb{R}^n .

Example

Which of the following sets are bases of \mathbb{R}^3 ?

- (i) $\{(1, 2, 1), (-1, 0, 3)\}$
- (ii) $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$

Answer:

- (i) No, since any basis of \mathbb{R}^3 must have exactly three vectors.
- (ii) Yes. We have

Theorem

Any set consist of n linearly independent vectors in \mathbb{R}^n is a basis of \mathbb{R}^n .

Example

Which of the following sets are bases of \mathbb{R}^3 ?

- (i) $\{(1, 2, 1), (-1, 0, 3)\}$
- (ii) $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$

Answer:

- (i) No, since any basis of \mathbb{R}^3 must have exactly three vectors.
- (ii) Yes. We have

$$A = \begin{bmatrix} -1 & 1 & 0 \\ 4 & 0 & 1 \\ 3 & 2 & -5 \end{bmatrix} \xrightarrow[4R_1+R_2]{3R_1+R_3} \begin{bmatrix} -1 & 1 & 0 \\ 0 & 4 & 1 \\ 0 & 5 & -5 \end{bmatrix} \xrightarrow[1/4R_2]{-5/4R_2+R_3} \begin{bmatrix} -1 & 1 & 0 \\ 0 & 1 & 1/4 \\ 0 & 0 & -25/4 \end{bmatrix}$$

See that $\text{rank}(A) = 3$.

Theorem

Any set consist of n linearly independent vectors in \mathbb{R}^n is a basis of \mathbb{R}^n .

Example

Which of the following sets are bases of \mathbb{R}^3 ?

- (i) $\{(1, 2, 1), (-1, 0, 3)\}$
- (ii) $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$

Answer:

- (i) No, since any basis of \mathbb{R}^3 must have exactly three vectors.
- (ii) Yes. We have

$$A = \begin{bmatrix} -1 & 1 & 0 \\ 4 & 0 & 1 \\ 3 & 2 & -5 \end{bmatrix} \xrightarrow[4R_1+R_2]{3R_1+R_3} \begin{bmatrix} -1 & 1 & 0 \\ 0 & 4 & 1 \\ 0 & 5 & -5 \end{bmatrix} \xrightarrow[1/4R_2]{-5/4R_2+R_3} \begin{bmatrix} -1 & 1 & 0 \\ 0 & 1 & 1/4 \\ 0 & 0 & -25/4 \end{bmatrix}$$

See that $\text{rank}(A) = 3$. Hence $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$ is linearly independent.

Theorem

Any set consist of n linearly independent vectors in \mathbb{R}^n is a basis of \mathbb{R}^n .

Example

Which of the following sets are bases of \mathbb{R}^3 ?

- (i) $\{(1, 2, 1), (-1, 0, 3)\}$
- (ii) $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$

Answer:

- (i) No, since any basis of \mathbb{R}^3 must have exactly three vectors.
- (ii) Yes. We have

$$A = \begin{bmatrix} -1 & 1 & 0 \\ 4 & 0 & 1 \\ 3 & 2 & -5 \end{bmatrix} \xrightarrow[4R_1+R_2]{3R_1+R_3} \begin{bmatrix} -1 & 1 & 0 \\ 0 & 4 & 1 \\ 0 & 5 & -5 \end{bmatrix} \xrightarrow[1/4R_2]{-5/4R_2+R_3} \begin{bmatrix} -1 & 1 & 0 \\ 0 & 1 & 1/4 \\ 0 & 0 & -25/4 \end{bmatrix}$$

See that $\text{rank}(A) = 3$. Hence $\{(-1, 4, 3), (1, 0, 2), (0, 1, -5)\}$ is linearly independent.

5.3 Orthogonality

Given $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ are two n -tuples (vectors) in \mathbb{R}^n .

Definition

- The **dot product** of \mathbf{x} and \mathbf{y} is defined as follows:

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n .$$

5.3 Orthogonality

Given $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ are two n -tuples (vectors) in \mathbb{R}^n .

Definition

- The **dot product** of \mathbf{x} and \mathbf{y} is defined as follows:

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n .$$

- The **length** of \mathbf{x} is

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} .$$

5.3 Orthogonality

Given $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ are two n -tuples (vectors) in \mathbb{R}^n .

Definition

- The **dot product** of \mathbf{x} and \mathbf{y} is defined as follows:

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n .$$

- The **length** of \mathbf{x} is

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} .$$

- A vector of length 1 is called a **unit vector**.

5.3 Orthogonality

Given $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ are two n -tuples (vectors) in \mathbb{R}^n .

Definition

- The **dot product** of \mathbf{x} and \mathbf{y} is defined as follows:

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n .$$

- The **length** of \mathbf{x} is

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} .$$

- A vector of length 1 is called a **unit vector**.

5.3 Orthogonality

Given $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ are two n -tuples (vectors) in \mathbb{R}^n .

Definition

- The **dot product** of \mathbf{x} and \mathbf{y} is defined as follows:

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n .$$

- The **length** of \mathbf{x} is

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} .$$

- A vector of length 1 is called a **unit vector**.

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$
3. $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$
3. $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$
4. $(a\mathbf{u}) \cdot \mathbf{v} = a(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (a\mathbf{v})$ for any scalar a .

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$
3. $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$
4. $(a\mathbf{u}) \cdot \mathbf{v} = a(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (a\mathbf{v})$ for any scalar a .
5. $\mathbf{u} \cdot (\mathbf{v} \pm \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} \pm \mathbf{u} \cdot \mathbf{w}$.

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$
3. $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$
4. $(a\mathbf{u}) \cdot \mathbf{v} = a(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (a\mathbf{v})$ for any scalar a .
5. $\mathbf{u} \cdot (\mathbf{v} \pm \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} \pm \mathbf{u} \cdot \mathbf{w}$.
6. $|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|$ (Cauchy inequality)

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$
3. $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$
4. $(a\mathbf{u}) \cdot \mathbf{v} = a(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (a\mathbf{v})$ for any scalar a .
5. $\mathbf{u} \cdot (\mathbf{v} \pm \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} \pm \mathbf{u} \cdot \mathbf{w}$.
6. $|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|$ (Cauchy inequality)
7. $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$ (Triangle inequality)

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$
3. $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$
4. $(a\mathbf{u}) \cdot \mathbf{v} = a(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (a\mathbf{v})$ for any scalar a .
5. $\mathbf{u} \cdot (\mathbf{v} \pm \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} \pm \mathbf{u} \cdot \mathbf{w}$.
6. $|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|$ (Cauchy inequality)
7. $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$ (Triangle inequality)

Example

If $\mathbf{u} = (1, 1, -1, 2)$ and $\mathbf{v} = (2, 0, 3, 1)$ in \mathbb{R}^4 then

$$\mathbf{u} \cdot \mathbf{v} = 1 \quad \|\mathbf{u}\| = \sqrt{1^2 + 1^2 + (-1)^2 + 2^2} = \sqrt{7}$$

Theorem

Let \mathbf{u}, \mathbf{v} and \mathbf{w} be vectors in \mathbb{R}^n .

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2. $\mathbf{u} \cdot \mathbf{0} = \mathbf{0} \cdot \mathbf{u} = 0$
3. $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$
4. $(a\mathbf{u}) \cdot \mathbf{v} = a(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (a\mathbf{v})$ for any scalar a .
5. $\mathbf{u} \cdot (\mathbf{v} \pm \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} \pm \mathbf{u} \cdot \mathbf{w}$.
6. $|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|$ (Cauchy inequality)
7. $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$ (Triangle inequality)

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

$$\mathbf{x}_i \cdot \mathbf{x}_j = 0 \text{ for all } i \neq j \text{ and } \mathbf{x}_i \neq \mathbf{0} \text{ for all } i.$$

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

$$\mathbf{x}_i \cdot \mathbf{x}_j = 0 \text{ for all } i \neq j \text{ and } \mathbf{x}_i \neq \mathbf{0} \text{ for all } i .$$

- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthonormal** if it is orthogonal and each \mathbf{x}_i is a unit vector $\|\mathbf{x}_i\| = 1$.

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

$$\mathbf{x}_i \cdot \mathbf{x}_j = 0 \text{ for all } i \neq j \text{ and } \mathbf{x}_i \neq \mathbf{0} \text{ for all } i.$$

- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthonormal** if it is orthogonal and each \mathbf{x}_i is a unit vector $\|\mathbf{x}_i\| = 1$.

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

$$\mathbf{x}_i \cdot \mathbf{x}_j = 0 \text{ for all } i \neq j \text{ and } \mathbf{x}_i \neq \mathbf{0} \text{ for all } i .$$

- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthonormal** if it is orthogonal and each \mathbf{x}_i is a unit vector $\|\mathbf{x}_i\| = 1$.

Example

The standard basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is an orthonormal set in \mathbb{R}^n .

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

$$\mathbf{x}_i \cdot \mathbf{x}_j = 0 \text{ for all } i \neq j \text{ and } \mathbf{x}_i \neq \mathbf{0} \text{ for all } i.$$

- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthonormal** if it is orthogonal and each \mathbf{x}_i is a unit vector $\|\mathbf{x}_i\| = 1$.

Example

The standard basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is an orthonormal set in \mathbb{R}^n .

Example

The set $\left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$ is orthogonal in \mathbb{R}^2 ,

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

$$\mathbf{x}_i \cdot \mathbf{x}_j = 0 \text{ for all } i \neq j \text{ and } \mathbf{x}_i \neq \mathbf{0} \text{ for all } i.$$

- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthonormal** if it is orthogonal and each \mathbf{x}_i is a unit vector $\|\mathbf{x}_i\| = 1$.

Example

The standard basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is an orthonormal set in \mathbb{R}^n .

Example

The set $\left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$ is orthogonal in \mathbb{R}^2 , but not orthonormal.

Orthogonal Sets

Definition

- Two vectors \mathbf{x} and \mathbf{y} in \mathbb{R}^n are **orthogonal** if $\mathbf{x} \cdot \mathbf{y} = 0$.
- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthogonal set** if

$$\mathbf{x}_i \cdot \mathbf{x}_j = 0 \text{ for all } i \neq j \text{ and } \mathbf{x}_i \neq \mathbf{0} \text{ for all } i.$$

- A set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ of vectors in \mathbb{R}^n is called an **orthonormal** if it is orthogonal and each \mathbf{x}_i is a unit vector $\|\mathbf{x}_i\| = 1$.

Example

The standard basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is an orthonormal set in \mathbb{R}^n .

Example

The set $\left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$ is orthogonal in \mathbb{R}^2 , but not orthonormal.

Example

Find the number c such that the set

$$\left\{ \mathbf{u}_1 = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ 3 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 2 \\ -1 \\ 0 \\ -1 \end{bmatrix}, \mathbf{u}_4 = \begin{bmatrix} a \\ b \\ c \\ 1 \end{bmatrix} \right\} \text{ is orthogonal.}$$

Answer:

Example

Find the number c such that the set

$$\left\{ \mathbf{u}_1 = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ 3 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 2 \\ -1 \\ 0 \\ -1 \end{bmatrix}, \mathbf{u}_4 = \begin{bmatrix} a \\ b \\ c \\ 1 \end{bmatrix} \right\} \text{ is orthogonal.}$$

Answer: Solve the system

$$\begin{cases} \mathbf{u}_1 \cdot \mathbf{u}_4 = a + 2b + c = 0 \\ \mathbf{u}_2 \cdot \mathbf{u}_4 = a - b + c + 3 = 0 \\ \mathbf{u}_3 \cdot \mathbf{u}_4 = 2a - b - 1 = 0 \end{cases}$$

Example

Find the number c such that the set

$$\left\{ \mathbf{u}_1 = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ 3 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 2 \\ -1 \\ 0 \\ -1 \end{bmatrix}, \mathbf{u}_4 = \begin{bmatrix} a \\ b \\ c \\ 1 \end{bmatrix} \right\} \text{ is orthogonal.}$$

Answer: Solve the system

$$\begin{cases} \mathbf{u}_1 \cdot \mathbf{u}_4 = a + 2b + c = 0 \\ \mathbf{u}_2 \cdot \mathbf{u}_4 = a - b + c + 3 = 0 \\ \mathbf{u}_3 \cdot \mathbf{u}_4 = 2a - b - 1 = 0 \end{cases}$$

we get $a = 1, b = 1, c = -3$.

Example

Find the number c such that the set

$$\left\{ \mathbf{u}_1 = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ 3 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 2 \\ -1 \\ 0 \\ -1 \end{bmatrix}, \mathbf{u}_4 = \begin{bmatrix} a \\ b \\ c \\ 1 \end{bmatrix} \right\} \text{ is orthogonal.}$$

Answer: Solve the system

$$\begin{cases} \mathbf{u}_1 \cdot \mathbf{u}_4 = a + 2b + c = 0 \\ \mathbf{u}_2 \cdot \mathbf{u}_4 = a - b + c + 3 = 0 \\ \mathbf{u}_3 \cdot \mathbf{u}_4 = 2a - b - 1 = 0 \end{cases}$$

we get $a = 1, b = 1, c = -3$.

Example

Find b so that $B = \left\{ \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right), \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right), (a, b, c) \right\}$ is an orthonormal set.

Answer:

Example

Find b so that $B = \left\{ \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right), \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right), (a, b, c) \right\}$ is an orthonormal set.

Answer: To B is an orthonormal set,

$$\begin{cases} \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right) \cdot (a, b, c) = \frac{1}{\sqrt{3}}(a + b + c) = 0 \\ \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right) \cdot (a, b, c) = \frac{-1}{\sqrt{2}}(a - b) = 0 \\ \|(a, b, c)\| = \sqrt{a^2 + b^2 + c^2} = 1 \end{cases}$$

Example

Find b so that $B = \left\{ \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right), \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right), (a, b, c) \right\}$ is an orthonormal set.

Answer: To B is an orthonormal set,

$$\begin{cases} \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right) \cdot (a, b, c) = \frac{1}{\sqrt{3}}(a + b + c) = 0 \\ \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right) \cdot (a, b, c) = \frac{-1}{\sqrt{2}}(a - b) = 0 \\ \|(a, b, c)\| = \sqrt{a^2 + b^2 + c^2} = 1 \end{cases}$$

Substitute a by b , c by $-2b$ into third equation we get

$$b^2 + b^2 + (-2b)^2 = 6b^2 = 1$$

Example

Find b so that $B = \left\{ \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right), \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right), (a, b, c) \right\}$ is an orthonormal set.

Answer: To B is an orthonormal set,

$$\begin{cases} \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right) \cdot (a, b, c) = \frac{1}{\sqrt{3}}(a + b + c) = 0 \\ \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right) \cdot (a, b, c) = \frac{-1}{\sqrt{2}}(a - b) = 0 \\ \|(a, b, c)\| = \sqrt{a^2 + b^2 + c^2} = 1 \end{cases}$$

Substitute a by b , c by $-2b$ into third equation we get

$$b^2 + b^2 + (-2b)^2 = 6b^2 = 1$$

Hence $b = 1/\sqrt{6}$ or $b = -1/\sqrt{6}$.

Example

Find b so that $B = \left\{ \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right), \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right), (a, b, c) \right\}$ is an orthonormal set.

Answer: To B is an orthonormal set,

$$\begin{cases} \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right) \cdot (a, b, c) = \frac{1}{\sqrt{3}}(a + b + c) = 0 \\ \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right) \cdot (a, b, c) = \frac{-1}{\sqrt{2}}(a - b) = 0 \\ \|(a, b, c)\| = \sqrt{a^2 + b^2 + c^2} = 1 \end{cases}$$

Substitute a by b , c by $-2b$ into third equation we get

$$b^2 + b^2 + (-2b)^2 = 6b^2 = 1$$

Hence $b = 1/\sqrt{6}$ or $b = -1/\sqrt{6}$.

Theorem (Pythagoras' Theorem)

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is an orthogonal set, then

$$\|\mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_k\|^2 = \|\mathbf{x}_1\|^2 + \|\mathbf{x}_2\|^2 + \dots + \|\mathbf{x}_k\|^2 .$$

Theorem

Every orthogonal set in \mathbb{R}^n is linearly independent.

Theorem (Pythagoras' Theorem)

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is an orthogonal set, then

$$\|\mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_k\|^2 = \|\mathbf{x}_1\|^2 + \|\mathbf{x}_2\|^2 + \dots + \|\mathbf{x}_k\|^2 .$$

Theorem

Every orthogonal set in \mathbb{R}^n is linearly independent.

Example

Suppose that $\{\mathbf{u}, \mathbf{v}\}$ is an orthogonal set and their length are 4 and 5 respectively. What is the length of $\mathbf{u} - 3\mathbf{v}$.

Theorem (Pythagoras' Theorem)

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is an orthogonal set, then

$$\|\mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_k\|^2 = \|\mathbf{x}_1\|^2 + \|\mathbf{x}_2\|^2 + \dots + \|\mathbf{x}_k\|^2 .$$

Theorem

Every orthogonal set in \mathbb{R}^n is linearly independent.

Example

Suppose that $\{\mathbf{u}, \mathbf{v}\}$ is an orthogonal set and their length are 4 and 5 respectively. What is the length of $\mathbf{u} - 3\mathbf{v}$.

Answer:

Theorem (Pythagoras' Theorem)

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is an orthogonal set, then

$$\|\mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_k\|^2 = \|\mathbf{x}_1\|^2 + \|\mathbf{x}_2\|^2 + \dots + \|\mathbf{x}_k\|^2 .$$

Theorem

Every orthogonal set in \mathbb{R}^n is linearly independent.

Example

Suppose that $\{\mathbf{u}, \mathbf{v}\}$ is an orthogonal set and their length are 4 and 5 respectively. What is the length of $\mathbf{u} - 3\mathbf{v}$.

Answer: By Pythagoras Theorem:

$$\|\mathbf{u} - 3\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|-3\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + 9\|\mathbf{v}\|^2 = 4^2 + 9 * 5^2 = 241 .$$

Theorem (Pythagoras' Theorem)

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is an orthogonal set, then

$$\|\mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_k\|^2 = \|\mathbf{x}_1\|^2 + \|\mathbf{x}_2\|^2 + \dots + \|\mathbf{x}_k\|^2 .$$

Theorem

Every orthogonal set in \mathbb{R}^n is linearly independent.

Example

Suppose that $\{\mathbf{u}, \mathbf{v}\}$ is an orthogonal set and their length are 4 and 5 respectively. What is the length of $\mathbf{u} - 3\mathbf{v}$.

Answer: By Pythagoras Theorem:

$$\|\mathbf{u} - 3\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|-3\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + 9\|\mathbf{v}\|^2 = 4^2 + 9 * 5^2 = 241 .$$

Hence $\|\mathbf{u} - 3\mathbf{v}\| = \sqrt{241}$.

Theorem (Pythagoras' Theorem)

If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ is an orthogonal set, then

$$\|\mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_k\|^2 = \|\mathbf{x}_1\|^2 + \|\mathbf{x}_2\|^2 + \dots + \|\mathbf{x}_k\|^2 .$$

Theorem

Every orthogonal set in \mathbb{R}^n is linearly independent.

Example

Suppose that $\{\mathbf{u}, \mathbf{v}\}$ is an orthogonal set and their length are 4 and 5 respectively. What is the length of $\mathbf{u} - 3\mathbf{v}$.

Answer: By Pythagoras Theorem:

$$\|\mathbf{u} - 3\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|-3\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + 9\|\mathbf{v}\|^2 = 4^2 + 9 * 5^2 = 241 .$$

Hence $\|\mathbf{u} - 3\mathbf{v}\| = \sqrt{241}$.

Example

Which of the following statements are true?

- (i) if $\{\mathbf{u}, \mathbf{v}\}$ is orthogonal in \mathbb{R}^n then $\{\mathbf{u}, \mathbf{u} + \mathbf{v}\}$ is also orthogonal.
- (ii) if $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is also orthogonal.
- (iii) if $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\{\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}\}$ is orthogonal.

Answer:

Example

Which of the following statements are true?

- (i) if $\{\mathbf{u}, \mathbf{v}\}$ is orthogonal in \mathbb{R}^n then $\{\mathbf{u}, \mathbf{u} + \mathbf{v}\}$ is also orthogonal.
- (ii) if $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is also orthogonal.
- (iii) if $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\{\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}\}$ is orthogonal.

Answer:

- (i) False. Since if $\{\mathbf{u}, \mathbf{v}\}$ then $\mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) = \|\mathbf{u}\|^2 \neq 0$, for any $\mathbf{u} \neq \mathbf{0}$.

Example

Which of the following statements are true?

- (i) if $\{\mathbf{u}, \mathbf{v}\}$ is orthogonal in \mathbb{R}^n then $\{\mathbf{u}, \mathbf{u} + \mathbf{v}\}$ is also orthogonal.
- (ii) if $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is also orthogonal.
- (iii) if $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\{\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}\}$ is orthogonal.

Answer:

- (i) False. Since if $\{\mathbf{u}, \mathbf{v}\}$ then $\mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) = \|\mathbf{u}\|^2 \neq 0$, for any $\mathbf{u} \neq \mathbf{0}$.
- (ii) False. For example $\mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\mathbf{z} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Then $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal

Example

Which of the following statements are true?

- (i) if $\{\mathbf{u}, \mathbf{v}\}$ is orthogonal in \mathbb{R}^n then $\{\mathbf{u}, \mathbf{u} + \mathbf{v}\}$ is also orthogonal.
- (ii) if $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is also orthogonal.
- (iii) if $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\{\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}\}$ is orthogonal.

Answer:

- (i) False. Since if $\{\mathbf{u}, \mathbf{v}\}$ then $\mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) = \|\mathbf{u}\|^2 \neq 0$, for any $\mathbf{u} \neq \mathbf{0}$.
- (ii) False. For example $\mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\mathbf{z} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Then $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal but $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is not orthogonal.

Example

Which of the following statements are true?

- (i) if $\{\mathbf{u}, \mathbf{v}\}$ is orthogonal in \mathbb{R}^n then $\{\mathbf{u}, \mathbf{u} + \mathbf{v}\}$ is also orthogonal.
- (ii) if $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is also orthogonal.
- (iii) if $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\{\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}\}$ is orthogonal.

Answer:

- (i) False. Since if $\{\mathbf{u}, \mathbf{v}\}$ then $\mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) = \|\mathbf{u}\|^2 \neq 0$, for any $\mathbf{u} \neq \mathbf{0}$.
- (ii) False. For example $\mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\mathbf{z} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Then $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal but $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is not orthogonal.
- (iii) True. If $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\mathbf{u} \cdot \mathbf{v} = 0$ and $\|\mathbf{u}\| = \|\mathbf{v}\| = 1$.
Hence

Example

Which of the following statements are true?

- (i) if $\{\mathbf{u}, \mathbf{v}\}$ is orthogonal in \mathbb{R}^n then $\{\mathbf{u}, \mathbf{u} + \mathbf{v}\}$ is also orthogonal.
- (ii) if $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is also orthogonal.
- (iii) if $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\{\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}\}$ is orthogonal.

Answer:

- (i) False. Since if $\{\mathbf{u}, \mathbf{v}\}$ then $\mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) = \|\mathbf{u}\|^2 \neq 0$, for any $\mathbf{u} \neq \mathbf{0}$.
- (ii) False. For example $\mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\mathbf{z} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Then $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal but $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is not orthogonal.
- (iii) True. If $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\mathbf{u} \cdot \mathbf{v} = 0$ and $\|\mathbf{u}\| = \|\mathbf{v}\| = 1$. Hence

$$(\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot \mathbf{u} - \mathbf{u} \cdot \mathbf{v} + \mathbf{v} \cdot \mathbf{u} - \mathbf{v} \cdot \mathbf{v} = \|\mathbf{u}\|^2 - \|\mathbf{v}\|^2 = 0.$$

Example

Which of the following statements are true?

- (i) if $\{\mathbf{u}, \mathbf{v}\}$ is orthogonal in \mathbb{R}^n then $\{\mathbf{u}, \mathbf{u} + \mathbf{v}\}$ is also orthogonal.
- (ii) if $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is also orthogonal.
- (iii) if $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\{\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}\}$ is orthogonal.

Answer:

- (i) False. Since if $\{\mathbf{u}, \mathbf{v}\}$ then $\mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) = \|\mathbf{u}\|^2 \neq 0$, for any $\mathbf{u} \neq \mathbf{0}$.
- (ii) False. For example $\mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\mathbf{z} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Then $\{\mathbf{u}, \mathbf{v}\}$ and $\{\mathbf{w}, \mathbf{z}\}$ are both orthogonal but $\{\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{z}\}$ is not orthogonal.
- (iii) True. If $\{\mathbf{u}, \mathbf{v}\}$ is orthonormal then $\mathbf{u} \cdot \mathbf{v} = 0$ and $\|\mathbf{u}\| = \|\mathbf{v}\| = 1$. Hence

$$(\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot \mathbf{u} - \mathbf{u} \cdot \mathbf{v} + \mathbf{v} \cdot \mathbf{u} - \mathbf{v} \cdot \mathbf{v} = \|\mathbf{u}\|^2 - \|\mathbf{v}\|^2 = 0.$$

Theorem

Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ be an orthogonal basis of U . Then any $\mathbf{u} \in U$:

$$\mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{x}_1}{\|\mathbf{x}_1\|^2} \mathbf{x}_1 + \frac{\mathbf{u} \cdot \mathbf{x}_2}{\|\mathbf{x}_2\|^2} \mathbf{x}_2 + \cdots + \frac{\mathbf{u} \cdot \mathbf{x}_m}{\|\mathbf{x}_m\|^2} \mathbf{x}_m$$

Theorem

Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ be an orthogonal basis of U . Then any $\mathbf{u} \in U$:

$$\mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{x}_1}{\|\mathbf{x}_1\|^2} \mathbf{x}_1 + \frac{\mathbf{u} \cdot \mathbf{x}_2}{\|\mathbf{x}_2\|^2} \mathbf{x}_2 + \cdots + \frac{\mathbf{u} \cdot \mathbf{x}_m}{\|\mathbf{x}_m\|^2} \mathbf{x}_m$$

Example

Let $\mathbf{x} = [a, b, c]$. Let U be the subspace spanned by the orthogonal basis $\{\mathbf{u} = [1, 1, 1], \mathbf{v} = [1, -1, 0], \mathbf{w} = [1, 1, -2]\}$.

Find the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$.

Theorem

Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ be an orthogonal basis of U . Then any $\mathbf{u} \in U$:

$$\mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{x}_1}{\|\mathbf{x}_1\|^2} \mathbf{x}_1 + \frac{\mathbf{u} \cdot \mathbf{x}_2}{\|\mathbf{x}_2\|^2} \mathbf{x}_2 + \cdots + \frac{\mathbf{u} \cdot \mathbf{x}_m}{\|\mathbf{x}_m\|^2} \mathbf{x}_m$$

Example

Let $\mathbf{x} = [a, b, c]$. Let U be the subspace spanned by the orthogonal basis $\{\mathbf{u} = [1, 1, 1], \mathbf{v} = [1, -1, 0], \mathbf{w} = [1, 1, -2]\}$.

Find the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$.

Answer:

Theorem

Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ be an orthogonal basis of U . Then any $\mathbf{u} \in U$:

$$\mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{x}_1}{\|\mathbf{x}_1\|^2} \mathbf{x}_1 + \frac{\mathbf{u} \cdot \mathbf{x}_2}{\|\mathbf{x}_2\|^2} \mathbf{x}_2 + \cdots + \frac{\mathbf{u} \cdot \mathbf{x}_m}{\|\mathbf{x}_m\|^2} \mathbf{x}_m$$

Example

Let $\mathbf{x} = [a, b, c]$. Let U be the subspace spanned by the orthogonal basis $\{\mathbf{u} = [1, 1, 1], \mathbf{v} = [1, -1, 0], \mathbf{w} = [1, 1, -2]\}$.

Find the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$.

Answer: We can see $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is an orthogonal basis of U . Hence the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is

$$\frac{\mathbf{x} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} = \frac{a - b}{2}$$

Theorem

Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ be an orthogonal basis of U . Then any $\mathbf{u} \in U$:

$$\mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{x}_1}{\|\mathbf{x}_1\|^2} \mathbf{x}_1 + \frac{\mathbf{u} \cdot \mathbf{x}_2}{\|\mathbf{x}_2\|^2} \mathbf{x}_2 + \cdots + \frac{\mathbf{u} \cdot \mathbf{x}_m}{\|\mathbf{x}_m\|^2} \mathbf{x}_m$$

Example

Let $\mathbf{x} = [a, b, c]$. Let U be the subspace spanned by the orthogonal basis $\{\mathbf{u} = [1, 1, 1], \mathbf{v} = [1, -1, 0], \mathbf{w} = [1, 1, -2]\}$.

Find the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$.

Answer: We can see $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is an orthogonal basis of U . Hence the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is

$$\frac{\mathbf{x} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} = \frac{a - b}{2}$$

(The question is: Write $\mathbf{x} = t_1\mathbf{u} + t_2\mathbf{v} + t_3\mathbf{w}$, find t_2 .)

Theorem

Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m\}$ be an orthogonal basis of U . Then any $\mathbf{u} \in U$:

$$\mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{x}_1}{\|\mathbf{x}_1\|^2} \mathbf{x}_1 + \frac{\mathbf{u} \cdot \mathbf{x}_2}{\|\mathbf{x}_2\|^2} \mathbf{x}_2 + \cdots + \frac{\mathbf{u} \cdot \mathbf{x}_m}{\|\mathbf{x}_m\|^2} \mathbf{x}_m$$

Example

Let $\mathbf{x} = [a, b, c]$. Let U be the subspace spanned by the orthogonal basis $\{\mathbf{u} = [1, 1, 1], \mathbf{v} = [1, -1, 0], \mathbf{w} = [1, 1, -2]\}$.

Find the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$.

Answer: We can see $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is an orthogonal basis of U . Hence the coefficient of \mathbf{v} when expressing \mathbf{x} as a linear combination of $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is

$$\frac{\mathbf{x} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} = \frac{a - b}{2}$$

(The question is: Write $\mathbf{x} = t_1\mathbf{u} + t_2\mathbf{v} + t_3\mathbf{w}$, find t_2 .)

Example

Express $\mathbf{X} = \begin{bmatrix} -3 & 2 & 2 & 7 \end{bmatrix}^T$ as a linear combination of the orthogonal basis of the subspace

$$U = \text{span} \left\{ \mathbf{u} = \begin{bmatrix} -1 & 2 & 0 & 3 \end{bmatrix}^T, \mathbf{v} = \begin{bmatrix} 1 & 2 & -2 & -1 \end{bmatrix}^T \right\}.$$

Answer:

Example

Express $\mathbf{X} = \begin{bmatrix} -3 & 2 & 2 & 7 \end{bmatrix}^T$ as a linear combination of the orthogonal basis of the subspace

$$U = \text{span} \left\{ \mathbf{u} = \begin{bmatrix} -1 & 2 & 0 & 3 \end{bmatrix}^T, \mathbf{v} = \begin{bmatrix} 1 & 2 & -2 & -1 \end{bmatrix}^T \right\}.$$

Answer: We have

$$\begin{aligned} \mathbf{X} &= \frac{\mathbf{X} \cdot \mathbf{u}}{\|\mathbf{u}\|^2} \mathbf{u} + \frac{\mathbf{X} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \mathbf{v} \\ &= \frac{28}{14} \mathbf{u} + \frac{-10}{10} \mathbf{v} \\ &= 2\mathbf{u} - \mathbf{v} \end{aligned}$$

Example

Express $\mathbf{X} = [-3 \ 2 \ 2 \ 7]^T$ as a linear combination of the orthogonal basis of the subspace

$$U = \text{span} \left\{ \mathbf{u} = [-1 \ 2 \ 0 \ 3]^T, \mathbf{v} = [1 \ 2 \ -2 \ -1]^T \right\}.$$

Answer: We have

$$\begin{aligned}\mathbf{X} &= \frac{\mathbf{X} \cdot \mathbf{u}}{\|\mathbf{u}\|^2} \mathbf{u} + \frac{\mathbf{X} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \mathbf{v} \\ &= \frac{28}{14} \mathbf{u} + \frac{-10}{10} \mathbf{v} \\ &= 2\mathbf{u} - \mathbf{v}\end{aligned}$$

5.4 Rank of a Matrix

Let A be $m \times n$ matrix.

Definition

- The **column space**, $\text{col}(A)$, of A is the subspace of \mathbb{R}^m spanned by the columns of A .

5.4 Rank of a Matrix

Let A be $m \times n$ matrix.

Definition

- The **column space**, $\text{col}(A)$, of A is the subspace of \mathbb{R}^m spanned by the columns of A .
- The **row space**, $\text{row}(A)$, of A is the subspace of \mathbb{R}^n spanned by the rows of A .

5.4 Rank of a Matrix

Let A be $m \times n$ matrix.

Definition

- The **column space**, $\text{col}(A)$, of A is the subspace of \mathbb{R}^m spanned by the columns of A .
- The **row space**, $\text{row}(A)$, of A is the subspace of \mathbb{R}^n spanned by the rows of A .

5.4 Rank of a Matrix

Let A be $m \times n$ matrix.

Definition

- The **column space**, $\text{col}(A)$, of A is the subspace of \mathbb{R}^m spanned by the columns of A .
- The **row space**, $\text{row}(A)$, of A is the subspace of \mathbb{R}^n spanned by the rows of A .

Example

If $A = \begin{bmatrix} 1 & 2 & 3 \\ -2 & 0 & 5 \end{bmatrix}$ then

$\text{col}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 5 \end{bmatrix} \right\}$ is a subspace of \mathbb{R}^2

$$= \left\{ x \begin{bmatrix} 1 \\ -2 \end{bmatrix} + y \begin{bmatrix} 2 \\ 0 \end{bmatrix} + z \begin{bmatrix} 3 \\ 5 \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\} = \left\{ \begin{bmatrix} x + 2y + 3z \\ -2x + 5z \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\}$$

$\text{row}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 5 \end{bmatrix} \right\}$ is a subspace of \mathbb{R}^3

$$= \left\{ x \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + y \begin{bmatrix} -2 \\ 0 \\ 5 \end{bmatrix} \mid x, y \in \mathbb{R} \right\} = \left\{ \begin{bmatrix} x - 2y \\ 2x \\ 3x + 5y \end{bmatrix} \mid x, y \in \mathbb{R} \right\}$$

Example

If $A = \begin{bmatrix} 1 & 2 & 3 \\ -2 & 0 & 5 \end{bmatrix}$ then

$\text{col}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 5 \end{bmatrix} \right\}$ is a subspace of \mathbb{R}^2

$$= \left\{ x \begin{bmatrix} 1 \\ -2 \end{bmatrix} + y \begin{bmatrix} 2 \\ 0 \end{bmatrix} + z \begin{bmatrix} 3 \\ 5 \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\} = \left\{ \begin{bmatrix} x + 2y + 3z \\ -2x + 5z \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\}$$

$\text{row}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 5 \end{bmatrix} \right\}$ is a subspace of \mathbb{R}^3

$$= \left\{ x \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + y \begin{bmatrix} -2 \\ 0 \\ 5 \end{bmatrix} \mid x, y \in \mathbb{R} \right\} = \left\{ \begin{bmatrix} x - 2y \\ 2x \\ 3x + 5y \end{bmatrix} \mid x, y \in \mathbb{R} \right\}$$

Lemma

Let A and B denote $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Lemma

Let A and B denote $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Lemma

Let A and B denote $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Theorem

If matrix R is a row-echelon form of A , then

1. The nonzero rows of R are a basis of $\text{row}(A)$.

Lemma

Let A and B denote $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Theorem

If matrix R is a row-echelon form of A , then

1. The nonzero rows of R are a basis of $\text{row}(A)$.
2. The columns of A corresponding to the columns of R that contain the leading 1 are a basis of $\text{col}(A)$.

Lemma

Let A and B denote $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Theorem

If matrix R is a row-echelon form of A , then

1. The nonzero rows of R are a basis of $\text{row}(A)$.
2. The columns of A corresponding to the columns of R that contain the leading 1 are a basis of $\text{col}(A)$.

Lemma

Let A and B denote $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Theorem

If matrix R is a row-echelon form of A , then

1. The nonzero rows of R are a basis of $\text{row}(A)$.
2. The columns of A corresponding to the columns of R that contain the leading 1 are a basis of $\text{col}(A)$.

Example

Let $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 3 & 0 & 3 \end{bmatrix}$. Find basis for $\text{row}(A)$ and $\text{col}(A)$.

Answer:

Example

Let $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 3 & 0 & 3 \end{bmatrix}$. Find basis for $\text{row}(A)$ and $\text{col}(A)$.

Answer:

$$A \xrightarrow[\begin{smallmatrix} -2R_1+R_2 \\ -3R_1+R_3 \end{smallmatrix}]{\begin{smallmatrix} -R_2+R_3 \\ 1/3R_2 \end{smallmatrix}} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 3 & 0 \\ 0 & 3 & 0 \end{bmatrix} \xrightarrow[\begin{smallmatrix} -R_2+R_3 \\ 1/3R_2 \end{smallmatrix}]{\begin{smallmatrix} -R_2+R_3 \\ 1/3R_2 \end{smallmatrix}} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 3 & 0 & 3 \end{bmatrix}$. Find basis for $\text{row}(A)$ and $\text{col}(A)$.

Answer:

$$A \xrightarrow[-3R_1+R_3]{-2R_1+R_2} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 3 & 0 \\ 0 & 3 & 0 \end{bmatrix} \xrightarrow[1/3R_2]{-R_2+R_3} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

A basis of $\text{row}(A)$ is $\{(1, -1, 1), (0, 1, 0)\}$.

Example

Let $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 3 & 0 & 3 \end{bmatrix}$. Find basis for $\text{row}(A)$ and $\text{col}(A)$.

Answer:

$$A \xrightarrow[-3R_1+R_3]{-2R_1+R_2} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 3 & 0 \\ 0 & 3 & 0 \end{bmatrix} \xrightarrow[1/3R_2]{-R_2+R_3} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

A basis of $\text{row}(A)$ is $\{(1, -1, 1), (0, 1, 0)\}$.

A basis of $\text{col}(A)$ is $\{(1, 2, 3)^T, (-1, 1, 0)^T\}$.

Example

Let $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 3 & 0 & 3 \end{bmatrix}$. Find basis for $\text{row}(A)$ and $\text{col}(A)$.

Answer:

$$A \xrightarrow[-3R_1+R_3]{-2R_1+R_2} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 3 & 0 \\ 0 & 3 & 0 \end{bmatrix} \xrightarrow[1/3R_2]{-R_2+R_3} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

A basis of $\text{row}(A)$ is $\{(1, -1, 1), (0, 1, 0)\}$.

A basis of $\text{col}(A)$ is $\{(1, 2, 3)^T, (-1, 1, 0)^T\}$.

Theorem

Let A be $m \times n$ matrix. Then

- $\text{rank}(A) \leq \min\{m, n\}$.
- $\text{col}(A) = \text{im}(A)$.

Theorem

Let A be $m \times n$ matrix. Then

- $\text{rank}(A) \leq \min\{m, n\}$.
- $\text{col}(A) = \text{im}(A)$.
- $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \dim(\text{im}(A)) = \text{rank}(A)$.

Theorem

Let A be $m \times n$ matrix. Then

- $\text{rank}(A) \leq \min\{m, n\}$.
- $\text{col}(A) = \text{im}(A)$.
- $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \dim(\text{im}(A)) = \text{rank}(A)$.
- $\dim(\text{null}(A)) = n - \text{rank}(A) = \text{the number of parameters (of the basic solutions) of the homogeneous system } A\mathbf{X} = \mathbf{0}$.

Theorem

Let A be $m \times n$ matrix. Then

- $\text{rank}(A) \leq \min\{m, n\}$.
- $\text{col}(A) = \text{im}(A)$.
- $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \dim(\text{im}(A)) = \text{rank}(A)$.
- $\dim(\text{null}(A)) = n - \text{rank}(A) = \text{the number of parameters (of the basic solutions) of the homogeneous system } A\mathbf{X} = \mathbf{0}$.

Theorem

Let A be $m \times n$ matrix. Then

- $\text{rank}(A) \leq \min\{m, n\}$.
- $\text{col}(A) = \text{im}(A)$.
- $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \dim(\text{im}(A)) = \text{rank}(A)$.
- $\dim(\text{null}(A)) = n - \text{rank}(A) = \text{the number of parameters (of the basic solutions) of the homogeneous system } A\mathbf{X} = \mathbf{0}$.

Example

Find the dimensions of the null space and the column space of the given matrix.

$$A = \begin{bmatrix} -1 & 2 & 1 & 3 \\ 2 & -3 & -2 & 1 \\ 3 & -5 & -3 & -2 \end{bmatrix}$$

Answer:

Example

Find the dimensions of the null space and the column space of the given matrix.

$$A = \begin{bmatrix} -1 & 2 & 1 & 3 \\ 2 & -3 & -2 & 1 \\ 3 & -5 & -3 & -2 \end{bmatrix}$$

Answer: We have

$$A \xrightarrow{\substack{2R_1+R_2 \\ 3R_1+R_3}} \begin{bmatrix} -1 & 2 & 1 & 3 \\ 0 & 1 & 0 & 7 \\ 0 & 1 & 0 & 7 \end{bmatrix} \xrightarrow{\substack{-R_2+R_3 \\ -R_1}} \begin{bmatrix} 1 & -2 & -1 & -3 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Example

Find the dimensions of the null space and the column space of the given matrix.

$$A = \begin{bmatrix} -1 & 2 & 1 & 3 \\ 2 & -3 & -2 & 1 \\ 3 & -5 & -3 & -2 \end{bmatrix}$$

Answer: We have

$$A \xrightarrow{\substack{2R_1+R_2 \\ 3R_1+R_3}} \begin{bmatrix} -1 & 2 & 1 & 3 \\ 0 & 1 & 0 & 7 \\ 0 & 1 & 0 & 7 \end{bmatrix} \xrightarrow{\substack{-R_2+R_3 \\ -R_1}} \begin{bmatrix} 1 & -2 & -1 & -3 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Hence $\text{rank}(A) = 2$, and

$$\dim(\text{null}(A)) = 4 - \text{rank}(A) = 2, \quad \dim(\text{col}(A)) = \text{rank}(A) = 2.$$

Example

Find the dimensions of the null space and the column space of the given matrix.

$$A = \begin{bmatrix} -1 & 2 & 1 & 3 \\ 2 & -3 & -2 & 1 \\ 3 & -5 & -3 & -2 \end{bmatrix}$$

Answer: We have

$$A \xrightarrow{\substack{2R_1+R_2 \\ 3R_1+R_3}} \begin{bmatrix} -1 & 2 & 1 & 3 \\ 0 & 1 & 0 & 7 \\ 0 & 1 & 0 & 7 \end{bmatrix} \xrightarrow{\substack{-R_2+R_3 \\ -R_1}} \begin{bmatrix} 1 & -2 & -1 & -3 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Hence $\text{rank}(A) = 2$, and

$$\dim(\text{null}(A)) = 4 - \text{rank}(A) = 2, \quad \dim(\text{col}(A)) = \text{rank}(A) = 2.$$

Example

Find the dimension of the subspace

$$U = \left\{ [x, y, z, w] \mid x - 2y + 3z + w = 0, 3x - 5y + z + 8w = 0 \right\}$$

Answer:

Example

Find the dimension of the subspace

$$U = \left\{ [x, y, z, w] \mid x - 2y + 3z + w = 0, 3x - 5y + z + 8w = 0 \right\}$$

Answer: See that $U = \text{null} \left(\begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} \right)$.

Example

Find the dimension of the subspace

$$U = \{[x, y, z, w] \mid x - 2y + 3z + w = 0, 3x - 5y + z + 8w = 0\}$$

Answer: See that $U = \text{null} \left(\begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} \right)$. We have

$$\text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} \stackrel{-3R_1 + R_2}{=} \text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 0 & 1 & -8 & 5 \end{bmatrix} = 2.$$

Example

Find the dimension of the subspace

$$U = \{[x, y, z, w] \mid x - 2y + 3z + w = 0, 3x - 5y + z + 8w = 0\}$$

Answer: See that $U = \text{null} \left(\begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} \right)$. We have

$$\text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} \stackrel{-3R_1+R_2}{=} \text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 0 & 1 & -8 & 5 \end{bmatrix} = 2.$$

$$\text{Hence } \dim(U) = 4 - \text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} = 4 - 2 = 2.$$

Example

Find the dimension of the subspace

$$U = \{[x, y, z, w] \mid x - 2y + 3z + w = 0, 3x - 5y + z + 8w = 0\}$$

Answer: See that $U = \text{null} \left(\begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} \right)$. We have

$$\text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} \stackrel{-3R_1+R_2}{=} \text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 0 & 1 & -8 & 5 \end{bmatrix} = 2.$$

$$\text{Hence } \dim(U) = 4 - \text{rank} \begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -5 & 1 & 8 \end{bmatrix} = 4 - 2 = 2.$$

Example

Find the dimension of

$$U = \left\{ (x + y, x - 2y, 3x) \mid x, y \in \mathbb{R} \right\}$$

Answer:

Example

Find the dimension of

$$U = \left\{ (x + y, x - 2y, 3x) \mid x, y \in \mathbb{R} \right\}$$

Answer: See that $U = \text{im} \left(\begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \right) = \text{col} \left(\begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \right)$.

Example

Find the dimension of

$$U = \left\{ (x + y, x - 2y, 3x) \mid x, y \in \mathbb{R} \right\}$$

Answer: See that $U = \text{im} \left(\begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \right) = \text{col} \left(\begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \right)$. We have

$$\dim(U) = \text{rank} \begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \xrightarrow[-3R_1+R_3]{-R_1+R_2} \text{rank} \begin{bmatrix} 1 & 1 \\ 0 & -3 \\ 0 & -3 \end{bmatrix} \xrightarrow[-R_2+R_3]{-1/3R_2} \text{rank} \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} = 2.$$

Example

Find the dimension of

$$U = \left\{ (x + y, x - 2y, 3x) \mid x, y \in \mathbb{R} \right\}$$

Answer: See that $U = \text{im} \left(\begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \right) = \text{col} \left(\begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \right)$. We have

$$\dim(U) = \text{rank} \begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 3 & 0 \end{bmatrix} \stackrel{-R_1+R_2}{\stackrel{-3R_1+R_3}{=}} \text{rank} \begin{bmatrix} 1 & 1 \\ 0 & -3 \\ 0 & -3 \end{bmatrix} \stackrel{-1/3R_2}{\stackrel{-R_2+R_3}{=}} \text{rank} \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} = 2.$$

Example

Let A be a 3×6 matrix and let $\dim(\text{null}(A)) = 2$. Which of the following statements are true?

- (i) All bases of the $\text{col}(A)$ have four vectors.
- (ii) $\dim(\text{row}(A)) = 1$.

Example

Let A be a 3×6 matrix and let $\dim(\text{null}(A)) = 2$. Which of the following statements are true?

- (i) All bases of the $\text{col}(A)$ have four vectors.
- (ii) $\dim(\text{row}(A)) = 1$.

Answer:

Example

Let A be a 3×6 matrix and let $\dim(\text{null}(A)) = 2$. Which of the following statements are true?

- (i) All bases of the $\text{col}(A)$ have four vectors.
- (ii) $\dim(\text{row}(A)) = 1$.

Answer: We have $\dim(\text{null}(A)) = 6 - \text{rank}(A) = 2$. Hence $\text{rank}(A) = 4$.

- (i) True. Since $\dim(\text{col}(A)) = \text{rank}(A) = 4$. Hence all bases of the $\text{col}(A)$ have four vectors.

Example

Let A be a 3×6 matrix and let $\dim(\text{null}(A)) = 2$. Which of the following statements are true?

- (i) All bases of the $\text{col}(A)$ have four vectors.
- (ii) $\dim(\text{row}(A)) = 1$.

Answer: We have $\dim(\text{null}(A)) = 6 - \text{rank}(A) = 2$. Hence $\text{rank}(A) = 4$.

- (i) True. Since $\dim(\text{col}(A)) = \text{rank}(A) = 4$. Hence all bases of the $\text{col}(A)$ have four vectors.
- (ii) False. Since $\dim(\text{row}(A)) = \text{rank}(A) = 4$.

Example

Let A be a 3×6 matrix and let $\dim(\text{null}(A)) = 2$. Which of the following statements are true?

- (i) All bases of the $\text{col}(A)$ have four vectors.
- (ii) $\dim(\text{row}(A)) = 1$.

Answer: We have $\dim(\text{null}(A)) = 6 - \text{rank}(A) = 2$. Hence $\text{rank}(A) = 4$.

- (i) True. Since $\dim(\text{col}(A)) = \text{rank}(A) = 4$. Hence all bases of the $\text{col}(A)$ have four vectors.
- (ii) False. Since $\dim(\text{row}(A)) = \text{rank}(A) = 4$.

Example

Let A is a 100×350 matrix. Which of the following statements are true?

- (i) $\dim(\text{Null}(A))$ must be at least 250.
- (ii) $\dim(\text{row}(A)) + \dim(\text{col}(A)) = 450$.

Answer:

Example

Let A is a 100×350 matrix. Which of the following statements are true?

- (i) $\dim(\text{Null}(A))$ must be at least 250.
- (ii) $\dim(\text{row}(A)) + \dim(\text{col}(A)) = 450$.

Answer:

- (i) True. Since $\dim(\text{Null}(A)) = 350 - \text{rank}(A) = 4$. Moreover, $\text{rank}(A) \leq \min\{100, 350\} = 100$. Thus,

$$\dim(\text{Null}(A)) = 350 - \text{rank}(A) \geq 250 .$$

Example

Let A is a 100×350 matrix. Which of the following statements are true?

- (i) $\dim(\text{Null}(A))$ must be at least 250.
- (ii) $\dim(\text{row}(A)) + \dim(\text{col}(A)) = 450$.

Answer:

- (i) True. Since $\dim(\text{Null}(A)) = 350 - \text{rank}(A) = 4$. Moreover, $\text{rank}(A) \leq \min\{100, 350\} = 100$. Thus,

$$\dim(\text{Null}(A)) = 350 - \text{rank}(A) \geq 250 .$$

- (ii) False. Since $\dim(\text{row}(A)) + \dim(\text{col}(A)) \leq 200$.

Example

Let A is a 100×350 matrix. Which of the following statements are true?

- (i) $\dim(\text{Null}(A))$ must be at least 250.
- (ii) $\dim(\text{row}(A)) + \dim(\text{col}(A)) = 450$.

Answer:

- (i) True. Since $\dim(\text{Null}(A)) = 350 - \text{rank}(A) = 4$. Moreover, $\text{rank}(A) \leq \min\{100, 350\} = 100$. Thus,

$$\dim(\text{Null}(A)) = 350 - \text{rank}(A) \geq 250 .$$

- (ii) False. Since $\dim(\text{row}(A)) + \dim(\text{col}(A)) \leq 200$.

Exercises

Section 5.1: 1, 2, 3, 4 (page 267-268)

Section 5.2: 1, 2, 3, 4, 6, 7abcfg (page 278)

Section 5.3: 1, 2, 3, 4, 5, 6, 7, 12 (page 286-287)

Section 5.4: 1, 2, 3, 7 (page 294-295)