

Chapter 3. Determinants and Diagonalization

- 1 The Cofactor Expansion
- 2 Determinants and Matrix Inverse
- 3 Diagonalization and Eigenvectors

3.1 The Cofactor Expansion

Let $A = [a_{ij}]_{n \times n}$ be a square matrix.

Definition

- A_{ij} is the $(n-1) \times (n-1)$ matrix obtained from A by deleting row i and column j .
- (i, j) -cofactor of A

$$c_{ij}(A) = (-1)^{i+j} \det(A_{ij}).$$

3.1 The Cofactor Expansion

Let $A = [a_{ij}]_{n \times n}$ be a square matrix.

Definition

- A_{ij} is the $(n-1) \times (n-1)$ matrix obtained from A by deleting row i and column j .
- (i, j) -cofactor of A

$$c_{ij}(A) = (-1)^{i+j} \det(A_{ij}).$$

- The **determinant** of A , $\det(A)$ or $|A|$ is defined recursively:

$$\det(A) = a_{i1}c_{i1} + a_{i2}c_{i2} + \cdots + a_{in}c_{in}.$$

3.1 The Cofactor Expansion

Let $A = [a_{ij}]_{n \times n}$ be a square matrix.

Definition

- A_{ij} is the $(n-1) \times (n-1)$ matrix obtained from A by deleting row i and column j .
- (i, j) -cofactor of A

$$c_{ij}(A) = (-1)^{i+j} \det(A_{ij}).$$

- The **determinant** of A , $\det(A)$ or $|A|$ is defined recursively:

$$\det(A) = a_{i1}c_{i1} + a_{i2}c_{i2} + \cdots + a_{in}c_{in}.$$

3.1 The Cofactor Expansion

Let $A = [a_{ij}]_{n \times n}$ be a square matrix.

Definition

- A_{ij} is the $(n-1) \times (n-1)$ matrix obtained from A by deleting row i and column j .
- (i, j) -cofactor of A

$$c_{ij}(A) = (-1)^{i+j} \det(A_{ij}).$$

- The **determinant** of A , $\det(A)$ or $|A|$ is defined recursively:

$$\det(A) = a_{i1}c_{i1} + a_{i2}c_{i2} + \cdots + a_{in}c_{in}.$$

Example

- $\det(a) = a$
- $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$

Example

- $\det(a) = a$
- $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$
-

$$\begin{aligned} & \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \\ &= a_{11}(-1)^{1+1} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} + a_{12}(-1)^{1+2} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} \\ & \quad + a_{13}(-1)^{1+3} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \end{aligned}$$

Example

- $\det(a) = a$
- $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$
-

$$\begin{aligned} & \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \\ &= a_{11}(-1)^{1+1} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} + a_{12}(-1)^{1+2} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} \\ & \quad + a_{13}(-1)^{1+3} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \end{aligned}$$

Sarrus rule

Example

We have

$$\det \begin{bmatrix} 1 & -1 & 2 \\ 3 & 0 & 2 \\ -2 & 1 & -1 \end{bmatrix} = \begin{vmatrix} 1 & -1 & 2 \\ 3 & 0 & 2 \\ -2 & 1 & -1 \end{vmatrix} \begin{vmatrix} 1 & -1 \\ 3 & 0 \\ -2 & 1 \end{vmatrix}$$
$$= 1 \cdot 0 \cdot (-1) + (-1) \cdot 2 \cdot (-2) + 2 \cdot 3 \cdot 1 - 2 \cdot 0 \cdot (-2) - 1 \cdot 2 \cdot 1 - (-1) \cdot 3 \cdot (-1) = 5.$$

Sarrus rule

Example

We have

$$\det \begin{bmatrix} 1 & -1 & 2 \\ 3 & 0 & 2 \\ -2 & 1 & -1 \end{bmatrix} = \begin{vmatrix} 1 & -1 & 2 \\ 3 & 0 & 2 \\ -2 & 1 & -1 \end{vmatrix} \begin{vmatrix} 1 & -1 \\ 3 & 0 \\ -2 & 1 \end{vmatrix}$$
$$= 1 \cdot 0 \cdot (-1) + (-1) \cdot 2 \cdot (-2) + 2 \cdot 3 \cdot 1 - 2 \cdot 0 \cdot (-2) - 1 \cdot 2 \cdot 1 - (-1) \cdot 3 \cdot (-1) = 5.$$

Example

Find the determinant of $A = \begin{bmatrix} 1 & 3 & -1 & 0 \\ -1 & 1 & 0 & 5 \\ 2 & 0 & 1 & 0 \\ 3 & 0 & -2 & 0 \end{bmatrix}$.

Answer:

Example

Find the determinant of $A = \begin{bmatrix} 1 & 3 & -1 & 0 \\ -1 & 1 & 0 & 5 \\ 2 & 0 & 1 & 0 \\ 3 & 0 & -2 & 0 \end{bmatrix}$.

Answer: Expand along the fourth column of A :

Example

Find the determinant of $A = \begin{bmatrix} 1 & 3 & -1 & 0 \\ -1 & 1 & 0 & 5 \\ 2 & 0 & 1 & 0 \\ 3 & 0 & -2 & 0 \end{bmatrix}$.

Answer: Expand along the fourth column of A :

$$|A| = 5 \cdot (-1)^{2+4} \begin{vmatrix} 1 & 3 & -1 \\ 2 & 0 & 1 \\ 3 & 0 & -2 \end{vmatrix}$$

expand along the second column

$$= 5 \cdot 3 \cdot (-1)^{1+2} \begin{vmatrix} 2 & 1 \\ 3 & -2 \end{vmatrix}$$
$$= -15 \cdot [2 \cdot (-2) - 3 \cdot 1] = 105.$$

Example

Find the determinant of $A = \begin{bmatrix} 1 & 3 & -1 & 0 \\ -1 & 1 & 0 & 5 \\ 2 & 0 & 1 & 0 \\ 3 & 0 & -2 & 0 \end{bmatrix}$.

Answer: Expand along the fourth column of A :

$$|A| = 5 \cdot (-1)^{2+4} \begin{vmatrix} 1 & 3 & -1 \\ 2 & 0 & 1 \\ 3 & 0 & -2 \end{vmatrix}$$

expand along the second column $\underline{=}$

$$5 \cdot 3 \cdot (-1)^{1+2} \begin{vmatrix} 2 & 1 \\ 3 & -2 \end{vmatrix}$$
$$= -15 \cdot [2 \cdot (-2) - 3 \cdot 1] = 105 .$$

Example

Let $A = \begin{bmatrix} 4 & -5 & 7 \\ 3 & 6 & -2 \\ 1 & 8 & -9 \end{bmatrix}$. Find $(2, 3)$ -cofactor of A .

Answer:

Example

Let $A = \begin{bmatrix} 4 & -5 & 7 \\ 3 & 6 & -2 \\ 1 & 8 & -9 \end{bmatrix}$. Find $(2, 3)$ -cofactor of A .

Answer: $(2, 3)$ -cofactor of A is

$$(-1)^{2+3} \begin{vmatrix} 4 & -5 \\ 1 & 8 \end{vmatrix} = -[4 * 8 - 1 * (-5)] = -37.$$

Example

Let $A = \begin{bmatrix} 4 & -5 & 7 \\ 3 & 6 & -2 \\ 1 & 8 & -9 \end{bmatrix}$. Find $(2, 3)$ -cofactor of A .

Answer: $(2, 3)$ -cofactor of A is

$$(-1)^{2+3} \begin{vmatrix} 4 & -5 \\ 1 & 8 \end{vmatrix} = -[4 * 8 - 1 * (-5)] = -37.$$

Elementary operations and Determinant

- Interchange two rows (or column), the determinant change the sign.
- Multiply a constant c to a row (or column), the determinant is multiple by the constant.

Elementary operations and Determinant

- Interchange two rows (or column), the determinant change the sign.
- Multiply a constant c to a row (or column), the determinant is multiple by the constant.
- Add a multiple of a row (or column) to a different row (or column), the determinant does not change.

Elementary operations and Determinant

- Interchange two rows (or column), the determinant change the sign.
- Multiply a constant c to a row (or column), the determinant is multiple by the constant.
- Add a multiple of a row (or column) to a different row (or column), the determinant does not change.

Example

$$\begin{vmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 4 & 5 \end{vmatrix} \xrightarrow{R_1 \leftrightarrow R_3} - \begin{vmatrix} 3 & 4 & 5 \\ 2 & 4 & 6 \\ 1 & 2 & 3 \end{vmatrix} \xrightarrow{1/2 R_2} -2 \begin{vmatrix} 3 & 4 & 5 \\ 1 & 2 & 3 \\ 1 & 2 & 3 \end{vmatrix} \xrightarrow{-R_2 + R_3} -2 \begin{vmatrix} 3 & 4 & 5 \\ 1 & 2 & 3 \\ 0 & 0 & 0 \end{vmatrix} = 0$$

Elementary operations and Determinant

- Interchange two rows (or column), the determinant change the sign.
- Multiply a constant c to a row (or column), the determinant is multiple by the constant.
- Add a multiple of a row (or column) to a different row (or column), the determinant does not change.

Example

$$\begin{vmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 4 & 5 \end{vmatrix} \xrightarrow{R_1 \leftrightarrow R_3} - \begin{vmatrix} 3 & 4 & 5 \\ 2 & 4 & 6 \\ 1 & 2 & 3 \end{vmatrix} \xrightarrow{1/2 R_2} -2 \begin{vmatrix} 3 & 4 & 5 \\ 1 & 2 & 3 \\ 1 & 2 & 3 \end{vmatrix} \xrightarrow{-R_2 + R_3} -2 \begin{vmatrix} 3 & 4 & 5 \\ 1 & 2 & 3 \\ 0 & 0 & 0 \end{vmatrix} = 0$$

Example

Let $\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 5$. Find $\begin{vmatrix} p+x & q+y & r+z \\ 3a-2x & 3b-2y & 3c-2z \\ 3x & 3y & 3z \end{vmatrix}$.

Answer:

Example

Let $\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 5$. Find $\begin{vmatrix} p+x & q+y & r+z \\ 3a-2x & 3b-2y & 3c-2z \\ 3x & 3y & 3z \end{vmatrix}$.

Answer: We have

$$\begin{vmatrix} p+x & q+y & r+z \\ 3a-2x & 3b-2y & 3c-2z \\ 3x & 3y & 3z \end{vmatrix} \xrightarrow{1/3 R_3} 3 \begin{vmatrix} p+x & q+y & r+z \\ 3a-2x & 3b-2y & 3c-2z \\ x & y & z \end{vmatrix}$$

$$\xrightarrow{-R_3 + R_1, 2R_3 + R_2} 3 \begin{vmatrix} p & q & r \\ 3a & 3b & 3c \\ x & y & z \end{vmatrix} \xrightarrow{1/3 R_2} 9 \begin{vmatrix} p & q & r \\ a & b & c \\ x & y & z \end{vmatrix}$$

$$\xrightarrow{R_1 \leftrightarrow R_2} -9 \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = -9 * 5 = -45.$$

Example

Let $\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 5$. Find $\begin{vmatrix} p+x & q+y & r+z \\ 3a-2x & 3b-2y & 3c-2z \\ 3x & 3y & 3z \end{vmatrix}$.

Answer: We have

$$\begin{vmatrix} p+x & q+y & r+z \\ 3a-2x & 3b-2y & 3c-2z \\ 3x & 3y & 3z \end{vmatrix} \xrightarrow{1/3 R_3} 3 \begin{vmatrix} p+x & q+y & r+z \\ 3a-2x & 3b-2y & 3c-2z \\ x & y & z \end{vmatrix}$$

$$\xrightarrow{-R_3 + R_1, 2R_3 + R_2} 3 \begin{vmatrix} p & q & r \\ 3a & 3b & 3c \\ x & y & z \end{vmatrix} \xrightarrow{1/3 R_2} 9 \begin{vmatrix} p & q & r \\ a & b & c \\ x & y & z \end{vmatrix}$$

$$\xrightarrow{R_1 \leftrightarrow R_2} -9 \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = -9 * 5 = -45.$$

Example

Let $\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 10$. Find $\begin{vmatrix} x & 0 & y & z \\ p & 0 & q & r \\ a & 0 & b & c \\ 2 & 3 & 4 & 5 \end{vmatrix}$.

Answer:

Example

Let $\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 10$. Find $\begin{vmatrix} x & 0 & y & z \\ p & 0 & q & r \\ a & 0 & b & c \\ 2 & 3 & 4 & 5 \end{vmatrix}$.

Answer: Expand the determinant along the second column,

Example

Let $\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 10$. Find $\begin{vmatrix} x & 0 & y & z \\ p & 0 & q & r \\ a & 0 & b & c \\ 2 & 3 & 4 & 5 \end{vmatrix}$.

Answer: Expand the determinant along the second column,

$$\begin{vmatrix} x & 0 & y & z \\ p & 0 & q & r \\ a & 0 & b & c \\ 2 & 3 & 4 & 5 \end{vmatrix} = 3 * (-1)^{4+2} \begin{vmatrix} x & y & z \\ p & q & r \\ a & b & c \end{vmatrix} \stackrel{R_1 \leftrightarrow R_3}{=} -3 \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = -30$$

Example

Let $\begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 10$. Find $\begin{vmatrix} x & 0 & y & z \\ p & 0 & q & r \\ a & 0 & b & c \\ 2 & 3 & 4 & 5 \end{vmatrix}$.

Answer: Expand the determinant along the second column,

$$\begin{vmatrix} x & 0 & y & z \\ p & 0 & q & r \\ a & 0 & b & c \\ 2 & 3 & 4 & 5 \end{vmatrix} = 3 * (-1)^{4+2} \begin{vmatrix} x & y & z \\ p & q & r \\ a & b & c \end{vmatrix} \stackrel{R_1 \leftrightarrow R_3}{=} -3 \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = -30$$

Properties of Determinant

- $\det(AB) = \det(A) \det(B)$. In particular $\det(A^m) = [\det(A)]^m$, for $m \in \mathbb{N}^*$.
- $\det(A^{-1}) = 1 / \det(A)$
- $\det(kA) = k^n \det(A)$, for any $n \times n$ matrix A .

Properties of Determinant

- $\det(AB) = \det(A) \det(B)$. In particular $\det(A^m) = [\det(A)]^m$, for $m \in \mathbb{N}^*$.
- $\det(A^{-1}) = 1 / \det(A)$
- $\det(kA) = k^n \det(A)$, for any $n \times n$ matrix A .
- $\det(A^T) = \det(A)$

Properties of Determinant

- $\det(AB) = \det(A) \det(B)$. In particular $\det(A^m) = [\det(A)]^m$, for $m \in \mathbb{N}^*$.
- $\det(A^{-1}) = 1 / \det(A)$
- $\det(kA) = k^n \det(A)$, for any $n \times n$ matrix A .
- $\det(A^T) = \det(A)$
- If A is a square **triangular matrix**, then $\det(A)$ is the product of the entries on the **main diagonal**.

Properties of Determinant

- $\det(AB) = \det(A) \det(B)$. In particular $\det(A^m) = [\det(A)]^m$, for $m \in \mathbb{N}^*$.
- $\det(A^{-1}) = 1 / \det(A)$
- $\det(kA) = k^n \det(A)$, for any $n \times n$ matrix A .
- $\det(A^T) = \det(A)$
- If A is a square **triangular matrix**, then $\det(A)$ is the product of the entries on the **main diagonal**.

Example

$$\begin{vmatrix} 3a & 3b \\ 3c & 3d \end{vmatrix} = 9 \begin{vmatrix} a & b \\ c & d \end{vmatrix}, \quad \begin{vmatrix} 2 & 4 & -8 & 9 \\ 0 & -1 & 5 & 6 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 0 & -5 \end{vmatrix} = 2 * (-1) * 4 * (-5) = 40$$

Properties of Determinant

- $\det(AB) = \det(A) \det(B)$. In particular $\det(A^m) = [\det(A)]^m$, for $m \in \mathbb{N}^*$.
- $\det(A^{-1}) = 1 / \det(A)$
- $\det(kA) = k^n \det(A)$, for any $n \times n$ matrix A .
- $\det(A^T) = \det(A)$
- If A is a square **triangular matrix**, then $\det(A)$ is the product of the entries on the **main diagonal**.

Example

$$\begin{vmatrix} 3a & 3b \\ 3c & 3d \end{vmatrix} = 9 \begin{vmatrix} a & b \\ c & d \end{vmatrix}, \quad \begin{vmatrix} 2 & 4 & -8 & 9 \\ 0 & -1 & 5 & 6 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 0 & -5 \end{vmatrix} = 2 * (-1) * 4 * (-5) = 40$$

Example

Let A and B be 4×4 matrices with $\det(A) = 8$ and $\det(B) = 6$. Find $\det(3B^{-2}A^2B^T)$.

Answer:

Example

Let A and B be 4×4 matrices with $\det(A) = 8$ and $\det(B) = 6$. Find $\det(3B^{-2}A^2B^T)$.

Answer: We have

$$\begin{aligned}\det(3B^{-2}A^2B^T) &= 3^4 \det(B)^{-2} \det(A)^2 \det(B^T) \\ &= 3^4 \det(A)^2 \det(B)^{-1} = 3^4 \cdot 8^2 / 6 = 864.\end{aligned}$$

Example

Let A and B be 4×4 matrices with $\det(A) = 8$ and $\det(B) = 6$. Find $\det(3B^{-2}A^2B^T)$.

Answer: We have

$$\begin{aligned}\det(3B^{-2}A^2B^T) &= 3^4 \det(B)^{-2} \det(A)^2 \det(B^T) \\ &= 3^4 \det(A)^2 \det(B)^{-1} = 3^4 \cdot 8^2 / 6 = 864.\end{aligned}$$

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$.

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$.
(F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$.
(F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A . (T)
- $\det(-A) = -\det(A)$, for any square matrix A .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A . (T)
- $\det(-A) = -\det(A)$, for any square matrix A .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A . (T)
- $\det(-A) = -\det(A)$, for any square matrix A . (F)
- $\det(-A) = -\det(A)$, for any 3×3 matrix A .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A . (T)
- $\det(-A) = -\det(A)$, for any square matrix A . (F)
- $\det(-A) = -\det(A)$, for any 3×3 matrix A .

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A . (T)
- $\det(-A) = -\det(A)$, for any square matrix A . (F)
- $\det(-A) = -\det(A)$, for any 3×3 matrix A . (T)

Quiz

Determine the following statements are True or False.

- Let A and B be $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$. (F)
- $\det(kA) = k\det(A)$, for any $n \times n$ matrix A and scalar k . (F)
- $\det(AB) = \det(A)\det(B)$, for any $n \times n$ matrices A, B . (T)
- $\det(A^T A) = [\det(A)]^2$, for any square matrix A . (T)
- $\det(-A) = -\det(A)$, for any square matrix A . (F)
- $\det(-A) = -\det(A)$, for any 3×3 matrix A . (T)

Example

Let A be a 4×4 matrix, $\det(A) = 2$.

B is the matrix obtained from A by interchanging R_3 and R_2 .

C is the matrix obtained from B by adding 12 times R_3 to R_2 .

D is the matrix obtained from C by multiplying R_3 by 5.

Find $\det(BCD)$.

Answer:

Example

Let A be a 4×4 matrix, $\det(A) = 2$.

B is the matrix obtained from A by interchanging R_3 and R_2 .

C is the matrix obtained from B by adding 12 times R_3 to R_2 .

D is the matrix obtained from C by multiplying R_3 by 5.

Find $\det(BCD)$.

Answer: We have

$$\det(B) = -\det(A) = -2$$

$$\det(C) = \det(B) = -2$$

$$\det(D) = 5\det(C) = -10.$$

Example

Let A be a 4×4 matrix, $\det(A) = 2$.

B is the matrix obtained from A by interchanging R_3 and R_2 .

C is the matrix obtained from B by adding 12 times R_3 to R_2 .

D is the matrix obtained from C by multiplying R_3 by 5.

Find $\det(BCD)$.

Answer: We have

$$\det(B) = -\det(A) = -2$$

$$\det(C) = \det(B) = -2$$

$$\det(D) = 5\det(C) = -10.$$

Hence

$$\det(BCD) = \det(B)\det(C)\det(D) = -2 * (-2) * (-10) = -40.$$

Example

Let A be a 4×4 matrix, $\det(A) = 2$.

B is the matrix obtained from A by interchanging R_3 and R_2 .

C is the matrix obtained from B by adding 12 times R_3 to R_2 .

D is the matrix obtained from C by multiplying R_3 by 5.

Find $\det(BCD)$.

Answer: We have

$$\det(B) = -\det(A) = -2$$

$$\det(C) = \det(B) = -2$$

$$\det(D) = 5\det(C) = -10.$$

Hence

$$\det(BCD) = \det(B)\det(C)\det(D) = -2 * (-2) * (-10) = -40.$$

3.2 Determinants and Matrix Inverse

Let A be a square $n \times n$ matrix.

Definition

The **adjugate** of A , denoted $\text{adj}(A)$:

$$\text{adj}(A) = [c_{ij}]^T = \begin{bmatrix} c_{11} & c_{21} & \cdots & c_{n1} \\ c_{12} & c_{22} & \cdots & c_{n2} \\ \cdots & \cdots & \ddots & \cdots \\ c_{1n} & c_{2n} & \cdots & c_{nn} \end{bmatrix}$$

3.2 Determinants and Matrix Inverse

Let A be a square $n \times n$ matrix.

Definition

The **adjugate** of A , denoted $\text{adj}(A)$:

$$\text{adj}(A) = [c_{ij}]^T = \begin{bmatrix} c_{11} & c_{21} & \cdots & c_{n1} \\ c_{12} & c_{22} & \cdots & c_{n2} \\ \cdots & \cdots & \ddots & \cdots \\ c_{1n} & c_{2n} & \cdots & c_{nn} \end{bmatrix}$$

where c_{ij} is (i, j) -cofactor of A .

3.2 Determinants and Matrix Inverse

Let A be a square $n \times n$ matrix.

Definition

The **adjugate** of A , denoted $\text{adj}(A)$:

$$\text{adj}(A) = [c_{ij}]^T = \begin{bmatrix} c_{11} & c_{21} & \cdots & c_{n1} \\ c_{12} & c_{22} & \cdots & c_{n2} \\ \cdots & \cdots & \ddots & \cdots \\ c_{1n} & c_{2n} & \cdots & c_{nn} \end{bmatrix}$$

where c_{ij} is (i, j) -cofactor of A .

Theorem

Matrix A is invertible if and only if $\det(A) \neq 0$, and

$$A^{-1} = \frac{1}{\det(A)} \operatorname{adj}(A)$$

Example

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

Theorem

Matrix A is invertible if and only if $\det(A) \neq 0$, and

$$A^{-1} = \frac{1}{\det(A)} \operatorname{adj}(A)$$

Example

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then A is invertible if $\det(A) = ad - bc \neq 0$ and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Theorem

Matrix A is invertible if and only if $\det(A) \neq 0$, and

$$A^{-1} = \frac{1}{\det(A)} \operatorname{adj}(A)$$

Example

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then A is invertible if $\det(A) = ad - bc \neq 0$ and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

For example,

$$\begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}^{-1} = \frac{1}{10} \begin{bmatrix} 4 & -1 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0.4 & -0.1 \\ -0.2 & 0.3 \end{bmatrix}$$

Theorem

Matrix A is invertible if and only if $\det(A) \neq 0$, and

$$A^{-1} = \frac{1}{\det(A)} \operatorname{adj}(A)$$

Example

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then A is invertible if $\det(A) = ad - bc \neq 0$ and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

For example,

$$\begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}^{-1} = \frac{1}{10} \begin{bmatrix} 4 & -1 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0.4 & -0.1 \\ -0.2 & 0.3 \end{bmatrix}$$

Find the inverse matrix of $A = \begin{bmatrix} 2 & 0 & 1 \\ -1 & 3 & 4 \\ 1 & 5 & 7 \end{bmatrix}$.

Answer: We have

$$\det(A) = 2 * (-1)^{1+1} \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} + 1 * (-1)^{1+3} \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = 2 * 1 + (-8) = -6.$$

Find the inverse matrix of $A = \begin{bmatrix} 2 & 0 & 1 \\ -1 & 3 & 4 \\ 1 & 5 & 7 \end{bmatrix}$.

Answer: We have

$$\det(A) = 2 * (-1)^{1+1} \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} + 1 * (-1)^{1+3} \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = 2 * 1 + (-8) = -6.$$

All (i, j) -cofactors of A :

Find the inverse matrix of $A = \begin{bmatrix} 2 & 0 & 1 \\ -1 & 3 & 4 \\ 1 & 5 & 7 \end{bmatrix}$.

Answer: We have

$$\det(A) = 2 * (-1)^{1+1} \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} + 1 * (-1)^{1+3} \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = 2 * 1 + (-8) = -6.$$

All (i,j) -cofactors of A :

$$c_{11} = \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} = 1, \quad c_{12} = - \begin{vmatrix} -1 & 4 \\ 1 & 7 \end{vmatrix} = 11, \quad c_{13} = \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = -8$$

$$c_{21} = - \begin{vmatrix} 0 & 1 \\ 5 & 7 \end{vmatrix} = 5, \quad c_{22} = \begin{vmatrix} 2 & 1 \\ 1 & 7 \end{vmatrix} = 13, \quad c_{23} = - \begin{vmatrix} 2 & 0 \\ 1 & 5 \end{vmatrix} = -10$$

$$c_{31} = \begin{vmatrix} 0 & 1 \\ 3 & 4 \end{vmatrix} = -3, \quad c_{32} = - \begin{vmatrix} 2 & 1 \\ -1 & 4 \end{vmatrix} = -9, \quad c_{33} = \begin{vmatrix} 2 & 0 \\ -1 & 3 \end{vmatrix} = 6$$

Find the inverse matrix of $A = \begin{bmatrix} 2 & 0 & 1 \\ -1 & 3 & 4 \\ 1 & 5 & 7 \end{bmatrix}$.

Answer: We have

$$\det(A) = 2 * (-1)^{1+1} \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} + 1 * (-1)^{1+3} \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = 2 * 1 + (-8) = -6.$$

All (i, j) -cofactors of A :

$$c_{11} = \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} = 1, \quad c_{12} = - \begin{vmatrix} -1 & 4 \\ 1 & 7 \end{vmatrix} = 11, \quad c_{13} = \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = -8$$

$$c_{21} = - \begin{vmatrix} 0 & 1 \\ 5 & 7 \end{vmatrix} = 5, \quad c_{22} = \begin{vmatrix} 2 & 1 \\ 1 & 7 \end{vmatrix} = 13, \quad c_{23} = - \begin{vmatrix} 2 & 0 \\ 1 & 5 \end{vmatrix} = -10$$

$$c_{31} = \begin{vmatrix} 0 & 1 \\ 3 & 4 \end{vmatrix} = -3, \quad c_{32} = - \begin{vmatrix} 2 & 1 \\ -1 & 4 \end{vmatrix} = -9, \quad c_{33} = \begin{vmatrix} 2 & 0 \\ -1 & 3 \end{vmatrix} = 6$$

$$\text{Thus, } A^{-1} = \frac{1}{-6} \begin{bmatrix} 1 & 5 & -3 \\ 11 & 13 & -9 \\ -8 & -10 & 6 \end{bmatrix} = \begin{bmatrix} -1/6 & -5/6 & 1/2 \\ -11/6 & -13/6 & 3/2 \\ 4/3 & 5/3 & -1 \end{bmatrix}$$

Find the inverse matrix of $A = \begin{bmatrix} 2 & 0 & 1 \\ -1 & 3 & 4 \\ 1 & 5 & 7 \end{bmatrix}$.

Answer: We have

$$\det(A) = 2 * (-1)^{1+1} \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} + 1 * (-1)^{1+3} \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = 2 * 1 + (-8) = -6.$$

All (i, j) -cofactors of A :

$$c_{11} = \begin{vmatrix} 3 & 4 \\ 5 & 7 \end{vmatrix} = 1, \quad c_{12} = - \begin{vmatrix} -1 & 4 \\ 1 & 7 \end{vmatrix} = 11, \quad c_{13} = \begin{vmatrix} -1 & 3 \\ 1 & 5 \end{vmatrix} = -8$$

$$c_{21} = - \begin{vmatrix} 0 & 1 \\ 5 & 7 \end{vmatrix} = 5, \quad c_{22} = \begin{vmatrix} 2 & 1 \\ 1 & 7 \end{vmatrix} = 13, \quad c_{23} = - \begin{vmatrix} 2 & 0 \\ 1 & 5 \end{vmatrix} = -10$$

$$c_{31} = \begin{vmatrix} 0 & 1 \\ 3 & 4 \end{vmatrix} = -3, \quad c_{32} = - \begin{vmatrix} 2 & 1 \\ -1 & 4 \end{vmatrix} = -9, \quad c_{33} = \begin{vmatrix} 2 & 0 \\ -1 & 3 \end{vmatrix} = 6$$

$$\text{Thus, } A^{-1} = \frac{1}{-6} \begin{bmatrix} 1 & 5 & -3 \\ 11 & 13 & -9 \\ -8 & -10 & 6 \end{bmatrix} = \begin{bmatrix} -1/6 & -5/6 & 1/2 \\ -11/6 & -13/6 & 3/2 \\ 4/3 & 5/3 & -1 \end{bmatrix}$$

Properties

- $\det(A^{-1}) = 1/\det(A)$.
- $(AB)^{-1} = B^{-1}A^{-1}$, for any invertible matrices A, B .

Properties

- $\det(A^{-1}) = 1/\det(A)$.
- $(AB)^{-1} = B^{-1}A^{-1}$, for any invertible matrices A, B .
- $I^{-1} = I$.

Properties

- $\det(A^{-1}) = 1/\det(A)$.
- $(AB)^{-1} = B^{-1}A^{-1}$, for any invertible matrices A, B .
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$.

Properties

- $\det(A^{-1}) = 1/\det(A)$.
- $(AB)^{-1} = B^{-1}A^{-1}$, for any invertible matrices A, B .
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$.
- $(aA)^{-1} = 1/aA^{-1}$.

Properties

- $\det(A^{-1}) = 1/\det(A)$.
- $(AB)^{-1} = B^{-1}A^{-1}$, for any invertible matrices A, B .
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$.
- $(aA)^{-1} = 1/aA^{-1}$.
- $(A^T)^{-1} = (A^{-1})^T$.

Properties

- $\det(A^{-1}) = 1/\det(A)$.
- $(AB)^{-1} = B^{-1}A^{-1}$, for any invertible matrices A, B .
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$.
- $(aA)^{-1} = 1/aA^{-1}$.
- $(A^T)^{-1} = (A^{-1})^T$.
- $\det(\text{adj}(A)) = [\det(A)]^{n-1}$, for any invertible $n \times n$ matrix A .

Properties

- $\det(A^{-1}) = 1/\det(A)$.
- $(AB)^{-1} = B^{-1}A^{-1}$, for any invertible matrices A, B .
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$.
- $(aA)^{-1} = 1/aA^{-1}$.
- $(A^T)^{-1} = (A^{-1})^T$.
- $\det(\text{adj}(A)) = [\det(A)]^{n-1}$, for any invertible $n \times n$ matrix A .

Example

Find the $(1, 3)$ - entry of A^{-1} provided that

$$A = \begin{bmatrix} 2 & -3 & 1 \\ 0 & 3 & 0 \\ 5 & -2 & 4 \end{bmatrix}$$

Answer:

Example

Find the $(1, 3)$ - entry of A^{-1} provided that

$$A = \begin{bmatrix} 2 & -3 & 1 \\ 0 & 3 & 0 \\ 5 & -2 & 4 \end{bmatrix}$$

Answer: First, compute $\det(A)$

$$\det(A) = 3 * (-1)^{2+2} \begin{vmatrix} 2 & 1 \\ 5 & 4 \end{vmatrix} = 9$$

The $(3, 1)$ -cofactor of A :

Example

Find the $(1, 3)$ - entry of A^{-1} provided that

$$A = \begin{bmatrix} 2 & -3 & 1 \\ 0 & 3 & 0 \\ 5 & -2 & 4 \end{bmatrix}$$

Answer: First, compute $\det(A)$

$$\det(A) = 3 * (-1)^{2+2} \begin{vmatrix} 2 & 1 \\ 5 & 4 \end{vmatrix} = 9$$

The $(3, 1)$ -cofactor of A :

$$c_{31} = (-1)^{3+1} \begin{vmatrix} -3 & 1 \\ 3 & 0 \end{vmatrix} = -3$$

Example

Find the $(1, 3)$ -entry of A^{-1} provided that

$$A = \begin{bmatrix} 2 & -3 & 1 \\ 0 & 3 & 0 \\ 5 & -2 & 4 \end{bmatrix}$$

Answer: First, compute $\det(A)$

$$\det(A) = 3 * (-1)^{2+2} \begin{vmatrix} 2 & 1 \\ 5 & 4 \end{vmatrix} = 9$$

The $(3, 1)$ -cofactor of A :

$$c_{31} = (-1)^{3+1} \begin{vmatrix} -3 & 1 \\ 3 & 0 \end{vmatrix} = -3$$

Hence, $(1, 3)$ -entry of A^{-1} is $-3/9 = -1/3$.

Example

Find the $(1, 3)$ -entry of A^{-1} provided that

$$A = \begin{bmatrix} 2 & -3 & 1 \\ 0 & 3 & 0 \\ 5 & -2 & 4 \end{bmatrix}$$

Answer: First, compute $\det(A)$

$$\det(A) = 3 * (-1)^{2+2} \begin{vmatrix} 2 & 1 \\ 5 & 4 \end{vmatrix} = 9$$

The $(3, 1)$ -cofactor of A :

$$c_{31} = (-1)^{3+1} \begin{vmatrix} -3 & 1 \\ 3 & 0 \end{vmatrix} = -3$$

Hence, $(1, 3)$ -entry of A^{-1} is $-3/9 = -1/3$.

Example

Let $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 3 & 0 \\ -5 & 0 & x \end{bmatrix}$. Find the values of x so that A is invertible.

Answer:

Example

Let $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 3 & 0 \\ -5 & 0 & x \end{bmatrix}$. Find the values of x so that A is invertible.

Answer: We have

$$\det(A) = 3 * (-1)^{2+2} \begin{vmatrix} 1 & 2 \\ -5 & x \end{vmatrix} = 3(x + 10)$$

Example

Let $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 3 & 0 \\ -5 & 0 & x \end{bmatrix}$. Find the values of x so that A is invertible.

Answer: We have

$$\det(A) = 3 * (-1)^{2+2} \begin{vmatrix} 1 & 2 \\ -5 & x \end{vmatrix} = 3(x + 10)$$

Hence A is invertible if and only if $\det(A) \neq 0 \Leftrightarrow x \neq -10$.

Example

Let $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 3 & 0 \\ -5 & 0 & x \end{bmatrix}$. Find the values of x so that A is invertible.

Answer: We have

$$\det(A) = 3 * (-1)^{2+2} \begin{vmatrix} 1 & 2 \\ -5 & x \end{vmatrix} = 3(x + 10)$$

Hence A is invertible if and only if $\det(A) \neq 0 \Leftrightarrow x \neq -10$.

Example

Let $A = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 3 & 4 \\ -5 & 2 & x \end{bmatrix}$. Find the $(1, 2)$ -entry of A^{-1} .

Answer:

Example

Let $A = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 3 & 4 \\ -5 & 2 & x \end{bmatrix}$. Find the $(1, 2)$ -entry of A^{-1} .

Answer: $\det(A) = 2 * (-1)^{1+3} \begin{vmatrix} 1 & 3 \\ -5 & 2 \end{vmatrix} = 2 * 17 = 34$. The $(2, 1)$ -cofactor of A is

Example

Let $A = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 3 & 4 \\ -5 & 2 & x \end{bmatrix}$. Find the $(1, 2)$ -entry of A^{-1} .

Answer: $\det(A) = 2 * (-1)^{1+3} \begin{vmatrix} 1 & 3 \\ -5 & 2 \end{vmatrix} = 2 * 17 = 34$. The $(2, 1)$ -cofactor of A is

$$c_{2,1} = (-1)^{2+1} \begin{vmatrix} 0 & 2 \\ 2 & x \end{vmatrix} = 4.$$

Example

Let $A = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 3 & 4 \\ -5 & 2 & x \end{bmatrix}$. Find the $(1, 2)$ -entry of A^{-1} .

Answer: $\det(A) = 2 * (-1)^{1+3} \begin{vmatrix} 1 & 3 \\ -5 & 2 \end{vmatrix} = 2 * 17 = 34$. The $(2, 1)$ -cofactor of A is

$$c_{2,1} = (-1)^{2+1} \begin{vmatrix} 0 & 2 \\ 2 & x \end{vmatrix} = 4.$$

Hence $(1, 2)$ -entry of A^{-1} is $4/34 = 2/17$.

Example

Let $A = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 3 & 4 \\ -5 & 2 & x \end{bmatrix}$. Find the $(1, 2)$ -entry of A^{-1} .

Answer: $\det(A) = 2 * (-1)^{1+3} \begin{vmatrix} 1 & 3 \\ -5 & 2 \end{vmatrix} = 2 * 17 = 34$. The $(2, 1)$ -cofactor of A is

$$c_{2,1} = (-1)^{2+1} \begin{vmatrix} 0 & 2 \\ 2 & x \end{vmatrix} = 4.$$

Hence $(1, 2)$ -entry of A^{-1} is $4/34 = 2/17$.

Cramer's rule

Theorem

If A is an invertible $n \times n$ matrix, the solution to the system

$$Ax = b$$

of n equations in the variables x_1, x_2, \dots, x_n is given by

$$x_1 = \frac{\det A_1}{\det A}$$

$$x_2 = \frac{\det A_2}{\det A}$$

\dots

$$x_n = \frac{\det A_n}{\det A}$$

where A_k is the matrix obtained from A by replacing column k by b .

Example

Find y satisfying the following system:

$$4x - y + 3z = 1$$

$$6x + 2y - z = 0$$

$$3x + 3y + 2z = -1$$

Example

Find y satisfying the following system:

$$4x - y + 3z = 1$$

$$6x + 2y - z = 0$$

$$3x + 3y + 2z = -1$$

Answer:

Example

Find y satisfying the following system:

$$\begin{array}{rcrcrcrcrcl} 4x & - & y & + & 3z & = & 1 \\ 6x & + & 2y & - & z & = & 0 \\ 3x & + & 3y & + & 2z & = & -1 \end{array}$$

Answer: We have

$$\begin{vmatrix} 4 & -1 & 3 \\ 6 & 2 & -1 \\ 3 & 3 & 2 \end{vmatrix} = 79, \quad \begin{vmatrix} 4 & 1 & 3 \\ 6 & 0 & -1 \\ 3 & -1 & 2 \end{vmatrix} = -37.$$

Example

Find y satisfying the following system:

$$\begin{array}{rcrcrcrcrcl} 4x & - & y & + & 3z & = & 1 \\ 6x & + & 2y & - & z & = & 0 \\ 3x & + & 3y & + & 2z & = & -1 \end{array}$$

Answer: We have

$$\begin{vmatrix} 4 & -1 & 3 \\ 6 & 2 & -1 \\ 3 & 3 & 2 \end{vmatrix} = 79, \quad \begin{vmatrix} 4 & 1 & 3 \\ 6 & 0 & -1 \\ 3 & -1 & 2 \end{vmatrix} = -37.$$

Hence $y = -37/79$.

Example

Find y satisfying the following system:

$$\begin{array}{rcrcrcrcrcl} 4x & - & y & + & 3z & = & 1 \\ 6x & + & 2y & - & z & = & 0 \\ 3x & + & 3y & + & 2z & = & -1 \end{array}$$

Answer: We have

$$\begin{vmatrix} 4 & -1 & 3 \\ 6 & 2 & -1 \\ 3 & 3 & 2 \end{vmatrix} = 79, \quad \begin{vmatrix} 4 & 1 & 3 \\ 6 & 0 & -1 \\ 3 & -1 & 2 \end{vmatrix} = -37.$$

Hence $y = -37/79$.

3.3 Diagonalization and Eigenvectors

How to compute A^k when A is a square matrix and n large?

Definition

A diagonal matrix is
$$\begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$$

3.3 Diagonalization and Eigenvectors

How to compute A^k when A is a square matrix and n large?

Definition

A diagonal matrix is $\begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$

We can see

$$\begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}^k = \begin{bmatrix} \lambda_1^k & 0 & \cdots & 0 \\ 0 & \lambda_2^k & \cdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \cdots & \lambda_n^k \end{bmatrix}$$

3.3 Diagonalization and Eigenvectors

How to compute A^k when A is a square matrix and n large?

Definition

A diagonal matrix is $\begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$

We can see

$$\begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}^k = \begin{bmatrix} \lambda_1^k & 0 & \cdots & 0 \\ 0 & \lambda_2^k & \cdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \cdots & \lambda_n^k \end{bmatrix}$$

Theorem

Suppose that $A = PDP^{-1}$ with D is a diagonal matrix then

$$A^k = PD^kP^{-1}.$$

Definition

If A is an $n \times n$ matrix, a number λ is called an **eigenvalue** of A if

$$AX = \lambda X \text{ for some column } X \neq 0 \text{ in } \mathbb{R}^n.$$

Theorem

Suppose that $A = PDP^{-1}$ with D is a diagonal matrix then

$$A^k = PD^kP^{-1}.$$

Definition

If A is an $n \times n$ matrix, a number λ is called an **eigenvalue** of A if

$$AX = \lambda X \text{ for some column } X \neq 0 \text{ in } \mathbb{R}^n.$$

In this case X is called an **eigenvector** of A corresponding to the eigenvalue λ , or a λ -eigenvector.

Theorem

Suppose that $A = PDP^{-1}$ with D is a diagonal matrix then

$$A^k = PD^kP^{-1}.$$

Definition

If A is an $n \times n$ matrix, a number λ is called an **eigenvalue** of A if

$$AX = \lambda X \text{ for some column } X \neq 0 \text{ in } \mathbb{R}^n.$$

In this case X is called an **eigenvector** of A corresponding to the eigenvalue λ , or a λ -eigenvector.

To find eigenvalues of A , we solve the equation

$$\det(\lambda I - A) = 0$$

Theorem

The *characteristic polynomial* $c_A(x)$ of an $n \times n$ matrix A is defined by

$$c_A(x) = \det(xI - A).$$

To find eigenvalues of A , we solve the equation

$$\det(\lambda I - A) = 0$$

Theorem

The *characteristic polynomial* $c_A(x)$ of an $n \times n$ matrix A is defined by

$$c_A(x) = \det(xI - A).$$

Example

Find all eigenvalues of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

To find eigenvalues of A , we solve the equation

$$\det(\lambda I - A) = 0$$

Theorem

The *characteristic polynomial* $c_A(x)$ of an $n \times n$ matrix A is defined by

$$c_A(x) = \det(xI - A).$$

Example

Find all eigenvalues of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

The characteristic polynomial of A is

$$c_A(x) = \begin{vmatrix} x-1 & -8 \\ -1 & x-3 \end{vmatrix} = (x-1)(x-3) - 8 = x^2 - 4x - 5.$$

To find eigenvalues of A , we solve the equation

$$\det(\lambda I - A) = 0$$

Theorem

The *characteristic polynomial* $c_A(x)$ of an $n \times n$ matrix A is defined by

$$c_A(x) = \det(xI - A).$$

Example

Find all eigenvalues of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

The characteristic polynomial of A is

$$c_A(x) = \begin{vmatrix} x-1 & -8 \\ -1 & x-3 \end{vmatrix} = (x-1)(x-3) - 8 = x^2 - 4x - 5.$$

$c_A(x) = 0 \Rightarrow x = -1$ or $x = 5$. Hence A has two eigenvalues -1 and 5 .

To find eigenvalues of A , we solve the equation

$$\det(\lambda I - A) = 0$$

Theorem

The *characteristic polynomial* $c_A(x)$ of an $n \times n$ matrix A is defined by

$$c_A(x) = \det(xI - A).$$

Example

Find all eigenvalues of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

The characteristic polynomial of A is

$$c_A(x) = \begin{vmatrix} x-1 & -8 \\ -1 & x-3 \end{vmatrix} = (x-1)(x-3) - 8 = x^2 - 4x - 5.$$

$c_A(x) = 0 \Rightarrow x = -1$ or $x = 5$. Hence A has two eigenvalues -1 and 5 .

Example

Find all eigenvectors of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

Answer: As above example, A has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

Example

Find all eigenvectors of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

Answer: As above example, A has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

- -1 -eigenvectors $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ is nonzero solution of the system

$$\begin{bmatrix} -2 & -8 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow x_1 + 4x_2 = 0 \Rightarrow x = \begin{bmatrix} 4t \\ -t \end{bmatrix} = t \begin{bmatrix} 4 \\ -1 \end{bmatrix}, \quad t \neq 0$$

Example

Find all eigenvectors of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

Answer: As above example, A has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

- -1 -eigenvectors $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ is nonzero solution of the system

$$\begin{bmatrix} -2 & -8 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow x_1 + 4x_2 = 0 \Rightarrow x = \begin{bmatrix} 4t \\ -t \end{bmatrix} = t \begin{bmatrix} 4 \\ -1 \end{bmatrix}, \quad t \neq 0.$$

- 5 -eigenvectors $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ is nonzero solution of the system

$$\begin{bmatrix} 4 & -8 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow x_1 - 2x_2 = 0 \Rightarrow X = \begin{bmatrix} 2t \\ t \end{bmatrix} = t \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad t \neq 0.$$

Example

Find all eigenvectors of $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$.

Answer: As above example, A has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

- -1 -eigenvectors $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ is nonzero solution of the system

$$\begin{bmatrix} -2 & -8 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow x_1 + 4x_2 = 0 \Rightarrow x = \begin{bmatrix} 4t \\ -t \end{bmatrix} = t \begin{bmatrix} 4 \\ -1 \end{bmatrix}, \quad t \neq 0.$$

- 5 -eigenvectors $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ is nonzero solution of the system

$$\begin{bmatrix} 4 & -8 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Leftrightarrow x_1 - 2x_2 = 0 \Rightarrow X = \begin{bmatrix} 2t \\ t \end{bmatrix} = t \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad t \neq 0.$$

Theorem

An $n \times n$ matrix A is *diagonalizable* if and only if it has eigenvectors x_1, x_2, \dots, x_n such that the matrix $P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$ is invertible.

$$A = P \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) P^{-1}$$

Theorem

An $n \times n$ matrix A is *diagonalizable* if and only if it has eigenvectors x_1, x_2, \dots, x_n such that the matrix $P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$ is invertible.

$$A = P \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) P^{-1}$$

where λ_i is the eigenvalue of A corresponding to x_i .

Theorem

An $n \times n$ matrix A is *diagonalizable* if and only if it has eigenvectors x_1, x_2, \dots, x_n such that the matrix $P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$ is invertible.

$$A = P \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) P^{-1}$$

where λ_i is the eigenvalue of A corresponding to x_i .

Example

The matrix $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$ has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

Theorem

An $n \times n$ matrix A is *diagonalizable* if and only if it has eigenvectors x_1, x_2, \dots, x_n such that the matrix $P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$ is invertible.

$$A = P \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) P^{-1}$$

where λ_i is the eigenvalue of A corresponding to x_i .

Example

The matrix $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$ has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

$x_1 = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$ is an eigenvector corresponding to λ_1 .

Theorem

An $n \times n$ matrix A is *diagonalizable* if and only if it has eigenvectors x_1, x_2, \dots, x_n such that the matrix $P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$ is invertible.

$$A = P \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) P^{-1}$$

where λ_i is the eigenvalue of A corresponding to x_i .

Example

The matrix $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$ has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

$x_1 = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$ is an eigenvector corresponding to λ_1 .

$x_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvector corresponding to λ_2 .

Theorem

An $n \times n$ matrix A is **diagonalizable** if and only if it has eigenvectors x_1, x_2, \dots, x_n such that the matrix $P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$ is invertible.

$$A = P \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) P^{-1}$$

where λ_i is the eigenvalue of A corresponding to x_i .

Example

The matrix $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$ has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

$x_1 = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$ is an eigenvector corresponding to λ_1 .

$x_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvector corresponding to λ_2 .

Hence:

$$A = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 5 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}^{-1}.$$

Theorem

An $n \times n$ matrix A is **diagonalizable** if and only if it has eigenvectors x_1, x_2, \dots, x_n such that the matrix $P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$ is invertible.

$$A = P \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) P^{-1}$$

where λ_i is the eigenvalue of A corresponding to x_i .

Example

The matrix $A = \begin{bmatrix} 1 & 8 \\ 1 & 3 \end{bmatrix}$ has two eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$.

$x_1 = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$ is an eigenvector corresponding to λ_1 .

$x_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvector corresponding to λ_2 .

Hence:

$$A = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 5 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}^{-1}.$$

Example

Let A be a 2×2 matrix with eigenvalues 2 and 3, and with corresponding eigenvectors $\begin{bmatrix} 2 & 1 \end{bmatrix}^T$ and $\begin{bmatrix} 5 & 3 \end{bmatrix}^T$. Find the $(2, 1)$ -entry of A .

Answer:

Example

Let A be a 2×2 matrix with eigenvalues 2 and 3, and with corresponding eigenvectors $\begin{bmatrix} 2 & 1 \end{bmatrix}^T$ and $\begin{bmatrix} 5 & 3 \end{bmatrix}^T$. Find the $(2, 1)$ -entry of A .

Answer: We have

$$A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}^{-1} = \begin{bmatrix} 4 & 15 \\ 2 & 9 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix}$$

Hence $(2, 1)$ -entry of A is $2 * 3 + 9 * (-1) = -3$.

Example

Let A be a 2×2 matrix with eigenvalues 2 and 3, and with corresponding eigenvectors $[2 \ 1]^T$ and $[5 \ 3]^T$. Find the $(2, 1)$ -entry of A .

Answer: We have

$$A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}^{-1} = \begin{bmatrix} 4 & 15 \\ 2 & 9 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix}$$

Hence $(2, 1)$ -entry of A is $2 * 3 + 9 * (-1) = -3$.

Example

Let $A = \begin{bmatrix} 1 & 1 & -1 \\ 0 & 0 & -1 \\ 0 & 2 & -3 \end{bmatrix}$. Given that $\lambda = -1$ is an eigenvalue of A . Which of the following are basic eigenvectors corresponding to λ ?

Answer:

$$(\lambda I_3 - A)x = 0 \Leftrightarrow \begin{bmatrix} -2 & -1 & 1 \\ 0 & -1 & 1 \\ 0 & -2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad (*)$$

Example

Let $A = \begin{bmatrix} 1 & 1 & -1 \\ 0 & 0 & -1 \\ 0 & 2 & -3 \end{bmatrix}$. Given that $\lambda = -1$ is an eigenvalue of A . Which of the following are basic eigenvectors corresponding to λ ?

Answer:

$$(\lambda I_3 - A)x = 0 \Leftrightarrow \begin{bmatrix} -2 & -1 & 1 \\ 0 & -1 & 1 \\ 0 & -2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad (*)$$

The system $(*)$ is equivalent to
$$\begin{cases} -2x_1 - x_2 + x_3 = 0 \\ -x_2 + x_3 = 0 \\ -2x_2 + 2x_3 = 0 \end{cases} \Leftrightarrow \begin{cases} x_1 = 0 \\ x_2 = t \\ x_3 = t \end{cases}.$$

Hence $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ is basic eigenvector corresponding to $\lambda = -1$.

Example

Let $A = \begin{bmatrix} 1 & 1 & -1 \\ 0 & 0 & -1 \\ 0 & 2 & -3 \end{bmatrix}$. Given that $\lambda = -1$ is an eigenvalue of A . Which of the following are basic eigenvectors corresponding to λ ?

Answer:

$$(\lambda I_3 - A)x = 0 \Leftrightarrow \begin{bmatrix} -2 & -1 & 1 \\ 0 & -1 & 1 \\ 0 & -2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad (*)$$

The system $(*)$ is equivalent to
$$\begin{cases} -2x_1 - x_2 + x_3 = 0 \\ -x_2 + x_3 = 0 \\ -2x_2 + 2x_3 = 0 \end{cases} \Leftrightarrow \begin{cases} x_1 = 0 \\ x_2 = t \\ x_3 = t \end{cases}.$$

Hence $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ is basic eigenvector corresponding to $\lambda = -1$.

Find the eigenvalues of the matrix

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 1 & 1 \\ 0 & 4 & -2 \end{bmatrix}.$$

Answer: The characteristic polynomial of A is

Find the eigenvalues of the matrix

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 1 & 1 \\ 0 & 4 & -2 \end{bmatrix}.$$

Answer: The characteristic polynomial of A is

$$\begin{aligned} c_A(x) &= \begin{vmatrix} x-2 & -1 & 1 \\ 0 & x-1 & -1 \\ 0 & -4 & x+2 \end{vmatrix} = (x-2) \begin{vmatrix} x-1 & -1 \\ -4 & x+2 \end{vmatrix} \\ &= (x-2)(x^2 + x - 6) = (x-2)^2(x+3). \end{aligned}$$

Find the eigenvalues of the matrix

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 1 & 1 \\ 0 & 4 & -2 \end{bmatrix}.$$

Answer: The characteristic polynomial of A is

$$\begin{aligned} c_A(x) &= \begin{vmatrix} x-2 & -1 & 1 \\ 0 & x-1 & -1 \\ 0 & -4 & x+2 \end{vmatrix} = (x-2) \begin{vmatrix} x-1 & -1 \\ -4 & x+2 \end{vmatrix} \\ &= (x-2)(x^2 + x - 6) = (x-2)^2(x+3). \end{aligned}$$

Hence, A has two eigenvalues $\lambda_1 = -3$ and $\lambda_2 = 2$.

Find the eigenvalues of the matrix

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 1 & 1 \\ 0 & 4 & -2 \end{bmatrix}.$$

Answer: The characteristic polynomial of A is

$$\begin{aligned} c_A(x) &= \begin{vmatrix} x-2 & -1 & 1 \\ 0 & x-1 & -1 \\ 0 & -4 & x+2 \end{vmatrix} = (x-2) \begin{vmatrix} x-1 & -1 \\ -4 & x+2 \end{vmatrix} \\ &= (x-2)(x^2 + x - 6) = (x-2)^2(x+3). \end{aligned}$$

Hence, A has two eigenvalues $\lambda_1 = -3$ and $\lambda_2 = 2$.

Exercises

Section 3.1 : $1bci, 5bd, 7, 9, 10, 16a$ (page 155 – 157)

Section 3.2 : $1ac, 2bdf, 3, 5, 6, 7, 9, 10, 20, 33$ (page 169 – 171)

Section 3.3 : $1ace, 8a, 12, 13, 14$ (page 189 – 190)