

Chapter 2. Matrix Algebra

1 Matrix Addition, Scalar Multiplication, Transposition

2 Multiplication

3 Matrix Inverse

4 Linear Transformations

2.1 Addition, Scalar Multiplication, Transpose

Definition

- An $m \times n$ matrix A is an array of m row and n column:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

2.1 Addition, Scalar Multiplication, Transpose

Definition

- An $m \times n$ matrix A is an array of m row and n column:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- The (i, j) -entry of A is the number lying simultaneously in row i and column j .

2.1 Addition, Scalar Multiplication, Transpose

Definition

- An $m \times n$ matrix A is an array of m row and n column:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- The (i,j) -entry of A is the number lying simultaneously in row i and column j .

Example

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{bmatrix}$$

2.1 Addition, Scalar Multiplication, Transpose

Definition

- An $m \times n$ matrix A is an array of m row and n column:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- The (i,j) -entry of A is the number lying simultaneously in row i and column j .

Example

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{bmatrix}$$

This is a 3×4 matrix.

2.1 Addition, Scalar Multiplication, Transpose

Definition

- An $m \times n$ matrix A is an array of m row and n column:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- The (i, j) -entry of A is the number lying simultaneously in row i and column j .

Example

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{bmatrix}$$

This is a 3×4 matrix.

The $(2, 3)$ -entry of A is 7.

2.1 Addition, Scalar Multiplication, Transpose

Definition

- An $m \times n$ matrix A is an array of m row and n column:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- The (i, j) -entry of A is the number lying simultaneously in row i and column j .

Example

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{bmatrix}$$

This is a 3×4 matrix.

The $(2, 3)$ -entry of A is 7.

Definition

- An $m \times m$ matrix is called a **square matrix** of size m .
- The zero matrix of size $m \times n$, denoted by $0_{m \times n}$ is the matrix that its all entries are 0.

Definition

- An $m \times m$ matrix is called a **square matrix** of size m .
- The zero matrix of size $m \times n$, denoted by $0_{m \times n}$ is the matrix that its all entries are 0.
- If $A = [a_{ij}]$ is an $m \times n$ matrix then $-A$ the **negative matrix** of A and defined by $-A = [-a_{ij}]$.

Definition

- An $m \times m$ matrix is called a **square matrix** of size m .
- The zero matrix of size $m \times n$, denoted by $0_{m \times n}$ is the matrix that its all entries are 0.
- If $A = [a_{ij}]$ is an $m \times n$ matrix then $-A$ the **negative matrix** of A and defined by $-A = [-a_{ij}]$.

Definition

An **identity matrix** I_n is a square matrix with 1s on the main diagonal and zeros elsewhere.

Definition

- An $m \times m$ matrix is called a **square matrix** of size m .
- The zero matrix of size $m \times n$, denoted by $0_{m \times n}$ is the matrix that its all entries are 0.
- If $A = [a_{ij}]$ is an $m \times n$ matrix then $-A$ the **negative matrix** of A and defined by $-A = [-a_{ij}]$.

Definition

An **identity matrix** I_n is a square matrix with 1s on the main diagonal and zeros elsewhere.

For example,

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \qquad I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Definition

- An $m \times m$ matrix is called a **square matrix** of size m .
- The zero matrix of size $m \times n$, denoted by $0_{m \times n}$ is the matrix that its all entries are 0.
- If $A = [a_{ij}]$ is an $m \times n$ matrix then $-A$ the **negative matrix** of A and defined by $-A = [-a_{ij}]$.

Definition

An **identity matrix** I_n is a square matrix with 1s on the main diagonal and zeros elsewhere.

For example,

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \qquad I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Let $A = [a_{ij}]$, $B = [b_{ij}]$ be two $m \times n$ matrices and k be a real numbers.

Definition

- **Addition** $A + B = [a_{ij} + b_{ij}]_{m \times n}$.

Let $A = [a_{ij}]$, $B = [b_{ij}]$ be two $m \times n$ matrices and k be a real numbers.

Definition

- **Addition** $A + B = [a_{ij} + b_{ij}]_{m \times n}$.
- **Scalar multiplication** $kA = [ka_{ij}]_{m \times n}$.

Let $A = [a_{ij}]$, $B = [b_{ij}]$ be two $m \times n$ matrices and k be a real numbers.

Definition

- **Addition** $A + B = [a_{ij} + b_{ij}]_{m \times n}$.
- **Scalar multiplication** $kA = [ka_{ij}]_{m \times n}$.

Example

$$2 \begin{bmatrix} -2 & 3 & 5 \\ 0 & 4 & 2 \\ 1 & -1 & 6 \end{bmatrix} - \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 4 & -1 & 3 \end{bmatrix} = \begin{bmatrix} -5 & 6 & 12 \\ 2 & 7 & 3 \\ -2 & -1 & 9 \end{bmatrix}$$

Let $A = [a_{ij}]$, $B = [b_{ij}]$ be two $m \times n$ matrices and k be a real numbers.

Definition

- **Addition** $A + B = [a_{ij} + b_{ij}]_{m \times n}$.
- **Scalar multiplication** $kA = [ka_{ij}]_{m \times n}$.

Example

$$2 \begin{bmatrix} -2 & 3 & 5 \\ 0 & 4 & 2 \\ 1 & -1 & 6 \end{bmatrix} - \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 4 & -1 & 3 \end{bmatrix} = \begin{bmatrix} -5 & 6 & 12 \\ 2 & 7 & 3 \\ -2 & -1 & 9 \end{bmatrix}$$

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$
- $A + (-A) = 0$ (zero matrix of size $m \times n$)

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$
- $A + (-A) = 0$ (zero matrix of size $m \times n$)
- $(A + B) + C = A + (B + C)$

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$
- $A + (-A) = 0$ (zero matrix of size $m \times n$)
- $(A + B) + C = A + (B + C)$
- $k(hA) = (kh)A$

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$
- $A + (-A) = 0$ (zero matrix of size $m \times n$)
- $(A + B) + C = A + (B + C)$
- $k(hA) = (kh)A$
- $(k + h)A = kA + hA$

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$
- $A + (-A) = 0$ (zero matrix of size $m \times n$)
- $(A + B) + C = A + (B + C)$
- $k(hA) = (kh)A$
- $(k + h)A = kA + hA$
- $k(A + B) = kA + kB$

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$
- $A + (-A) = 0$ (zero matrix of size $m \times n$)
- $(A + B) + C = A + (B + C)$
- $k(hA) = (kh)A$
- $(k + h)A = kA + hA$
- $k(A + B) = kA + kB$

Properties

For all $m \times n$ matrices A, B, C and all $k, h \in \mathbb{R}$,

- $A + B = B + A$
- $A + (-A) = 0$ (zero matrix of size $m \times n$)
- $(A + B) + C = A + (B + C)$
- $k(hA) = (kh)A$
- $(k + h)A = kA + hA$
- $k(A + B) = kA + kB$

Definition

Let A be $m \times n$ matrix. The **transpose** of A , written $A^T = [a_{ji}]$ is an $n \times m$ matrix.

Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Definition

Let A be $m \times n$ matrix. The **transpose** of A , written $A^T = [a_{ji}]$ is an $n \times m$ matrix.

Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Properties:

Definition

Let A be $m \times n$ matrix. The **transpose** of A , written $A^T = [a_{ji}]$ is an $n \times m$ matrix.

Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Properties:

- $(A^T)^T = A$

Definition

Let A be $m \times n$ matrix. The **transpose** of A , written $A^T = [a_{ji}]$ is an $n \times m$ matrix.

Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Properties:

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$

Definition

Let A be $m \times n$ matrix. The **transpose** of A , written $A^T = [a_{ji}]$ is an $n \times m$ matrix.

Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Properties:

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(kA)^T = kA^T$

Definition

Let A be $m \times n$ matrix. The **transpose** of A , written $A^T = [a_{ji}]$ is an $n \times m$ matrix.

Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Properties:

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(kA)^T = kA^T$

Definition

Let A be $m \times n$ matrix. The **transpose** of A , written $A^T = [a_{ji}]$ is an $n \times m$ matrix.

Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Properties:

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(kA)^T = kA^T$

Example

Find the $(2, 1)$ -entry of the matrix A that satisfies

$$\left(2A^T - 3 \begin{bmatrix} 1 & 0 \\ -2 & 2 \end{bmatrix}\right)^T = 4A - 7 \begin{bmatrix} 1 & 2 \\ -2 & 0 \end{bmatrix}$$

Answer:

Example

Find the $(2, 1)$ -entry of the matrix A that satisfies

$$\left(2A^T - 3 \begin{bmatrix} 1 & 0 \\ -2 & 2 \end{bmatrix}\right)^T = 4A - 7 \begin{bmatrix} 1 & 2 \\ -2 & 0 \end{bmatrix}$$

Answer: We have

$$2A - \begin{bmatrix} 3 & -6 \\ 0 & 6 \end{bmatrix} = 4A - \begin{bmatrix} 7 & 14 \\ -14 & 0 \end{bmatrix}$$

Example

Find the $(2, 1)$ -entry of the matrix A that satisfies

$$\left(2A^T - 3 \begin{bmatrix} 1 & 0 \\ -2 & 2 \end{bmatrix}\right)^T = 4A - 7 \begin{bmatrix} 1 & 2 \\ -2 & 0 \end{bmatrix}$$

Answer: We have

$$2A - \begin{bmatrix} 3 & -6 \\ 0 & 6 \end{bmatrix} = 4A - \begin{bmatrix} 7 & 14 \\ -14 & 0 \end{bmatrix}$$

Hence

$$2A = \begin{bmatrix} 4 & 20 \\ -14 & -6 \end{bmatrix}$$

Example

Find the $(2, 1)$ -entry of the matrix A that satisfies

$$\left(2A^T - 3 \begin{bmatrix} 1 & 0 \\ -2 & 2 \end{bmatrix}\right)^T = 4A - 7 \begin{bmatrix} 1 & 2 \\ -2 & 0 \end{bmatrix}$$

Answer: We have

$$2A - \begin{bmatrix} 3 & -6 \\ 0 & 6 \end{bmatrix} = 4A - \begin{bmatrix} 7 & 14 \\ -14 & 0 \end{bmatrix}$$

Hence

$$2A = \begin{bmatrix} 4 & 20 \\ -14 & -6 \end{bmatrix}$$

Thus, $(2, 1)$ -entry of A is $-14/2 = -7$.

Example

Find the $(2, 1)$ -entry of the matrix A that satisfies

$$\left(2A^T - 3 \begin{bmatrix} 1 & 0 \\ -2 & 2 \end{bmatrix}\right)^T = 4A - 7 \begin{bmatrix} 1 & 2 \\ -2 & 0 \end{bmatrix}$$

Answer: We have

$$2A - \begin{bmatrix} 3 & -6 \\ 0 & 6 \end{bmatrix} = 4A - \begin{bmatrix} 7 & 14 \\ -14 & 0 \end{bmatrix}$$

Hence

$$2A = \begin{bmatrix} 4 & 20 \\ -14 & -6 \end{bmatrix}$$

Thus, $(2, 1)$ -entry of A is $-14/2 = -7$.

Symmetric matrix

Definition

A matrix A is **symmetric** if $A^T = A$ (it means that (i, j) -entry of $A = (j, i)$ -entry of A).

Example

$\begin{bmatrix} 2 & 3 & 1 \\ 3 & -1 & 5 \\ 1 & 5 & 6 \end{bmatrix}$ is symmetric.

Symmetric matrix

Definition

A matrix A is **symmetric** if $A^T = A$ (it means that (i, j) -entry of $A = (j, i)$ -entry of A).

Example

$\begin{bmatrix} 2 & 3 & 1 \\ 3 & -1 & 5 \\ 1 & 5 & 6 \end{bmatrix}$ is symmetric.

$\begin{bmatrix} 1 & 2 \\ -2 & 4 \end{bmatrix}$ is not symmetric.

Properties

Symmetric matrix

Definition

A matrix A is **symmetric** if $A^T = A$ (it means that (i, j) -entry of $A = (j, i)$ -entry of A).

Example

$\begin{bmatrix} 2 & 3 & 1 \\ 3 & -1 & 5 \\ 1 & 5 & 6 \end{bmatrix}$ is symmetric.

$\begin{bmatrix} 1 & 2 \\ -2 & 4 \end{bmatrix}$ is not symmetric.

Properties

$A_{m \times m}$ and $B_{m \times m}$ are symmetric $\Rightarrow kA + hB$ is symmetric, for all $k, h \in \mathbb{R}$.

Symmetric matrix

Definition

A matrix A is **symmetric** if $A^T = A$ (it means that (i, j) -entry of $A = (j, i)$ -entry of A).

Example

$\begin{bmatrix} 2 & 3 & 1 \\ 3 & -1 & 5 \\ 1 & 5 & 6 \end{bmatrix}$ is symmetric.

$\begin{bmatrix} 1 & 2 \\ -2 & 4 \end{bmatrix}$ is not symmetric.

Properties

$A_{m \times m}$ and $B_{m \times m}$ are symmetric $\Rightarrow kA + hB$ is symmetric, for all $k, h \in \mathbb{R}$.

Example

Determine a and b such that the following matrix is symmetric.

$$\begin{bmatrix} -3 & a+b & 8 \\ 7 & 5 & -1 \\ 8 & a-b & ab \end{bmatrix}$$

Answer:

Example

Determine a and b such that the following matrix is symmetric.

$$\begin{bmatrix} -3 & a+b & 8 \\ 7 & 5 & -1 \\ 8 & a-b & ab \end{bmatrix}$$

Answer: Find a, b such that

$$\begin{cases} a+b=7 \\ a-b=-1 \end{cases} \Rightarrow a=3, b=4$$

Example

Determine a and b such that the following matrix is symmetric.

$$\begin{bmatrix} -3 & a+b & 8 \\ 7 & 5 & -1 \\ 8 & a-b & ab \end{bmatrix}$$

Answer: Find a, b such that

$$\begin{cases} a+b=7 \\ a-b=-1 \end{cases} \Rightarrow a=3, b=4$$

2.3 Multiplication

Definition (Dot product)

$$\begin{bmatrix} a_1 & a_2 & \cdots & a_m \end{bmatrix} \times \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = a_1 b_1 + a_2 b_2 + \cdots + a_m b_m$$

Definition

Let $A_{m \times n}, B_{n \times p}$ be two matrices. The multiplication of A and B

$$AB = [c_{ij}]_{m \times p},$$

2.3 Multiplication

Definition (Dot product)

$$\begin{bmatrix} a_1 & a_2 & \cdots & a_m \end{bmatrix} \times \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = a_1 b_1 + a_2 b_2 + \cdots + a_m b_m$$

Definition

Let $A_{m \times n}, B_{n \times p}$ be two matrices. The multiplication of A and B

$$AB = [c_{ij}]_{m \times p},$$

where c_{ij} is the dot product of i^{th} row of A and j^{th} column of B .

2.3 Multiplication

Definition (Dot product)

$$\begin{bmatrix} a_1 & a_2 & \cdots & a_m \end{bmatrix} \times \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = a_1 b_1 + a_2 b_2 + \cdots + a_m b_m$$

Definition

Let $A_{m \times n}$, $B_{n \times p}$ be two matrices. The multiplication of A and B

$$AB = [c_{ij}]_{m \times p},$$

where c_{ij} is the dot product of i^{th} row of A and j^{th} column of B .

2.3 Multiplication

Definition (Dot product)

$$\begin{bmatrix} a_1 & a_2 & \cdots & a_m \end{bmatrix} \times \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = a_1 b_1 + a_2 b_2 + \cdots + a_m b_m$$

Definition

Let $A_{m \times n}$, $B_{n \times p}$ be two matrices. The multiplication of A and B

$$AB = [c_{ij}]_{m \times p},$$

where c_{ij} is the dot product of i^{th} row of A and j^{th} column of B .

Example

$$\begin{bmatrix} 1 & 2 & -3 \\ -2 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 & 5 \\ -1 & 1 & 4 & 6 \\ 2 & 4 & 0 & -1 \end{bmatrix} = \begin{bmatrix} -7 & -10 & 7 & 20 \\ -7 & -1 & 14 & 9 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 3 & 2 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -3 & 2 \\ 0 & -1 & 3 \end{bmatrix}$.

Find $(2, 1)$ -entry of $AB - B^T A$.

Example

$$\begin{bmatrix} 1 & 2 & -3 \\ -2 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 & 5 \\ -1 & 1 & 4 & 6 \\ 2 & 4 & 0 & -1 \end{bmatrix} = \begin{bmatrix} -7 & -10 & 7 & 20 \\ -7 & -1 & 14 & 9 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 3 & 2 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -3 & 2 \\ 0 & -1 & 3 \end{bmatrix}$.

Find $(2, 1)$ -entry of $AB - B^T A$.

Answer:

Example

$$\begin{bmatrix} 1 & 2 & -3 \\ -2 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 & 5 \\ -1 & 1 & 4 & 6 \\ 2 & 4 & 0 & -1 \end{bmatrix} = \begin{bmatrix} -7 & -10 & 7 & 20 \\ -7 & -1 & 14 & 9 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 3 & 2 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -3 & 2 \\ 0 & -1 & 3 \end{bmatrix}$.

Find $(2, 1)$ -entry of $AB - B^T A$.

Answer:

- The $(2, 1)$ -entry of AB is the dot product of the second row of A and the first column of B and equal to $-2 * 2 + 1 * 4 + 1 * 0 = 0$.

Example

$$\begin{bmatrix} 1 & 2 & -3 \\ -2 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 & 5 \\ -1 & 1 & 4 & 6 \\ 2 & 4 & 0 & -1 \end{bmatrix} = \begin{bmatrix} -7 & -10 & 7 & 20 \\ -7 & -1 & 14 & 9 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 3 & 2 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -3 & 2 \\ 0 & -1 & 3 \end{bmatrix}$.

Find $(2, 1)$ -entry of $AB - B^T A$.

Answer:

- The $(2, 1)$ -entry of AB is the dot product of the second row of A and the first column of B and equal to $-2 * 2 + 1 * 4 + 1 * 0 = 0$.
- The $(2, 1)$ -entry of $B^T A$ is the dot product of the second column of B and the first column of A , is $1 * 1 + (-3) * (-2) + (-1) * 3 = 4$.

Example

$$\begin{bmatrix} 1 & 2 & -3 \\ -2 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 & 5 \\ -1 & 1 & 4 & 6 \\ 2 & 4 & 0 & -1 \end{bmatrix} = \begin{bmatrix} -7 & -10 & 7 & 20 \\ -7 & -1 & 14 & 9 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 3 & 2 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -3 & 2 \\ 0 & -1 & 3 \end{bmatrix}$.

Find $(2, 1)$ -entry of $AB - B^T A$.

Answer:

- The $(2, 1)$ -entry of AB is the dot product of the second row of A and the first column of B and equal to $-2 * 2 + 1 * 4 + 1 * 0 = 0$.
- The $(2, 1)$ -entry of $B^T A$ is the dot product of the second column of B and the first column of A , is $1 * 1 + (-3) * (-2) + (-1) * 3 = 4$.

Hence $(2, 1)$ -entry of $AB - B^T A$ is $0 - 4 = -4$.

Example

$$\begin{bmatrix} 1 & 2 & -3 \\ -2 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 & 5 \\ -1 & 1 & 4 & 6 \\ 2 & 4 & 0 & -1 \end{bmatrix} = \begin{bmatrix} -7 & -10 & 7 & 20 \\ -7 & -1 & 14 & 9 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & 0 & -2 \\ -2 & 1 & 1 \\ 3 & 2 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -3 & 2 \\ 0 & -1 & 3 \end{bmatrix}$.

Find $(2, 1)$ -entry of $AB - B^T A$.

Answer:

- The $(2, 1)$ -entry of AB is the dot product of the second row of A and the first column of B and equal to $-2 * 2 + 1 * 4 + 1 * 0 = 0$.
- The $(2, 1)$ -entry of $B^T A$ is the dot product of the second column of B and the first column of A , is $1 * 1 + (-3) * (-2) + (-1) * 3 = 4$.

Hence $(2, 1)$ -entry of $AB - B^T A$ is $0 - 4 = -4$.

Example

Find the second entry of the first row of the matrix A that satisfies

$$A^T - \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$$

Answer:

Example

Find the second entry of the first row of the matrix A that satisfies

$$A^T - \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$$

Answer: We have

$$A^T = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -1 & -2 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} = \begin{bmatrix} 2 & 5 \\ 1 & 2 \\ 5 & 10 \end{bmatrix}$$

Example

Find the second entry of the first row of the matrix A that satisfies

$$A^T - \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$$

Answer: We have

$$A^T = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -1 & -2 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} = \begin{bmatrix} 2 & 5 \\ 1 & 2 \\ 5 & 10 \end{bmatrix}$$

Hence the second entry of the first row of the matrix A is 1.

Example

Find the second entry of the first row of the matrix A that satisfies

$$A^T - \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$$

Answer: We have

$$A^T = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -1 & -2 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} = \begin{bmatrix} 2 & 5 \\ 1 & 2 \\ 5 & 10 \end{bmatrix}$$

Hence the second entry of the first row of the matrix A is 1.

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$
- $AI_m = I_m A = A$, for $A_{m \times m}$

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$
- $AI_m = I_m A = A$, for $A_{m \times m}$
- $(AB)^T = B^T A^T$

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$
- $AI_m = I_m A = A$, for $A_{m \times m}$
- $(AB)^T = B^T A^T$
- In general, $AB \neq BA$

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$
- $AI_m = I_m A = A$, for $A_{m \times m}$
- $(AB)^T = B^T A^T$
- In general, $AB \neq BA$

Example

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix} \neq \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} -1 & -2 \\ 3 & 5 \end{bmatrix}$$

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$
- $AI_m = I_m A = A$, for $A_{m \times m}$
- $(AB)^T = B^T A^T$
- In general, $AB \neq BA$

Example

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix} \neq \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} -1 & -2 \\ 3 & 5 \end{bmatrix}$$

Definition

We say A and B are **commute** if $AB = BA$.

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$
- $AI_m = I_m A = A$, for $A_{m \times m}$
- $(AB)^T = B^T A^T$
- In general, $AB \neq BA$

Example

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix} \neq \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} -1 & -2 \\ 3 & 5 \end{bmatrix}$$

Definition

We say A and B are **commute** if $AB = BA$.

Properties

- $A(BC) = (AB)C$, for $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$
- $A(B + C) = AB + AC$, for $A_{m \times n}$, $B_{n \times p}$, $C_{n \times p}$
- $(A + B)C = AC + BC$, for $A_{m \times n}$, $B_{m \times n}$, $C_{n \times p}$
- $AI_m = I_m A = A$, for $A_{m \times m}$
- $(AB)^T = B^T A^T$
- In general, $AB \neq BA$

Example

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix} \neq \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} -1 & -2 \\ 3 & 5 \end{bmatrix}$$

Definition

We say A and B are **commute** if $AB = BA$.

Quiz

Decide the following statements are True or False.

(a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix).

Quiz

Decide the following statements are True or False.

(a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix).

Quiz

Decide the following statements are True or False.

(a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F

(b) $(A + B)(A - B) = A^2 - B^2$.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric. T
- (h) If A, B are symmetric matrices of the same size, then AB is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric. T
- (h) If A, B are symmetric matrices of the same size, then AB is symmetric.

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric. T
- (h) If A, B are symmetric matrices of the same size, then AB is symmetric. F
- (i) Let A is 3×5 matrix and B is 6×7 matrix, and ACB can be performed. What is the size of C ?

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric. T
- (h) If A, B are symmetric matrices of the same size, then AB is symmetric. F
- (i) Let A is 3×5 matrix and B is 6×7 matrix, and ACB can be performed. What is the size of C ?

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric. T
- (h) If A, B are symmetric matrices of the same size, then AB is symmetric. F
- (i) Let A is 3×5 matrix and B is 6×7 matrix, and ACB can be performed. What is the size of C ? 5×6

Quiz

Decide the following statements are True or False.

- (a) If $AB = 0$ then either $A = 0$ or $B = 0$, (where 0 is zero matrix). F
- (b) $(A + B)(A - B) = A^2 - B^2$. F
- (c) If A^2 can be formed then A must be square matrix. T
- (d) If A is symmetric and square, then A^2 is symmetric. T
- (e) If A is square then AA^T is symmetric. T
- (f) If AB and BA can be formed then A and B must be square matrices. F
- (g) If A, B are symmetric matrices of the same size, then $A + B$ is symmetric. T
- (h) If A, B are symmetric matrices of the same size, then AB is symmetric. F
- (i) Let A is 3×5 matrix and B is 6×7 matrix, and ACB can be performed. What is the size of C ? 5×6

Multiplication matrix with system

A system of m linear equations and n variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

can be rewritten in the matrix form

$$AX = b,$$

Multiplication matrix with system

A system of m linear equations and n variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \dots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

can be rewritten in the matrix form

$$AX = b,$$

where $X = \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix}$ is the variables,

Multiplication matrix with system

A system of m linear equations and n variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

can be rewritten in the matrix form

$$AX = b,$$

where $X = \begin{bmatrix} x_1 \\ x_2 \\ \cdots \\ x_n \end{bmatrix}$ is the variables, $b = \begin{bmatrix} b_1 \\ b_2 \\ \cdots \\ b_n \end{bmatrix}$ and

Multiplication matrix with system

A system of m linear equations and n variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

can be rewritten in the matrix form

$$AX = b,$$

where $X = \begin{bmatrix} x_1 \\ x_2 \\ \cdots \\ x_n \end{bmatrix}$ is the variables, $b = \begin{bmatrix} b_1 \\ b_2 \\ \cdots \\ b_n \end{bmatrix}$ and A is the coefficient matrix.

Multiplication matrix with system

A system of m linear equations and n variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \dots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

can be rewritten in the matrix form

$$AX = b,$$

where $X = \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix}$ is the variables, $b = \begin{bmatrix} b_1 \\ b_2 \\ \dots \\ b_n \end{bmatrix}$ and A is the coefficient matrix.

Example

The system of linear equations

$$\begin{array}{rrcrcl} 2x & + & 3y & - & 5z & = & 9 \\ -x & - & 2y & + & 4z & = & 11 \end{array}$$

can be written in the matrix form

$$\begin{bmatrix} 2 & 3 & -5 \\ -1 & -2 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 11 \end{bmatrix}$$

Example

The system of linear equations

$$\begin{array}{rrcrcl} 2x & + & 3y & - & 5z & = & 9 \\ -x & - & 2y & + & 4z & = & 11 \end{array}$$

can be written in the matrix form

$$\begin{bmatrix} 2 & 3 & -5 \\ -1 & -2 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 11 \end{bmatrix}$$

2.4 Matrix Inverse

Let A be a square $n \times n$ matrix.

Definition

A matrix B is called an **inverse** of A if and only if

2.4 Matrix Inverse

Let A be a square $n \times n$ matrix.

Definition

A matrix B is called an **inverse** of A if and only if

$$AB = I \text{ and } BA = I$$

2.4 Matrix Inverse

Let A be a square $n \times n$ matrix.

Definition

A matrix B is called an **inverse** of A if and only if

$$AB = I \text{ and } BA = I$$

A matrix A that has an inverse is called an **invertible** matrix

2.4 Matrix Inverse

Let A be a square $n \times n$ matrix.

Definition

A matrix B is called an **inverse** of A if and only if

$$AB = I \text{ and } BA = I$$

A matrix A that has an inverse is called an **invertible** matrix

Example

Find the inverse matrix of $A = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$.

Answer:

Example

Find the inverse matrix of $A = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

Example

Find the inverse matrix of $A = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} 2x + z & 2y + t \\ z & t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

This is equivalent to

Example

Find the inverse matrix of $A = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} 2x + z & 2y + t \\ z & t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

This is equivalent to

$$\begin{array}{rcl} 2x + z & = & 1 \\ 2y + t & = & 0 \\ z & = & 0 \\ t & = & 1 \end{array} \Leftrightarrow \begin{array}{rcl} x & = & 1/2 \\ y & = & -1/2 \\ z & = & 0 \\ t & = & 1 \end{array}.$$

Example

Find the inverse matrix of $A = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} 2x + z & 2y + t \\ z & t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

This is equivalent to

$$\begin{array}{rclcl} 2x + z & = & 1 & x & = 1/2 \\ 2y + t & = & 0 & y & = -1/2 \\ z & = & 0 & z & = 0 \\ t & = & 1 & t & = 1 \end{array} \Leftrightarrow$$

$$\text{Hence } A^{-1} = \begin{bmatrix} 1/2 & -1/2 \\ 0 & 1 \end{bmatrix}.$$

Example

Find the inverse matrix of $A = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} 2x + z & 2y + t \\ z & t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

This is equivalent to

$$\begin{array}{rclcl} 2x + z & = & 1 & x & = 1/2 \\ 2y + t & = & 0 & y & = -1/2 \\ z & = & 0 & z & = 0 \\ t & = & 1 & t & = 1 \end{array} \Leftrightarrow$$

$$\text{Hence } A^{-1} = \begin{bmatrix} 1/2 & -1/2 \\ 0 & 1 \end{bmatrix}.$$

Example

Show that $A = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix}$ is not invertible.

Answer:

Example

Show that $A = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix}$ is not invertible.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$.

Example

Show that $A = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix}$ is not invertible.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} x - 2z & y - 2t \\ -x + 2z & -y + 2t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Example

Show that $A = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix}$ is not invertible.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} x - 2z & y - 2t \\ -x + 2z & -y + 2t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

This is equivalent to

$$\begin{aligned} x - 2z &= 1 \\ y - 2t &= 0 \\ -x + 2z &= 0 \\ -y + 2t &= 1 \end{aligned}$$

Example

Show that $A = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix}$ is not invertible.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} x - 2z & y - 2t \\ -x + 2z & -y + 2t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

This is equivalent to

$$\begin{aligned} x - 2z &= 1 \\ y - 2t &= 0 \\ -x + 2z &= 0 \\ -y + 2t &= 1 \end{aligned}$$

This system has no solution. Hence, A is not invertible.

Example

Show that $A = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix}$ is not invertible.

Answer: Say $B = \begin{bmatrix} x & y \\ z & t \end{bmatrix}$. We have

$$AB = \begin{bmatrix} 1 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} x - 2z & y - 2t \\ -x + 2z & -y + 2t \end{bmatrix} = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

This is equivalent to

$$\begin{aligned} x - 2z &= 1 \\ y - 2t &= 0 \\ -x + 2z &= 0 \\ -y + 2t &= 1 \end{aligned}$$

This system has no solution. Hence, A is not invertible.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then A is invertible if $\det(A) = ad - bc \neq 0$ and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Example

$$\begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}^{-1} = \frac{1}{10} \begin{bmatrix} 4 & -1 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0.4 & -0.1 \\ -0.2 & 0.3 \end{bmatrix}$$

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then A is invertible if $\det(A) = ad - bc \neq 0$ and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Example

$$\begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}^{-1} = \frac{1}{10} \begin{bmatrix} 4 & -1 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0.4 & -0.1 \\ -0.2 & 0.3 \end{bmatrix}$$

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1},$

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1},$
- $(A^{-1})^{-1} = A,$

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1}$,
- $(A^{-1})^{-1} = A$,
- $I^{-1} = I$.

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1}$,
- $(A^{-1})^{-1} = A$,
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$,

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1}$,
- $(A^{-1})^{-1} = A$,
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$,
- $(aA)^{-1} = 1/aA^{-1}$, for any $a \neq 0$.

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1}$,
- $(A^{-1})^{-1} = A$,
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$,
- $(aA)^{-1} = 1/aA^{-1}$, for any $a \neq 0$.
- $(A^T)^{-1} = (A^{-1})^T$.

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1}$,
- $(A^{-1})^{-1} = A$,
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$,
- $(aA)^{-1} = 1/aA^{-1}$, for any $a \neq 0$.
- $(A^T)^{-1} = (A^{-1})^T$.

Properties

For any invertible matrices A, B of the same size:

- $(AB)^{-1} = B^{-1}A^{-1}$,
- $(A^{-1})^{-1} = A$,
- $I^{-1} = I$.
- $[A^k]^{-1} = (A^{-1})^k$,
- $(aA)^{-1} = 1/aA^{-1}$, for any $a \neq 0$.
- $(A^T)^{-1} = (A^{-1})^T$.

Find $(2, 1)$ -entry of the matrix A such that

$$(2A^{-1} + I)^{-1} = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}.$$

Answer:

Find (2, 1)-entry of the matrix A such that

$$(2A^{-1} + I)^{-1} = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}.$$

Answer: We have

$$2A^{-1} + I = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}^{-1} = \frac{1}{3 * 7 - 4 * 5} \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix}.$$

Hence

Find (2, 1)-entry of the matrix A such that

$$(2A^{-1} + I)^{-1} = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}.$$

Answer: We have

$$2A^{-1} + I = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}^{-1} = \frac{1}{3 * 7 - 4 * 5} \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix}.$$

Hence

$$2A^{-1} = \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 6 & -4 \\ -5 & 2 \end{bmatrix}.$$

$$\Rightarrow A = \begin{bmatrix} 3 & -2 \\ -5/2 & 1 \end{bmatrix}^{-1} = \frac{1}{3 * 1 - (-2) * (-5/2)} \begin{bmatrix} 1 & 2 \\ 5/2 & 3 \end{bmatrix} = \begin{bmatrix} -1/2 & -1 \\ -5/4 & -3/2 \end{bmatrix}$$

Find $(2, 1)$ -entry of the matrix A such that

$$(2A^{-1} + I)^{-1} = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}.$$

Answer: We have

$$2A^{-1} + I = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}^{-1} = \frac{1}{3 * 7 - 4 * 5} \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix}.$$

Hence

$$2A^{-1} = \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 6 & -4 \\ -5 & 2 \end{bmatrix}.$$

$$\Rightarrow A = \begin{bmatrix} 3 & -2 \\ -5/2 & 1 \end{bmatrix}^{-1} = \frac{1}{3 * 1 - (-2) * (-5/2)} \begin{bmatrix} 1 & 2 \\ 5/2 & 3 \end{bmatrix} = \begin{bmatrix} -1/2 & -1 \\ -5/4 & -3/2 \end{bmatrix}$$

Thus, the $(2, 1)$ -entry of A is $-5/4$.

Find $(2, 1)$ -entry of the matrix A such that

$$(2A^{-1} + I)^{-1} = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}.$$

Answer: We have

$$2A^{-1} + I = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}^{-1} = \frac{1}{3 * 7 - 4 * 5} \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix}.$$

Hence

$$2A^{-1} = \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 6 & -4 \\ -5 & 2 \end{bmatrix}.$$

$$\Rightarrow A = \begin{bmatrix} 3 & -2 \\ -5/2 & 1 \end{bmatrix}^{-1} = \frac{1}{3 * 1 - (-2) * (-5/2)} \begin{bmatrix} 1 & 2 \\ 5/2 & 3 \end{bmatrix} = \begin{bmatrix} -1/2 & -1 \\ -5/4 & -3/2 \end{bmatrix}$$

Thus, the $(2, 1)$ -entry of A is $-5/4$.

Find the first entry of the second row of the matrix A that satisfies:

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A \right)^{-1} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Answer:

Find the first entry of the second row of the matrix A that satisfies:

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A \right)^{-1} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Answer: We have

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}^{-1} = \frac{1}{1 * (-1) - 0 * 2} \begin{bmatrix} -1 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Find the first entry of the second row of the matrix A that satisfies:

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A \right)^{-1} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Answer: We have

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}^{-1} = \frac{1}{1 * (-1) - 0 * 2} \begin{bmatrix} -1 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Hence,

Find the first entry of the second row of the matrix A that satisfies:

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A \right)^{-1} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Answer: We have

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}^{-1} = \frac{1}{1 * (-1) - 0 * 2} \begin{bmatrix} -1 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Hence,

$$\begin{aligned} A &= \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} = \frac{1}{-1} \begin{bmatrix} 0 & -1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \end{aligned}$$

Find the first entry of the second row of the matrix A that satisfies:

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A \right)^{-1} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Answer: We have

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}^{-1} = \frac{1}{1 * (-1) - 0 * 2} \begin{bmatrix} -1 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Hence,

$$\begin{aligned} A &= \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} = \frac{1}{-1} \begin{bmatrix} 0 & -1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \end{aligned}$$

The first entry of the second row of the matrix A is $1 * 1 + (-1) * 0 = 1$.

Find the first entry of the second row of the matrix A that satisfies:

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A \right)^{-1} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Answer: We have

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}^{-1} = \frac{1}{1 * (-1) - 0 * 2} \begin{bmatrix} -1 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Hence,

$$\begin{aligned} A &= \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} = \frac{1}{-1} \begin{bmatrix} 0 & -1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \end{aligned}$$

The first entry of the second row of the matrix A is $1 * 1 + (-1) * 0 = 1$.

Matrix inverse algorithm

If A is an invertible matrix, there exists a sequence of elementary row operations that

$$[A|I] \rightarrow [I|A^{-1}]$$

where the row operations on A and I are carried out simultaneously.

Matrix inverse algorithm

If A is an invertible matrix, there exists a sequence of elementary row operations that

$$[A|I] \rightarrow [I|A^{-1}]$$

where the row operations on A and I are carried out simultaneously.

Example

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 0 & 1 & 0 & 0 \\ 0 & 2 & 3 & 0 & 1 & 0 \\ 1 & 3 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{-R_1+R_3} \left[\begin{array}{ccc|ccc} 1 & 2 & 0 & 1 & 0 & 0 \\ 0 & 2 & 3 & 0 & 1 & 0 \\ 0 & 1 & 1 & -1 & 0 & 1 \end{array} \right]$$

$$\xrightarrow[\begin{array}{c} 1/2R_2 \\ -1/2R_2+R_3 \end{array}]{\begin{array}{c} 1/2R_2 \\ -1/2R_2+R_3 \end{array}} \left[\begin{array}{ccc|ccc} 1 & 2 & 0 & 1 & 0 & 0 \\ 0 & 1 & 3/2 & 0 & 1/2 & 0 \\ 0 & 0 & -1/2 & -1 & -1/2 & 1 \end{array} \right]$$

$$\xrightarrow[\begin{array}{c} -2R_3 \\ 3R_3+R_2 \end{array}]{\begin{array}{c} -2R_3 \\ 3R_3+R_2 \end{array}} \left[\begin{array}{ccc|ccc} 1 & 2 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & -3 & -1 & 3 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right]$$

$$\xrightarrow{-2R_2+R_1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 7 & 2 & -6 \\ 0 & 1 & 0 & -3 & -1 & 3 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right] \Rightarrow \left[\begin{array}{ccc} 1 & 2 & 0 \\ 0 & 2 & 3 \\ 1 & 3 & 1 \end{array} \right]^{-1} = \left[\begin{array}{ccc} 7 & 2 & -6 \\ -3 & -1 & 3 \\ 2 & 1 & -2 \end{array} \right]$$

2.6 Linear transformation

Definition

Let A be $m \times n$ matrix. The transformation $T_A : \mathbb{R}^n \times \mathbb{R}^m$ defined by

$$T_A(x) = Ax$$

is called the **matrix transformation induced by A** .

Example

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \end{bmatrix}$.

2.6 Linear transformation

Definition

Let A be $m \times n$ matrix. The transformation $T_A : \mathbb{R}^n \times \mathbb{R}^m$ defined by

$$T_A(x) = Ax$$

is called the **matrix transformation induced by A** .

Example

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \end{bmatrix}$. Then the matrix transformation induced by A is

$T_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T_A \left(\begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x + 2y + 3z \\ 2x - y + 4z \end{bmatrix}.$$

2.6 Linear transformation

Definition

Let A be $m \times n$ matrix. The transformation $T_A : \mathbb{R}^n \times \mathbb{R}^m$ defined by

$$T_A(x) = Ax$$

is called the **matrix transformation induced by A** .

Example

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \end{bmatrix}$. Then the matrix transformation induced by A is

$T_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T_A \left(\begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x + 2y + 3z \\ 2x - y + 4z \end{bmatrix}.$$

Example

Consider a transformation

$T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ given by

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x - 3y \\ 5y \\ -x + 6y \end{bmatrix}.$$

Then T is a matrix transformation induced by $A = \begin{bmatrix} 1 & -3 \\ 0 & 5 \\ -1 & 6 \end{bmatrix}$.

Example

Consider a transformation

$T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ given by

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x - 3y \\ 5y \\ -x + 6y \end{bmatrix}.$$

Then T is a matrix transformation induced by $A = \begin{bmatrix} 1 & -3 \\ 0 & 5 \\ -1 & 6 \end{bmatrix}$.

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a **linear transformation** if it satisfies the following two conditions for all vectors x and y in \mathbb{R}^n and all scalars $a \in \mathbb{R}$:

(1) $T(x + y) = T(x) + T(y)$

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a **linear transformation** if it satisfies the following two conditions for all vectors x and y in \mathbb{R}^n and all scalars $a \in \mathbb{R}$:

(1) $T(x + y) = T(x) + T(y)$

(2) $T(ax) = aT(x)$.

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a **linear transformation** if it satisfies the following two conditions for all vectors x and y in \mathbb{R}^n and all scalars $a \in \mathbb{R}$:

- (1) $T(x + y) = T(x) + T(y)$
- (2) $T(ax) = aT(x)$.

Note.

1. If T is a linear transformation, then $T(0) = 0$ (T preserves the zero vector).

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a **linear transformation** if it satisfies the following two conditions for all vectors x and y in \mathbb{R}^n and all scalars $a \in \mathbb{R}$:

- (1) $T(x + y) = T(x) + T(y)$
- (2) $T(ax) = aT(x)$.

Note.

1. If T is a linear transformation, then $T(0) = 0$ (T preserves the zero vector).
2. The conditions (1) and (2) are equivalent to

$$T(ax + by) = aT(x) + bT(y), \quad \text{for all } x, y \in \mathbb{R}^n; a, b \in \mathbb{R}.$$

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a **linear transformation** if it satisfies the following two conditions for all vectors x and y in \mathbb{R}^n and all scalars $a \in \mathbb{R}$:

- (1) $T(x + y) = T(x) + T(y)$
- (2) $T(ax) = aT(x)$.

Note.

1. If T is a linear transformation, then $T(0) = 0$ (T preserves the zero vector).
2. The conditions (1) and (2) are equivalent to

$$T(ax + by) = aT(x) + bT(y), \quad \text{for all } x, y \in \mathbb{R}^n; a, b \in \mathbb{R}.$$

Example

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(x, y) = (x, x + y)$ is a linear transformation. Indeed,

$$\begin{aligned}(1) \quad T\left(\begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} x' \\ y' \end{bmatrix}\right) &= T\left(\begin{bmatrix} x + x' \\ y + y' \end{bmatrix}\right) = \begin{bmatrix} x + x' \\ x + x' + y + y' \end{bmatrix} \\ &= \begin{bmatrix} x \\ x + y \end{bmatrix} + \begin{bmatrix} x' \\ x' + y' \end{bmatrix} = T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) + T\left(\begin{bmatrix} x' \\ y' \end{bmatrix}\right).\end{aligned}$$

$$(2) \quad T\left(a \begin{bmatrix} x \\ y \end{bmatrix}\right) = T\left(\begin{bmatrix} ax \\ ay \end{bmatrix}\right) = \begin{bmatrix} ax \\ ax + ay \end{bmatrix} = a \begin{bmatrix} x \\ x + y \end{bmatrix} = aT\left(\begin{bmatrix} x \\ y \end{bmatrix}\right).$$

Example

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ given by $T(x, y) = (x - y, 0, y^2)$ is not a linear transformation since

Example

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(x, y) = (x, x + y)$ is a linear transformation. Indeed,

$$\begin{aligned}(1) \quad T\left(\begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} x' \\ y' \end{bmatrix}\right) &= T\left(\begin{bmatrix} x + x' \\ y + y' \end{bmatrix}\right) = \begin{bmatrix} x + x' \\ x + x' + y + y' \end{bmatrix} \\ &= \begin{bmatrix} x \\ x + y \end{bmatrix} + \begin{bmatrix} x' \\ x' + y' \end{bmatrix} = T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) + T\left(\begin{bmatrix} x' \\ y' \end{bmatrix}\right).\end{aligned}$$

$$(2) \quad T\left(a \begin{bmatrix} x \\ y \end{bmatrix}\right) = T\left(\begin{bmatrix} ax \\ ay \end{bmatrix}\right) = \begin{bmatrix} ax \\ ax + ay \end{bmatrix} = a \begin{bmatrix} x \\ x + y \end{bmatrix} = aT\left(\begin{bmatrix} x \\ y \end{bmatrix}\right).$$

Example

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ given by $T(x, y) = (x - y, 0, y^2)$ is not a linear transformation since

$$T\left(2 \begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = T\left(\begin{bmatrix} 2 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \\ 4 \end{bmatrix} \neq 2T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = 2 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix}$$

Example

$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(x, y) = (x, x + y)$ is a linear transformation. Indeed,

$$\begin{aligned}(1) \quad T\left(\begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} x' \\ y' \end{bmatrix}\right) &= T\left(\begin{bmatrix} x + x' \\ y + y' \end{bmatrix}\right) = \begin{bmatrix} x + x' \\ x + x' + y + y' \end{bmatrix} \\ &= \begin{bmatrix} x \\ x + y \end{bmatrix} + \begin{bmatrix} x' \\ x' + y' \end{bmatrix} = T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) + T\left(\begin{bmatrix} x' \\ y' \end{bmatrix}\right).\end{aligned}$$

$$(2) \quad T\left(a \begin{bmatrix} x \\ y \end{bmatrix}\right) = T\left(\begin{bmatrix} ax \\ ay \end{bmatrix}\right) = \begin{bmatrix} ax \\ ax + ay \end{bmatrix} = a \begin{bmatrix} x \\ x + y \end{bmatrix} = aT\left(\begin{bmatrix} x \\ y \end{bmatrix}\right).$$

Example

$T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ given by $T(x, y) = (x - y, 0, y^2)$ is not a linear transformation since

$$T\left(2 \begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = T\left(\begin{bmatrix} 2 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \\ 4 \end{bmatrix} \neq 2T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = 2 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix}$$

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then for each $k = 1, 2, \dots$:

$$T(a_1x_1 + a_2x_2 + \cdots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \cdots + a_kT(x_k),$$

for all scalars a_i and all vectors x_i in \mathbb{R}^n .

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then for each $k = 1, 2, \dots$:

$$T(a_1x_1 + a_2x_2 + \cdots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \cdots + a_kT(x_k),$$

for all scalars a_i and all vectors x_i in \mathbb{R}^n .

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that

$$T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -3 \end{bmatrix}. \quad \text{Find } T \begin{bmatrix} 0 \\ 5 \end{bmatrix}.$$

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then for each $k = 1, 2, \dots$:

$$T(a_1x_1 + a_2x_2 + \cdots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \cdots + a_kT(x_k),$$

for all scalars a_i and all vectors x_i in \mathbb{R}^n .

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that

$$T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -3 \end{bmatrix}. \quad \text{Find } T \begin{bmatrix} 0 \\ 5 \end{bmatrix}.$$

Answer:

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then for each $k = 1, 2, \dots$:

$$T(a_1x_1 + a_2x_2 + \cdots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \cdots + a_kT(x_k),$$

for all scalars a_i and all vectors x_i in \mathbb{R}^n .

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that

$$T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -3 \end{bmatrix}. \quad \text{Find } T \begin{bmatrix} 0 \\ 5 \end{bmatrix}.$$

Answer: Find a, b : $\begin{bmatrix} 0 \\ 5 \end{bmatrix} = a \begin{bmatrix} 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then for each $k = 1, 2, \dots$:

$$T(a_1x_1 + a_2x_2 + \cdots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \cdots + a_kT(x_k),$$

for all scalars a_i and all vectors x_i in \mathbb{R}^n .

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that

$$T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -3 \end{bmatrix}. \quad \text{Find } T \begin{bmatrix} 0 \\ 5 \end{bmatrix}.$$

Answer: Find a, b : $\begin{bmatrix} 0 \\ 5 \end{bmatrix} = a \begin{bmatrix} 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} -1 \\ 1 \end{bmatrix}$. Solve this system: $\begin{cases} a = 1 \\ b = 2 \end{cases}$.

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then for each $k = 1, 2, \dots$:

$$T(a_1x_1 + a_2x_2 + \cdots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \cdots + a_kT(x_k),$$

for all scalars a_i and all vectors x_i in \mathbb{R}^n .

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that

$$T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -3 \end{bmatrix}. \quad \text{Find } T \begin{bmatrix} 0 \\ 5 \end{bmatrix}.$$

Answer: Find a, b : $\begin{bmatrix} 0 \\ 5 \end{bmatrix} = a \begin{bmatrix} 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} -1 \\ 1 \end{bmatrix}$. Solve this system: $\begin{cases} a = 1 \\ b = 2 \end{cases}$.

$$\text{Hence } T \begin{bmatrix} 0 \\ 5 \end{bmatrix} = 1T \begin{bmatrix} 2 \\ 3 \end{bmatrix} + 2T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} + 2 \begin{bmatrix} 4 \\ -3 \end{bmatrix} = \begin{bmatrix} 10 \\ -1 \end{bmatrix}.$$

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then for each $k = 1, 2, \dots$:

$$T(a_1x_1 + a_2x_2 + \cdots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \cdots + a_kT(x_k),$$

for all scalars a_i and all vectors x_i in \mathbb{R}^n .

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that

$$T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -3 \end{bmatrix}. \quad \text{Find } T \begin{bmatrix} 0 \\ 5 \end{bmatrix}.$$

Answer: Find a, b : $\begin{bmatrix} 0 \\ 5 \end{bmatrix} = a \begin{bmatrix} 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} -1 \\ 1 \end{bmatrix}$. Solve this system: $\begin{cases} a = 1 \\ b = 2 \end{cases}$.

$$\text{Hence } T \begin{bmatrix} 0 \\ 5 \end{bmatrix} = 1T \begin{bmatrix} 2 \\ 3 \end{bmatrix} + 2T \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \end{bmatrix} + 2 \begin{bmatrix} 4 \\ -3 \end{bmatrix} = \begin{bmatrix} 10 \\ -1 \end{bmatrix}.$$

Example

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a linear transformation and u, v be vectors such that $T(u + v) = 1$ and $T(u - v) = 5$. Find $T(-u + 2v)$.

Answer:

Example

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a linear transformation and u, v be vectors such that $T(u + v) = 1$ and $T(u - v) = 5$. Find $T(-u + 2v)$.

Answer: We have

$$\begin{cases} T(u + v) = T(u) + T(v) = 1 \\ T(u - v) = T(u) - T(v) = 5 \end{cases} \Rightarrow \begin{cases} T(u) = 3 \\ T(v) = -2 \end{cases}.$$

Hence

$$T(-u + 2v) = -T(u) + 2T(v) = -3 + 2(-2) = -7.$$

Example

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a linear transformation and u, v be vectors such that $T(u + v) = 1$ and $T(u - v) = 5$. Find $T(-u + 2v)$.

Answer: We have

$$\begin{cases} T(u + v) = T(u) + T(v) = 1 \\ T(u - v) = T(u) - T(v) = 5 \end{cases} \Rightarrow \begin{cases} T(u) = 3 \\ T(v) = -2 \end{cases}.$$

Hence

$$T(-u + 2v) = -T(u) + 2T(v) = -3 + 2(-2) = -7.$$

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a transformation. Then T is linear if and only if it is a matrix transformation induced by an $m \times n$ matrix A , given in terms of its columns by

$$A = \begin{bmatrix} T(e_1) & T(e_2) & \cdots & T(e_n) \end{bmatrix}$$

where $\{e_1, e_2, \dots, e_n\}$ is the *standard basis* of \mathbb{R}^n .

Example

Consider a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$: $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - 3y \\ x + 4y \end{bmatrix}$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a transformation. Then T is linear if and only if it is a matrix transformation induced by an $m \times n$ matrix A , given in terms of its columns by

$$A = \begin{bmatrix} T(e_1) & T(e_2) & \cdots & T(e_n) \end{bmatrix}$$

where $\{e_1, e_2, \dots, e_n\}$ is the *standard basis* of \mathbb{R}^n .

Example

Consider a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$: $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - 3y \\ x + 4y \end{bmatrix}$.

Then T is also a matrix transformation induced by

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a transformation. Then T is linear if and only if it is a matrix transformation induced by an $m \times n$ matrix A , given in terms of its columns by

$$A = \begin{bmatrix} T(e_1) & T(e_2) & \cdots & T(e_n) \end{bmatrix}$$

where $\{e_1, e_2, \dots, e_n\}$ is the *standard basis* of \mathbb{R}^n .

Example

Consider a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$: $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - 3y \\ x + 4y \end{bmatrix}$.

Then T is also a matrix transformation induced by

$$A = \begin{bmatrix} T(e_1) & T(e_2) \end{bmatrix} = \begin{bmatrix} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} & T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 2 & -3 \\ 1 & 4 \end{bmatrix}.$$

We can see that

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a transformation. Then T is linear if and only if it is a matrix transformation induced by an $m \times n$ matrix A , given in terms of its columns by

$$A = \begin{bmatrix} T(e_1) & T(e_2) & \cdots & T(e_n) \end{bmatrix}$$

where $\{e_1, e_2, \dots, e_n\}$ is the **standard basis** of \mathbb{R}^n .

Example

Consider a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$: $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - 3y \\ x + 4y \end{bmatrix}$.

Then T is also a matrix transformation induced by

$$A = \begin{bmatrix} T(e_1) & T(e_2) \end{bmatrix} = \begin{bmatrix} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} & T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 2 & -3 \\ 1 & 4 \end{bmatrix}.$$

We can see that $T \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 & -3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - 3y \\ x + 4y \end{bmatrix}$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a transformation. Then T is linear if and only if it is a matrix transformation induced by an $m \times n$ matrix A , given in terms of its columns by

$$A = \begin{bmatrix} T(e_1) & T(e_2) & \cdots & T(e_n) \end{bmatrix}$$

where $\{e_1, e_2, \dots, e_n\}$ is the **standard basis** of \mathbb{R}^n .

Example


Consider a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$: $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - 3y \\ x + 4y \end{bmatrix}$.

Then T is also a matrix transformation induced by

$$A = \begin{bmatrix} T(e_1) & T(e_2) \end{bmatrix} = \begin{bmatrix} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} & T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 2 & -3 \\ 1 & 4 \end{bmatrix}.$$

We can see that $T \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 & -3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - 3y \\ x + 4y \end{bmatrix}$.


Rotation




Given an angle θ , let

$$R_\theta: \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

denote **counterclockwise rotation** of \mathbb{R}^2 about the origin through the angle θ .



Rotation




Given an angle θ , let

$$R_\theta: \mathbb{R}^2 \rightarrow \mathbb{R}^2$$


denote **counterclockwise rotation** of \mathbb{R}^2 about the origin through the angle θ .

Then R_θ is a linear transformation induced by matrix

$$\begin{bmatrix} R_\theta(e_1) & R_\theta(e_2) \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$



Rotation




Given an angle θ , let

$$R_\theta: \mathbb{R}^2 \rightarrow \mathbb{R}^2$$


denote **counterclockwise rotation** of \mathbb{R}^2 about the origin through the angle θ .

Then R_θ is a linear transformation induced by matrix

$$\begin{bmatrix} R_\theta(e_1) & R_\theta(e_2) \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$



Reflection




The line through the origin with slope m has equation $y = mx$, and we let $Q_m: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote **reflection in the line $y = mx$** .

Then Q_m is a linear transformation induced by matrix

$$\frac{1}{1 + m^2} \begin{bmatrix} 1 - m^2 & 2m \\ 2m & m^2 - 1 \end{bmatrix}.$$

Reflection




The line through the origin with slope m has equation $y = mx$, and we let $Q_m: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote **reflection in the line $y = mx$** .

Then Q_m is a linear transformation induced by matrix

$$\frac{1}{1 + m^2} \begin{bmatrix} 1 - m^2 & 2m \\ 2m & m^2 - 1 \end{bmatrix}.$$

Projection




Let $P_m: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote **projection on the line $y = mx$** .

Then P_m is a linear transformation with matrix

$$\frac{1}{1+m^2} \begin{bmatrix} 1 & m \\ m & m^2 \end{bmatrix}.$$

Projection



Let $P_m: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote **projection on the line $y = mx$** .

Then P_m is a linear transformation with matrix

$$\frac{1}{1+m^2} \begin{bmatrix} 1 & m \\ m & m^2 \end{bmatrix}.$$

Composition

Given two “linked” transformations

$$\begin{array}{ccccc} \mathbb{R}^k & \xrightarrow{T} & \mathbb{R}^n & \xrightarrow{S} & \mathbb{R}^m \\ x & \mapsto & T(x) & \mapsto & S[T(x)] \end{array}$$

The **composite** of S and T :

$$S \circ T: \mathbb{R}^k \rightarrow \mathbb{R}^m$$

defined by

Composition

Given two “linked” transformations

$$\begin{array}{ccccc} \mathbb{R}^k & \xrightarrow{T} & \mathbb{R}^n & \xrightarrow{S} & \mathbb{R}^m \\ x & \mapsto & T(x) & \mapsto & S[T(x)] \end{array}$$

The **composite** of S and T :

$$S \circ T: \mathbb{R}^k \rightarrow \mathbb{R}^m$$

defined by

$$(S \circ T)(x) = S[T(x)] \quad \text{for all } x \in \mathbb{R}^k.$$

Composition

Given two “linked” transformations

$$\begin{array}{ccccc} \mathbb{R}^k & \xrightarrow{T} & \mathbb{R}^n & \xrightarrow{S} & \mathbb{R}^m \\ x & \mapsto & T(x) & \mapsto & S[T(x)] \end{array}$$

The **composite** of S and T :

$$S \circ T: \mathbb{R}^k \rightarrow \mathbb{R}^m$$

defined by

$$(S \circ T)(x) = S[T(x)] \quad \text{for all } x \in \mathbb{R}^k.$$

Theorem

Let $\mathbb{R}^k \xrightarrow[A_{n \times k}]{T} \mathbb{R}^n \xrightarrow[B_{m \times n]{S}} \mathbb{R}^m$ be linear transformations, and let A and B be the matrices of T and S respectively. Then $S \circ T$ is linear with matrix BA .

Composition

Given two “linked” transformations

$$\begin{array}{ccccc} \mathbb{R}^k & \xrightarrow{T} & \mathbb{R}^n & \xrightarrow{S} & \mathbb{R}^m \\ x & \mapsto & T(x) & \mapsto & S[T(x)] \end{array}$$

The **composite** of S and T :

$$S \circ T: \mathbb{R}^k \rightarrow \mathbb{R}^m$$

defined by

$$(S \circ T)(x) = S[T(x)] \quad \text{for all } x \in \mathbb{R}^k.$$

Theorem

Let $\mathbb{R}^k \xrightarrow[A_{n \times k}]{T} \mathbb{R}^n \xrightarrow[B_{m \times n]{S}} \mathbb{R}^m$ be linear transformations, and let A and B be the matrices of T and S respectively. Then $S \circ T$ is linear with matrix BA .

Example

Let T and S be linear transformations given by:

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x + 3y \\ -x + 2y \end{bmatrix}$$

$$S: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad S \begin{bmatrix} x \\ y \end{bmatrix} = x + 5y$$

Then $S \circ T: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by:

Example

Let T and S be linear transformations given by:

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x + 3y \\ -x + 2y \end{bmatrix}$$

$$S: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad S \begin{bmatrix} x \\ y \end{bmatrix} = x + 5y$$

Then $S \circ T: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by:

$$\begin{aligned} S \circ T \begin{bmatrix} x \\ y \end{bmatrix} &= S \left(T \begin{bmatrix} x \\ y \end{bmatrix} \right) = S \begin{bmatrix} 2x + 3y \\ -x + 2y \end{bmatrix} \\ &= (2x + 3y) + 5(-x + 2y) = -3x + 13y. \end{aligned}$$

Example

Let T and S be linear transformations given by:

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x + 3y \\ -x + 2y \end{bmatrix}$$

$$S: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad S \begin{bmatrix} x \\ y \end{bmatrix} = x + 5y$$

Then $S \circ T: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by:

$$\begin{aligned} S \circ T \begin{bmatrix} x \\ y \end{bmatrix} &= S \left(T \begin{bmatrix} x \\ y \end{bmatrix} \right) = S \begin{bmatrix} 2x + 3y \\ -x + 2y \end{bmatrix} \\ &= (2x + 3y) + 5(-x + 2y) = -3x + 13y. \end{aligned}$$

We can see that the matrices of T , S and $S \circ T$ respectively are

$$A = \begin{bmatrix} 2 & 3 \\ -1 & 2 \end{bmatrix}, B = \begin{bmatrix} 1 & 5 \end{bmatrix}, \text{ and } BA = \begin{bmatrix} -3 & 13 \end{bmatrix}.$$

Example

Let T and S be linear transformations given by:

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x + 3y \\ -x + 2y \end{bmatrix}$$

$$S: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad S \begin{bmatrix} x \\ y \end{bmatrix} = x + 5y$$

Then $S \circ T: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by:

$$\begin{aligned} S \circ T \begin{bmatrix} x \\ y \end{bmatrix} &= S \left(T \begin{bmatrix} x \\ y \end{bmatrix} \right) = S \begin{bmatrix} 2x + 3y \\ -x + 2y \end{bmatrix} \\ &= (2x + 3y) + 5(-x + 2y) = -3x + 13y. \end{aligned}$$

We can see that the matrices of T , S and $S \circ T$ respectively are

$$A = \begin{bmatrix} 2 & 3 \\ -1 & 2 \end{bmatrix}, B = [1 \quad 5], \text{ and } BA = [-3 \quad 13].$$

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

1st way. The matrix of the rotation through $\frac{\pi}{2}$ is

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

1st **way.** The matrix of the rotation through $\frac{\pi}{2}$ is

$$\begin{bmatrix} \cos \frac{\pi}{2} & -\sin \frac{\pi}{2} \\ \sin \frac{\pi}{2} & \cos \frac{\pi}{2} \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

The matrix of the reflection in the line $y = -x$ is

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

1st **way.** The matrix of the rotation through $\frac{\pi}{2}$ is

$$\begin{bmatrix} \cos \frac{\pi}{2} & -\sin \frac{\pi}{2} \\ \sin \frac{\pi}{2} & \cos \frac{\pi}{2} \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

The matrix of the reflection in the line $y = -x$ is

$$\frac{1}{1 + (-1)^2} \begin{bmatrix} 1 - (-1)^2 & 2 \cdot (-1) \\ 2 \cdot (-1) & (-1)^2 - 1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

Hence, the matrix of T is

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

1st **way.** The matrix of the rotation through $\frac{\pi}{2}$ is

$$\begin{bmatrix} \cos \frac{\pi}{2} & -\sin \frac{\pi}{2} \\ \sin \frac{\pi}{2} & \cos \frac{\pi}{2} \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

The matrix of the reflection in the line $y = -x$ is

$$\frac{1}{1 + (-1)^2} \begin{bmatrix} 1 - (-1)^2 & 2 \cdot (-1) \\ 2 \cdot (-1) & (-1)^2 - 1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

Hence, the matrix of T is $\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$.

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

1st **way.** The matrix of the rotation through $\frac{\pi}{2}$ is

$$\begin{bmatrix} \cos \frac{\pi}{2} & -\sin \frac{\pi}{2} \\ \sin \frac{\pi}{2} & \cos \frac{\pi}{2} \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

The matrix of the reflection in the line $y = -x$ is

$$\frac{1}{1 + (-1)^2} \begin{bmatrix} 1 - (-1)^2 & 2 \cdot (-1) \\ 2 \cdot (-1) & (-1)^2 - 1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

Hence, the matrix of T is $\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$.

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

2nd way. We can see

$$\begin{aligned} e_1 &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} \xrightarrow{\text{rotation through } \pi/2} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \xrightarrow{\text{reflection in } y=-x} \begin{bmatrix} -1 \\ 0 \end{bmatrix} \\ e_2 &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} \xrightarrow{\text{rotation through } \pi/2} \begin{bmatrix} -1 \\ 0 \end{bmatrix} \xrightarrow{\text{reflection in } y=-x} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{aligned}$$

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

2nd way. We can see

$$\begin{aligned} e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} &\xrightarrow{\text{rotation through } \pi/2} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \xrightarrow{\text{reflection in } y=-x} \begin{bmatrix} -1 \\ 0 \end{bmatrix} \\ e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} &\xrightarrow{\text{rotation through } \pi/2} \begin{bmatrix} -1 \\ 0 \end{bmatrix} \xrightarrow{\text{reflection in } y=-x} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{aligned}$$

Hence, the matrix of T is

$$\begin{bmatrix} T(e_1) & T(e_2) \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be rotation through $\frac{\pi}{2}$ followed by reflection in the line $y = -x$. Find the matrix of T .

2nd way. We can see

$$\begin{aligned} e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} &\xrightarrow{\text{rotation through } \pi/2} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \xrightarrow{\text{reflection in } y=-x} \begin{bmatrix} -1 \\ 0 \end{bmatrix} \\ e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} &\xrightarrow{\text{rotation through } \pi/2} \begin{bmatrix} -1 \\ 0 \end{bmatrix} \xrightarrow{\text{reflection in } y=-x} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{aligned}$$

Hence, the matrix of T is

$$\begin{bmatrix} T(e_1) & T(e_2) \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the matrix transformation induced by

$$A = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}. \text{ Find } (T \circ T) \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$

Answer:

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the matrix transformation induced by

$$A = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}. \text{ Find } (T \circ T) \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$

Answer: We have

$$\begin{aligned} (T \circ T) \begin{bmatrix} -1 \\ 2 \end{bmatrix} &= A^2 \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -5 \\ 2 \end{bmatrix}. \end{aligned}$$

Example

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the matrix transformation induced by

$$A = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}. \text{ Find } (T \circ T) \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$

Answer: We have

$$\begin{aligned} (T \circ T) \begin{bmatrix} -1 \\ 2 \end{bmatrix} &= A^2 \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -5 \\ 2 \end{bmatrix}. \end{aligned}$$

Section 2.1 : $1ad, 2aef, 3acd, 4, 14, 15, 19$ (page 45, 46)

Section 2.3 : $1adef, 2, 3, 4, 6, 7, 12, 22$ (page 76 – 79)

Section 2.4 : $1bd, 2aeg, 3acd, 4, 5, 6, 9, 12$ (page 91 – 93)

Section 2.6 : $1, 3, 7, 8, 9, 12$ (page 115, 116)