

# Chapter 1. Systems of Linear Equations

1 Solutions and Elementary Operations

2 Gaussian Elimination

3 Homogeneous Systems

# 1.1 Solutions and Elementary Operations

## Definition

A system of  $m$  linear equations and  $n$  variables ( unknowns):

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases} \quad (\star) \text{ where } a_{ij}, b_i \in \mathbb{R}$$

We say  $A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$  the coefficient matrix

# 1.1 Solutions and Elementary Operations

## Definition

A system of  $m$  linear equations and  $n$  variables ( unknowns):

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases} \quad (\star) \text{ where } a_{ij}, b_i \in \mathbb{R}$$

We say  $A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$  the coefficient matrix ,  $b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$ ,

# 1.1 Solutions and Elementary Operations

## Definition

A system of  $m$  linear equations and  $n$  variables ( unknowns):

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases} \quad (*) \text{ where } a_{ij}, b_i \in \mathbb{R}$$

We say  $A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$  the coefficient matrix ,  $b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$ ,

$\bar{A} = [A|b] = A = \left[ \begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{array} \right]$  the augmented matrix.

# 1.1 Solutions and Elementary Operations

## Definition

A system of  $m$  linear equations and  $n$  variables ( unknowns):

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases} \quad (\star) \text{ where } a_{ij}, b_i \in \mathbb{R}$$

We say  $A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$  the coefficient matrix ,  $b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$ ,

$\bar{A} = [A|b] = A = \left[ \begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{array} \right]$  the augmented matrix.

## Example

$$\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$$

is a system of 2 linear equations and 2 variables with

the augmented matrix:

## Example

$$\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$$
 is a system of 2 linear equations and 2 variables with the augmented matrix:

$$\bar{A} = \left[ \begin{array}{cc|c} 1 & 1 & 36 \\ 2 & 4 & 100 \end{array} \right]$$



## Example

$$\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$$
 is a system of 2 linear equations and 2 variables with the augmented matrix:

$$\bar{A} = \left[ \begin{array}{cc|c} 1 & 1 & 36 \\ 2 & 4 & 100 \end{array} \right]$$

## Example

$$\begin{cases} x_1 - 3x_2 + 3x_3 + x_4 = 9 \\ 4x_2 - x_3 = 6 \end{cases}$$

## Example

$$\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$$
 is a system of 2 linear equations and 2 variables with the augmented matrix:

$$\bar{A} = \left[ \begin{array}{cc|c} 1 & 1 & 36 \\ 2 & 4 & 100 \end{array} \right]$$

## Example

$$\begin{cases} x_1 - 3x_2 + 3x_3 + x_4 = 9 \\ 4x_2 - x_3 = 6 \end{cases}$$
 is a system of 2 equations and 4 variables, with the augmented matrix

## Example

$$\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$$
 is a system of 2 linear equations and 2 variables with the augmented matrix:

$$\bar{A} = \left[ \begin{array}{cc|c} 1 & 1 & 36 \\ 2 & 4 & 100 \end{array} \right]$$

## Example

$$\begin{cases} x_1 - 3x_2 + 3x_3 + x_4 = 9 \\ 4x_2 - x_3 = 6 \end{cases}$$
 is a system of 2 equations and 4 variables, with the augmented matrix

$$\bar{A} = \left[ \begin{array}{cccc|c} 1 & -3 & 3 & 1 & 9 \\ 0 & 4 & -1 & 0 & 6 \end{array} \right]$$

## Example

$$\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$$
 is a system of 2 linear equations and 2 variables with the augmented matrix:

$$\bar{A} = \left[ \begin{array}{cc|c} 1 & 1 & 36 \\ 2 & 4 & 100 \end{array} \right]$$

## Example

$$\begin{cases} x_1 - 3x_2 + 3x_3 + x_4 = 9 \\ 4x_2 - x_3 = 6 \end{cases}$$
 is a system of 2 equations and 4 variables, with the augmented matrix

$$\bar{A} = \left[ \begin{array}{cccc|c} 1 & -3 & 3 & 1 & 9 \\ 0 & 4 & -1 & 0 & 6 \end{array} \right]$$

- A **solution** to one equation is a sequence of real numbers  $s_1, s_2, \dots, s_n$  such that  $a_1s_1 + a_2s_2 + \dots + a_ns_n = b$ .
- $s_1, s_2, \dots, s_n$  is called a solution to a system of linear equations if  $s_1, s_2, \dots, s_n$  is a solution to every equation of the system.
- Two systems are said to be **equivalent** if they have the same set of solutions.

A system can have **no solution** or **unique solution**, or **infinite solution**.

### Definition

A system has a solution is called **consistent**.

A system can have **no solution** or **unique solution**, or **infinite solution**.

## Definition

A system has a solution is called **consistent**.

A system has no solution is called **inconsistent**.

A system can have **no solution** or **unique solution**, or **infinite solution**.

## Definition

A system has a solution is called **consistent**.

A system has no solution is called **inconsistent**.

## Example

•  $\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$  is consistent (unique solution  $\begin{cases} x = 22 \\ y = 14 \end{cases}$ ).



A system can have **no solution** or **unique solution**, or **infinite solution**.

### Definition

A system has a solution is called **consistent**.

A system has no solution is called **inconsistent**.

### Example

- $\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$  is consistent (unique solution  $\begin{cases} x = 22 \\ y = 14 \end{cases}$ ).
- $\begin{cases} x + y = 36 \\ 2x + 2y = 100 \end{cases}$  is inconsistent (no solution).

A system can have **no solution** or **unique solution**, or **infinite solution**.

## Definition

A system has a solution is called **consistent**.

A system has no solution is called **inconsistent**.

## Example

- $\begin{cases} x + y = 36 \\ 2x + 4y = 100 \end{cases}$  is consistent (unique solution  $\begin{cases} x = 22 \\ y = 14 \end{cases}$ ).
- $\begin{cases} x + y = 36 \\ 2x + 2y = 100 \end{cases}$  is inconsistent (no solution).

## Example

Find the solution of 
$$\begin{cases} x + y - z = 1 \\ 2x - y + z = 5 \\ -x + 2y - 2z = -4 \end{cases} .$$

The system is **equivalent to** (by using operations  $R'_2 = -2R_1 + R_2$  ,  $R'_3 = R_1 + R_3$ ) the following system

## Example

Find the solution of 
$$\begin{cases} x + y - z = 1 \\ 2x - y + z = 5 \\ -x + 2y - 2z = -4 \end{cases}.$$

The system is **equivalent to** (by using operations  $R'_2 = -2R_1 + R_2$ ,  $R'_3 = R_1 + R_3$ ) the following system

$$\begin{cases} x + y - z = 1 \\ 0x - 3y + 3z = 3 \\ 0x + 3y - 3z = -3 \end{cases} \Leftrightarrow \begin{cases} x + y - z = 1 \\ y - z = -1 \end{cases}$$

which has infinite solutions of the form

## Example

Find the solution of 
$$\begin{cases} x + y - z = 1 \\ 2x - y + z = 5 \\ -x + 2y - 2z = -4 \end{cases}.$$

The system is **equivalent to** (by using operations  $R'_2 = -2R_1 + R_2$ ,  $R'_3 = R_1 + R_3$ ) the following system

$$\begin{cases} x + y - z = 1 \\ 0x - 3y + 3z = 3 \\ 0x + 3y - 3z = -3 \end{cases} \Leftrightarrow \begin{cases} x + y - z = 1 \\ y - z = -1 \end{cases}$$

which has infinite solutions of the form 
$$\begin{cases} x = 2 \\ y = -1 + z = -1 + t \\ z = t \end{cases}.$$

## Example

Find the solution of 
$$\begin{cases} x + y - z = 1 \\ 2x - y + z = 5 \\ -x + 2y - 2z = -4 \end{cases}.$$

The system is **equivalent to** (by using operations  $R'_2 = -2R_1 + R_2$ ,  $R'_3 = R_1 + R_3$ ) the following system

$$\begin{cases} x + y - z = 1 \\ 0x - 3y + 3z = 3 \\ 0x + 3y - 3z = -3 \end{cases} \Leftrightarrow \begin{cases} x + y - z = 1 \\ y - z = -1 \end{cases}$$

which has infinite solutions of the form 
$$\begin{cases} x = 2 \\ y = -1 + z = -1 + t \\ z = t \end{cases}$$

## Definition

The following operations, called **elementary operations**, can routinely be performed on systems of linear equations (rows of matrix) to produce equivalent systems.

1. Interchange two equations/rows.

## Definition

The following operations, called **elementary operations**, can routinely be performed on systems of linear equations (rows of matrix) to produce equivalent systems.

1. Interchange two equations/rows.
2. Multiply one equation/row by a nonzero number.



## Definition

The following operations, called **elementary operations**, can routinely be performed on systems of linear equations (rows of matrix) to produce equivalent systems.

1. Interchange two equations/rows.
2. Multiply one equation/row by a nonzero number.
3. Add a multiple of one equation/row to a different equation/row.

## Definition

The following operations, called **elementary operations**, can routinely be performed on systems of linear equations (rows of matrix) to produce equivalent systems.

1. Interchange two equations/rows.
2. Multiply one equation/row by a nonzero number.
3. Add a multiple of one equation/row to a different equation/row.

## Definition

The following operations, called **elementary operations**, can routinely be performed on systems of linear equations (rows of matrix) to produce equivalent systems.

1. Interchange two equations/rows.
2. Multiply one equation/row by a nonzero number.
3. Add a multiple of one equation/row to a different equation/row.

## Theorem

*Suppose that a sequence of elementary operations is performed on a system of linear equations. Then the resulting system has the same set of solutions as the original, so the two systems are equivalent.*

## Definition

The following operations, called **elementary operations**, can routinely be performed on systems of linear equations (rows of matrix) to produce equivalent systems.

1. Interchange two equations/rows.
2. Multiply one equation/row by a nonzero number.
3. Add a multiple of one equation/row to a different equation/row.

## Theorem

*Suppose that a sequence of elementary operations is performed on a system of linear equations. Then the resulting system has the same set of solutions as the original, so the two systems are equivalent.*

## Example

Find the solution of the system  $\begin{cases} x + 2y = 1 \\ 2x + 3y = 5 \end{cases}$ .

The augmented matrix

$$\begin{aligned} \bar{A} &= \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 2 & 3 & 5 \end{array} \right] \xrightarrow{-2R_1+R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & -1 & 3 \end{array} \right] \\ &\xrightarrow{-R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & 1 & -3 \end{array} \right]. \end{aligned}$$

## Example

Find the solution of the system  $\begin{cases} x + 2y = 1 \\ 2x + 3y = 5 \end{cases}$ .

The augmented matrix

$$\begin{aligned} \bar{A} &= \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 2 & 3 & 5 \end{array} \right] \xrightarrow{-2R_1+R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & -1 & 3 \end{array} \right] \\ &\xrightarrow{-R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & 1 & -3 \end{array} \right]. \end{aligned}$$

Hence the system is equivalent to the following system

## Example

Find the solution of the system  $\begin{cases} x + 2y = 1 \\ 2x + 3y = 5 \end{cases}$ .

The augmented matrix

$$\begin{aligned} \bar{A} &= \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 2 & 3 & 5 \end{array} \right] \xrightarrow{-2R_1+R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & -1 & 3 \end{array} \right] \\ &\xrightarrow{-R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & 1 & -3 \end{array} \right]. \end{aligned}$$

Hence the system is equivalent to the following system

$$\begin{cases} x + 2y = 1 \\ y = -3 \end{cases}$$

## Example

Find the solution of the system  $\begin{cases} x + 2y = 1 \\ 2x + 3y = 5 \end{cases}$ .

The augmented matrix

$$\begin{aligned} \bar{A} &= \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 2 & 3 & 5 \end{array} \right] \xrightarrow{-2R_1+R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & -1 & 3 \end{array} \right] \\ &\xrightarrow{-R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & 1 & -3 \end{array} \right]. \end{aligned}$$

Hence the system is equivalent to the following system

$$\begin{cases} x + 2y = 1 \\ y = -3 \end{cases}$$

which has unique solution  $\begin{cases} x = 7 \\ y = -3 \end{cases}$ .



## Example

Find the solution of the system  $\begin{cases} x + 2y = 1 \\ 2x + 3y = 5 \end{cases}$ .

The augmented matrix

$$\begin{aligned} \bar{A} &= \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 2 & 3 & 5 \end{array} \right] \xrightarrow{-2R_1+R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & -1 & 3 \end{array} \right] \\ &\xrightarrow{-R_2} \left[ \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & 1 & -3 \end{array} \right]. \end{aligned}$$

Hence the system is equivalent to the following system

$$\begin{cases} x + 2y = 1 \\ y = -3 \end{cases}$$

which has unique solution  $\begin{cases} x = 7 \\ y = -3 \end{cases}$ .

Find the solution of the system 
$$\begin{cases} x + y - z &= 2 \\ -x + 2y + 3z &= 5 \\ 2x - y - 4z &= 6 \end{cases}$$

The augmented matrix

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ -1 & 2 & 3 & 5 \\ 2 & -1 & -4 & 6 \end{array} \right] \xrightarrow[\begin{smallmatrix} -2R_1+R_3 \end{smallmatrix}]{\begin{smallmatrix} R_1+R_2 \end{smallmatrix}} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & -3 & -2 & 2 \end{array} \right]$$
$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & 0 & 0 & 9 \end{array} \right]$$

Find the solution of the system 
$$\begin{cases} x + y - z &= 2 \\ -x + 2y + 3z &= 5 \\ 2x - y - 4z &= 6 \end{cases}.$$

The augmented matrix

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ -1 & 2 & 3 & 5 \\ 2 & -1 & -4 & 6 \end{array} \right] \xrightarrow[-2R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & -3 & -2 & 2 \end{array} \right]$$
$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & 0 & 0 & 9 \end{array} \right]$$

We obtain an equivalent system 
$$\begin{cases} x + y - z &= 2 \\ 3y + 2z &= 7 \\ 0x + 0y + 0z &= 9 \end{cases}$$

Find the solution of the system 
$$\begin{cases} x + y - z &= 2 \\ -x + 2y + 3z &= 5 \\ 2x - y - 4z &= 6 \end{cases}.$$

The augmented matrix

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ -1 & 2 & 3 & 5 \\ 2 & -1 & -4 & 6 \end{array} \right] \xrightarrow[-2R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & -3 & -2 & 2 \end{array} \right]$$
$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & 0 & 0 & 9 \end{array} \right]$$

We obtain an equivalent system 
$$\begin{cases} x + y - z &= 2 \\ 3y + 2z &= 7 \\ 0x + 0y + 0z &= 9 \end{cases}$$
 which is

inconsistent.

Find the solution of the system 
$$\begin{cases} x + y - z &= 2 \\ -x + 2y + 3z &= 5 \\ 2x - y - 4z &= 6 \end{cases}.$$

The augmented matrix

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ -1 & 2 & 3 & 5 \\ 2 & -1 & -4 & 6 \end{array} \right] \xrightarrow[-2R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & -3 & -2 & 2 \end{array} \right]$$
$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & -1 & 2 \\ 0 & 3 & 2 & 7 \\ 0 & 0 & 0 & 9 \end{array} \right]$$

We obtain an equivalent system 
$$\begin{cases} x + y - z &= 2 \\ 3y + 2z &= 7 \\ 0x + 0y + 0z &= 9 \end{cases}$$
 which is

inconsistent.

# 1.2 Gaussian Elimination

## Definition

A matrix is said to be in **row-echelon** if it satisfies the following three conditions:

- (i) All zero rows (consisting entirely of zeros) are at the bottom.

## 1.2 Gaussian Elimination

### Definition

A matrix is said to be in **row-echelon** if it satisfies the following three conditions:

- (i) All zero rows (consisting entirely of zeros) are at the bottom.
- (ii) The first nonzero entry from the left in each nonzero row is a 1, called **the leading 1** for that row.

## 1.2 Gaussian Elimination

### Definition

A matrix is said to be in **row-echelon** if it satisfies the following three conditions:

- (i) All zero rows (consisting entirely of zeros) are at the bottom.
- (ii) The first nonzero entry from the left in each nonzero row is a 1, called **the leading 1** for that row.
- (iii) Each leading 1 is to the right of all leading 1's in all rows above it.



## 1.2 Gaussian Elimination

### Definition

A matrix is said to be in **row-echelon** if it satisfies the following three conditions:

- (i) All zero rows (consisting entirely of zeros) are at the bottom.
- (ii) The first nonzero entry from the left in each nonzero row is a 1, called **the leading 1** for that row.
- (iii) Each leading 1 is to the right of all leading 1's in all rows above it.

A row-echelon matrix is **reduced row-echelon** if it satisfies:

## 1.2 Gaussian Elimination

### Definition

A matrix is said to be in **row-echelon** if it satisfies the following three conditions:

- (i) All zero rows (consisting entirely of zeros) are at the bottom.
- (ii) The first nonzero entry from the left in each nonzero row is a 1, called **the leading 1** for that row.
- (iii) Each leading 1 is to the right of all leading 1's in all rows above it.

A row-echelon matrix is **reduced row-echelon** if it satisfies:

- (iv) Each leading 1 is the only nonzero entry in its column.

## 1.2 Gaussian Elimination

### Definition

A matrix is said to be in **row-echelon** if it satisfies the following three conditions:

- (i) All zero rows (consisting entirely of zeros) are at the bottom.
- (ii) The first nonzero entry from the left in each nonzero row is a 1, called **the leading 1** for that row.
- (iii) Each leading 1 is to the right of all leading 1's in all rows above it.

A row-echelon matrix is **reduced row-echelon** if it satisfies:

- (iv) Each leading 1 is the only nonzero entry in its column.

## 1.2 Gaussian Elimination

### Definition

A matrix is said to be in **row-echelon** if it satisfies the following three conditions:

- (i) All zero rows (consisting entirely of zeros) are at the bottom.
- (ii) The first nonzero entry from the left in each nonzero row is a 1, called **the leading 1** for that row.
- (iii) Each leading 1 is to the right of all leading 1's in all rows above it.

A row-echelon matrix is **reduced row-echelon** if it satisfies:

- (iv) Each leading 1 is the only nonzero entry in its column.

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon



## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon

but not reduced

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon

but not reduced

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon

but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon

but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon

but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon  
but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & -6 & 8 \\ 0 & 0 & 1 & 3 \\ 0 & 1 & -5 & 6 \end{bmatrix}$$

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon  
but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & -6 & 8 \\ 0 & 0 & 1 & 3 \\ 0 & 1 & -5 & 6 \end{bmatrix}$$

is not in row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon  
but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & -6 & 8 \\ 0 & 0 & 1 & 3 \\ 0 & 1 & -5 & 6 \end{bmatrix}$$

is not in row-echelon.



## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon  
but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & -6 & 8 \\ 0 & 0 & 1 & 3 \\ 0 & 1 & -5 & 6 \end{bmatrix}$$

is not in row-echelon.

$$\begin{bmatrix} 1 & 0 & -6 & 5 \\ 0 & 1 & 2 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon  
but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & -6 & 8 \\ 0 & 0 & 1 & 3 \\ 0 & 1 & -5 & 6 \end{bmatrix}$$

is not in row-echelon.

$$\begin{bmatrix} 1 & 0 & -6 & 5 \\ 0 & 1 & 2 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

is in reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 2 & 3 \end{bmatrix}$$

not row-echelon

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

row-echelon  
but not reduced

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced row-echelon.

## Example

$$\begin{bmatrix} 1 & 2 & -6 & 8 \\ 0 & 0 & 1 & 3 \\ 0 & 1 & -5 & 6 \end{bmatrix}$$

is not in row-echelon.

$$\begin{bmatrix} 1 & 0 & -6 & 5 \\ 0 & 1 & 2 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

is in reduced row-echelon.

## Theorem

*Every matrix can be brought to (reduced) row-echelon form by a sequence of elementary row operations.*

## Example

Consider the matrix  $B = \begin{bmatrix} 2 & 1 & 3 & -3 \\ 1 & -1 & 2 & 5 \\ -2 & 5 & 1 & 4 \end{bmatrix}$ .

## Theorem

*Every matrix can be brought to (reduced) row-echelon form by a sequence of elementary row operations.*

## Example

Consider the matrix  $B = \begin{bmatrix} 2 & 1 & 3 & -3 \\ 1 & -1 & 2 & 5 \\ -2 & 5 & 1 & 4 \end{bmatrix}$ .

The matrix  $B$  can be brought to row-echelon as:

## Theorem

*Every matrix can be brought to (reduced) row-echelon form by a sequence of elementary row operations.*

## Example

Consider the matrix  $B = \begin{bmatrix} 2 & 1 & 3 & -3 \\ 1 & -1 & 2 & 5 \\ -2 & 5 & 1 & 4 \end{bmatrix}$ .

The matrix  $B$  can be brought to row-echelon as:

$$\begin{aligned} B &\xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 2 & 1 & 3 & -3 \\ -2 & 5 & 1 & 4 \end{bmatrix} \xrightarrow[\substack{-2R_1 + R_2 \\ 2R_1 + R_3}]{\phantom{0}} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 0 & 3 & -1 & -13 \\ 0 & 3 & 5 & 14 \end{bmatrix} \\ &\xrightarrow{-R_2 + R_3} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 0 & 3 & -1 & -13 \\ 0 & 0 & 6 & 27 \end{bmatrix} \xrightarrow[\substack{1/3R_2 \\ 1/6R_3}]{\phantom{0}} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 0 & 1 & -1/3 & -13/3 \\ 0 & 0 & 1 & 27/6 \end{bmatrix}. \end{aligned}$$

## Theorem

*Every matrix can be brought to (reduced) row-echelon form by a sequence of elementary row operations.*

## Example

Consider the matrix  $B = \begin{bmatrix} 2 & 1 & 3 & -3 \\ 1 & -1 & 2 & 5 \\ -2 & 5 & 1 & 4 \end{bmatrix}$ .

The matrix  $B$  can be brought to row-echelon as:

$$\begin{aligned} B &\xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 2 & 1 & 3 & -3 \\ -2 & 5 & 1 & 4 \end{bmatrix} \xrightarrow{\substack{-2R_1 + R_2 \\ 2R_1 + R_3}} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 0 & 3 & -1 & -13 \\ 0 & 3 & 5 & 14 \end{bmatrix} \\ &\xrightarrow{-R_2 + R_3} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 0 & 3 & -1 & -13 \\ 0 & 0 & 6 & 27 \end{bmatrix} \xrightarrow{\substack{1/3 R_2 \\ 1/6 R_3}} \begin{bmatrix} 1 & -1 & 2 & 5 \\ 0 & 1 & -1/3 & -13/3 \\ 0 & 0 & 1 & 27/6 \end{bmatrix}. \end{aligned}$$

## Example

Consider the matrix  $C = \begin{bmatrix} 1 & -2 & -1 & 3 \\ 2 & -4 & 1 & 0 \\ 1 & -2 & -1 & -1 \end{bmatrix}$ .

The matrix  $C$  can be brought to reduced row-echelon form as:



## Example

Consider the matrix  $C = \begin{bmatrix} 1 & -2 & -1 & 3 \\ 2 & -4 & 1 & 0 \\ 1 & -2 & -1 & -1 \end{bmatrix}$ .

The matrix  $C$  can be brought to reduced row-echelon form as:

$$\begin{aligned} C &\xrightarrow[-R_1+R_3]{-2R_1+R_2} \begin{bmatrix} 1 & -2 & -1 & 3 \\ 0 & 0 & 3 & -6 \\ 0 & 0 & 0 & -4 \end{bmatrix} \xrightarrow[-1/4R_3]{1/3R_2} \begin{bmatrix} 1 & -2 & -1 & 3 \\ 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 1 \end{bmatrix} \\ &\xrightarrow[-3R_3+R_1]{2R_3+R_2} \begin{bmatrix} 1 & -2 & -1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_2+R_1} \begin{bmatrix} 1 & -2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

## Example

Consider the matrix  $C = \begin{bmatrix} 1 & -2 & -1 & 3 \\ 2 & -4 & 1 & 0 \\ 1 & -2 & -1 & -1 \end{bmatrix}$ .

The matrix  $C$  can be brought to reduced row-echelon form as:

$$\begin{aligned} C &\xrightarrow[-R_1+R_3]{-2R_1+R_2} \begin{bmatrix} 1 & -2 & -1 & 3 \\ 0 & 0 & 3 & -6 \\ 0 & 0 & 0 & -4 \end{bmatrix} \xrightarrow[-1/4R_3]{1/3R_2} \begin{bmatrix} 1 & -2 & -1 & 3 \\ 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 1 \end{bmatrix} \\ &\xrightarrow[-3R_3+R_1]{2R_3+R_2} \begin{bmatrix} 1 & -2 & -1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_2+R_1} \begin{bmatrix} 1 & -2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

# Rank of a matrix

- The reduced row-echelon form of a matrix  $A$  is uniquely determined by  $A$ , but the row-echelon form of  $A$  is not unique.
- The number  $r$  of leading 1's is the same in each of the different row-echelon matrices.
- As  $r$  depends only on  $A$  and not on the row-echelon forms, it is called the rank of the matrix  $A$ , denoted by  $\text{rank}(A)$ .

# Rank of a matrix

## Definition

The **rank** of matrix  $A$  is the number of leading 1s in any row-echelon matrix to which  $A$  can be carried by row operations.

## Example

Find the rank of  $A = \begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix}$ .

# Rank of a matrix

## Definition

The **rank** of matrix  $A$  is the number of leading 1s in any row-echelon matrix to which  $A$  can be carried by row operations.

## Example

Find the rank of  $A = \begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix}$ .

Answer:

# Rank of a matrix

## Definition

The **rank** of matrix  $A$  is the number of leading 1s in any row-echelon matrix to which  $A$  can be carried by row operations.

## Example

Find the rank of  $A = \begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix}$ .

**Answer:**

$$A \xrightarrow[-R_1+R_3]{R_1+R_2} \begin{bmatrix} 1 & 1 & -1 & 3 \\ 0 & 5 & 4 & 1 \\ 0 & 5 & 4 & 1 \end{bmatrix} \xrightarrow{1/5 R_2} \begin{bmatrix} 1 & 1 & -1 & 3 \\ 0 & 1 & 4/5 & 1/5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

# Rank of a matrix

## Definition

The **rank** of matrix  $A$  is the number of leading 1s in any row-echelon matrix to which  $A$  can be carried by row operations.

## Example

Find the rank of  $A = \begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix}$ .

**Answer:**

$$A \xrightarrow[-R_1+R_3]{R_1+R_2} \begin{bmatrix} 1 & 1 & -1 & 3 \\ 0 & 5 & 4 & 1 \\ 0 & 5 & 4 & 1 \end{bmatrix} \xrightarrow[1/5R_2]{-R_2+R_3} \begin{bmatrix} 1 & 1 & -1 & 3 \\ 0 & 1 & 4/5 & 1/5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Hence  $\text{rank}(A) = 2$ .

# Rank of a matrix

## Definition

The **rank** of matrix  $A$  is the number of leading 1s in any row-echelon matrix to which  $A$  can be carried by row operations.

## Example

Find the rank of  $A = \begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix}$ .

**Answer:**

$$A \xrightarrow[\begin{smallmatrix} -R_1+R_3 \end{smallmatrix}]{\begin{smallmatrix} R_1+R_2 \end{smallmatrix}} \begin{bmatrix} 1 & 1 & -1 & 3 \\ 0 & 5 & 4 & 1 \\ 0 & 5 & 4 & 1 \end{bmatrix} \xrightarrow[1/5R_2]{-R_2+R_3} \begin{bmatrix} 1 & 1 & -1 & 3 \\ 0 & 1 & 4/5 & 1/5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Hence  $\text{rank}(A) = 2$ .



## Theorem

*A system of  $m$  equations and  $n$  variables with coefficient matrix  $A$  and augmented matrix  $\bar{A}$*

- The system is inconsistent if and only if  $\text{rank}(A) \neq \text{rank}(\bar{A})$ .*

## Theorem

*A system of  $m$  equations and  $n$  variables with coefficient matrix  $A$  and augmented matrix  $\bar{A}$*

- The system is inconsistent if and only if  $\text{rank}(A) \neq \text{rank}(\bar{A})$ .*
- The system has unique solution if and only if  $\text{rank}(A) = \text{rank}(\bar{A}) = n$ .*

## Theorem

*A system of  $m$  equations and  $n$  variables with coefficient matrix  $A$  and augmented matrix  $\bar{A}$*

- *The system is inconsistent if and only if  $\text{rank}(A) \neq \text{rank}(\bar{A})$ .*
- *The system has unique solution if and only if  $\text{rank}(A) = \text{rank}(\bar{A}) = n$ .*
- *The system has infinite solution if and only if  $\text{rank}(A) = \text{rank}(\bar{A}) < n$ . In this cases, the system has  $n - \text{rank}(A)$  general solutions, the set of solutions described in parametric form including  $n - \text{rank}(A)$  parameters.*

## Theorem

*A system of  $m$  equations and  $n$  variables with coefficient matrix  $A$  and augmented matrix  $\bar{A}$*

- The system is inconsistent if and only if  $\text{rank}(A) \neq \text{rank}(\bar{A})$ .*
- The system has unique solution if and only if  $\text{rank}(A) = \text{rank}(\bar{A}) = n$ .*
- The system has infinite solution if and only if  $\text{rank}(A) = \text{rank}(\bar{A}) < n$ . In this cases, the system has  $n - \text{rank}(A)$  general solutions, the set of solutions described in parametric form including  $n - \text{rank}(A)$  parameters.*

## Example

Find all values of  $m$  so that the system

$$\begin{cases} x + y + 3z = 1 \\ -x + 3y + 2z = 3 \\ 3x - y + 4z = m \end{cases}$$

has infinitely many solutions.

**Answer:** This is a system of 3 equations and 3 variables.

## Example

Find all values of  $m$  so that the system

$$\begin{cases} x + y + 3z = 1 \\ -x + 3y + 2z = 3 \\ 3x - y + 4z = m \end{cases}$$

has infinitely many solutions.

**Answer:** This is a system of 3 equations and 3 variables.

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ -1 & 3 & 2 & 3 \\ 3 & -1 & 4 & m \end{array} \right] \xrightarrow[-3R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & -4 & -5 & m-3 \end{array} \right]$$

## Example

Find all values of  $m$  so that the system

$$\begin{cases} x + y + 3z = 1 \\ -x + 3y + 2z = 3 \\ 3x - y + 4z = m \end{cases}$$

has infinitely many solutions.

**Answer:** This is a system of 3 equations and 3 variables.

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ -1 & 3 & 2 & 3 \\ 3 & -1 & 4 & m \end{array} \right] \xrightarrow[-3R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & -4 & -5 & m-3 \end{array} \right]$$

$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & 0 & 0 & m+1 \end{array} \right]$$

## Example

Find all values of  $m$  so that the system

$$\begin{cases} x + y + 3z = 1 \\ -x + 3y + 2z = 3 \\ 3x - y + 4z = m \end{cases}$$

has infinitely many solutions.

**Answer:** This is a system of 3 equations and 3 variables.

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ -1 & 3 & 2 & 3 \\ 3 & -1 & 4 & m \end{array} \right] \xrightarrow[-3R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & -4 & -5 & m-3 \end{array} \right]$$

$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & 0 & 0 & m+1 \end{array} \right]$$

$\text{rank}(A) = 2 < 3$  (variables).

$\text{rank}(\bar{A}) = 2$  if  $m = -1$



## Example

Find all values of  $m$  so that the system

$$\begin{cases} x + y + 3z = 1 \\ -x + 3y + 2z = 3 \\ 3x - y + 4z = m \end{cases}$$

has infinitely many solutions.

**Answer:** This is a system of 3 equations and 3 variables.

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ -1 & 3 & 2 & 3 \\ 3 & -1 & 4 & m \end{array} \right] \xrightarrow[-3R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & -4 & -5 & m-3 \end{array} \right]$$

$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & 0 & 0 & m+1 \end{array} \right]$$

$\text{rank}(A) = 2 < 3$  (variables).

$\text{rank}(\bar{A}) = 2$  if  $m = -1$

$\text{rank}(\bar{A}) = 3$  if  $m \neq -1$ .

## Example

Find all values of  $m$  so that the system

$$\begin{cases} x + y + 3z = 1 \\ -x + 3y + 2z = 3 \\ 3x - y + 4z = m \end{cases}$$

has infinitely many solutions.

**Answer:** This is a system of 3 equations and 3 variables.

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ -1 & 3 & 2 & 3 \\ 3 & -1 & 4 & m \end{array} \right] \xrightarrow[-3R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & -4 & -5 & m-3 \end{array} \right]$$

$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & 0 & 0 & m+1 \end{array} \right]$$

$\text{rank}(A) = 2 < 3$  (variables).

$\text{rank}(\bar{A}) = 2$  if  $m = -1$

$\text{rank}(\bar{A}) = 3$  if  $m \neq -1$ .

The system has infinite solution if and only if  $m = -1$ .

## Example

Find all values of  $m$  so that the system

$$\begin{cases} x + y + 3z = 1 \\ -x + 3y + 2z = 3 \\ 3x - y + 4z = m \end{cases}$$

has infinitely many solutions.

**Answer:** This is a system of 3 equations and 3 variables.

$$\bar{A} = \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ -1 & 3 & 2 & 3 \\ 3 & -1 & 4 & m \end{array} \right] \xrightarrow[-3R_1+R_3]{R_1+R_2} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & -4 & -5 & m-3 \end{array} \right]$$

$$\xrightarrow{R_2+R_3} \left[ \begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & 0 & 0 & m+1 \end{array} \right]$$

$\text{rank}(A) = 2 < 3$  (variables).

$\text{rank}(\bar{A}) = 2$  if  $m = -1$

$\text{rank}(\bar{A}) = 3$  if  $m \neq -1$ .

The system has infinite solution if and only if  $m = -1$ .

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & -19 \\ 0 & 1 & 4 & 4 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

Answer:

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & -19 \\ 0 & 1 & 4 & 4 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

**Answer:** We can see  $\text{rank}(\bar{A}) = 3, \text{rank}(A) = 2$ .

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & -19 \\ 0 & 1 & 4 & 4 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

**Answer:** We can see  $\text{rank}(\bar{A}) = 3, \text{rank}(A) = 2$ . The system has no solution (inconsistent).

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & -19 \\ 0 & 1 & 4 & 4 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

**Answer:** We can see  $\text{rank}(\bar{A}) = 3, \text{rank}(A) = 2$ . The system has no solution (inconsistent).

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & 8 \\ 0 & 1 & 4 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Answer:



## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & 8 \\ 0 & 1 & 4 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

**Answer:** The system has 3 equations and 3 variables and

$$\text{rank}(A) = \text{rank}(\bar{A}) = 2 < 3.$$

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & 8 \\ 0 & 1 & 4 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

**Answer:** The system has 3 equations and 3 variables and

$$\text{rank}(A) = \text{rank}(\bar{A}) = 2 < 3.$$

The system has infinite solution:

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & 8 \\ 0 & 1 & 4 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

**Answer:** The system has 3 equations and 3 variables and

$$\text{rank}(A) = \text{rank}(\bar{A}) = 2 < 3.$$

The system has infinite solution:

$$\begin{cases} x_1 + 2x_2 - 3x_3 = 8 \\ x_2 + 4x_3 = -3 \end{cases} \Leftrightarrow \begin{cases} x_1 = 8 + 3x_3 - 2x_2 = 14 + 7s \\ x_2 = -3 - 4x_3 = -3 - 4s \\ x_3 = s \end{cases}$$

## Example

The augmented matrix is given for a system of equations. If the system is consistent, find the general solution. Otherwise state that there is no solution.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & 8 \\ 0 & 1 & 4 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

**Answer:** The system has 3 equations and 3 variables and

$$\text{rank}(A) = \text{rank}(\bar{A}) = 2 < 3.$$

The system has infinite solution:

$$\begin{cases} x_1 + 2x_2 - 3x_3 = 8 \\ x_2 + 4x_3 = -3 \end{cases} \Leftrightarrow \begin{cases} x_1 = 8 + 3x_3 - 2x_2 = 14 + 7s \\ x_2 = -3 - 4x_3 = -3 - 4s \\ x_3 = s \end{cases}$$

## 1.3 Homogeneous Systems

### Definition

A **homogeneous system** consists of  $m$  linear equations and  $n$  variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \dots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases} \quad (\star) \text{ where } a_{ij} \in \mathbb{R}.$$

### Example

$$\begin{cases} 3x_1 - x_2 + x_3 = 0 \\ 2x_1 + 4x_2 - 5x_3 = 0 \\ x_1 + 2x_2 - x_3 = 0 \end{cases}$$

## 1.3 Homogeneous Systems

### Definition

A **homogeneous system** consists of  $m$  linear equations and  $n$  variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \dots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases} \quad (*) \text{ where } a_{ij} \in \mathbb{R}.$$

### Example

$$\begin{cases} 3x_1 - x_2 + x_3 = 0 \\ 2x_1 + 4x_2 - 5x_3 = 0 \\ x_1 + 2x_2 - x_3 = 0 \end{cases} \quad \text{is a homogeneous system of 3 equations and 3 variables.}$$

## 1.3 Homogeneous Systems

### Definition

A **homogeneous system** consists of  $m$  linear equations and  $n$  variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \dots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases} \quad (\star) \text{ where } a_{ij} \in \mathbb{R}.$$

### Example

$$\begin{cases} 3x_1 - x_2 + x_3 = 0 \\ 2x_1 + 4x_2 - 5x_3 = 0 \\ x_1 + 2x_2 - x_3 = 0 \end{cases} \quad \text{is a homogeneous system of 3 equations and 3 variables.}$$

## Note.

1. A homogeneous system always has a **trivial solution**  
 $(x_1, x_2, \dots, x_n) = (0, 0, \dots, 0)$ .
2. If it has a **non-trivial solution**, then it has infinite solution.



## Note.

1. A homogeneous system always has a **trivial solution**  
 $(x_1, x_2, \dots, x_n) = (0, 0, \dots, 0)$ .
2. If it has a **non-trivial solution**, then it has infinite solution.

## Theorem

*Given a homogeneous solution of  $m$ -equation and  $n$ -variables with the coefficient matrix  $A_{m \times n}$ .*

## Note.

1. A homogeneous system always has a **trivial solution**  
 $(x_1, x_2, \dots, x_n) = (0, 0, \dots, 0)$ .
2. If it has a **non-trivial solution**, then it has infinite solution.

## Theorem

*Given a homogeneous system of  $m$ -equation and  $n$ -variables with the coefficient matrix  $A_{m \times n}$ .*

- *If  $\text{rank}(A) < n$ , then the system has infinite solutions (solutions can be formed under  $n - \text{rank}(A)$  **basic solutions**).*

## Note.

1. A homogeneous system always has a **trivial solution**  
 $(x_1, x_2, \dots, x_n) = (0, 0, \dots, 0)$ .
2. If it has a **non-trivial solution**, then it has infinite solution.

## Theorem

*Given a homogeneous solution of  $m$ -equation and  $n$ -variables with the coefficient matrix  $A_{m \times n}$ .*

- *If  $\text{rank}(A) < n$ , then the system has infinite solutions (solutions can be formed under  $n - \text{rank}(A)$  **basic solutions**).*
- *If  $\text{rank}(A) = n$ , then the system has only trivial solution.*

## Note.

1. A homogeneous system always has a **trivial solution**  
 $(x_1, x_2, \dots, x_n) = (0, 0, \dots, 0)$ .
2. If it has a **non-trivial solution**, then it has infinite solution.

## Theorem

*Given a homogeneous solution of  $m$ -equation and  $n$ -variables with the coefficient matrix  $A_{m \times n}$ .*

- *If  $\text{rank}(A) < n$ , then the system has infinite solutions (solutions can be formed under  $n - \text{rank}(A)$  **basic solutions**).*
- *If  $\text{rank}(A) = n$ , then the system has only trivial solution.*

## Note.

1. A homogeneous system always has a **trivial solution**  
 $(x_1, x_2, \dots, x_n) = (0, 0, \dots, 0)$ .
2. If it has a **non-trivial solution**, then it has infinite solution.

## Theorem

*Given a homogeneous solution of  $m$ -equation and  $n$ -variables with the coefficient matrix  $A_{m \times n}$ .*

- *If  $\text{rank}(A) < n$ , then the system has infinite solutions (solutions can be formed under  $n - \text{rank}(A)$  **basic solutions**).*
- *If  $\text{rank}(A) = n$ , then the system has only trivial solution.*

## Example

Solve the following system

$$\begin{array}{rrrrrr} x_1 & +2x_2 & -x_3 & +2x_4 & +x_5 & = 0 \\ x_1 & +2x_2 & +2x_3 & & +x_5 & = 0 \\ 2x_1 & +4x_2 & -2x_3 & +3x_4 & +x_5 & = 0 \end{array} .$$

The coefficient matrix

$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 1 & 2 & 2 & 0 & 1 \\ 2 & 4 & -2 & 3 & 1 \end{bmatrix} \xrightarrow{\substack{-R_1+R_2 \\ -2R_1+R_3}} \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & 3 & -2 & 0 \\ 0 & 0 & 0 & -1 & -1 \end{bmatrix} .$$

## Example

Solve the following system

$$\begin{array}{rrrrrr} x_1 & +2x_2 & -x_3 & +2x_4 & +x_5 & = 0 \\ x_1 & +2x_2 & +2x_3 & & +x_5 & = 0 \\ 2x_1 & +4x_2 & -2x_3 & +3x_4 & +x_5 & = 0 \end{array}$$

The coefficient matrix

$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 1 & 2 & 2 & 0 & 1 \\ 2 & 4 & -2 & 3 & 1 \end{bmatrix} \xrightarrow{\substack{-R_1+R_2 \\ -2R_1+R_3}} \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & 3 & -2 & 0 \\ 0 & 0 & 0 & -1 & -1 \end{bmatrix}.$$

$$\rightarrow \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & 1 & -2/3 & 0 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

$$\text{Rank}(A) = 3.$$

## Example

Solve the following system

$$\begin{array}{rrrrrr} x_1 & +2x_2 & -x_3 & +2x_4 & +x_5 & = 0 \\ x_1 & +2x_2 & +2x_3 & & +x_5 & = 0 \\ 2x_1 & +4x_2 & -2x_3 & +3x_4 & +x_5 & = 0 \end{array}$$

The coefficient matrix

$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 1 & 2 & 2 & 0 & 1 \\ 2 & 4 & -2 & 3 & 1 \end{bmatrix} \xrightarrow{\substack{-R_1+R_2 \\ -2R_1+R_3}} \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & 3 & -2 & 0 \\ 0 & 0 & 0 & -1 & -1 \end{bmatrix}.$$

$$\rightarrow \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & 1 & -2/3 & 0 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

$$\text{Rank}(A) = 3.$$

The system has infinite solution:

$$\begin{cases} x_1 = -2x_2 + x_3 - 2x_4 - x_5 \\ \phantom{x_1} = -2t + 1/3s \\ x_2 = t \\ x_3 = 2/3x_4 = -2/3s \\ x_4 = -x_5 = -s \\ x_5 = s \end{cases}$$



## Example

Solve the following system

$$\begin{array}{rrrrrr} x_1 & +2x_2 & -x_3 & +2x_4 & +x_5 & = 0 \\ x_1 & +2x_2 & +2x_3 & & +x_5 & = 0 \\ 2x_1 & +4x_2 & -2x_3 & +3x_4 & +x_5 & = 0 \end{array}$$

The coefficient matrix

$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 1 & 2 & 2 & 0 & 1 \\ 2 & 4 & -2 & 3 & 1 \end{bmatrix} \xrightarrow{\substack{-R_1+R_2 \\ -2R_1+R_3}} \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & 3 & -2 & 0 \\ 0 & 0 & 0 & -1 & -1 \end{bmatrix}.$$

$$\rightarrow \begin{bmatrix} 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & 1 & -2/3 & 0 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

$$\text{Rank}(A) = 3.$$

The system has infinite solution:

$$\begin{cases} x_1 = -2x_2 + x_3 - 2x_4 - x_5 \\ \quad = -2t + 1/3s \\ x_2 = t \\ x_3 = 2/3x_4 = -2/3s \\ x_4 = -x_5 = -s \\ x_5 = s \end{cases}$$

## Example

Find all values of  $a$  such that the homogeneous system

$$\begin{cases} x - 2y + z = 0 \\ x - y + 3z = 0 \\ 2x + ay + 4z = 0 \end{cases}$$

has only the trivial solution.

**Answer:** The coefficient matrix

$$\begin{aligned} A &= \begin{bmatrix} 1 & -2 & 1 \\ 1 & -1 & 3 \\ 2 & a & 4 \end{bmatrix} \xrightarrow[-2R_1+R_3]{-R_1+R_2} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 2 \\ 0 & a+4 & 2 \end{bmatrix} \\ &\xrightarrow{-(a+4)R_2+R_3} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 2-2(a+4) \end{bmatrix}. \end{aligned}$$

We need  $\text{rank}(A) = 3$ . This implies that  $2 - 2(a + 4) \neq 0$ , i.e.,  $a \neq -3$ .

# Exercises

1.1 : 10, 14, 18, 19 (page 8, 9).

1.2 : 1, 2a, 3, 4d, 5ace, 7ab, 8ad, 9a, 11acd, 12 (page 17-19).

1.3 : 1, 2ac, 3bc, 5ab (page 25, 26).