

# Discrete Mathematics

## Chapter 2: Basic Structures: Sets, Functions, Sequences and Sums

Department of Mathematics  
The FPT university

# 2.1 Sets

## 2.1 Sets

### Definition

## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.

## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.

## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.
- The **cardinality** of the set  $A$  is the number of distinct elements of  $A$ , denoted by  $|A|$ .

## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.
- The **cardinality** of the set  $A$  is the number of distinct elements of  $A$ , denoted by  $|A|$ .
- The **empty** set, denoted by  $\emptyset$ , is the set whose cardinality is 0.

## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.
- The **cardinality** of the set  $A$  is the number of distinct elements of  $A$ , denoted by  $|A|$ .
- The **empty** set, denoted by  $\emptyset$ , is the set whose cardinality is 0.

**Example.**



## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.
- The **cardinality** of the set  $A$  is the number of distinct elements of  $A$ , denoted by  $|A|$ .
- The **empty** set, denoted by  $\emptyset$ , is the set whose cardinality is 0.

### Example.

- The set  $\{a, cat, catches, a, mouse\}$  has

## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.
- The **cardinality** of the set  $A$  is the number of distinct elements of  $A$ , denoted by  $|A|$ .
- The **empty** set, denoted by  $\emptyset$ , is the set whose cardinality is 0.

### Example.

- The set  $\{a, cat, catches, a, mouse\}$  has 4 elements.

## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.
- The **cardinality** of the set  $A$  is the number of distinct elements of  $A$ , denoted by  $|A|$ .
- The **empty** set, denoted by  $\emptyset$ , is the set whose cardinality is 0.

### Example.

- The set  $\{a, cat, catches, a, mouse\}$  has 4 elements.
- The set  $\{a, b, \{a, b\}, c, \{a, b, c, \}, \emptyset\}$  has


## 2.1 Sets

### Definition

- A **set** is an unordered collection of objects.
- An object of a set is called an **element**, or a **member**, of that set.
- The **cardinality** of the set  $A$  is the number of distinct elements of  $A$ , denoted by  $|A|$ .
- The **empty** set, denoted by  $\emptyset$ , is the set whose cardinality is 0.

### Example.

- The set  $\{a, cat, catches, a, mouse\}$  has 4 elements.
- The set  $\{a, b, \{a, b\}, c, \{a, b, c, \}, \emptyset\}$  has 6 elements



- If  $x$  is an element of  $A$  we write  $x \in A$ .

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .



- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\}$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\}$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\}$$



- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\}$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\}$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\} \quad (\text{T})$$

$$\emptyset \in \{\emptyset\}$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\emptyset \in \{\emptyset\} \quad (\text{T})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \text{ (T)}$$

$$x \subseteq \{x\} \text{ (F)}$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \text{ (T)}$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \text{ (T)}$$

$$\emptyset \subseteq \{\emptyset\} \text{ (T)}$$

$$\emptyset \in \{\emptyset\} \text{ (T)}$$

$$\{a, b, c\} \subseteq \{a, b, c\}$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\} \quad (\text{T})$$

$$\emptyset \in \{\emptyset\} \quad (\text{T})$$

$$\{a, b, c\} \subseteq \{a, b, c\} \quad (\text{T})$$



- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\} \quad (\text{T})$$

$$\emptyset \in \{\emptyset\} \quad (\text{T})$$

$$\{a, b, c\} \subseteq \{a, b, c\} \quad (\text{T})$$

$$\{a, b, c\} \in \{a, b, c\}$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\} \quad (\text{T})$$

$$\emptyset \in \{\emptyset\} \quad (\text{T})$$

$$\{a, b, c\} \subseteq \{a, b, c\} \quad (\text{T})$$

$$\{a, b, c\} \in \{a, b, c\} \quad (\text{F})$$

- If  $x$  is an element of  $A$  we write  $x \in A$ . If  $x$  is not an element of  $A$  we write  $x \notin A$ .
- If all elements of  $A$  are also elements of  $B$  we write  $A \subseteq B$ , and  $A$  is called a **subset** of  $B$ .
- If  $A$  is a **proper** subset of  $B$ , meaning  $A \subseteq B$  and  $A \neq B$ , we write  $A \subset B$ .
- The empty set  $\emptyset$  is a subset of any set.

**Example.** Which statements are true?

$$x \in \{x\} \quad (\text{T})$$

$$x \subseteq \{x\} \quad (\text{F})$$

$$\{a, b\} \subseteq \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\{a, b\} \in \{a, b, \{a, b\}, c\} \quad (\text{T})$$

$$\emptyset \subseteq \{\emptyset\} \quad (\text{T})$$

$$\emptyset \in \{\emptyset\} \quad (\text{T})$$

$$\{a, b, c\} \subseteq \{a, b, c\} \quad (\text{T})$$

$$\{a, b, c\} \in \{a, b, c\} \quad (\text{F})$$



## Definitions

## Definitions

- The **Cartesian product** of two sets  $A$  and  $B$ , denoted by  $A \times B$ , is the set of all ordered pairs  $(a, b)$  where  $a \in A$  and  $b \in B$ .

## Definitions

- The **Cartesian product** of two sets  $A$  and  $B$ , denoted by  $A \times B$ , is the set of all ordered pairs  $(a, b)$  where  $a \in A$  and  $b \in B$ .
- The **power set** of the set  $A$ , denoted by  $P(A)$ , is the set of all subsets of  $A$ .

## Definitions

- The **Cartesian product** of two sets  $A$  and  $B$ , denoted by  $A \times B$ , is the set of all ordered pairs  $(a, b)$  where  $a \in A$  and  $b \in B$ .
- The **power set** of the set  $A$ , denoted by  $P(A)$ , is the set of all subsets of  $A$ .

**Note.**



## Definitions

- The **Cartesian product** of two sets  $A$  and  $B$ , denoted by  $A \times B$ , is the set of all ordered pairs  $(a, b)$  where  $a \in A$  and  $b \in B$ .
- The **power set** of the set  $A$ , denoted by  $P(A)$ , is the set of all subsets of  $A$ .

### Note.

If  $|A| = m$  and  $|B| = n$  then  $|A \times B| = mn$ .

## Definitions

- The **Cartesian product** of two sets  $A$  and  $B$ , denoted by  $A \times B$ , is the set of all ordered pairs  $(a, b)$  where  $a \in A$  and  $b \in B$ .
- The **power set** of the set  $A$ , denoted by  $P(A)$ , is the set of all subsets of  $A$ .

### Note.

If  $|A| = m$  and  $|B| = n$  then  $|A \times B| = mn$ .

If  $|A| = n$  then  $|P(A)| = 2^n$ .

## 2.2 Set Operations

## 2.2 Set Operations

Let  $A$  and  $B$  be two sets.

## 2.2 Set Operations

Let  $A$  and  $B$  be two sets.

- **Union** of  $A$  and  $B$ :  $A \cup B = \{x \mid (x \in A) \vee (x \in B)\}$

## 2.2 Set Operations

Let  $A$  and  $B$  be two sets.

- **Union** of  $A$  and  $B$ :  $A \cup B = \{x \mid (x \in A) \vee (x \in B)\}$
- **Intersection** of  $A$  and  $B$ :  $A \cap B = \{x \mid (x \in A) \wedge (x \in B)\}$

## 2.2 Set Operations

Let  $A$  and  $B$  be two sets.

- **Union** of  $A$  and  $B$ :  $A \cup B = \{x \mid (x \in A) \vee (x \in B)\}$
- **Intersection** of  $A$  and  $B$ :  $A \cap B = \{x \mid (x \in A) \wedge (x \in B)\}$
- **Difference** of  $A$  and  $B$ :  $A - B = \{x \mid (x \in A) \wedge (x \notin B)\}$

## 2.2 Set Operations

Let  $A$  and  $B$  be two sets.

- **Union** of  $A$  and  $B$ :  $A \cup B = \{x \mid (x \in A) \vee (x \in B)\}$
- **Intersection** of  $A$  and  $B$ :  $A \cap B = \{x \mid (x \in A) \wedge (x \in B)\}$
- **Difference** of  $A$  and  $B$ :  $A - B = \{x \mid (x \in A) \wedge (x \notin B)\}$
- **Symmetric difference** of  $A$  and  $B$ :  $A \oplus B = \{x \mid (x \in A) \oplus (x \in B)\}$



## 2.2 Set Operations

Let  $A$  and  $B$  be two sets.

- **Union** of  $A$  and  $B$ :  $A \cup B = \{x \mid (x \in A) \vee (x \in B)\}$
- **Intersection** of  $A$  and  $B$ :  $A \cap B = \{x \mid (x \in A) \wedge (x \in B)\}$
- **Difference** of  $A$  and  $B$ :  $A - B = \{x \mid (x \in A) \wedge (x \notin B)\}$
- **Symmetric difference** of  $A$  and  $B$ :  $A \oplus B = \{x \mid (x \in A) \oplus (x \in B)\}$
- **Complement** of  $A$  with respect to the universal set  $U$ :  $\overline{A} = U - A$

# Set Identities

# Set Identities

Name	Identity

# Set Identities

Name	Identity
Complementation law	$\overline{(\overline{A})} = A$

# Set Identities

Name	Identity
Complementation law	$\overline{(\overline{A})} = A$
Identity laws	$A \cup \emptyset = A, A \cap U = A$

# Set Identities

Name	Identity
Complementation law	$\overline{\overline{A}} = A$
Identity laws	$A \cup \emptyset = A, \quad A \cap U = A$
Domination laws	$A \cup U = U, \quad A \cap \emptyset = \emptyset$

# Set Identities

Name	Identity
Complementation law	$\overline{\overline{A}} = A$
Identity laws	$A \cup \emptyset = A, A \cap U = A$
Domination laws	$A \cup U = U, A \cap \emptyset = \emptyset$
Complement laws	$A \cup \overline{A} = U, A \cap \overline{A} = \emptyset$

# Set Identities

Name	Identity
Complementation law	$\overline{(\overline{A})} = A$
Identity laws	$A \cup \emptyset = A, A \cap U = A$
Domination laws	$A \cup U = U, A \cap \emptyset = \emptyset$
Complement laws	$A \cup \overline{A} = U, A \cap \overline{A} = \emptyset$
Idempotent laws	$A \cup A = A, A \cap A = A$



# Set Identities

Name	Identity
Complementation law	$\overline{\overline{A}} = A$
Identity laws	$A \cup \emptyset = A, A \cap U = A$
Domination laws	$A \cup U = U, A \cap \emptyset = \emptyset$
Complement laws	$A \cup \overline{A} = U, A \cap \overline{A} = \emptyset$
Idempotent laws	$A \cup A = A, A \cap A = A$
Commutative laws	$A \cup B = B \cup A, A \cap B = B \cap A$

# Set Identities

Name	Identity
Complementation law	$\overline{(\overline{A})} = A$
Identity laws	$A \cup \emptyset = A, A \cap U = A$
Domination laws	$A \cup U = U, A \cap \emptyset = \emptyset$
Complement laws	$A \cup \overline{A} = U, A \cap \overline{A} = \emptyset$
Idempotent laws	$A \cup A = A, A \cap A = A$
Commutative laws	$A \cup B = B \cup A, A \cap B = B \cap A$
Associative laws	$(A \cup B) \cup C = A \cup (B \cup C)$ $(A \cap B) \cap C = A \cap (B \cap C)$

# Set Identities

Name	Identity
Complementation law	$\overline{(\overline{A})} = A$
Identity laws	$A \cup \emptyset = A, A \cap U = A$
Domination laws	$A \cup U = U, A \cap \emptyset = \emptyset$
Complement laws	$A \cup \overline{A} = U, A \cap \overline{A} = \emptyset$
Idempotent laws	$A \cup A = A, A \cap A = A$
Commutative laws	$A \cup B = B \cup A, A \cap B = B \cap A$
Associative laws	$(A \cup B) \cup C = A \cup (B \cup C)$ $(A \cap B) \cap C = A \cap (B \cap C)$
Distributive laws	$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

# Set Identities

Name	Identity
Complementation law	$\overline{(\overline{A})} = A$
Identity laws	$A \cup \emptyset = A, A \cap U = A$
Domination laws	$A \cup U = U, A \cap \emptyset = \emptyset$
Complement laws	$A \cup \overline{A} = U, A \cap \overline{A} = \emptyset$
Idempotent laws	$A \cup A = A, A \cap A = A$
Commutative laws	$A \cup B = B \cup A, A \cap B = B \cap A$
Associative laws	$(A \cup B) \cup C = A \cup (B \cup C)$ $(A \cap B) \cap C = A \cap (B \cap C)$
Distributive laws	$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
De Morgan's laws	$\overline{A \cup B} = \overline{A} \cap \overline{B}$ $\overline{A \cap B} = \overline{A} \cup \overline{B}$

Two methods to show the equality of two sets.

Two methods to show the equality of two sets.

- Show that each set is a subset of the other.

Two methods to show the equality of two sets.

- Show that each set is a subset of the other.
- Use Membership table, similar to the method of using truth table to establish propositional equivalences.

# Computer Representation of Sets



# Computer Representation of Sets

Let  $U$  be a universal set. Fix an ordering of elements of  $U$  as  $a_1, a_2, \dots, a_n$ .

# Computer Representation of Sets

Let  $U$  be a universal set. Fix an ordering of elements of  $U$  as  $a_1, a_2, \dots, a_n$ .

If  $A$  is a subset of  $U$ , represent  $A$  with a bit string of length  $n$ , where the  $i$ th bit is 1 if  $a_i$  is in  $A$ , and is 0 if  $a_i$  is not in  $A$ .

# Computer Representation of Sets

Let  $U$  be a universal set. Fix an ordering of elements of  $U$  as  $a_1, a_2, \dots, a_n$ .

If  $A$  is a subset of  $U$ , represent  $A$  with a bit string of length  $n$ , where the  $i$ th bit is 1 if  $a_i$  is in  $A$ , and is 0 if  $a_i$  is not in  $A$ .

**Example.**

# Computer Representation of Sets

Let  $U$  be a universal set. Fix an ordering of elements of  $U$  as  $a_1, a_2, \dots, a_n$ .

If  $A$  is a subset of  $U$ , represent  $A$  with a bit string of length  $n$ , where the  $i$ th bit is 1 if  $a_i$  is in  $A$ , and is 0 if  $a_i$  is not in  $A$ .

**Example.** Let  $U = \{1, 2, 3, 4, 5, 6, 7, 8\}$ .

# Computer Representation of Sets

Let  $U$  be a universal set. Fix an ordering of elements of  $U$  as  $a_1, a_2, \dots, a_n$ .


If  $A$  is a subset of  $U$ , represent  $A$  with a bit string of length  $n$ , where the  $i$ th bit is 1 if  $a_i$  is in  $A$ , and is 0 if  $a_i$  is not in  $A$ .

**Example.** Let  $U = \{1, 2, 3, 4, 5, 6, 7, 8\}$ . Then the subset  $A = \{1, 3, 4, 6\}$  is represented as the bit string 10110100.

## 2.3 Functions


## 2.3 Functions

## 2.3 Functions






## 2.3 Functions




## 2.3 Functions




The set  $A$  is called **domain** and  $B$  is called **codomain** of  $f$ .

## 2.3 Functions



The set  $A$  is called **domain** and  $B$  is called **codomain** of  $f$ .  
If  $f(a) = b$  we say  $b$  is the **image** of  $a$  and  $a$  is a **preimage** of  $b$ .

## 2.3 Functions




The set  $A$  is called **domain** and  $B$  is called **codomain** of  $f$ .  
If  $f(a) = b$  we say  $b$  is the **image** of  $a$  and  $a$  is a **preimage** of  $b$ .

Let  $S$  be a subset of  $A$ . The set

$$f(S) = \{b \in B \mid \exists a \in A (f(a) = b)\}$$

is called the image of  $S$ ,

## 2.3 Functions



The set  $A$  is called **domain** and  $B$  is called **codomain** of  $f$ .  
If  $f(a) = b$  we say  $b$  is the **image** of  $a$  and  $a$  is a **preimage** of  $b$ .

Let  $S$  be a subset of  $A$ . The set


$$f(S) = \{b \in B \mid \exists a \in A (f(a) = b)\}$$

is called the image of  $S$ , and the set

$$f^{-1}(S) = \{a \in A \mid f(a) \in S\}$$

is called the preimage of  $S$ .

## 2.3 Functions



The set  $A$  is called **domain** and  $B$  is called **codomain** of  $f$ .  
If  $f(a) = b$  we say  $b$  is the **image** of  $a$  and  $a$  is a **preimage** of  $b$ .

Let  $S$  be a subset of  $A$ . The set

$$f(S) = \{b \in B \mid \exists a \in A (f(a) = b)\}$$

is called the image of  $S$ , and the set

$$f^{-1}(S) = \{a \in A \mid f(a) \in S\}$$

is called the preimage of  $S$ . The set  $f(A)$  is called the **range** of  $f$ .

# Some Important Functions

# Some Important Functions

- Floor function:  $\lfloor x \rfloor$



# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor =$  the greatest integer that is not greater than  $x$ .

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

**Note.** For all real numbers  $x$  we have

$$x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

**Note.** For all real numbers  $x$  we have

$$x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

**Example.** Which statements are true for all real numbers  $x, y$  and all integers  $n$ ?

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

**Note.** For all real numbers  $x$  we have

$$x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

**Example.** Which statements are true for all real numbers  $x, y$  and all integers  $n$ ?

$$\lceil x + y \rceil = \lceil x \rceil + \lceil y \rceil$$

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

**Note.** For all real numbers  $x$  we have

$$x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

**Example.** Which statements are true for all real numbers  $x, y$  and all integers  $n$ ?

$$\lceil x + y \rceil = \lceil x \rceil + \lceil y \rceil$$

$$\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$$

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

**Note.** For all real numbers  $x$  we have

$$x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

**Example.** Which statements are true for all real numbers  $x, y$  and all integers  $n$ ?

$$\lceil x + y \rceil = \lceil x \rceil + \lceil y \rceil$$

$$\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$$

$$\lceil x + n \rceil = \lceil x \rceil + n$$



# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

**Note.** For all real numbers  $x$  we have

$$x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

**Example.** Which statements are true for all real numbers  $x, y$  and all integers  $n$ ?

$$\lceil x + y \rceil = \lceil x \rceil + \lceil y \rceil$$

$$\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$$

$$\lceil x + n \rceil = \lceil x \rceil + n$$

$$\lfloor x + n \rfloor = \lfloor x \rfloor + n$$

# Some Important Functions

- **Floor** function:  $\lfloor x \rfloor$  = the greatest integer that is not greater than  $x$ .
- **Ceiling** function:  $\lceil x \rceil$  = the smallest integer that is not smaller than  $x$ .

**Note.** For all real numbers  $x$  we have

$$x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

**Example.** Which statements are true for all real numbers  $x, y$  and all integers  $n$ ?

$$\lceil x + y \rceil = \lceil x \rceil + \lceil y \rceil$$

$$\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$$

$$\lceil x + n \rceil = \lceil x \rceil + n$$

$$\lfloor x + n \rfloor = \lfloor x \rfloor + n$$


# One-to-One, Onto, and Bijection

# One-to-One, Onto, and Bijection

The function  $f : A \rightarrow B$  is **one-to-one** if  $f(a_1) \neq f(a_2)$  for all  $a_1 \neq a_2$  in  $A$ .

# One-to-One, Onto, and Bijection

The function  $f : A \rightarrow B$  is **one-to-one** if  $f(a_1) \neq f(a_2)$  for all  $a_1 \neq a_2$  in  $A$ .





**Example.** Which functions are one-to-one?

**Example.** Which functions are one-to-one?

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$



**Example.** Which functions are one-to-one?

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R}^+ \rightarrow \mathbb{R}; \quad f(x) = x^2$

**Example.** Which functions are one-to-one?

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R}^+ \rightarrow \mathbb{R}; \quad f(x) = x^2$

(c)  $f : \mathbb{Z} \rightarrow \mathbb{Z}; \quad f(n) = \lfloor \frac{n+1}{2} \rfloor$

**Example.** Which functions are one-to-one?

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R}^+ \rightarrow \mathbb{R}; \quad f(x) = x^2$

(c)  $f : \mathbb{Z} \rightarrow \mathbb{Z}; \quad f(n) = \lfloor \frac{n+1}{2} \rfloor$


(d)  $f : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}; \quad f(m, n) = m + n$



The function  $f : A \rightarrow B$  is **onto** if for each  $b$  in  $B$  there is  $a$  in  $A$  such that  $f(a) = b$ .

The function  $f : A \rightarrow B$  is **onto** if for each  $b$  in  $B$  there is  $a$  in  $A$  such that  $f(a) = b$ . In other words, the function  $f : A \rightarrow B$  is onto if  $f(A) = B$ .

The function  $f : A \rightarrow B$  is **onto** if for each  $b$  in  $B$  there is  $a$  in  $A$  such that  $f(a) = b$ . In other words, the function  $f : A \rightarrow B$  is onto if  $f(A) = B$ .







**Example.** Which functions are onto:

**Example.** Which functions are onto:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

**Example.** Which functions are onto:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

**Example.** Which functions are onto:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

(c)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = 2\lfloor x \rfloor$

**Example.** Which functions are onto:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

(c)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = 2\lfloor x \rfloor$

(d)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = \lfloor 2x \rfloor$

**Example.** Which functions are onto:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

(c)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = 2\lfloor x \rfloor$

(d)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = \lfloor 2x \rfloor$

(e)  $f : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}; \quad f(m, n) = m + n$




The function  $f : A \rightarrow B$  is a **bijection** (or **one-to-one correspondence**) if it is both one-to-one and onto.



The function  $f : A \rightarrow B$  is a **bijection** (or **one-to-one correspondence**) if it is both one-to-one and onto.

The function  $f : A \rightarrow B$  is a **bijection** (or **one-to-one correspondence**) if it is both one-to-one and onto.





**Example 1.** Which functions are bijection:

**Example 1.** Which functions are bijection:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

**Example 1.** Which functions are bijection:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

**Example 1.** Which functions are bijection:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

(c)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = \lfloor 2x \rfloor$

**Example 1.** Which functions are bijection:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

(c)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = \lfloor 2x \rfloor$

(d)  $f : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}; \quad f(m, n) = m + n$



**Example 1.** Which functions are bijection:

(a)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2$

(b)  $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^3$

(c)  $f : \mathbb{R} \rightarrow \mathbb{Z}; \quad f(x) = \lfloor 2x \rfloor$

(d)  $f : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}; \quad f(m, n) = m + n$

(e)  $f : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z} \times \mathbb{Z}; \quad f(m, n) = (m, m + n)$



**Example 2.** Does there exist:

**Example 2.** Does there exist:

- (a) A bijection /one-to-one/onto function from a set of 7 elements to a set of 5 elements?

**Example 2.** Does there exist:

- (a) A bijection /one-to-one/onto function from a set of 7 elements to a set of 5 elements? From a set of 5 elements to a set of 7 elements?

**Example 2.** Does there exist:

- (a) A bijection /one-to-one/onto function from a set of 7 elements to a set of 5 elements? From a set of 5 elements to a set of 7 elements?
- (b) A bijection from the set of even integers to the set of odd integers?

**Example 2.** Does there exist:

- (a) A bijection /one-to-one/onto function from a set of 7 elements to a set of 5 elements? From a set of 5 elements to a set of 7 elements?
- (b) A bijection from the set of even integers to the set of odd integers?
- (c) A bijection from the set of odd integers to the set of all integers?

**Example 2.** Does there exist:


- (a) A bijection /one-to-one/onto function from a set of 7 elements to a set of 5 elements? From a set of 5 elements to a set of 7 elements?
- (b) A bijection from the set of even integers to the set of odd integers?
- (c) A bijection from the set of odd integers to the set of all integers?
- (d) A bijection from the set of all real numbers to the set of positive real numbers?




# Compositions and Inverses

## Composition

## Composition




## Composition






# Inverse

# Inverse




## Inverse



**Note.**



## Inverse



**Note.** The function  $f : A \rightarrow B$  has an inverse if and only if  $f$  is a bijection.

## 2.4 Sequences and Summations

## 2.4 Sequences and Summations

### Sequences

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list.

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

**Example.**

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

**Example.** Find a general formula for  $a_n$  of each sequence:

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

**Example.** Find a general formula for  $a_n$  of each sequence:

(a)  $\frac{1}{2}, -\frac{2}{3}, \frac{3}{4}, -\frac{4}{5}, \dots$



## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

**Example.** Find a general formula for  $a_n$  of each sequence:

(a)  $\frac{1}{2}, -\frac{2}{3}, \frac{3}{4}, -\frac{4}{5}, \dots$

(b)  $-2, 1, 4, 7, 10, \dots$

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

**Example.** Find a general formula for  $a_n$  of each sequence:

(a)  $\frac{1}{2}, -\frac{2}{3}, \frac{3}{4}, -\frac{4}{5}, \dots$

(b)  $-2, 1, 4, 7, 10, \dots$  (an arithmetic progression)

(c)  $1, 2, 2, 3, 3, 3, 4, 4, 4, 4, \dots$

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

**Example.** Find a general formula for  $a_n$  of each sequence:

- (a)  $\frac{1}{2}, -\frac{2}{3}, \frac{3}{4}, -\frac{4}{5}, \dots$
- (b)  $-2, 1, 4, 7, 10, \dots$  (an arithmetic progression)
- (c)  $1, 2, 2, 3, 3, 3, 4, 4, 4, 4, \dots$
- (d)  $1, 1, 2, 3, 5, 8, 13, 21, \dots$

## 2.4 Sequences and Summations

### Sequences

Sequence is a discrete structure used to represent an order list. It is usually denoted as  $\{a_1, a_2, \dots\} = \{a_n, n = 1, 2, \dots\}$ .

**Example.** Find a general formula for  $a_n$  of each sequence:

- (a)  $\frac{1}{2}, -\frac{2}{3}, \frac{3}{4}, -\frac{4}{5}, \dots$
- (b)  $-2, 1, 4, 7, 10, \dots$  (an arithmetic progression)
- (c)  $1, 2, 2, 3, 3, 3, 4, 4, 4, 4, \dots$
- (d)  $1, 1, 2, 3, 5, 8, 13, 21, \dots$  (Fibonacci sequence)

# Summations

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ .

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

$$\sum_{i=1}^{100} a_i = a_1 + a_2 + \cdots + a_{100}$$



# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

$$\sum_{i=1}^{100} a_i = a_1 + a_2 + \cdots + a_{100}$$

$$\sum_{i=1}^{100} a_{2i+1} = a_3 + a_5 + \cdots + a_{201}$$

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

$$\sum_{i=1}^{100} a_i = a_1 + a_2 + \cdots + a_{100}$$

$$\sum_{i=1}^{100} a_{2i+1} = a_3 + a_5 + \cdots + a_{201}$$

$$\sum_{i=1}^{100} 1 = 1 + 1 + \cdots + 1 \text{ ( 100 terms )}$$

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

$$\sum_{i=1}^{100} a_i = a_1 + a_2 + \cdots + a_{100}$$

$$\sum_{i=1}^{100} a_{2i+1} = a_3 + a_5 + \cdots + a_{201}$$

$$\sum_{i=1}^{100} 1 = 1 + 1 + \cdots + 1 \text{ ( 100 terms )}$$

## Properties

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

$$\sum_{i=1}^{100} a_i = a_1 + a_2 + \dots + a_{100}$$

$$\sum_{i=1}^{100} a_{2i+1} = a_3 + a_5 + \dots + a_{201}$$

$$\sum_{i=1}^{100} 1 = 1 + 1 + \dots + 1 \text{ ( 100 terms )}$$

## Properties

$$\sum_{i=1}^n (a_i + b_i) = \sum_{i=1}^n a_i + \sum_{i=1}^n b_i$$

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

$$\sum_{i=1}^{100} a_i = a_1 + a_2 + \dots + a_{100}$$

$$\sum_{i=1}^{100} a_{2i+1} = a_3 + a_5 + \dots + a_{201}$$

$$\sum_{i=1}^{100} 1 = 1 + 1 + \dots + 1 \text{ ( 100 terms )}$$

## Properties

$$\sum_{i=1}^n (a_i + b_i) = \sum_{i=1}^n a_i + \sum_{i=1}^n b_i \qquad \sum_{i=1}^n k a_i = k \sum_{i=1}^n a_i$$

# Summations

Given a sequence  $\{a_n, n = 1, 2, \dots\}$ . The summation notation,  $\sum$ , is used as follows:

$$\sum_{i=1}^{100} a_i = a_1 + a_2 + \dots + a_{100}$$

$$\sum_{i=1}^{100} a_{2i+1} = a_3 + a_5 + \dots + a_{201}$$

$$\sum_{i=1}^{100} 1 = 1 + 1 + \dots + 1 \text{ ( 100 terms )}$$

## Properties

$$\sum_{i=1}^n (a_i + b_i) = \sum_{i=1}^n a_i + \sum_{i=1}^n b_i \qquad \sum_{i=1}^n ka_i = k \sum_{i=1}^n a_i$$

# Some Important Sums

# Some Important Sums

$$\sum_{i=1}^n i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$



# Some Important Sums

$$\sum_{i=1}^n i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

# Some Important Sums

$$\sum_{i=1}^n i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = 1^3 + 2^3 + \cdots + n^3 = \left( \frac{n(n+1)}{2} \right)^2$$

# Some Important Sums

$$\sum_{i=1}^n i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = 1^3 + 2^3 + \cdots + n^3 = \left( \frac{n(n+1)}{2} \right)^2$$

$$\sum_{i=0}^n r^i = 1 + r + r^2 + \cdots + r^n = \frac{1 - r^{n+1}}{1 - r}$$



## Example 1.

**Example 1.** Find:

**Example 1.** Find:

(a) 
$$\sum_{i=1}^{100} \frac{3^i}{4^{i+1}}$$

**Example 1.** Find:

(a)  $\sum_{i=1}^{100} \frac{3^i}{4^{i+1}}$

(b)  $\sum_{i=1}^{100} \frac{1}{i(i+1)}$  (Telescoping sum)

**Example 2.**



**Example 1.** Find:

(a)  $\sum_{i=1}^{100} \frac{3^i}{4^{i+1}}$

(b)  $\sum_{i=1}^{100} \frac{1}{i(i+1)}$  (Telescoping sum)

**Example 2.** Find double summations:

**Example 1.** Find:

(a)  $\sum_{i=1}^{100} \frac{3^i}{4^{i+1}}$

(b)  $\sum_{i=1}^{100} \frac{1}{i(i+1)}$  (Telescoping sum)

**Example 2.** Find double summations:

(a)  $\sum_{i=1}^2 \sum_{j=0}^2 (i + 2j)$

**Example 1.** Find:

$$(a) \sum_{i=1}^{100} \frac{3^i}{4^{i+1}}$$

$$(b) \sum_{i=1}^{100} \frac{1}{i(i+1)} \text{ (Telescoping sum)}$$

**Example 2.** Find double summations:

$$(a) \sum_{i=1}^2 \sum_{j=0}^2 (i + 2j)$$

$$(b) \sum_{i=1}^{10} \sum_{j=1}^{100} ij$$