

Discrete Mathematics

Chapter 9: Graphs

Department of Mathematics
The FPT university

Chapter 9: Introduction

Chapter 9: Introduction

Topics covered:

Chapter 9: Introduction

Topics covered:

9.1 Graphs and Graph Models

Chapter 9: Introduction

Topics covered:

9.1 Graphs and Graph Models

9.2 Graph Terminologies and Special Types of Graphs

Topics covered:

9.1 Graphs and Graph Models

9.2 Graph Terminologies and Special Types of Graphs

9.3 Representing Graphs and Graph Isomorphism

Topics covered:

9.1 Graphs and Graph Models

9.2 Graph Terminologies and Special Types of Graphs

9.3 Representing Graphs and Graph Isomorphism

9.4 Connectivity

Chapter 9: Introduction

Topics covered:

- 9.1 Graphs and Graph Models
- 9.2 Graph Terminologies and Special Types of Graphs
- 9.3 Representing Graphs and Graph Isomorphism
- 9.4 Connectivity
- 9.5 Euler and Hamilton Paths


Chapter 9: Introduction

Topics covered:

- 9.1 Graphs and Graph Models
- 9.2 Graph Terminologies and Special Types of Graphs
- 9.3 Representing Graphs and Graph Isomorphism
- 9.4 Connectivity
- 9.5 Euler and Hamilton Paths
- 9.6 Shortest-Path Problem

9.1 Graphs and Graph Models

9.1 Graphs and Graph Models



9.1 Graphs and Graph Models

A **graph** $G = (V, E)$ consists of V a non-empty set of **vertices** (or **nodes**) and E , a set of **edges**. Each edge has one or two vertices associated with it, called **endpoints**.

9.1 Graphs and Graph Models

A **graph** $G = (V, E)$ consists of V a non-empty set of **vertices** (or **nodes**) and E , a set of **edges**. Each edge has one or two vertices associated with it, called **endpoints**.

Graph classification:

9.1 Graphs and Graph Models

A **graph** $G = (V, E)$ consists of V a non-empty set of **vertices** (or **nodes**) and E , a set of **edges**. Each edge has one or two vertices associated with it, called **endpoints**.

Graph classification:

- **Undirected graphs:**

9.1 Graphs and Graph Models

A **graph** $G = (V, E)$ consists of V a non-empty set of **vertices** (or **nodes**) and E , a set of **edges**. Each edge has one or two vertices associated with it, called **endpoints**.

Graph classification:

- **Undirected graphs:** Simple graphs, Multigraphs and Pseudographs

9.1 Graphs and Graph Models

A **graph** $G = (V, E)$ consists of V a non-empty set of **vertices** (or **nodes**) and E , a set of **edges**. Each edge has one or two vertices associated with it, called **endpoints**.

Graph classification:

- **Undirected graphs:** Simple graphs, Multigraphs and Pseudographs
- **Directed graphs:**

9.1 Graphs and Graph Models

A **graph** $G = (V, E)$ consists of V a non-empty set of **vertices** (or **nodes**) and E , a set of **edges**. Each edge has one or two vertices associated with it, called **endpoints**.

Graph classification:

- **Undirected graphs:** Simple graphs, Multigraphs and Pseudographs
- **Directed graphs:** Simple directed graphs, Directed multigraphs

Undirected graphs

Undirected graphs


Simple graphs:

Undirected graphs

Simple graphs: For any two vertices there is at most one edge connecting them, and there are no loops.

Undirected graphs


Simple graphs: For any two vertices there is at most one edge connecting them, and there are no loops.



Multigraphs:

Multigraphs: There are possibly **multiple edges**, but no loops.


Multigraphs: There are possibly **multiple edges**, but no loops.



Pseudographs:

Pseudographs: There are possibly multiple edges and loops.

Pseudographs: There are possibly multiple edges and loops.



Directed graphs

Directed graphs


Simple directed graphs:

Directed graphs

Simple directed graphs: There are no loops and no multiple directed edges.

Directed graphs


Simple directed graphs: There are no loops and no multiple directed edges.



Directed graphs

Simple directed graphs: There are no loops and no multiple directed edges.


Directed multigraphs:



Directed graphs


Simple directed graphs: There are no loops and no multiple directed edges.

Directed multigraphs: There are possibly multiple directed edges and loops.




Directed graphs

Simple directed graphs: There are no loops and no multiple directed edges.




Directed multigraphs: There are possibly multiple directed edges and loops.




Directed graphs

Simple directed graphs: There are no loops and no multiple directed edges.



Directed multigraphs: There are possibly multiple directed edges and loops.



9.2 Graph Terminologies and Special Types of Graphs

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

The Handshaking Theorem for Undirected graphs

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

The Handshaking Theorem for Undirected graphs

In an undirected graph G with e edges,

$$\sum \deg(v) = 2e$$

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

The Handshaking Theorem for Undirected graphs

In an undirected graph G with e edges,

$$\sum \deg(v) = 2e$$

Therefore, the number of vertices of odd degrees is an even number.

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

The Handshaking Theorem for Undirected graphs

In an undirected graph G with e edges,

$$\sum \deg(v) = 2e$$

Therefore, the number of vertices of odd degrees is an even number.

Example. Do there exist simple graphs with 5 vertices of degrees:

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

The Handshaking Theorem for Undirected graphs

In an undirected graph G with e edges,

$$\sum \deg(v) = 2e$$

Therefore, the number of vertices of odd degrees is an even number.

Example. Do there exist simple graphs with 5 vertices of degrees:

(a) 1, 2, 3, 3, 4

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

The Handshaking Theorem for Undirected graphs

In an undirected graph G with e edges,

$$\sum \deg(v) = 2e$$

Therefore, the number of vertices of odd degrees is an even number.

Example. Do there exist simple graphs with 5 vertices of degrees:

(a) 1, 2, 3, 3, 4 (b) 1, 2, 3, 3, 3

9.2 Graph Terminologies and Special Types of Graphs

- In an undirected graph, two vertices u and v are called **adjacent** if they are endpoints of an edge e , and e is called **incident with** u and v .
- The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex.

The Handshaking Theorem for Undirected graphs

In an undirected graph G with e edges,

$$\sum \deg(v) = 2e$$

Therefore, the number of vertices of odd degrees is an even number.

Example. Do there exist simple graphs with 5 vertices of degrees:

- (a) 1, 2, 3, 3, 4 (b) 1, 2, 3, 3, 3 (c) 1, 2, 3, 4, 4

- If $e = (u, v)$ is an edge of a directed graph, u is said to be adjacent to v and v is adjacent from u . The vertex u is called the **initial** and v is called the **terminal** or **end** vertex of e .

- If $e = (u, v)$ is an edge of a directed graph, u is said to be adjacent to v and v is adjacent from u . The vertex u is called the **initial** and v is called the **terminal** or **end** vertex of e .
- The **in-degree** of a vertex v in a directed graph, denoted by $\deg^-(v)$, is the number of edges with v as their terminal vertex. The **out-degree** of u , denoted by $\deg^+(u)$, is the number of edges with u as their initial vertex.

- If $e = (u, v)$ is an edge of a directed graph, u is said to be adjacent to v and v is adjacent from u . The vertex u is called the **initial** and v is called the **terminal** or **end** vertex of e .
- The **in-degree** of a vertex v in a directed graph, denoted by $\deg^-(v)$, is the number of edges with v as their terminal vertex. The **out-degree** of u , denoted by $\deg^+(u)$, is the number of edges with u as their initial vertex.

The Handshaking Theorem for directed graphs

- If $e = (u, v)$ is an edge of a directed graph, u is said to be adjacent to v and v is adjacent from u . The vertex u is called the **initial** and v is called the **terminal** or **end** vertex of e .
- The **in-degree** of a vertex v in a directed graph, denoted by $\deg^-(v)$, is the number of edges with v as their terminal vertex. The **out-degree** of u , denoted by $\deg^+(u)$, is the number of edges with u as their initial vertex.

The Handshaking Theorem for directed graphs

In a directed graph G ,

$$\sum \deg^+(v) = \sum \deg^-(v)$$

- If $e = (u, v)$ is an edge of a directed graph, u is said to be adjacent to v and v is adjacent from u . The vertex u is called the **initial** and v is called the **terminal** or **end** vertex of e .
- The **in-degree** of a vertex v in a directed graph, denoted by $\deg^-(v)$, is the number of edges with v as their terminal vertex. The **out-degree** of u , denoted by $\deg^+(u)$, is the number of edges with u as their initial vertex.

The Handshaking Theorem for directed graphs

In a directed graph G ,

$$\sum \deg^+(v) = \sum \deg^-(v)$$

- If $e = (u, v)$ is an edge of a directed graph, u is said to be adjacent to v and v is adjacent from u . The vertex u is called the **initial** and v is called the **terminal** or **end** vertex of e .
- The **in-degree** of a vertex v in a directed graph, denoted by $\deg^-(v)$, is the number of edges with v as their terminal vertex. The **out-degree** of u , denoted by $\deg^+(u)$, is the number of edges with u as their initial vertex.

The Handshaking Theorem for directed graphs

In a directed graph G ,

$$\sum \deg^+(v) = \sum \deg^-(v)$$

Some Special Simple Graphs

Some Special Simple Graphs

Complete Graphs K_n , $n \geq 1$:

Some Special Simple Graphs

Complete Graphs K_n , $n \geq 1$: n vertices, any two distinct vertices are connected by only one edge.


Some Special Simple Graphs

Complete Graphs K_n , $n \geq 1$: n vertices, any two distinct vertices are connected by only one edge.


K_1



K_2



K_3




K_4

Cycles C_n , $n \geq 3$:


Cycles C_n , $n \geq 3$: n vertices v_1, v_2, \dots, v_n and edges

$v_1 v_2, v_2 v_3, \dots, v_{n-1} v_n, v_n v_1$.


Cycles C_n , $n \geq 3$: n vertices v_1, v_2, \dots, v_n and edges $v_1 v_2, v_2 v_3, \dots, v_{n-1} v_n, v_n v_1$.



C_3



C_4




C_5


Wheels W_n , $n \geq 3$:

Wheels W_n , $n \geq 3$: Add one vertex to C_n and connect it with the remaining vertices.


Wheels W_n , $n \geq 3$: Add one vertex to C_n and connect it with the remaining vertices.



W_3



W_4





W_5

n -Cubes Q_n , $n \geq 1$:

n -Cubes Q_n , $n \geq 1$: 2^n vertices, and the edges are drawn by the following rule: represent each vertex by a bit string of length n , and two vertices are connected if their bit strings differ in exactly one position.

n -Cubes Q_n , $n \geq 1$: 2^n vertices, and the edges are drawn by the following rule: represent each vertex by a bit string of length n , and two vertices are connected if their bit strings differ in exactly one position.



Question. How many edges each of the graphs K_n , C_n , W_n , Q_n has?

Bipartite graphs

Bipartite graphs



Bipartite graphs


A simple graph G is called **bipartite** if the vertex set can be divided in two disjoint subsets such that each edge connects one vertex from one of these two subsets to another vertex of the other subset.

Bipartite graphs

A simple graph G is called **bipartite** if the vertex set can be divided in two disjoint subsets such that each edge connects one vertex from one of these two subsets to another vertex of the other subset.


Bipartite graphs

A simple graph G is called **bipartite** if the vertex set can be divided in two disjoint subsets such that each edge connects one vertex from one of these two subsets to another vertex of the other subset.




Bipartite graphs

A simple graph G is called **bipartite** if the vertex set can be divided in two disjoint subsets such that each edge connects one vertex from one of these two subsets to another vertex of the other subset.




Question. Which graphs K_n , C_n , W_n , Q_n are bipartite?




The **complete bipartite** graph K_{mn} is the graph whose vertex set is divided to two disjoint subsets of m and n vertices, such that two vertices are connected if and only if they do not belong to the same subset.

The **complete bipartite** graph K_{mn} is the graph whose vertex set is divided to two disjoint subsets of m and n vertices, such that two vertices are connected if and only if they do not belong to the same subset.



$K_{3,4}$

The **complete bipartite** graph K_{mn} is the graph whose vertex set is divided to two disjoint subsets of m and n vertices, such that two vertices are connected if and only if they do not belong to the same subset.



$K_{3,4}$

Question. How many edges does the graph K_{mn} have?

New Graphs from Olds



New Graphs from Olds

The **union** of 2 graphs G and H is a new graph whose vertex set consists of vertices of G and H , and whose edge set consists of edges of G and H .

New Graphs from Olds

The **union** of 2 graphs G and H is a new graph whose vertex set consists of vertices of G and H , and whose edge set consists of edges of G and H .




Example. Union of 2 graphs



New Graphs from Olds

The **union** of 2 graphs G and H is a new graph whose vertex set consists of vertices of G and H , and whose edge set consists of edges of G and H .


Example. Union of 2 graphs



is the graph


A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .

A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .



Example. Given the graph


A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .



Example. Given the graph


Then:

A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .




Example. Given the graph

Then:




A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .




Example. Given the graph

Then:




is a subgraph of G

A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .




Example. Given the graph


Then:



is a subgraph of G




A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .




Example. Given the graph

Then:




is a subgraph of G




is not a subgraph of G

A **subgraph** of the graph G is a graph whose vertex set is a subset of the vertex set of G , and whose edge set is a subset of the edge set of G .




Example. Given the graph

Then:



is a subgraph of G



is not a subgraph of G

9.3 Representing Graphs and Graph Isomorphism

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for **Undirected graphs**.

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n .

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n ,

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G connecting v_i and v_j .

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G connecting v_i and v_j .

Adjacency matrices for Directed graphs.

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G connecting v_i and v_j .

Adjacency matrices for Directed graphs.

Let G be a directed graph with vertices v_1, v_2, \dots, v_n .

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G connecting v_i and v_j .

Adjacency matrices for Directed graphs.

Let G be a directed graph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n ,

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G connecting v_i and v_j .

Adjacency matrices for Directed graphs.

Let G be a directed graph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G connecting v_i and v_j .

Adjacency matrices for Directed graphs.

Let G be a directed graph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G whose initial vertex is v_i and whose end vertex is v_j .

9.3 Representing Graphs and Graph Isomorphism

Adjacency matrices for Undirected graphs.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G connecting v_i and v_j .

Adjacency matrices for Directed graphs.

Let G be a directed graph with vertices v_1, v_2, \dots, v_n . We can represent G by a square matrix $[a_{ij}]$ of order n , whose entries are determined as follows:

a_{ij} = the number of edges in G whose initial vertex is v_i and whose end vertex is v_j .

Incident matrices.

Incident matrices.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n and edges e_1, e_2, \dots, e_m .

Incident matrices.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n and edges e_1, e_2, \dots, e_m . We can represent G by an **incident matrix** $[a_{ij}]$ of size $n \times m$,

Incident matrices.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n and edges e_1, e_2, \dots, e_m . We can represent G by an **incident matrix** $[a_{ij}]$ of size $n \times m$, whose entries are determined as follows:

Incident matrices.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n and edges e_1, e_2, \dots, e_m . We can represent G by an **incident matrix** $[a_{ij}]$ of size $n \times m$, whose entries are determined as follows:

$a_{ij} = 1$ if e_j is incident to v_i ,

Incident matrices.

Let G be a pseudograph with vertices v_1, v_2, \dots, v_n and edges e_1, e_2, \dots, e_m . We can represent G by an **incident matrix** $[a_{ij}]$ of size $n \times m$, whose entries are determined as follows:

$a_{ij} = 1$ if e_j is incident to v_i , and

$a_{ij} = 0$ otherwise.

Graph Isomorphism

Graph Isomorphism





Graph Isomorphism

Two graphs G and H are **isomorphic** if there is a bijection f between the vertex sets of two graphs with the property that, u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in V .

Graph Isomorphism

Two graphs G and H are **isomorphic** if there is a bijection f between the vertex sets of two graphs with the property that, u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in V .



Example 1.



Graph Isomorphism

Two graphs G and H are **isomorphic** if there is a bijection f between the vertex sets of two graphs with the property that, u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in V .

Example 1.





Isomorphism:

Graph Isomorphism

Two graphs G and H are **isomorphic** if there is a bijection f between the vertex sets of two graphs with the property that, u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in V .

Example 1.





Isomorphism:

$$f(A) = X$$

Graph Isomorphism

Two graphs G and H are **isomorphic** if there is a bijection f between the vertex sets of two graphs with the property that, u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in V .

Example 1.



Isomorphism:



$$f(A) = X$$

$$f(B) = Y$$

Graph Isomorphism

Two graphs G and H are **isomorphic** if there is a bijection f between the vertex sets of two graphs with the property that, u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in V .

Example 1.



Isomorphism:

$$f(A) = X$$



$$f(B) = Y$$

$$f(C) = W$$

Graph Isomorphism

Two graphs G and H are **isomorphic** if there is a bijection f between the vertex sets of two graphs with the property that, u and v are adjacent in G if and only if $f(u)$ and $f(v)$ are adjacent in V .

Example 1.





Isomorphism:

$$\begin{aligned}f(A) &= X \\f(B) &= Y \\f(C) &= W \\f(D) &= Z\end{aligned}$$



Example 2.

Example 2. Determine if the two graphs are isomorphic:

Example 2. Determine if the two graphs are isomorphic:





Example 2. Determine if the two graphs are isomorphic:



Problem.

Example 2. Determine if the two graphs are isomorphic:



Problem. Find an algorithm to check if two graphs are isomorphic.

Graph Invariants

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

These properties are called **graph invariants**.

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

These properties are called [graph invariants](#). They can be:

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

These properties are called **graph invariants**. They can be:

- The number of vertices

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

These properties are called **graph invariants**. They can be:

- The number of vertices
- The number of edges

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

These properties are called **graph invariants**. They can be:

- The number of vertices
- The number of edges
- Degrees

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

These properties are called **graph invariants**. They can be:

- The number of vertices
- The number of edges
- Degrees
- Circuits, paths (to be studied later)

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.

These properties are called **graph invariants**. They can be:

- The number of vertices
- The number of edges
- Degrees
- Circuits, paths (to be studied later)
- ...

Graph Invariants

If there are no correspondences between two vertex sets preserving the adjacency, then the two graphs are not isomorphic.

In some cases, it is easy to conclude that the given graphs are not isomorphic if we can find some properties of the graphs that are not the same.



These properties are called **graph invariants**. They can be:

- The number of vertices
- The number of edges
- Degrees
- Circuits, paths (to be studied later)
- ...





Example.

Example. Are the following pairs of graph isomorphic?

Example. Are the following pairs of graph isomorphic?



Example. Are the following pairs of graph isomorphic?



9.4 Connectivity

9.4 Connectivity

- Let G be an undirected graph. A **path** of length n from u to v is a sequence of edges $x_0x_1, x_1x_2, \dots, x_{n-1}x_n$, in which x_i and x_j are not necessarily distinct, and $x_0 = u, x_n = v$.

9.4 Connectivity

- Let G be an undirected graph. A **path** of length n from u to v is a sequence of edges $x_0x_1, x_1x_2, \dots, x_{n-1}x_n$, in which x_i and x_j are not necessarily distinct, and $x_0 = u, x_n = v$.
- A path is called a **circuit** if it starts and ends at the same vertex, and has length greater than 0.

9.4 Connectivity

- Let G be an undirected graph. A **path** of length n from u to v is a sequence of edges $x_0x_1, x_1x_2, \dots, x_{n-1}x_n$, in which x_i and x_j are not necessarily distinct, and $x_0 = u, x_n = v$.
- A path is called a **circuit** if it starts and ends at the same vertex, and has length greater than 0.
- A path, or a circuit, is **simple** if it does not contain the same edge more than once.

9.4 Connectivity

- Let G be an undirected graph. A **path** of length n from u to v is a sequence of edges $x_0x_1, x_1x_2, \dots, x_{n-1}x_n$, in which x_i and x_j are not necessarily distinct, and $x_0 = u, x_n = v$.
- A path is called a **circuit** if it starts and ends at the same vertex, and has length greater than 0.
- A path, or a circuit, is **simple** if it does not contain the same edge more than once.

Connectedness in Undirected Graphs

Connectedness in Undirected Graphs

- An undirected graph is **connected** if there is a path between any pair of distinct vertices.

Connectedness in Undirected Graphs

- An undirected graph is **connected** if there is a path between any pair of distinct vertices.
- A **connected component** of an undirected graph is a maximal subgraph that is connected.

Connectedness in Undirected Graphs

- An undirected graph is **connected** if there is a path between any pair of distinct vertices.
- A **connected component** of an undirected graph is a maximal subgraph that is connected.
- A **cut vertex**, or an **articulation point**, is a vertex that if we remove it and all the edges incident with it we will obtain a subgraph having more connected components than the original graph.

Connectedness in Undirected Graphs

- An undirected graph is **connected** if there is a path between any pair of distinct vertices.
- A **connected component** of an undirected graph is a maximal subgraph that is connected.
- A **cut vertex**, or an **articulation point**, is a vertex that if we remove it and all the edges incident with it we will obtain a subgraph having more connected components than the original graph.
- A **cut edge**, or a **bridge**, is an edge that if we remove it we will obtain a subgraph having more connected components than the original graph.

Connectedness in Undirected Graphs

- An undirected graph is **connected** if there is a path between any pair of distinct vertices.
- A **connected component** of an undirected graph is a maximal subgraph that is connected.
- A **cut vertex**, or an **articulation point**, is a vertex that if we remove it and all the edges incident with it we will obtain a subgraph having more connected components than the original graph.
- A **cut edge**, or a **bridge**, is an edge that if we remove it we will obtain a subgraph having more connected components than the original graph.

Connectedness in Directed Graphs

Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa.

Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.

Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.
- A **strongly connected component** of a directed graph G is a maximal subgraph of G that is strongly connected.

Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.
- A **strongly connected component** of a directed graph G is a maximal subgraph of G that is strongly connected.

Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.
- A **strongly connected component** of a directed graph G is a maximal subgraph of G that is strongly connected.

Example.

Connectedness in Directed Graphs


- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.
- A **strongly connected component** of a directed graph G is a maximal subgraph of G that is strongly connected.

Example. Given the graph

Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.
- A **strongly connected component** of a directed graph G is a maximal subgraph of G that is strongly connected.


Example. Given the graph



Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.
- A **strongly connected component** of a directed graph G is a maximal subgraph of G that is strongly connected.

Example. Given the graph




Determine if the graph is strongly connected, weakly connected, and find the number of strongly connected components.

Connectedness in Directed Graphs

- A directed graph is **strongly connected** if for all pairs of vertices u and v there is a path from u to v and vice versa. A directed graph is **weakly connected** if the underlying undirected graph is connected.
- A **strongly connected component** of a directed graph G is a maximal subgraph of G that is strongly connected.



Example. Given the graph







Determine if the graph is strongly connected, weakly connected, and find the number of strongly connected components.

Use graph invariants of paths and circuits to check if the two graphs are isomorphic:

Use graph invariants of paths and circuits to check if the two graphs are isomorphic:



Use graph invariants of paths and circuits to check if the two graphs are isomorphic:



Counting Paths Between Vertices

Counting Paths Between Vertices

Counting Paths Between Vertices

Let G be a graph (undirected, directed) whose adjacency matrix with respect to the ordering of vertices v_1, v_2, \dots, v_n is A .

Counting Paths Between Vertices

Let G be a graph (undirected, directed) whose adjacency matrix with respect to the ordering of vertices v_1, v_2, \dots, v_n is A . The number of paths of length r from v_i to v_j is the (i, j) -entry of the matrix A^r .

Counting Paths Between Vertices


Let G be a graph (undirected, directed) whose adjacency matrix with respect to the ordering of vertices v_1, v_2, \dots, v_n is A . The number of paths of length r from v_i to v_j is the (i, j) -entry of the matrix A^r .

Example. Count the number of paths of length 3 between A and C in the graph

Counting Paths Between Vertices

Let G be a graph (undirected, directed) whose adjacency matrix with respect to the ordering of vertices v_1, v_2, \dots, v_n is A . The number of paths of length r from v_i to v_j is the (i, j) -entry of the matrix A^r .

Example. Count the number of paths of length 3 between A and C in the graph




9.5 Euler and Hamilton Paths

9.5 Euler and Hamilton Paths

The 7 bridges problem.


9.5 Euler and Hamilton Paths

The 7 bridges problem.



9.5 Euler and Hamilton Paths

The 7 bridges problem.



Question. Is this possible to start at some location, travel across all bridges without crossing any bridge twice, then return to the starting point?

Euler Paths and Circuits

Euler Paths and Circuits

Euler Paths and Circuits

A simple circuit containing all edges of a graph is called **Euler circuit**.

Euler Paths and Circuits

A simple circuit containing all edges of a graph is called **Euler circuit**. A simple path containing all edges of a graph is called **Euler path**.

Euler Paths and Circuits

A simple circuit containing all edges of a graph is called **Euler circuit**. A simple path containing all edges of a graph is called **Euler path**.

Conditions for existence of Euler paths and circuits

Euler Paths and Circuits

A simple circuit containing all edges of a graph is called **Euler circuit**. A simple path containing all edges of a graph is called **Euler path**.

Conditions for existence of Euler paths and circuits

- A connected multigraph G has Euler circuits if and only if every vertex has even degree.

Euler Paths and Circuits

A simple circuit containing all edges of a graph is called **Euler circuit**. A simple path containing all edges of a graph is called **Euler path**.

Conditions for existence of Euler paths and circuits

- A connected multigraph G has Euler circuits if and only if every vertex has even degree.
- If G does not have Euler circuits, then it has Euler paths if and only if it has exactly two vertices of odd degrees.

Euler Paths and Circuits

A simple circuit containing all edges of a graph is called **Euler circuit**. A simple path containing all edges of a graph is called **Euler path**.

Conditions for existence of Euler paths and circuits

- A connected multigraph G has Euler circuits if and only if every vertex has even degree.
- If G does not have Euler circuits, then it has Euler paths if and only if it has exactly two vertices of odd degrees.

Example. The answer to the 7 bridges problem is NO.


Euler Paths and Circuits

A simple circuit containing all edges of a graph is called **Euler circuit**. A simple path containing all edges of a graph is called **Euler path**.

Conditions for existence of Euler paths and circuits

- A connected multigraph G has Euler circuits if and only if every vertex has even degree.
- If G does not have Euler circuits, then it has Euler paths if and only if it has exactly two vertices of odd degrees.

Example. The answer to the 7 bridges problem is NO.



Question.

Question. For what values of m, n do the special graphs $K_n, C_n, W_n, Q_n, K_{mn}$ has Euler paths/circuits?

Question. For what values of m, n do the special graphs $K_n, C_n, W_n, Q_n, K_{mn}$ has Euler paths/circuits?

Problem 1. Find an algorithm to find Euler circuits/paths.

Question. For what values of m, n do the special graphs $K_n, C_n, W_n, Q_n, K_{mn}$ has Euler paths/circuits?

Problem 1. Find an algorithm to find Euler circuits/paths.

Problem 2. Find conditions for the existence of Euler paths/circuits in directed graphs.

Question. For what values of m, n do the special graphs $K_n, C_n, W_n, Q_n, K_{mn}$ has Euler paths/circuits?

Problem 1. Find an algorithm to find Euler circuits/paths.

Problem 2. Find conditions for the existence of Euler paths/circuits in directed graphs.

Hamilton Paths and Circuits

Hamilton Paths and Circuits



Hamilton Paths and Circuits

- A simple path that passes through all vertices exactly once is called **Hamilton path**.

Hamilton Paths and Circuits

- A simple path that passes through all vertices exactly once is called **Hamilton path**.
- A simple circuit that passes through all vertices exactly once is called **Hamilton circuit**.

Hamilton Paths and Circuits


- A simple path that passes through all vertices exactly once is called **Hamilton path**.
- A simple circuit that passes through all vertices exactly once is called **Hamilton circuit**.

Example. Find Hamilton paths/circuits of the graphs:

Hamilton Paths and Circuits

- A simple path that passes through all vertices exactly once is called **Hamilton path**.
- A simple circuit that passes through all vertices exactly once is called **Hamilton circuit**.


Example. Find Hamilton paths/circuits of the graphs:



Hamilton Paths and Circuits

- A simple path that passes through all vertices exactly once is called **Hamilton path**.
- A simple circuit that passes through all vertices exactly once is called **Hamilton circuit**.

Example. Find Hamilton paths/circuits of the graphs:



Question. For what values of m, n do the special graphs $K_n, C_n, W_n, Q_n, K_{mn}$ has Hamilton paths/circuits?

9.6 Shortest-Path Problem

9.6 Shortest-Path Problem

9.6 Shortest-Path Problem

A graph that has a number assigned to each edge is called a **weighted graph**.

9.6 Shortest-Path Problem

A graph that has a number assigned to each edge is called a **weighted graph**. The **length** of a path in a weighted graph is the sum of all weights of the edges of this path.

9.6 Shortest-Path Problem

A graph that has a number assigned to each edge is called a **weighted graph**. The **length** of a path in a weighted graph is the sum of all weights of the edges of this path.

Problem.

9.6 Shortest-Path Problem


A graph that has a number assigned to each edge is called a **weighted graph**. The **length** of a path in a weighted graph is the sum of all weights of the edges of this path.

Problem. Find the path of shortest length between two vertices in a weighted graph.

9.6 Shortest-Path Problem

A graph that has a number assigned to each edge is called a **weighted graph**. The **length** of a path in a weighted graph is the sum of all weights of the edges of this path.

Problem. Find the path of shortest length between two vertices in a weighted graph.



Dijkstra's Algorithm

Dijkstra's Algorithm

Let G be a weighted graph.

Dijkstra's Algorithm

Let G be a weighted graph. To find the shortest path between A and Z in G , Dijkstra's algorithm:

Dijkstra's Algorithm

Let G be a weighted graph. To find the shortest path between A and Z in G , Dijkstra's algorithm:

- Finds the length of the shortest path from A to the first vertex.

Dijkstra's Algorithm

Let G be a weighted graph. To find the shortest path between A and Z in G , Dijkstra's algorithm:

- Finds the length of the shortest path from A to the first vertex.
- Finds the length of the shortest path from A to the second vertex.

Dijkstra's Algorithm

Let G be a weighted graph. To find the shortest path between A and Z in G , Dijkstra's algorithm:

- Finds the length of the shortest path from A to the first vertex.
- Finds the length of the shortest path from A to the second vertex.
- Finds the length of the shortest path from A to the third vertex.

Dijkstra's Algorithm

Let G be a weighted graph. To find the shortest path between A and Z in G , Dijkstra's algorithm:

- Finds the length of the shortest path from A to the first vertex.
- Finds the length of the shortest path from A to the second vertex.
- Finds the length of the shortest path from A to the third vertex.
- ...

Dijkstra's Algorithm


Let G be a weighted graph. To find the shortest path between A and Z in G , Dijkstra's algorithm:


- Finds the length of the shortest path from A to the first vertex.
- Finds the length of the shortest path from A to the second vertex.
- Finds the length of the shortest path from A to the third vertex.
- ...
- Continue the process until Z is reached.



Example.




Example. Find the shortest path from A to Z in the weighted graph





Example. Find the shortest path from A to Z in the weighted graph
































Procedure Dijkstra(G : weighted connected simple graph with n vertices v_1, v_2, \dots, v_n)

for $i := 1$ **to** n

$L(v_i) := \infty$

$L(A) := 0$

$S := \emptyset$

while $Z \notin S$

begin

$u :=$ vertex not in S with minimum label

$S := S \cup \{u\}$

for all vertices v not in S

$L(v) := \min\{L(v), L(u) + \textit{distance}(u, v)\}$

end