

## Supplemental Material

Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines for *IFNL3* (*IL28B*)  
genotype and peginterferon alpha based regimens

Andrew J. Muir<sup>1</sup>, Li Gong<sup>2</sup>, Samuel G. Johnson<sup>3,4</sup>, Ming Ta Michael Lee<sup>5,6,7</sup>, Marc S. Williams<sup>8</sup>, Teri E. Klein<sup>2</sup>, Kelly E. Caudle<sup>9</sup>, David R. Nelson<sup>10</sup>

<sup>1</sup>Division of Gastroenterology, Department of Medicine, Duke Clinical Research Institute, Duke University School of Medicine, Durham, North Carolina, USA

<sup>2</sup>Department of Genetics, Stanford University, Palo Alto, California, USA

<sup>3</sup>Department of Clinical Pharmacy, Skaggs School of Pharmacy and Pharmaceutical Sciences, University of Colorado, Denver, Colorado, USA

<sup>4</sup>Clinical Pharmacy Services, Kaiser Permanente Colorado, Denver, Colorado, USA

<sup>5</sup>National Center for Genome Medicine, Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan

<sup>6</sup>School of Chinese Medicine, China Medical University, Taichung, Taiwan

<sup>7</sup>Laboratory for International Alliance, RIKEN Center for Genomic Medicine, Yokohama, Japan

<sup>8</sup>Genomic Medicine Institute, Geisinger Health System, Danville, Pennsylvania, USA

<sup>9</sup>Department of Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, Tennessee, USA

<sup>10</sup>Department of Medicine, University of Florida, Gainesville, Florida, USA

### Corresponding Author:

Andrew J. Muir

2400 Pratt Street, Terrace Level, Room 0311

Durham, NC 27705

Phone: 919.668.8557

Fax: 919.668.7164

Email: [andrew.muir@duke.edu](mailto:andrew.muir@duke.edu)

## Table of Contents

CPIC Updates.....	3
Focused Literature Review .....	3
Other genes affecting PEG-interferon response .....	3
Available Genetic Test Options .....	3
Levels of Evidence linking genotype to phenotype.....	4
Strength of Recommendations.....	4
Supplemental Table S1. Genes associated with HCV treatment response .....	5
Supplemental Table S2. Frequencies of genotypes/alleles in major race/ethnic groups .....	6
Supplemental Table S3. Frequency of genotypes/alleles by individual publications for rs12979860.....	7
Supplemental Table S4. Evidence linking genotype with phenotype.....	11
References.....	14

## **CPIC Updates**

Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines are published in full on the PharmGKB website ([www.pharmgkb.org](http://www.pharmgkb.org)). Relevant information will be periodically reviewed and updated guidelines will be published online.

## **Focused Literature Review**

A literature search of the PubMed (1966 to January 2013) and Ovid MEDLINE (1950 to January 2013) database using the keywords ((IL28B OR interleukin 28) AND (peginterferon OR pegylated-interferon alfa OR PEG/IFN) AND genotype) was performed. Only articles available in English were reviewed.

## **Other genes affecting PEG-interferon response**

Many studies over the last three decades have evaluated genetic predictors of Hepatitis C virus (HCV) treatment response. Although the *IFNL3* genotype has emerged as the strongest predictor of treatment response, a recent systematic review found that more than forty genes have been associated with treatment response or treatment-related adverse events (1). The review did not attempt to validate the individual studies suggesting associations but was meant to provide awareness of the research. Other than the *IFNL3* genotype, these studies have not reported a consistent genetic association with HCV treatment response. Gene Ontology analysis categorized these genes into major groups of genes associated with the immune response, the inflammatory response, and the response to the virus (Table S1).

## **Available Genetic Test Options**

Commercially available genetic testing options change over time. An updated and fully linked table is available at <http://www.pharmgkb.org/gene/PA134952671#tabview=tab0&subtab=34>. Furthermore, the Genetic Testing Registry (GTR) provides a central location for voluntary submission of genetic test information by providers and is available at <http://www.ncbi.nlm.nih.gov/gtr/>. However, at the time of the writing of this manuscript no genetic test information was available on the GTR website for *IFNL3*.

## **Levels of Evidence linking genotype to phenotype**

The evidence summarized in Supplemental Table S4 has been graded using the three tiered system required by the Clinical Pharmacogenetics Implementation Consortium:

High: Evidence includes consistent results from well-designed, well conducted studies.

Moderate: Evidence is sufficient to determine effects, but the strength of the evidence is limited by the number, quality, or consistency of the individual studies, generalizability to routine practice, or indirect nature of the evidence.

Weak: Evidence is insufficient to assess the effects on health outcomes because of limited number or power of studies, important flaws in their design or conduct, gaps in the chain of evidence, or lack of information.

Every effort was made to present evidence from high-quality studies, which provided the framework for the strength of therapeutic recommendations in Table 2.

## **Strength of Recommendations**

Overall, the dosing recommendations are simplified to allow rapid interpretation by clinicians. We chose to use a slight modification of a transparent and simple system for just three categories for recommendations adopted from the rating scale for evidence-based recommendations on the use of retroviral agents (<http://aidsinfo.nih.gov/contentfiles/AdultandAdolescentGL.pdf>): strong, where “the evidence is high quality and the desirable effects clearly outweigh the undesirable effects”; moderate, in which “there is a close or uncertain balance” as to whether the evidence is high quality and the desirable clearly outweigh the undesirable effects; and optional, in which the desirable effects are closely balanced with undesirable effects and there is room for differences in opinion as to the need for the recommended course of action.

Strong recommendation for the statement

Moderate recommendation for the statement

Optional recommendation for the statement

**Supplemental Table S1. Genes associated with HCV treatment response (1)**

<b>Immune response</b>	<b>Inflammatory response</b>	<b>Response to the virus</b>
<i>CCR5</i>	<i>APOE</i>	<i>IFNAR1</i>
<i>CTLA4</i>	<i>CCR5</i>	<i>IFNAR2</i>
<i>CYP27B1</i>	<i>IL6</i>	<i>IFN<math>\gamma</math></i>
<i>HLA-A</i>	<i>IL10</i>	<i>IL6</i>
<i>HLA-B</i>	<i>MBL2</i>	<i>CCL5</i>
<i>HLA-C</i>	<i>CCL5</i>	<i>TNF</i>
<i>HLA-DQB1</i>	<i>TGFB1</i>	<i>TLR7</i>
<i>HLA-DRB1</i>	<i>TNF</i>	<i>TNK2</i>
<i>HFE</i>	<i>TRL7</i>	<i>MAPKAPK3</i>
<i>IL6</i>		<i>ADAR</i>
<i>IL10</i>		<i>OASL</i>
<i>KIR2DL3</i>		<i>EIF2AK2</i>
<i>MBL2</i>		<i>MX1</i>
<i>TGFB1</i>		<i>PSMB8</i>
<i>TNF</i>		<i>SPP1</i>
<i>TLR7</i>		

**Supplemental Table S2. Frequencies of genotypes/alleles in major race/ethnic groups**

Genotype	Caucasian (n=2042)	Asian (n=1518)	African American (n=491)	Hispanic (n=191)
rs12979860 : CC	0.38 (n=772)	0.77 (n=1166)	0.15 (n=72)	0.31 (n=60)
rs12979860 : CT	0.50 (n=1029)	0.21 (n=323)	0.48 (n=237)	0.48 (n=91)
rs12979860 : TT	0.12 (n=241)	0.02 (n=29)	0.37 (n=182)	0.21 (n=40)
rs12979860: C allele	0.63	0.87	0.39	0.55

Frequencies were calculated using the genotype information from subjects in studies by *Ge D et al* (2) and *Thompson AJ et al* (3) for Caucasians, African Americans and Hispanics and study by *Kobayashi M* (4) for Asians.

**Supplemental Table S3. Frequency of genotypes/alleles by individual publications for rs12979860**

Reference	Race	Study subject	CC (%)	CT (%)	TT (%)	Total (N)
Ramos <i>et al.</i> (2012) (5)	European & African		21 (32)	29 (44)	16 (24)	66
Falleti <i>et al.</i> (2012) (6)	Caucasian		76 (37)	130 (63)		206
El-Awady <i>et al.</i> (2012) (7)	Egyptian	Healthy control	48 (48)	38 (38)	14 (14)	100
		Spontaneous clearance	72 (86)	6 (7)	6 (7)	84
		Chronic HCV	14 (13)	83 (75)	13 (12)	110
		End stage liver disease	0 (0)	88 (80)	30 (20)	118
Xie <i>et al.</i> (2012) (8)	Chinese	Healthy control	85 (85)	13 (13)	2 (2)	100
		CHC patient	122 (55.5)	80 (36.4)	18 (8.1)	220
Riva <i>et al.</i> (2012) (9)	Caucasian	Sustained virological response (SVR)	28 (28)	59 (59)	13 (13)	100
		Nonsponse (NR)	0	60 (85)	11 (15)	70
Ramos <i>et al.</i> (2012) (10)	Brazilian	Acute hepatitis C with viral clearance	16 (88.9)	0 (0.0)	2 (11.1)	18
		Chronic hepatitis C	36 (30.3)	61 (51.3)	22 (18.5)	119
Poordad <i>et al.</i> (2012) (11)		Untreated-PR48	64 (29)	116 (53)	37 (17)	217
		Untreated-BOC RGT	77 (35)	103 (46)	42 (19)	222
		Untreated-BOC/PR48	55 (26)	115 (54)	44 (21)	214
		Treatment-failure-PR48	13 (25)	29 (56)	10 (19)	52

		Treatment-failure-BOC RGT	28 (28)	62 (62)	11 (11)	101
		Treatment-failure-BOC/PR48	22 (21)	66 (62)	18(17)	106
Fabris <i>et al.</i> (2012) (12)	Caucasian	With minimal or no fibrosis	26 (28)	54 (58)	13 (14)	93
		With chronic HCV infection and abnormal levels of transaminase	52 (36)	74 (52)	17 (12)	143
Martin-Carbonero <i>et al.</i> (2012) (13)	Spain	Chronic hepatitis B	29 (59)	17 (35)	3 (6)	49
		Spontaneous HBV clearance	22 (45)	21 (42)	6 (12)	49
Asselah <i>et al.</i> (2012) (14)	Egyptian, European, Sub-Saharan African	All patients	43 (26.2)	85 (51.8)	36 (22)	164
		Treated patients	22 (26.8)	43 (52.4)	17 (20.8)	82
Cariani <i>et al.</i> (2011) (15)	Caucasian		36.50%	47.60%	15.90%	170
Golden-Mason <i>et al.</i> (2011) (16)	African American		15.63%	40.62%	43.75%	55
	Caucasian		50.00%	34.62%	15.38%	46
Luo <i>et al.</i> (2013) (17)		Severe Fibrosis	1 (5)	13(65)	4(30)	30
		Mild Fibrosis	5 (50)	2 (20)	3(30)	
De la Fuente <i>et al.</i> (2013) (18)	Egyptian	Drug: Responder	45 (71.4)	13 (20.6)	5 (7.9)	100
		Drug: Non-responder	9 (24.3)	21(56.7)	7(18.9)	



Shaker and Sakik (2012) (19)	Caucasian		17(27)	34(53)	13(20)	64
Ramos <i>et al.</i> (2012) (10)			17(27%)	39(63%)	6(10%)	68
Falleti <i>et al.</i> (2011) (20)		Control	11.20%	41.80%	47%	629
		Hepatitis Patients	14.60%	52.80%	32.60%	
Fonseca-Coronado <i>et al.</i> (2011) (21)		Donor	3	5	22	50
		Patient	16	0	4	
Bitetto <i>et al.</i> (2011) (22)	Caucasian	Drug: Responder	14(10.4%)	60(44.8%)	60 (44.8%)	211
		Drug: Non-responder	10(13.0%)	51(66.2%)	16(20.8%)	
Ito <i>et al.</i> (2011) (23)	Japanese		4(1.4%)	85(29.6%)	198(69%)	287
Neukam <i>et al.</i> (2011) (24)	German & Spanish	acute	7(8.8%)	35(43.8%)	38(47.5%)	587
		chronic	46(9.5%)	243(44.7%)	218(45.8%)	
Ochi <i>et al.</i> (2011)(25)	Asian	HCV 1b	31(1.9%)	367(23%)	1198(75.1%)	2432
		HCV 2a	10(1.8%)	92(16.6%)	452(81.6%)	
		control	2(0.7%)	44(15.6%)	236(83.7%)	
Fabris <i>et al.</i> (2011) (26)	Italian	control	35(10.2%)	145(42.1%)	164(47.7%)	756
		Viral cirrhosis	8(5.8%)	58(42.0%)	72(52.2%)	
		Non-viral cirrhosis	38(13.9%)	137(50%)	99(36.1%)	
Langhans <i>et al.</i> (2011)(27)	Caucasian	Control	5	36	59	400
		Acute hepatitis	16	37	47	
		Chronic hepatitis	11	44	45	
		Self	7	39	54	
Clausen <i>et al.</i> (2011) (28)	Danish	Virus Clearance	0	15 (32%)	32 (68%)	206
		infection	30 (19%)	67 (42%)	62 (39%)	

Mangia <i>et al.</i> (2010)(29)	Caucasian	HCV infection	40(15%)	128((48%)	100(37%)	446
		control	14(8%)	90(51%)	74(42%)	
Thompson <i>et al.</i> (2010)(3)	Caucasian		139	596	436	1171
	African American		42	146	112	300
	Hispanic		34	56	26	116
Ge <i>et al.</i> (2009) (2)	Caucasian		336	433	102	871
	African American		30	91	70	191
	Hispanic		26	35	14	75
Kobayashi <i>et al.</i> (2012)(4)	Asian		1166	323	29	1518
			CC (%)	CT/TT (%)		Total N
Rizk and Derbala (2012) (30)	White & Black		13 (40%)	20 (60%)		33
Yuan <i>et al.</i> (2012) (31)		Viral Type 1	15(20%)	25(33.33%)		75
		Viral Type 3	8(10.67%)	8(10.67%)		
		Viral Type 4	5(6.67%)	4(5.33%)		
		1y4	2(2.67%)	3(4%)		
			C allele (%)	T allele (%)		Total N
Shi <i>et al.</i> (2012) (32)	Han Chinese	Healthy control	325 (95)	17 (5.0)		171
		Persistent HCV infection	1008 (95.3)	50 (4.7)		529
		Spontaneously cleared the virus	383 (97.7)	9 (2.3)		196

**Supplemental Table S4. Evidence linking genotype with phenotype**

Type of experimental model	Major findings	References	Level of evidence*
Clinical	<i>IFNL3</i> rs12979860 (CC genotype) is associated with increased SVR rates in HCV genotype 1 patients treated with PEG-IFN $\alpha$ and RBV therapy	McCarthy <i>et al.</i> (2010) (33) Montes-Cano <i>et al.</i> (2010) (34) Stattermayer <i>et al.</i> (2011) (35) Liao <i>et al.</i> (2011) (36) Chen <i>et al.</i> (2011) (37) Galmozzi <i>et al.</i> (2011) (38) Sarrazin <i>et al.</i> (2011) (39) Reiberger <i>et al.</i> (2011) (40) Bitetto <i>et al.</i> (2011) (22) de Rueda <i>et al.</i> (2011) (41) Darling <i>et al.</i> (2011) (42) Hayes <i>et al.</i> (2011) (43) Lin <i>et al.</i> (2011) (44) Patel <i>et al.</i> (2011) (45) Shi <i>et al.</i> (2012) (46) Ladero <i>et al.</i> (2012)(47)	High
Clinical	<i>IFNL3</i> rs8099917 (TT genotype) is associated with increased SVR rates in HCV genotype 1 patients treated with PEG-IFN $\alpha$ and RBV therapy	Stattermayer <i>et al.</i> (2011) (35) Tanaka <i>et al.</i> (2009) (48) Suppiah <i>et al.</i> (2009) (49) Rauch <i>et al.</i> (2010) (50) Abe <i>et al.</i> (2010) (51) Sinn <i>et al.</i> (2011) (52) Hayes <i>et al.</i> (2011) (43) Fukuhara <i>et al.</i> (2010) (53) Kurosaki <i>et al.</i> (2011) (54)	High
Clinical	<i>IFNL3</i> rs8099917 (TT genotype) is associated with increased SVR rates in HCV genotype 1 patients treated with PEG- PEG-IFN $\alpha$ and RBV therapy	Stattermayer <i>et al.</i> (2011) (35) Tanaka <i>et al.</i> (2009) (48) Suppiah <i>et al.</i> (2009)(49) Rauch <i>et al.</i> (2010) (50) Abe <i>et al.</i> (2010) (51) Sinn <i>et al.</i> (2011) (52) Hayes <i>et al.</i> (2011) (43) Fukuhara <i>et al.</i> (2010) (53) Kurosaki <i>et al.</i> (2011) (54)	High
Clinical	<i>IFNL3</i> rs12980275 (AA genotype) is associated with increased SVR rates in HCV genotype 1 patients treated with PEG-IFN $\alpha$ and RBV therapy	Fattovich <i>et al.</i> (2011) (55)	Moderate

Clinical	<i>IFNL3</i> rs12979860 is associated with early viral kinetics in patients infected with HCV during PEG- PEG-IFN $\alpha$ and RBV therapy	Naggie <i>et al.</i> (2012) (56) Howell <i>et al.</i> (2012) (57) Lindh <i>et al.</i> (2011) (58) Bochud <i>et al.</i> (2011) (59) Chu <i>et al.</i> (2012) (60)	Moderate
Clinical	<i>IFNL3</i> rs12979860 (CC genotype) is associated with increased SVR rates in HCV genotype 2 patients treated with PEG-IFN $\alpha$ and RBV therapy	Mangia <i>et al.</i> (2010) (29) Sarrazin <i>et al.</i> (2011) (39) Bitetto <i>et al.</i> (2011) (22) de Rueda <i>et al.</i> (2011) (41)	Moderate
Clinical	<i>IFNL3</i> rs8099917 (TT genotype) is associated with increased SVR rates in HCV genotype 2 patients treated with PEG-IFN/RBV therapy	Kawaoka <i>et al.</i> (2011) (61) Sinn <i>et al.</i> (2011) (52) Sarrazin <i>et al.</i> (2011) (39) Shi <i>et al.</i> (2012) (46)	Moderate
Clinical	<i>IFNL3</i> rs12979860 (CC genotype) is associated with increased SVR rates in HCV genotype 3 patients treated with PEG-IFN $\alpha$ and RBV therapy	Mangia <i>et al.</i> (2010) (29) Scherzer <i>et al.</i> (2011) (62) Sarrazin <i>et al.</i> (2011) (39) Moghaddam <i>et al.</i> (2011) (63) Bitetto <i>et al.</i> (2011) (22) de Rueda <i>et al.</i> (2011) (41)	Moderate
Clinical	<i>IFNL3</i> rs8099917 (TT genotype) is associated with increased SVR rates in HCV genotype 3 patients treated with PEG- PEG-IFN $\alpha$ and RBV therapy	Scherzer <i>et al.</i> (2011) (62) Sarrazin <i>et al.</i> (2011) (39) Shi <i>et al.</i> (2012) (46)	Moderate
Clinical	<i>IFNL3</i> rs12979860 (CC genotype) is associated with increased SVR rates in HCV genotype 4 patients treated with PEG-IFN $\alpha$ and RBV therapy	Stattermayer <i>et al.</i> (2011) (35) Ge <i>et al.</i> (2009) (2) Kawaoka <i>et al.</i> (2011) (61) Reiberger <i>et al.</i> (2011) (40) Bitetto <i>et al.</i> (2011) (22) de Rueda <i>et al.</i> (2011) (41) Shi <i>et al.</i> (2012) (46) Asselah <i>et al.</i> (2012) (14)	High
Clinical	<i>IFNL3</i> rs8099917 (TT genotype) is associated with increased SVR rates in HCV genotype 4 patients treated with PEG-IFN $\alpha$ and RBV therapy	Stattermayer <i>et al.</i> (2011) (35)	Low

Clinical	<i>IFNL3</i> rs12979860 is associated with spontaneous clearance of acute HCV	Thomas <i>et al.</i> (2009)(64) Ruiz-Extremera <i>et al.</i> (2011) (65) Tillmann <i>et al.</i> (2010) (66) Knapp <i>et al.</i> (2011) (67)	Moderate
Clinical	<i>IFNL3</i> rs8099917 is associated with spontaneous clearance of acute HCV infection	Rauch <i>et al.</i> (2010) (50)	Low
In vitro	Effect of <i>IFNL3</i> variation on <i>IFNL3</i> gene expression	Ge <i>et al.</i> (2009) (2) Suppiah <i>et al.</i> (2009) (49) Honda <i>et al.</i> (2010) (68) Urban <i>et al.</i> (2010) (69)	Moderate
In vitro	Effect of <i>IFNL3</i> variation on intrahepatic interferon-stimulated genes (ISG) expression	Honda <i>et al.</i> (2010) (68) Urban <i>et al.</i> (2010) (69) Dill <i>et al.</i> (2011) (70) Abe <i>et al.</i> (2011) (71) McGilvray <i>et al.</i> (2012) (72)	Moderate
In vitro	IFN lamda-3 inhibits HCV replication	Ank <i>et al.</i> (2006) (73)	Low

Evidence Review Table: IL28B = interleukin-28-beta; SVR = Sustained Virologic Response; HCV = Hepatitis C Virus; PEG-IFN $\alpha$  = Pegylated Interferon; RBV = Ribavirin; ISG = Interferon-Stimulated-Genes.

## References

- (1) Schlecker, C., Ultsch, A., Geisslinger, G. & Lotsch, J. The pharmacogenetic background of hepatitis C treatment. *Mutat Res* **751**, 36-48 (2012).
- (2) Ge, D. *et al.* Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* **461**, 399-401 (2009).
- (3) Thompson, A.J. *et al.* Interleukin-28B polymorphism improves viral kinetics and is the strongest pretreatment predictor of sustained virologic response in genotype 1 hepatitis C virus. *Gastroenterology* **139**, 120-9 e18 (2010).
- (4) Kobayashi, M. *et al.* Association of two polymorphisms of the IL28B gene with viral factors and treatment response in 1,518 patients infected with hepatitis C virus. *Journal of gastroenterology* **47**, 596-605 (2012).
- (5) Ramos, J.A. *et al.* A single nucleotide polymorphism, rs129679860, in the IL28B locus is associated with the viral kinetics and a sustained virological response in a chronic, monoinfected hepatitis C virus genotype-1 Brazilian population treated with pegylated interferon-ribavirin. *Memorias do Instituto Oswaldo Cruz* **107**, 888-92 (2012).
- (6) Falletti, E. *et al.* Vitamin D binding protein gene polymorphisms and baseline vitamin D levels as predictors of antiviral response in chronic hepatitis C. *Hepatology* **56**, 1641-50 (2012).
- (7) El-Awady, M.K. *et al.* Association of IL28B SNP With Progression of Egyptian HCV Genotype 4 Patients to End Stage Liver Disease. *Hepatitis monthly* **12**, 271-7 (2012).
- (8) Xie, J.Q. *et al.* Relationship between the genetic variation in interleukin 28B and response to antiviral therapy in patients with chronic hepatitis C. *Chinese medical journal* **125**, 2334-8 (2012).
- (9) Riva, E. *et al.* Interleukin-28B (IL-28B) single-nucleotide polymorphisms and interferon plus ribavirin treatment outcome in Italian chronically HCV-infected patients. *Journal of viral hepatitis* **19**, 650-3 (2012).
- (10) Ramos, J.A. *et al.* Association of IL-10, IL-4, and IL-28B gene polymorphisms with spontaneous clearance of hepatitis C virus in a population from Rio de Janeiro. *BMC research notes* **5**, 508 (2012).
- (11) Poordad, F. *et al.* Factors that predict response of patients with hepatitis C virus infection to boceprevir. *Gastroenterology* **143**, 608-18 e1-5 (2012).
- (12) Fabris, C. *et al.* The interleukin 28B rs12979860 C/T polymorphism and serum cholesterol as predictors of fibrosis progression in patients with chronic hepatitis C and persistently normal transaminases. *Journal of medical virology* **84**, 747-55 (2012).
- (13) Martin-Carbonero, L., Rallon, N.I., Benito, J.M., Poveda, E., Gonzalez-Lahoz, J. & Soriano, V. Short communication: Does interleukin-28B single nucleotide polymorphisms influence the natural history of hepatitis B? *AIDS research and human retroviruses* **28**, 1262-4 (2012).
- (14) Asselah, T. *et al.* IL28B polymorphism is associated with treatment response in patients with genotype 4 chronic hepatitis C. *Journal of hepatology* **56**, 527-32 (2012).

- (15) Cariani, E., Critelli, R., Rota, C., Luongo, M., Trenti, T. & Villa, E. Interleukin 28B genotype determination using DNA from different sources: A simple and reliable tool for the epidemiological and clinical characterization of hepatitis C. *Journal of virological methods* **178**, 235-8 (2011).
- (16) Golden-Mason, L. *et al.* Natural killer inhibitory receptor expression associated with treatment failure and interleukin-28B genotype in patients with chronic hepatitis C. *Hepatology* **54**, 1559-69 (2011).
- (17) Luo, Y., Jin, C., Ling, Z., Mou, X., Zhang, Q. & Xiang, C. Association study of IL28B: rs12979860 and rs8099917 polymorphisms with SVR in patients infected with chronic HCV genotype 1 to PEG-INF/RBV therapy using systematic meta-analysis. *Gene* **513**, 292-6 (2013).
- (18) de la Fuente, C. *et al.* Interleukin 28B rs12979860 (CT/TT) genotype is associated with milder hepatic damage in the natural evolution of HCV/HIV coinfection. *Journal of interferon & cytokine research* **33**, 43-7 (2013).
- (19) Shaker, O.G. & Sadik, N.A. Polymorphisms in interleukin-10 and interleukin-28B genes in Egyptian patients with chronic hepatitis C virus genotype 4 and their effect on the response to pegylated interferon/ribavirin-therapy. *Journal of gastroenterology and hepatology* **27**, 1842-9 (2012).
- (20) Falletti, E. *et al.* Role of interleukin 28B rs12979860 C/T polymorphism on the histological outcome of chronic hepatitis C: relationship with gender and viral genotype. *Journal of clinical immunology* **31**, 891-9 (2011).
- (21) Fonseca-Coronado, S. *et al.* Interleukin-28B genotyping by melt-mismatch amplification mutation assay PCR analysis using single nucleotide polymorphisms rs12979860 and rs8099917, a useful tool for prediction of therapy response in hepatitis C patients. *Journal of clinical microbiology* **49**, 2706-10 (2011).
- (22) Bitetto, D. *et al.* Complementary role of vitamin D deficiency and the interleukin-28B rs12979860 C/T polymorphism in predicting antiviral response in chronic hepatitis C. *Hepatology* **53**, 1118-26 (2011).
- (23) Ito, K. *et al.* The rs8099917 polymorphism, when determined by a suitable genotyping method, is a better predictor for response to pegylated alpha interferon/ribavirin therapy in Japanese patients than other single nucleotide polymorphisms associated with interleukin-28B. *Journal of clinical microbiology* **49**, 1853-60 (2011).
- (24) Neukam, K. *et al.* Different distributions of hepatitis C virus genotypes among HIV-infected patients with acute and chronic hepatitis C according to interleukin-28B genotype. *HIV medicine* **12**, 487-93 (2011).
- (25) Ochi, H. *et al.* IL-28B predicts response to chronic hepatitis C therapy--fine-mapping and replication study in Asian populations. *The Journal of general virology* **92**, 1071-81 (2011).
- (26) Fabris, C. *et al.* IL-28B rs12979860 C/T allele distribution in patients with liver cirrhosis: role in the course of chronic viral hepatitis and the development of HCC. *Journal of hepatology* **54**, 716-22 (2011).
- (27) Langhans, B. *et al.* Interferon-lambda serum levels in hepatitis C. *Journal of hepatology* **54**, 859-65 (2011).

- (28) Clausen, L.N. *et al.* Interleukin-28B polymorphisms are associated with hepatitis C virus clearance and viral load in a HIV-1-infected cohort. *Journal of viral hepatitis* **18**, e66-74 (2011).
- (29) Mangia, A. *et al.* An IL28B polymorphism determines treatment response of hepatitis C virus genotype 2 or 3 patients who do not achieve a rapid virologic response. *Gastroenterology* **139**, 821-7, 7 e1 (2010).
- (30) Rizk, N.M. & Derbala, M.F. Genetic polymorphisms of ICAM 1 and IL28 as predictors of liver fibrosis severity and viral clearance in hepatitis C genotype 4. *Clinics and research in hepatology and gastroenterology* **751**, 36-48 (2012).
- (31) Yuan, H., Adams-Huet, B., Petersen, T., Attar, N., Lee, W.M. & Jain, M.K. A single nucleotide polymorphism in IL28B affects viral evolution of hepatitis C quasispecies after pegylated interferon and ribavirin therapy. *Journal of medical virology* **84**, 1913-9 (2012).
- (32) Shi, X. *et al.* IL28B genetic variation is associated with spontaneous clearance of hepatitis C virus, treatment response, serum IL-28B levels in Chinese population. *PloS one* **7**, e37054 (2012).
- (33) McCarthy, J.J. *et al.* Replicated association between an IL28B gene variant and a sustained response to pegylated interferon and ribavirin. *Gastroenterology* **138**, 2307-14 (2010).
- (34) Montes-Cano, M.A. *et al.* Interleukin-28B genetic variants and hepatitis virus infection by different viral genotypes. *Hepatology* **52**, 33-7 (2010).
- (35) Stattermayer, A.F. *et al.* Impact of IL28B genotype on the early and sustained virologic response in treatment-naïve patients with chronic hepatitis C. *Clinical gastroenterology and hepatology* **9**, 344-50 e2 (2011).
- (36) Liao, X.W. *et al.* Association of genetic variation in IL28B with hepatitis C treatment-induced viral clearance in the Chinese Han population. *Antiviral therapy* **16**, 141-7 (2011).
- (37) Chen, J.Y. *et al.* IL28B genetic variations are associated with high sustained virological response (SVR) of interferon-alpha plus ribavirin therapy in Taiwanese chronic HCV infection. *Genes and immunity* **12**, 300-9 (2011).
- (38) Galmozzi, E. *et al.* A tetra-primer amplification refractory mutation system polymerase chain reaction for the evaluation of rs12979860 IL28B genotype. *Journal of viral hepatitis* **18**, 628-30 (2011).
- (39) Sarrazin, C. *et al.* Importance of IL28B gene polymorphisms in hepatitis C virus genotype 2 and 3 infected patients. *Journal of hepatology* **54**, 415-21 (2011).
- (40) Reiberger, T. *et al.* Portal pressure predicts outcome and safety of antiviral therapy in cirrhotic patients with hepatitis C virus infection. *Clinical gastroenterology and hepatology* **9**, 602-8 e1 (2011).
- (41) de Rueda, P.M. *et al.* Importance of host genetic factors HLA and IL28B as predictors of response to pegylated interferon and ribavirin. *The American journal of gastroenterology* **106**, 1246-54 (2011).
- (42) Darling, J.M. *et al.* Quantitation of pretreatment serum interferon-gamma-inducible protein-10 improves the predictive value of an IL28B gene polymorphism for hepatitis C treatment response. *Hepatology* **53**, 14-22 (2011).
- (43) Hayes, C.N. *et al.* HCV substitutions and IL28B polymorphisms on outcome of peg-interferon plus ribavirin combination therapy. *Gut* **60**, 261-7 (2011).



- (44) Lin, C.Y. *et al.* IL28B SNP rs12979860 is a critical predictor for on-treatment and sustained virologic response in patients with hepatitis C virus genotype-1 infection. *PloS one* **6**, e18322 (2011).
- (45) Patel, K. *et al.* High predictive accuracy of an unbiased proteomic profile for sustained virologic response in chronic hepatitis C patients. *Hepatology* **53**, 1809-18 (2011).
- (46) Shi, K.Q., Liu, W.Y., Lin, X.F., Fan, Y.C., Chen, Y.P. & Zheng, M.H. Interleukin-28B polymorphisms on the SVR in the treatment of naive chronic hepatitis C with pegylated interferon-alpha plus ribavirin: a meta-analysis. *Gene* **507**, 27-35 (2012).
- (47) Ladero, J.M. *et al.* Predicting response to therapy in chronic hepatitis C: an approach combining interleukin-28B gene polymorphisms and clinical data. *Journal of gastroenterology and hepatology* **27**, 279-85 (2012).
- (48) Tanaka, Y. *et al.* Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nature genetics* **41**, 1105-9 (2009).
- (49) Suppiah, V. *et al.* IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nature genetics* **41**, 1100-4 (2009).
- (50) Rauch, A. *et al.* Genetic variation in IL28B is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology* **138**, 1338-45, 45 e1-7 (2010).
- (51) Abe, H. *et al.* Common variation of IL28 affects gamma-GTP levels and inflammation of the liver in chronically infected hepatitis C virus patients. *Journal of hepatology* **53**, 439-43 (2010).
- (52) Sinn, D.H. *et al.* Association of a single nucleotide polymorphism near the interleukin-28B gene with response to hepatitis C therapy in Asian patients. *Journal of gastroenterology and hepatology* **26**, 1374-9 (2011).
- (53) Fukuhara, T. *et al.* Variants in IL28B in liver recipients and donors correlate with response to peg-interferon and ribavirin therapy for recurrent hepatitis C. *Gastroenterology* **139**, 1577-85, 85 e1-3 (2010).
- (54) Kurosaki, M. *et al.* Pre-treatment prediction of response to pegylated-interferon plus ribavirin for chronic hepatitis C using genetic polymorphism in IL28B and viral factors. *Journal of hepatology* **54**, 439-48 (2011).
- (55) Fattovich, G. *et al.* IL28B polymorphisms, IP-10 and viral load predict virological response to therapy in chronic hepatitis C. *Alimentary pharmacology & therapeutics* **33**, 1162-72 (2011).
- (56) Naggie, S. *et al.* Dysregulation of innate immunity in hepatitis C virus genotype 1 IL28B-unfavorable genotype patients: impaired viral kinetics and therapeutic response. *Hepatology* **56**, 444-54 (2012).
- (57) Howell, C.D. *et al.* Single nucleotide polymorphism upstream of interleukin 28B associated with phase 1 and phase 2 of early viral kinetics in patients infected with HCV genotype 1. *Journal of hepatology* **56**, 557-63 (2012).
- (58) Lindh, M. *et al.* Interleukin 28B gene variation at rs12979860 determines early viral kinetics during treatment in patients carrying genotypes 2 or 3 of hepatitis C virus. *The Journal of infectious diseases* **203**, 1748-52 (2011).
- (59) Bochud, P.Y. *et al.* IL28B polymorphisms predict reduction of HCV RNA from the first day of therapy in chronic hepatitis C. *Journal of hepatology* **55**, 980-8 (2011).

- (60) Chu, T.W. *et al.* Effect of IL28B genotype on early viral kinetics during interferon-free treatment of patients with chronic hepatitis C. *Gastroenterology* **142**, 790-5 (2012).
- (61) Kawaoka, T. *et al.* Predictive value of the IL28B polymorphism on the effect of interferon therapy in chronic hepatitis C patients with genotypes 2a and 2b. *Journal of hepatology* **54**, 408-14 (2011).
- (62) Scherzer, T.M. *et al.* Early virologic response and IL28B polymorphisms in patients with chronic hepatitis C genotype 3 treated with peginterferon alfa-2a and ribavirin. *Journal of hepatology* **54**, 866-71 (2011).
- (63) Moghaddam, A. *et al.* IL28B genetic variation and treatment response in patients with hepatitis C virus genotype 3 infection. *Hepatology* **53**, 746-54 (2011).
- (64) Thomas, D.L. *et al.* Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature* **461**, 798-801 (2009).
- (65) Ruiz-Extremera, A. *et al.* Genetic variation in interleukin 28B with respect to vertical transmission of hepatitis C virus and spontaneous clearance in HCV-infected children. *Hepatology* **53**, 1830-8 (2011).
- (66) Tillmann, H.L. *et al.* A polymorphism near IL28B is associated with spontaneous clearance of acute hepatitis C virus and jaundice. *Gastroenterology* **139**, 1586-92, 92 e1 (2010).
- (67) Knapp, S. *et al.* A polymorphism in IL28B distinguishes exposed, uninfected individuals from spontaneous resolvers of HCV infection. *Gastroenterology* **141**, 320-5, 5 e1-2 (2011).
- (68) Honda, M. *et al.* Hepatic ISG expression is associated with genetic variation in interleukin 28B and the outcome of IFN therapy for chronic hepatitis C. *Gastroenterology* **139**, 499-509 (2010).
- (69) Urban, T.J. *et al.* IL28B genotype is associated with differential expression of intrahepatic interferon-stimulated genes in patients with chronic hepatitis C. *Hepatology* **52**, 1888-96 (2010).
- (70) Dill, M.T. *et al.* Interferon-induced gene expression is a stronger predictor of treatment response than IL28B genotype in patients with hepatitis C. *Gastroenterology* **140**, 1021-31 (2011).
- (71) Abe, H. *et al.* IL28 variation affects expression of interferon stimulated genes and peg-interferon and ribavirin therapy. *Journal of hepatology* **54**, 1094-101 (2011).
- (72) McGilvray, I. *et al.* Hepatic cell-type specific gene expression better predicts HCV treatment outcome than IL28B genotype. *Gastroenterology* **142**, 1122-31 e1 (2012).
- (73) Ank, N., West, H., Bartholdy, C., Eriksson, K., Thomsen, A.R. & Paludan, S.R. Lambda interferon (IFN-lambda), a type III IFN, is induced by viruses and IFNs and displays potent antiviral activity against select virus infections in vivo. *Journal of virology* **80**, 4501-9 (2006).