

## Supplemental Material

### **Clinical Pharmacogenetics Implementation Consortium (CPIC) Guidelines for Dihydropyrimidine Dehydrogenase Genotype and Fluoropyrimidine Dosing: 2017 update**

**Ursula Amstutz<sup>1</sup>, Linda M. Henricks<sup>2</sup>, Steven M. Offer<sup>3</sup>, Julia Barbarino<sup>4</sup>, Jan H.M. Schellens<sup>2,5</sup>, Jesse J. Swen<sup>6</sup>, Teri E. Klein<sup>4</sup>, Howard L. McLeod<sup>7</sup>, Kelly E. Caudle<sup>8</sup>, Robert B. Diasio<sup>3,9</sup>, Matthias Schwab<sup>10,11,12</sup>**

<sup>1</sup>University Institute of Clinical Chemistry, Inselspital Bern University Hospital, University of Bern, Bern, Switzerland

<sup>2</sup>Department of Clinical Pharmacology, Division of Medical Oncology and Division of Pharmacology, the Netherlands Cancer Institute, Amsterdam, the Netherlands

<sup>3</sup>Department of Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic, Rochester, Minnesota, USA

<sup>4</sup>Department of Biomedical Data Science, Stanford University, Stanford, California, USA

<sup>5</sup>Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Utrecht, the Netherlands

<sup>6</sup>Department of Clinical Pharmacy & Toxicology, Leiden University Medical Center, Leiden, the Netherlands

<sup>7</sup>DeBartolo Family Personalized Medicine Institute and the Department of Population Sciences, H. Lee Moffitt Cancer Center & Research Institute, Tampa, Florida, USA

<sup>8</sup>Department of Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, Tennessee, USA

<sup>9</sup>Mayo Clinic Cancer Center, Mayo Clinic, Rochester, Minnesota, USA

<sup>10</sup>Dr Margarete Fischer-Bosch-Institute of Clinical Pharmacology, Stuttgart, Germany

<sup>11</sup>Department of Clinical Pharmacology, University Hospital, Tuebingen, Germany

<sup>12</sup>Department of Pharmacy and Biochemistry, University of Tuebingen, Tuebingen, Germany

## Table of Contents

Guideline Updates.....	3
Literature Review.....	3
Genetic Test Interpretation .....	4
Other considerations .....	6
Level of Evidence .....	7
Strength of Recommendation .....	7
Resources to Incorporate Pharmacogenetics into an Electronic Health Record with Clinical Decision Support .....	8
Supplemental Table S1. Evidence linking <i>DPYD</i> genotype with DPD phenotype and dihydropyrimidine toxicity .....	11
References.....	65

## **Guideline Updates**

The Clinical Pharmacogenetics Implementation Consortium (CPIC) guideline for *DPYD* genotypes and the dosing of fluoropyrimidines is published in full on [https://cpicpgx.org/guidelines/guideline-for-fluoropyrimidines-and-dpyd/\(1\)](https://cpicpgx.org/guidelines/guideline-for-fluoropyrimidines-and-dpyd/(1)) and at [pharmgkb.org](http://pharmgkb.org). Relevant information will be reviewed periodically and updated guidelines published online.

## **Literature Review**

### **2013 guideline**

A literature search of the PubMed® database (1966 to March 2013) using the keywords ((DPD OR DPYD OR Dihydropyrimidine Dehydrogenase) AND (fluorouracil OR 5-FU OR fluoropyrimidines OR capecitabine OR tegafur) AND genotype) was performed and results were limited to those available in English. Further articles were found via the reference sections of reviews. Using these search terms, 104 publications were identified. Study inclusion criteria included publications that included analyses for the association between *DPYD* genotypes (c.1905+1G>A , c.1679T>G, and c.2846A>T ) and metabolism of dihydropyrimidines and adverse drug events or clinical outcomes. Non-English manuscripts were excluded. Following application of these inclusion criteria, 30 publications were reviewed and included in the evidence tables

### **2017 guideline**

We searched PubMed® database as described above between 1966 and March 2017. The 2013 literature review was repeated to include all known *DPYD* genotypes. Using these search terms, 150 publications were identified. Following application of the inclusion criteria, 49 publications were reviewed and included in the evidence tables. An additional 43 studies were identified from the reference sections of reviews and other published paper, and included in the evidence tables, bringing the total included studies to 92 (**Supplemental Table S1**).

## Genetic Test Interpretation

While some *DPYD* variants have been assigned a (\*) allele, this nomenclature has not been updated to include more recently identified decreased function variants. As a consequence, only a minority of *DPYD* variants has a (\*) allele designation. Furthermore, the (\*) allele nomenclature is used for other drug metabolizing enzymes to designate haplotypes. Due to the size of *DPYD*, the gene encompasses several haplotype blocks (2, 3) with low linkage disequilibrium between variants located in different haplotype blocks. As a consequence, it is not possible to reliably determine full haplotypes that incorporate genotypes for common polymorphisms (e.g. c.85T>C, c.2194G>A) across the entire gene. Therefore, any (\*) alleles used for *DPYD* generally do not refer to haplotypes but only to a genotype at one specific SNP locus. To avoid confusion with (\*) allele nomenclature used for haplotypes of other drug metabolizing enzymes, the preferred nomenclature for *DPYD* variants is therefore the use of rs# or HGVS nomenclature.

Test results for *DPYD* do not report a diplotype for the entire gene, but genotypes for individual SNP loci. Importantly, however, all currently established toxicity-associated decreased/no function *DPYD* variants have a low population frequency (<5%) and are observed most frequently in individuals without a second decreased/no function variant. Therefore, in patients who carry two different decreased/no function variants, for the test interpretation, it is assumed that the two variants with an impact on DPD activity are located on different gene copies. For patients, in whom novel *DPYD* variants with suspected deleterious impact are detected in combination with known decreased/no function variants, this assumption may not be correct. In such a case, a phenotyping test may be helpful to determine enzyme activity, or genotyping of relatives (parents, siblings, and offspring) to determine segregation patterns. In addition, a genetic test may also include genotyping of other, common *DPYD* variants (e.g. c.85T>C, c.1627A>G, c.2194G>A). If this is case, a patient may be heterozygous for multiple of these variants and it cannot be determined which alleles are located on the same gene copy. However, based on current data, none of these common variants have a clinically relevant impact in the context of 5-fluorouracil related toxicity. The exact haplotype configuration of these

normal function variants is thus not required for the test interpretation. Therefore, to calculate the *DPYD* gene activity score, only the variant activity scores for the two variants with the lowest activity score is considered. For example, if a patient is a heterozygous carrier of a decreased function variant (e.g. c.1129–5923C>G) and two normal function variants (e.g. c.85T>C and c.1627A>G), the variant activity score of 0.5 for c.1129–5923C>G would be considered for one gene copy, and an activity score of 1 for the second gene copy, resulting in a total gene activity score of 1.5.

The dosing recommendations in this guideline are specific for variant alleles in which there are clear data linking the *DPYD* genotype to fluoropyrimidine toxicity (c.1905+1G>A, c.1679T>G, c.2846A>T, c.1129–5923C>G) (**Supplementary Table S1**). Several other variants have been reported to be associated with reduced enzyme activity and/or linked to fluoropyrimidine toxicity, albeit with somewhat weaker evidence (see ***DPYD* Allele Functionality Table (4)**, “moderate evidence supporting function”). While most of these variants are rare (see ***DPYD* Allele Frequency Table (4)**), the decreased function variant rs115232898 (c.557T>C, p.Y186C) is relatively common in individuals of African ancestry and has been observed in case reports of patients with severe 5-fluorouracil related toxicity (5, 6).

On the other hand, several *DPYD* variants that are relatively common in the population have strong or moderate evidence that they do not impact DPD function in a clinically relevant manner in the context of 5-fluorouracil related toxicity. For rs1801159 (\*5, c.1627A>G, p.I543V) and rs1801265 (\*9A, c.85T>C, p.C29R) none of the large cohort and case-control studies observed a significant association with toxicity or reduced DPD activity (see **Supplemental Table S1**). For other variants, associations with toxicity have been observed in single studies, but could not be reproduced in a majority of studies (rs1801160, \*6, c.2194G>A, p.V732I; rs2297595, c.496T>C, p.M166V) or by meta-analysis (rs1801158, \*4, c.1601G>A, p.S534N) (see **Supplemental Table S1**). Based on current knowledge, a fluorouracil dose adaptation in carriers of these variants is thus not warranted.

Many of the variants listed in the “in vitro data only and/or limited clinical/ex vivo data” category (see **DPYD Allele Functionality Table (4)**) as decreased or no function variants have a very low (<0.5%) allele frequency in the populations studied (see **DPYD Allele Frequency Table (4)**) and to date, there are no studies linking these variant alleles directly to toxicity related to fluoropyrimidines. Their functional effect was evaluated by comparison of their in vitro activity to the *in vitro* activity of known toxicity-associated *DPYD* variants: All variants with *in vitro* activity similar to c.1905+1G>A and c.1679T>G were categorized as “no function” variants; variants with *in vitro* activity greater than that of known “no function” variants but equal to or lower than the *in vitro* activity of c.2846A>T were classified as “decreased function” variants.

Several variants listed in the “unclear or conflicting data supporting function” category had *in vitro* DPD activity (i.e. homozygous expression of the variant) that was significantly lower than wildtype activity, but the magnitude of the decrease was smaller than for any established toxicity-associated variant. For these variants, it is currently not known if the decrease in DPD activity observed *in vitro* has a clinically relevant impact on 5-fluorouracil toxicity. At the time of writing, these variants would thus not be actionable for a reduction of the starting dose in fluoropyrimidine-based therapies.

### **Other considerations**

Several other genes may influence responses to 5-fluorouracil (7, 8), in particular genes of the folate pathway. The most well-studied of these are *MTHFR* and *TYMS*, although to date for *TYMS* the underlying causal variants of associations (9) and their clinical utility (10) are unclear, and associations have been inconsistent for *MTHFR* (10). Therefore, predictive dosing strategies for these genes have yet to be successfully applied. Similarly, a recently identified association of a variant (rs17822471) in *ABCC11*, a transporter of 5-fluorouracil metabolites, with fluoropyrimidine-related leukopenia requires further investigation (11, 12). In the context of capecitabine-based therapies, genes in the capecitabine activation pathway have also been studied, most notably *CDA*, *CES1* and *CES2* (13, 14). While some associations have been reported, these results have not been sufficiently replicated to determine potential genotype-based therapeutic strategies.

Furthermore, Fernandez-Rozadilla *et al* performed a genome-wide association study on 221 colorectal cancer patients (including a validation set of 791 patients) that had been treated with a 5-fluorouracil-based regimen (15). Seven SNPs (rs16857540 (*NLGN1*), rs2465403 (*COLEC10*), rs10876844 (*OR10AE3P*, *PSMB2P*), rs10784749, rs17626122 (*PARD3B*), rs7325568 and rs4243761) showed evidence of association with adverse drug reactions. They also evaluated the association signals for seven SNPs that had been linked to 5-fluorouracil-related toxicity in the literature (rs1801159 and rs1801265 (*DPYD*), rs18010919 (*UMPS*), rs1801133 (*MTHFR*), rs34743033, rs34489327 (*TYMS*), rs1695 (*GSTP1*)). Four of these variants had good proxy SNPs in the study, but none of them showed a statistically significant association. Some of the identified associations underscore the potential importance of other genes that may contribute increased risk of toxicity of 5-fluorouracil, although further studies are needed to determine their clinical utility.

### **Level of Evidence**

The evidence summarized in **Supplemental Table S1** is graded using a scaled modified slightly from Valdes *et al* (16).

High: Evidence includes consistent results from well-designed, well-conducted studies.

Moderate: Evidence is sufficient to determine effects, but the strength of the evidence is limited by the number, quality, or consistency of the individual studies; generalizability to routine practice; or indirect nature of the evidence.

Weak: Evidence is insufficient to assess the effects on health outcomes because of limited number or power of studies, important flaws in their design or conduct, gaps in the chain of evidence, or lack of information

### **Strength of Recommendation**

CPIC's dosing recommendations (**Table 2, main manuscript**) are based on weighting the evidence from a combination of preclinical functional and clinical data, as well as on some existing disease-specific consensus guidelines. Some of the factors that are taken

into account include *in vivo* clinical outcome for reference drug, *in vivo* PK/PD for reference drug, and *in vitro* enzyme activity with probe substrate only.

Overall, the dosing recommendations are simplified to allow rapid interpretation by clinicians. We chose to use a slight modification of a transparent and simple system for just three categories for recommendations adopted from the rating scale for evidence-based recommendations on the use of retroviral agents

(<http://aidsinfo.nih.gov/contentfiles/AdultandAdolescentGL.pdf>):

**Strong** recommendation for the statement: “The evidence is high quality and the desirable effects clearly outweigh the undesirable effects.”

**Moderate** recommendation for the statement: “There is a close or uncertain balance” as to whether the evidence is high quality and the desirable clearly outweigh the undesirable effects.

**Optional** recommendation for the statement: The desirable effects are closely balanced with undesirable effects, or the evidence is weak or based on extrapolations. There is room for differences in opinion as to the need for the recommended course of action.

**No recommendation:** There is insufficient evidence, confidence, or agreement to provide a recommendation to guide clinical practice at this time

The strength of the 5-fluorouracil dosing recommendations (**Table 2, main manuscript**) is based on the fact that some variants (c.1905+1G>A, c.1679T>G, c.2846A>T, c.1129–5923C>G) clearly affect DPD activity, and DPD activity is clearly related to 5-fluorouracil clearance, and 5-fluorouracil exposure is associated with its toxic effects. Therefore, reduction of 5-fluorouracil dosage in patients with these variants can prevent severe and possibly life-threatening toxicities, as has been demonstrated for c.1905+1G>A (17). The strength of the capecitabine dosing recommendations is based on the fact that this prodrug of 5-fluorouracil is metabolized by DPD in the same manner.

### **Resources to Incorporate Pharmacogenetics into an Electronic Health Record with Clinical Decision Support**

Clinical decision support (CDS) tools integrated within electronic health records (EHRs) can help guide clinical pharmacogenetics at the point of care (18-22). See



<https://cpicpgx.org/guidelines/guideline-for-fluoropyrimidines-and-dpyd/> for resources to support the adoption of CPIC guidelines within an EHR. Based on the capabilities of various EHRs and local preferences, we recognize that approaches may vary across organizations. Our intent is to synthesize foundational knowledge that provides a common starting point for incorporating the use of *DPYD* genotype results to guide fluoropyrimidine dosing in an EHR.

Effectively incorporating pharmacogenetic information into an EHR to optimize drug therapy should have some key attributes. Pharmacogenetic results, an interpreted phenotype, and a concise interpretation or summary of the result must be documented in the EHR (23, 24). To incorporate a phenotype in the EHR in a standardized manner, genotype test results provided by the laboratory must be consistently translated into an interpreted phenotype (**Table 1, main manuscript**). Because clinicians must be able to easily find the information, the interpreted phenotype may be documented as a problem list entry or in a patient summary section; these phenotypes are best stored in the EHR at the “person level” rather than at the date-centric “encounter level”. Additionally, results should be entered as standardized and discrete terms to facilitate using them to provide point-of-care CDS (18, 25).

Because pharmacogenetic results have lifetime implications and clinical significance, results should be placed into a section of the EHR that is accessible independent of the test result date to allow clinicians to quickly find the result at any time after it is initially placed in the EHR. To facilitate this process, CPIC is providing gene-specific information figures and tables that include full diplotype to phenotype tables, diagram(s) that illustrate how *DPYD* pharmacogenetic test results could be entered into an EHR, example EHR consultation/genetic test interpretation language and widely used nomenclature systems for genes relevant to the CPIC guideline (see <https://cpicpgx.org/guidelines/guideline-for-fluoropyrimidines-and-dpyd/>) (26).

Point-of-care CDS should be designed to effectively notify clinicians of prescribing implications at any time after the test result is entered into the EHR. CPIC is also

providing gene-drug specific tables that provide guidance to achieve these objectives with diagrams that illustrate how point-of-care CDS should be entered into the EHR, example pre- and post-test alert language, and widely used nomenclature systems for drugs relevant to the CPIC guideline (see <https://cpicpgx.org/guidelines/guideline-for-fluoropyrimidines-and-dpyd/>).

**Supplemental Table S1. Evidence linking *DPYD* genotype with DPD phenotype and dihydropyrimidine toxicity**

<i>DPYD</i> *allele rsID nucleotide change <sup>a</sup> protein change <sup>b</sup>	Parameter	Major findings	References	Type of experimental model	Level of Evidence <sup>c,d</sup>
*2A  rs3918290  c.1905+1G>A	Activity	AG is associated with decreased DPD activity as compared to GG	<b>Supports Statement: Statistically Significant:</b> Wei, <i>et al.</i> (1996)(27) Kuilenburg, <i>et al.</i> (2016)(28) Nie, <i>et al.</i> (2017)(29) <b>Same Direction of Association:</b> Sistonen, <i>et al.</i> (2014)(30)	Clinical, Ex vivo	<b>Moderate</b>
		Within cell lines, AA is associated with decreased DPD activity as compared to GG	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(31)	In vitro	<b>High</b>
		AA + AG were observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Holopainen, <i>et al.</i> (1997)(32) Vreken, <i>et al.</i> (1997)(33) Van Kuilenburg, <i>et al.</i> (1997)(34) Ridge, <i>et al.</i> (1998)(35) Van Kuilenburg, <i>et al.</i> (1999)(36) van Kuilenburg, <i>et al.</i> (2001)(37) Johnson, <i>et al.</i> (2002)(38) Maring, <i>et al.</i> (2002)(39) van Kuilenburg, <i>et al.</i> (2002)(40) Van Kuilenburg, <i>et al.</i> (2002)(41) Al-Sanna'a, <i>et al.</i> (2005)(42)	Clinical, Ex vivo	<b>High</b>

			Ezzeldin, <i>et al.</i> (2005)(43) Largillier, <i>et al.</i> (2006)(44) Magne, <i>et al.</i> (2007)(45) Loganayagam, <i>et al.</i> (2010)(46) van Kuilenburg, <i>et al.</i> (2010)(47) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Loganayagam, <i>et al.</i> (2010)(46)		
	Dose	AG is associated with decreased capecitabine dose as compared to GG	<b>Supports Statement: Statistically Significant:</b> Deenen, <i>et al.</i> (2011)(50)	Clinical	<b>Moderate</b>
		Individuals with AG received a decreased dose of capecitabine	<b>Supports Statement:</b> Joerger, <i>et al.</i> (2015)(51)	Clinical	<b>Weak</b>
	Efficacy	AG is associated with a decreased acute lymphoblastic leukemia complete remission rate as compared to GG	<b>Supports Statement: Statistically Significant:</b> Zhao, <i>et al.</i> (2016)(52)	Clinical	<b>Weak</b>
		AA + AG are not associated with progression-free, disease-free, event-free or overall survival times or confirmed response rate as compared to GG	McLeod, <i>et al.</i> (2010)(53) Deenen, <i>et al.</i> (2011)(50) Cai, <i>et al.</i> (2014)(54) Zhao, <i>et al.</i> (2016)(52)	Clinical	<b>Moderate</b>
	Metabolism	AG is associated with decreased metabolism of fluorouracil as compared to GG	<b>Supports Statement: Statistically Significant:</b> Boisdron-Celle, <i>et al.</i> (2007)(55) van Kuilenburg, <i>et al.</i> (2008)(56) Gentile, <i>et al.</i> (2016)(57)	Clinical, Ex vivo	<b>High</b>

		Individuals with AG were observed to have decreased metabolism of fluorouracil	<b>Supports Statement:</b> Maring, <i>et al.</i> (2002)(58) Joerger, <i>et al.</i> (2015)(51)	Clinical	<b>Moderate</b>
	Toxicity	AA + AG are associated with increased risk or severity of fluoropyrimidine toxicity as compared to GG	<b>Supports Statement: Statistically Significant (overall toxicity):</b> Van Kuilenburg, <i>et al.</i> (2002)(59) Salgueiro, <i>et al.</i> (2004)(60) Boisdron-Celle, <i>et al.</i> (2007)(55) Schwab, <i>et al.</i> (2008)(61) Deenen, <i>et al.</i> (2011)(50) Lee, <i>et al.</i> (2014)(62) Toffoli, <i>et al.</i> (2015)(63) <b>Statistically Significant (myelosuppression):</b> Schwab, <i>et al.</i> (2008)(61) Kleibl, <i>et al.</i> (2009)(64) Kristensen, <i>et al.</i> (2010)(65) Rosmarin, <i>et al.</i> (2014)(66) Cai, <i>et al.</i> (2014)(54) <b>Statistically Significant (hand-foot syndrome):</b> Cai, <i>et al.</i> (2014)(54) <b>Statistically Significant (diarrhea):</b> Deenen, <i>et al.</i> (2011)(50) Cai, <i>et al.</i> (2014)(54) <b>Statistically Significant (mucositis):</b> Schwab, <i>et al.</i> (2008)(61)	Clinical	<b>High</b>

			<p>Kleibl, <i>et al.</i> (2009)(64)  <b>Statistically Significant</b>  <b>(hepatotoxicity):</b>  Zhao, <i>et al.</i> (2016)(52)  <b>Statistically Significant</b>  <b>(infection):</b>  Zhao, <i>et al.</i> (2016)(52)  <b>Same Direction of Association</b>  <b>(overall toxicity):</b>  Braun, <i>et al.</i> (2009)(67)  Dhawan, <i>et al.</i> (2013)(68)  Rosmarin, <i>et al.</i> (2014)(66)  Rosmarin, <i>et al.</i> (2015)(9)  Froehlich, <i>et al.</i> (2015)(3)  Boige, <i>et al.</i> (2016)(69)  <b>Same Direction of Association</b>  <b>(neutropenia):</b>  McLeod, <i>et al.</i> (2010)(53)  <b>Does Not Support Statement:</b>  Amstutz, <i>et al.</i> (2009)(2)  McLeod, <i>et al.</i> (2010)(53)</p>		
--	--	--	--	--	--

		AA + AG were observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Wei, <i>et al.</i> (1996)(27) Van Kuilenburg, <i>et al.</i> (1997)(34) van Kuilenburg, <i>et al.</i> (2000)(70) van Kuilenburg, <i>et al.</i> (2001)(37) Raida, <i>et al.</i> (2001)(71) Johnson, <i>et al.</i> (2002)(38) Maring, <i>et al.</i> (2002)(58) Van Kuilenburg, <i>et al.</i> (2002)(41) Steiner, <i>et al.</i> (2005)(72) Ezzeldin, <i>et al.</i> (2005)(43) Largillier, <i>et al.</i> (2006)(44) Morel, <i>et al.</i> (2006)(73) Saif, <i>et al.</i> (2007)(74) Salgado, <i>et al.</i> (2007)(75) Magne, <i>et al.</i> (2007)(45) Sulzyc-Bielicka, <i>et al.</i> (2008)(76) Gross, <i>et al.</i> (2008)(77) Loganayagam, <i>et al.</i> (2010)(46) Boige, <i>et al.</i> (2010)(78) Ceric, <i>et al.</i> (2010)(79) van Kuilenburg, <i>et al.</i> (2010)(47) Savva-Bordalo, <i>et al.</i> (2010)(80) Cellier, <i>et al.</i> (2011)(147) Loganayagam, <i>et al.</i> (2013)(13) Suarez Martinez-Falero, <i>et al.</i> (2014)(81) Joerger, <i>et al.</i> (2015)(51) Thomas, <i>et al.</i> (2016)(48) Roberto, <i>et al.</i> (2017)(82)	Clinical	<b>High</b>
--	--	---	--	----------	-------------

<p>*13</p> <p>rs55886062</p> <p>c.1679T&gt;G</p> <p>p.I560S</p>	Activity	GT is associated with decreased DPD activity as compared to TT	<b>Supports Statement: <i>Statistically Significant:</i></b> Offer, <i>et al.</i> (2013)(5) Nie, <i>et al.</i> (2017)(29) <b><i>Same Direction of Association:</i></b> Sistonen, <i>et al.</i> (2014)(30)	Clinical, Ex vivo	<b>Moderate</b>
		Within cell lines, GG is associated with decreased DPD activity was compared to TT	<b>Supports Statement: <i>Statistically Significant:</i></b> Offer, <i>et al.</i> (2013)(31)	In vitro	<b>High</b>
		GT was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Johnson, <i>et al.</i> (2002)(38) van Kuilenburg, <i>et al.</i> (2002)(40) Ezzeldin, <i>et al.</i> (2005)(43) Thomas, <i>et al.</i> (2016)(48)	Clinical, Ex vivo	<b>High</b>
	Toxicity	GT is associated with increased risk or severity of fluoropyrimidine toxicity as compared to TT	<b>Supports Statement: <i>Same Direction of Association (overall toxicity):</i></b> Rosmarin, <i>et al.</i> (2015)(9) Froehlich, <i>et al.</i> (2015)(3) Lee, <i>et al.</i> (2014)(62) Toffoli, <i>et al.</i> (2015)(63) <b>Does Not Support Statement:</b> Amstutz, <i>et al.</i> (2009)(2) Deenen, <i>et al.</i> (2011)(50) Boige, <i>et al.</i> (2016)(69)	Clinical	<b>High</b>
		GT was observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Johnson, <i>et al.</i> (2002)(38) Ezzeldin, <i>et al.</i> (2005)(43)	Clinical	<b>High</b>



			Morel, <i>et al.</i> (2006)(73) Loganayagam, <i>et al.</i> (2010)(46) Cellier, <i>et al.</i> (2011)(83) Loganayagam, <i>et al.</i> (2013)(13) Dhelens, <i>et al.</i> (2016)(84) Thomas, <i>et al.</i> (2016)(48)		
rs67376798  c.2846A>T  p.D949V	Activity	The AT genotype/the T allele is associated with decreased DPD activity as compared to the AA genotype	<b>Supports Statement: Statistically Significant:</b> Seck, <i>et al.</i> (2005)(85) <b>Same Direction of Association:</b> Sistonen, <i>et al.</i> (2014)(30) Kuilenburg, <i>et al.</i> (2016)(28) Nie, <i>et al.</i> (2017)(29) <b>Does Not Support Statement:</b> Offer, <i>et al.</i> (2013)(5)	Clinical, Ex vivo	<b>Moderate</b>
		Within cell lines, the T allele is associated with decreased DPD activity as compared to AA	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86) Kuilenburg, <i>et al.</i> (2016)(28)	In vitro	<b>High</b>
		The AT genotype/the T allele was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2002)((40) Loganayagam, <i>et al.</i> (2010)(46) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Loganayagam, <i>et al.</i> (2010)(46)	Clinical, Ex vivo	<b>Moderate</b>

	Dose	The AT genotype is associated with decreased capecitabine dose as compared to the AA genotype	<b>Supports Statement: <i>Statistically Significant:</i></b> Deenen, <i>et al.</i> (2011)(50)	Clinical	<b>Moderate</b>
	Efficacy	The AT genotype is not associated with disease-free survival as compared to the AA genotype	Lee, <i>et al.</i> (2014)(62)	Clinical	<b>Moderate</b>
	Metabolism	The AT genotype is associated with decreased metabolism of fluorouracil as compared to the AA genotype	<b>Supports Statement: <i>Statistically Significant:</i></b> Boisdron-Celle, <i>et al.</i> (2007)(55) <b><i>Same Direction of Association:</i></b> Gentile, <i>et al.</i> (2016)(57)	Clinical, Ex vivo	<b>Moderate</b>
	Toxicity	The AT genotype/the T allele is associated with increased risk or severity of fluoropyrimidine toxicity as compared to the AA genotype	<b>Supports Statement: <i>Statistically Significant (overall toxicity):</i></b> Boisdron-Celle, <i>et al.</i> (2007)(55) Schwab, <i>et al.</i> (2008)(61) Deenen, <i>et al.</i> (2011)(50) Rosmarin, <i>et al.</i> (2014)(66) Lee, <i>et al.</i> (2014)(62) Toffoli, <i>et al.</i> (2015)(63) Boige, <i>et al.</i> (2016)(69) <b><i>Statistically Significant (diarrhea):</i></b> Deenen, <i>et al.</i> (2011)(50) Joerger, <i>et al.</i> (2015)(51)	Clinical	<b>High</b>

			<b>Same Direction of Association (overall toxicity):</b> Rosmarin, <i>et al.</i> (2015)(9) Froehlich, <i>et al.</i> (2015)(3)		
		The AT genotype/the T allele was observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2000)(70) Morel, <i>et al.</i> (2006)(73) Gross, <i>et al.</i> (2008)(77) Loganayagam, <i>et al.</i> (2010)(46) Obi, <i>et al.</i> (2011)(87) Cellier, <i>et al.</i> (2011)(83) Loganayagam, <i>et al.</i> (2013)(13) Thomas, <i>et al.</i> (2016)(48) Kuilenburg, <i>et al.</i> (2016)(28) <b>Does Not Support Statement:</b> Kristensen, <i>et al.</i> (2010)(65)	Clinical	<b>High</b>
HapB3  rs75017182 + rs56038477 + rs56276561  c.1129-5923C>G + c.1236G>A (p.E412E) + c.483+18G>A	Activity	HapB3 is associated with decreased DPD activity	<b>Supports Statement: Statistically Significant:</b> Sistonen, <i>et al.</i> (2014)(30) Nie, <i>et al.</i> (2017)(29) <b>Same Direction of Association:</b> Offer, <i>et al.</i> (2013)(5)	Ex vivo, Clinical	<b>Moderate</b>
		HapB3 was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2010)(47) Meulendijks, <i>et al.</i> (2016)(88) <b>Does Not Support Statement:</b> Seck, <i>et al.</i> (2005)(85)	Ex vivo	<b>Moderate</b>

	Dose	HapB3 was observed in individuals who required a fluoropyrimidine dose reduction	<b>Supports Statement:</b> Meulendijks, <i>et al.</i> (2016)(88) <b>Does Not Support Statement:</b> Meulendijks, <i>et al.</i> (2016)(88)	Clinical	<b>Weak</b>
	Toxicity	HapB3 is associated with increased risk or severity of fluoropyrimidine toxicity	<b>Supports Statement: <i>Statistically Significant (overall toxicity):</i></b> Amstutz, <i>et al.</i> (2009)(2) van Kuilenburg, <i>et al.</i> (2010)(47) Froehlich, <i>et al.</i> (2015)(3) <b><i>Statistically Significant (diarrhea):</i></b> Deenen, <i>et al.</i> (2011)(50) <b><i>Statistically Significant (neutropenia):</i></b> Lee, <i>et al.</i> (2016)(89) <b><i>Same Direction of Association (overall toxicity):</i></b> Schwab, <i>et al.</i> (2008)(61) Deenen, <i>et al.</i> (2011)(50) Rosmarin, <i>et al.</i> (2014)(66) Rosmarin, <i>et al.</i> (2015)(9) Lee, <i>et al.</i> (2016)(89) Boige, <i>et al.</i> (2016)(69) <b>Does Not Support Statement:</b> Kleibl, <i>et al.</i> (2009)(64) Loganayagam, <i>et al.</i> (2013)(13) Falvella, <i>et al.</i> (2015)(90)	Clinical	<b>High</b>
		HapB3 was observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2010)(47) Meulendijks, <i>et al.</i> (2016)(88)	Clinical	<b>Weak</b>

			<b>Does Not Support Statement:</b> Meulendijks, <i>et al.</i> (2016)(88)		
*2A (rs3918290, c.1905+1G>A) + rs67376798 (c.2846A>T, p.D949V) + *13 (rs55886062, c.1679T>G, p.I560S)	Toxicity	When the 1905+1 A allele and 2846 T allele are assessed together, with or without the 1679 G allele, they are associated with increased risk or severity of fluoropyrimidine toxicity	<b>Supports Statement: Statistically Significant (overall toxicity):</b> Morel, <i>et al.</i> (2006)(73) Saif, <i>et al.</i> (2013)(91) <b>Statistically Significant (gastrointestinal events):</b> Capitain, <i>et al.</i> (2008)(92)	Clinical	<b>High</b>
*2A (rs3918290, c.1905+1G>A) + rs67376798 (c.2846A>T, p.D949V) + *13 (rs55886062, c.1679T>G, p.I560S) + HapB3	Toxicity	When the 1905+1 A allele, the rs67376798 T allele and the HapB3 haplotype are assessed together, with or without the 1679 G allele, they are associated with increased risk or severity of fluoropyrimidine toxicity	<b>Supports Statement: Statistically Significant (overall toxicity):</b> Jennings, <i>et. al</i> (2013)(93) Froehlich, <i>et al</i> (2015)(3)	Clinical	<b>High</b>
*2A (rs3918290, c.1905+1G>A) +	Toxicity	When the 1905+1 A allele, 2846 T allele, 1679 G allele and 1601 A allele are assessed	<b>Supports Statement: Statistically Significant (overall toxicity):</b> Loganayagam, <i>et al.</i> (2013)(13)	Clinical	<b>Moderate</b>

rs67376798 (c.2846A>T, p.D949V) + *13 (rs55886062, c.1679T>G, p.I560S) + *4 (rs1801158, c.1601G>A, p.S534N)		together they are associated with increased risk or severity of fluoropyrimidine toxicity			
*2A (rs3918290, c.1905+1G>A) + rs67376798 (c.2846A>T, p.D949V) + HapB3	Activity	When the 1905+1 A allele, 2846 T allele and the HapB3 haplotype are assessed together they are associated with decreased DPD activity	<b>Supports Statement: <i>Statistically Significant:</i></b> Sistonen, <i>et al.</i> (2014)(30) Kuilenburg, <i>et al.</i> (2016)(28)	Clinical, Ex vivo	<b>Moderate</b>
*4  rs1801158  c.1601G>A  p.S534N	Activity	AG/the A allele is associated with decreased DPD activity as compared to GG	<b>Supports Statement: <i>Statistically Significant:</i></b> Seck, <i>et al.</i> (2005)(85) <b><i>Same Direction of Association:</i></b> Sistonen, <i>et al.</i> (2014)(30) Kuilenburg, <i>et al.</i> (2016)(28) <b>Does Not Support Statement:</b> Offer, <i>et al</i> (2013)(5)	Clinical, Ex vivo	<b>Weak</b>

		AG was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Gross, <i>et al.</i> (2003)(94) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Ridge, <i>et al.</i> (1998)(35)	Clinical, Ex vivo	<b>Weak</b>
		Within cell lines, the A allele is associated with decreased DPD activity as compared to GG	<b>Supports Statement: Statistically Significant:</b> Kuilenburg, <i>et al.</i> (2016)(28)	In vitro	<b>Weak</b>
		Within cell lines AA is associated with increased DPD activity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(31)	In vitro	<b>Weak</b>
	Metabolism	AG is not associated with altered fluorouracil metabolism as compared to GG	Gentile, <i>et al.</i> (2016)(57)	Ex vivo	<b>Weak</b>
	Toxicity	The A allele is associated with increased risk or severity of fluoropyrimidine toxicity as compared to GG	<b>Supports Statement: Same Direction of Association:</b> Froehlich, <i>et al.</i> (2015)(3) Loganayagam, <i>et al.</i> (2013)(13) Rosmarin, <i>et al.</i> (2014)(66) Rosmarin, <i>et al.</i> (2015)(9) <b>Does Not Support Statement:</b> Schwab, <i>et al.</i> (2008)(61) Kleibl, <i>et al.</i> (2009)(64) Amstutz, <i>et al.</i> (2009)(2) Deenen, <i>et al.</i> (2011)(50) Froehlich, <i>et al.</i> (2015)(3)	Clinical	<b>Weak</b>

			Toffoli, <i>et al.</i> (2015)(63) Boige, <i>et al.</i> (2016)(69)		
		AG was observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Gross, <i>et al.</i> (2003)(94) Lazar, <i>et al.</i> (2004)(95) van Kuilenburg, <i>et al.</i> (2010)(47)	Clinical	<b>Weak</b>
*5  rs1801159  c.1627A>G  p.I543V	Activity	AG + GG are not associated with altered DPD activity as compared to AA	He, <i>et al.</i> (2008)(96) Offer, <i>et al.</i> (2013)(5) Sistonen, <i>et al.</i> (2014)(30) Kuilenburg, <i>et al.</i> (2016)(28)	Clinical, Ex vivo	<b>High</b>
		Within cell lines, the GG or the G allele are not associated with altered DPD activity as compared to AA	Offer, <i>et al.</i> (2013)(31) Kuilenburg, <i>et al.</i> (2016)(28)	In vitro	<b>High</b>
		AG + GG were observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Gross, <i>et al.</i> (2003)(94) Ezzeldin, <i>et al.</i> (2005)(43) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Ridge, <i>et al.</i> (1998)(35) Collie-Duguid, <i>et al.</i> (2000)(49) Seck, <i>et al.</i> (2005)(85) Ezzeldin, <i>et al.</i> (2005)(43)	Clinical, Ex vivo	<b>Weak</b>



	Metabolism	AG + GG are associated with decreased metabolism of fluorouracil as compared to AA	<b>Supports Statement: Statistically Significant:</b> Zhang, <i>et al.</i> (2007)(97) Teh, <i>et al.</i> (2013)(98) <b>Same Direction of Association:</b> Gentile, <i>et al.</i> (2016)(57) <b>Does Not Support Statement:</b> Rudek, <i>et al.</i> (2013)(99)	Clinical, Ex vivo	<b>Weak</b>
	Efficacy	AG + GG are associated with increased overall survival time, or increased response to fluoropyrimidine treatment as compared to AA	<b>Supports Statement: Statistically Significant:</b> Grau, <i>et al.</i> (2008)(100) Joerger, <i>et al.</i> (2015)(51) <b>Does Not Support Statement:</b> McLeod, <i>et al.</i> (2010)(53)	Clinical	<b>Weak</b>
		AG + GG are associated with decreased response to fluoropyrimidine treatment as compared to AA	<b>Supports Statement: Statistically Significant:</b> Zhang, <i>et al.</i> (2012)(101)	Clinical	<b>Weak</b>
		AG + GG are not associated with progression-free survival time as compared to AA	McLeod, <i>et al.</i> (2010)(53) Farina-Sarasqueta, <i>et al.</i> (2010)(102)	Clinical	<b>Weak</b>

	Toxicity	AG + GG are associated with risk or severity of fluoropyrimidine toxicity as compared to AA	<b>Supports Statement: <i>Statistically Significant (nausea/vomiting):</i></b> Zhang, <i>et al.</i> (2007)(97) <b><i>Statistically Significant (leukopenia):</i></b> Zhang, <i>et al.</i> (2007)(97) <b><i>Same Direction of Association:</i></b> Rosmarin, <i>et al.</i> (2014)(66) <b>Does Not Support Statement:</b> Gross, <i>et al.</i> (2003)(94) Cho, <i>et al.</i> (2007)(103) Schwab, <i>et al.</i> (2008)(61) Kleibl, <i>et al.</i> (2009)(64) Amstutz, <i>et al.</i> (2009)(2) McLeod, <i>et al.</i> (2010)(53) Deenen, <i>et al.</i> (2011)(50) Zhang, <i>et al.</i> (2012)(101) Teh, <i>et al.</i> (2013)(98) Rosmarin, <i>et al.</i> (2015)(9) Froehlich, <i>et al.</i> (2015)(3) Joerger, <i>et al.</i> (2015)(51) Toffoli, <i>et al.</i> (2015)(63) Boige, <i>et al.</i> (2016)(69)	Clinical	<b>Weak</b>
		AG + GG were observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) van Kuilenburg, <i>et al.</i> (2000)(70) Lazar, <i>et al.</i> (2004)(95) Ezzeldin, <i>et al.</i> (2005)(43) Kim, <i>et al.</i> (2010)(104) van Kuilenburg, <i>et al.</i> (2010)(47) Zaanan, <i>et al.</i> (2014)(105)	Clinical	<b>Weak</b>

			Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49)		
*6  rs1801160  c.2194G>A  p.V732I	Activity	AG/the A allele is associated with decreased DPD activity as compared to GG	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(5) <b>Same Direction of Association:</b> Kuilenburg, <i>et al.</i> (2016)(28)	Ex vivo	<b>weak</b>
		Within cell lines, the AA genotype or the A allele are associated with decreased DPD activity as compared to GG	<b>Supports Statement: Statistically Significant:</b> Kuilenburg, <i>et al.</i> (2016)(28) <b>Does Not Support Statement:</b> Offer, <i>et al.</i> (2013)(31)	In vitro	<b>weak</b>
		AA + AG were observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Seck, <i>et al.</i> (2005)(85)	Clinical, Ex vivo	<b>weak</b>
	Metabolism	AA + AG are associated with decreased metabolism of fluorouracil as compared to GG	<b>Supports Statement: Statistically Significant:</b> Gentile, <i>et al.</i> (2016)(57)	Ex vivo	<b>weak</b>

	Efficacy	AA + AG are not associated with complete remission rate, event-free survival or response to fluoropyrimidine treatment as compared to GG	Zhang, <i>et al.</i> (2012)(101) Zhao, <i>et al.</i> (2016)(52)	Clinical	<b>weak</b>
	Toxicity	AA + AG are associated with increased risk or severity of fluoropyrimidine toxicity as compared to GG	<b>Supports Statement: Statistically Significant (overall toxicity):</b> Boige, <i>et al.</i> (2016)(69) <b>Statistically Significant (myelosuppression):</b> Kleibl, <i>et al.</i> (2009)(64) Boige, <i>et al.</i> (2016)(69) <b>Statistically Significant (diarrhea):</b> Deenen, <i>et al.</i> (2011)(50) <b>Same Direction of Association:</b> <b>Does Not Support Statement:</b> Schwab, <i>et al.</i> (2008)(61) Amstutz, <i>et al.</i> (2009)(2) Deenen, <i>et al.</i> (2011)(50) Zhang, <i>et al.</i> (2012)(101) Rosmarin, <i>et al.</i> (2015)(9) Froehlich, <i>et al.</i> (2015)(3) Toffoli, <i>et al.</i> (2015)(63) Zhao, <i>et al.</i> (2016)(52)	Clinical	<b>weak</b>
		AA + AG were observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) van Kuilenburg, <i>et al.</i> (2000)(70) Thomas, <i>et al.</i> (2016)(48)	Clinical	<b>weak</b>

			Del Re, <i>et al.</i> (2015)(106) <b>Does Not Support Statement:</b> Thomas, <i>et al.</i> (2016)(48)		
*9A  rs1801265  c.85T>C  p.C29R	Activity	The CC + CT genotypes are associated with increased DPD activity as compared to TT	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(5) Sistonen, <i>et al.</i> (2014)(30) <b>Does Not Support Statement:</b> He, <i>et al.</i> (2008)(96) Kuilenburg, <i>et al.</i> (2016)(28)	Clinical, Ex vivo	<b>weak</b>
		Within cell lines, CC is associated with increased DPD activity as compared to TT	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(31)	In vitro	<b>weak</b>
		Within cell lines, the C allele is associated with decreased DPD activity as compared to TT	<b>Supports Statement: Statistically Significant:</b> Kuilenburg, <i>et al.</i> (2016)(28)	In vitro	<b>weak</b>
		CC + CT were observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Vreken, <i>et al.</i> (1997)(33) Van Kuilenburg, <i>et al.</i> (1999)(36) Van Kuilenburg, <i>et al.</i> (1999)(107) Collie-Duguid, <i>et al.</i> (2000)(49) van Kuilenburg, <i>et al.</i> (2002)(40) Gross, <i>et al.</i> (2003)(94) Ezzeldin, <i>et al.</i> (2005)(43) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Johnson, <i>et al.</i> (2002)(38)	Clinical, Ex vivo	<b>weak</b>

			Gross, <i>et al.</i> (2003)(94) Seck, <i>et al.</i> (2005)(85) Ezzeldin, <i>et al.</i> (2005)(43)		
	Metabolism	CC is associated with decreased metabolism of fluorouracil as compared to CT + TT	<b>Supports Statement:</b> Gentile, <i>et al.</i> (2016)(57) <b>Does Not Support Statement:</b> Boisdron-Celle, <i>et al.</i> (2007)(55) Zhang, <i>et al.</i> (2007)(97)	Clinical, Ex vivo	<b>weak</b>
	Efficacy	CC + CT are associated with decreased event-free survival time and decreased response to fluorouracil treatment as compared to TT	<b>Supports Statement: Statistically Significant:</b> Zhao, <i>et al.</i> (2016)(52)	Clinical	<b>weak</b>
		CC + CT is not associated with overall survival, progression-free survival or response to fluoropyrimidine treatment as compared to TT	Grau, <i>et al.</i> (2008)(100) McLeod, <i>et al.</i> (2010)(53) Joerger, <i>et al.</i> (2015)(51)	Clinical	<b>moderate</b>

	Toxicity	CC are associated with risk or severity of fluoropyrimidine toxicity as compared to TT	<p><b>Supports Statement: <i>Statistically Significant (nausea/vomiting):</i></b> Zhang, <i>et al.</i> (2007)(97)</p> <p><b><i>Statistically Significant (hand-foot syndrome):</i></b> Joerger, <i>et al.</i> (2015)(51)</p> <p><b><i>Statistically Significant (diarrhea):</i></b> Joerger, <i>et al.</i> (2015)(51)</p> <p><b><i>Statistically Significant (infection):</i></b> Zhao, <i>et al.</i> (2016)(52)</p> <p><b><i>Statistically Significant (nephrotoxicity):</i></b> Zhao, <i>et al.</i> (2016)(52)</p> <p><b><i>Statistically Significant (hepatotoxicity):</i></b> Zhao, <i>et al.</i> (2016)(52)</p> <p><b><i>Same Direction of Association (overall toxicity):</i></b> Froehlich, <i>et al.</i> (2015)(3)</p> <p><b>Does Not Support Statement:</b> Gross, <i>et al.</i> (2003)(94) Boisdron-Celle, <i>et al.</i> (2007)(55) Morel, <i>et al.</i>(2006)(73) Schwab, <i>et al.</i> (2008)(61) Amstutz, <i>et al.</i> (2009)(2) McLeod, <i>et al.</i> (2010)(53) Deenen, <i>et al.</i> (2011)(50) Dhawan, <i>et al.</i> (2013)(68) Rosmarin, <i>et al.</i> (2014)(66)</p>	Clinical	<b>weak</b>
--	----------	--	--	----------	-------------

			Rosmarin, <i>et al.</i> (2015)(9) Boige, <i>et al.</i> (2016)(69)		
		CC + CT are associated with decreased risk of gastrointestinal toxicity as compared to TT	<b>Supports Statement: Statistically Significant:</b> Kleibl, <i>et al.</i> (2009)(64)	Clinical	<b>weak</b>
		CC + CT were observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) van Kuilenburg, <i>et al.</i> (2000)(70) Lazar, <i>et al.</i> (2004)(95) Kim, <i>et al.</i> (2010)(104) van Kuilenburg, <i>et al.</i> (2010)(47) Kristensen, <i>et al.</i> (2010)(65) Zaanan, <i>et al.</i> (2014)(105) Saif, <i>et al.</i> (2014)(6) Baskin, <i>et al.</i> (2015)(108) Thomas, <i>et al.</i> (2016)(48) Del Re, <i>et al.</i> (2015)(106) <b>Does Not Support Statement:</b> Kristensen, <i>et al.</i> (2010)(65) Thomas, <i>et al.</i> (2016)(48)	Clinical	<b>weak</b>



rs2297595 c.496A>G p.M166V	Activity	The AG + GG genotypes are not associated with altered DPD activity as compared to the AA genotype	Seck, <i>et al.</i> (2005)(85) Offer, <i>et al.</i> (2013)(5) Kuilenburg, <i>et al.</i> (2016)(28)	Ex vivo	<b>weak</b>
		Within cell lines, the G allele is associated with decreased DPD activity as compared to the AA genotype	<b>Supports Statement: Statistically Significant:</b> Kuilenburg, <i>et al.</i> (2016)(28)	In vitro	<b>weak</b>
		Within cell lines, the G allele is associated with increased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant :</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The AG genotype was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Gross, <i>et al.</i> (2003)(94) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Johnson, <i>et al.</i> (2002)(38)	Clinical, Ex vivo	<b>weak</b>
	Metabolism	The AG + GG genotypes are associated with decreased metabolism of fluorouracil as compared to the AA genotype	<b>Supports Statement: Statistically Significant:</b> Gentile, <i>et al.</i> (2016)(57)	Ex vivo	<b>weak</b>

	Efficacy	The AG genotype is not associated with response to fluoropyrimidine treatment as compared to the AA genotype	Zhang, <i>et al.</i> (2012)(101)	Clinical	<b>weak</b>
	Toxicity	The AG + GG genotypes are associated with risk or severity of fluoropyrimidine toxicity as compared to the AA genotype	<b>Supports Statement: Statistically Significant (overall toxicity):</b> Gross, <i>et al.</i> (2008)(77) Falvella, <i>et al.</i> (2015)(90) <b>Statistically Significant (diarrhea):</b> Deenen, <i>et al.</i> (2011)(50) <b>Statistically Significant (hand-foot syndrome):</b> Deenen, <i>et al.</i> (2011)(50) <b>Same Direction of Association:</b> Deenen, <i>et al.</i> (2011)(50) <b>Does Not Support Statement for overall toxicity:</b> Schwab, <i>et al.</i> (2008)(61) Amstutz, <i>et al.</i> (2009)(2) Zhang, <i>et al.</i> (2012)(101) Loganayagam, <i>et al.</i> (2013)(13) Rosmarin, <i>et al.</i> (2014)(66) Rosmarin, <i>et al.</i> (2015)(9) Froehlich, <i>et al.</i> (2015)(3) Toffoli, <i>et al.</i> (2015)(63) Boige, <i>et al.</i> (2016)(69)	Clinical	<b>weak</b>

		The AG + GG genotypes are associated with a decreased risk of neutropenia as compared to the AA genotype	<b>Supports Statement: Statistically Significant:</b> Kleibl, <i>et al.</i> (2009)(64)	Clinical	<b>weak</b>
		The AG genotype/the G allele was observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2000)(70) Gross, <i>et al.</i> (2003)(94) van Kuilenburg, <i>et al.</i> (2010)(47) Kristensen, <i>et al.</i> (2010)(65) Thomas, <i>et al.</i> (2016)(48) <b>Does Not Support Statement:</b> Kristensen, <i>et al.</i> (2010)(65) Thomas, <i>et al.</i> (2016)(48)	Clinical	<b>weak</b>
rs115232898  c.557A>G  p.Y186C	Activity	The AG genotype is associated with decreased DPD activity as compared to the AA genotype	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>Moderate</b>
		Within cell lines, the GG genotype/the G allele is associated with decreased DPD activity as compared to the AA genotype	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(109) Offer, <i>et al.</i> (2014)(86)	In vitro	<b>Moderate</b>
		The AG genotype/the G allele was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Ezzeldin, <i>et al.</i> (2005)(43) Zaanan, <i>et al.</i> (2014)(105)	Clinical, Ex vivo	<b>Moderate</b>

	Toxicity	The AG genotype was observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Zaanan, <i>et al.</i> (2014)(105) Saif, <i>et al.</i> (2014)(6)	Clinical	<b>Weak</b>
rs61622928 c.1218G>A p.M406I	Activity	The AG genotype/the A allele is not associated with altered DPD activity as compared to the GG genotype	Offer, <i>et al.</i> (2013)(5) Kuilenburg, <i>et al.</i> (2016)(28)	Ex vivo	<b>weak</b>
		Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele	Offer, <i>et al.</i> (2014)(86) Kuilenburg, <i>et al.</i> (2016)(28)	In vitro	<b>weak</b>
		The AG genotype/the A allele was observed in an individual with decreased DPD activity	<b>Supports Statement:</b> Ezzeldin, <i>et al.</i> (2005)(43) Thomas, <i>et al.</i> (2016)(48)	Clinical, Ex vivo	<b>weak</b>
rs17376848 c.1896T>C p.F632F	Activity	The CC + CT genotypes are not associated with altered DPD activity as compared to the TT genotype	He, <i>et al.</i> (2008)(96) Offer, <i>et al.</i> (2013)(5)	Clinical, Ex vivo	<b>weak</b>
		The CT genotype/the C allele were observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49) Ezzeldin, <i>et al.</i> (2005)(43) <b>Does Not Support Statement:</b> Collie-Duguid, <i>et al.</i> (2000)(49)	Ex vivo	<b>weak</b>

	Metabolism	The CC + CT genotypes are associated with decreased metabolism of fluorouracil as compared to the TT genotype	<b>Supports Statement: <i>Statistically Significant:</i></b> Teh, <i>et al.</i> (2013)(98)	Clinical	<b>weak</b>
	Toxicity	The CT genotype/the C allele is associated with risk or severity of fluoropyrimidine toxicity as compared to the TT genotype	<b>Supports Statement: <i>Statistically Significant (overall toxicity):</i></b> Kristensen, <i>et al.</i> (2010)(65) Falvella, <i>et al.</i> (2015)(90) <b><i>Statistically Significant (neutropenia):</i></b> Teh, <i>et al.</i> (2013)(98) <b><i>Statistically Significant (stomatitis):</i></b> Joerger, <i>et al.</i> (2015)(51) <b>Does Not Support Statement:</b> Schwab, <i>et al.</i> (2008)(61) Kleibl, <i>et al.</i> (2009)(64) Amstutz, <i>et al.</i> (2009)(2) Deenen, <i>et al.</i> (2011)(50) Froehlich, <i>et al.</i> (2015)(3) Toffoli, <i>et al.</i> (2015)(63) Boige, <i>et al.</i> (2016)(69)	Clinical	<b>weak</b>
		The CT genotype was observed in individuals with fluoropyrimidine toxicity	<b>Supports Statement:</b> Gross, <i>et al.</i> (2003)(94)	Clinical	<b>weak</b>

*2B  rs1801159 + rs3918290  1627A>G (I543V) + 1905+1G>A	Activity	The *2B/*4 genotype was observed in an individual with decreased DPD activity	<b>Supports Statement:</b> Ridge, <i>et al.</i> (1998)(35)	Ex vivo	<b>weak</b>
	Toxicity	The *2B/*4 genotype was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Ridge, <i>et al.</i> (1998)(35)	Clinical	<b>weak</b>
*3  rs72549303  c.1898delC  p.P633Qfs	Activity	Within cell lines, the del allele is associated with decreased DPD activity as compared the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The C/del + del/del genotypes were observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Vreken, <i>et al.</i> (1997)(110) Vreken, <i>et al.</i> (1997)(33) Van Kuilenburg, <i>et al.</i> (1999)(36)	Clinical, Ex vivo	<b>weak</b>
*7  rs72549309  c.295_298delTCAT  p.F100Sfs	Activity	Within cell lines, the del allele is associated with decreased DPD activity as compared to the TCAT allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The del/del genotype was observed in individuals with DPYD deficiency	<b>Supports Statement:</b> Van Kuilenburg, <i>et al.</i> (1999)(36)	Clinical	<b>weak</b>

<p>*8</p> <p>rs1801266</p> <p>c.703C&gt;T</p> <p>p.R235W</p>	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The T allele was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Vreken, <i>et al.</i> (1997)(110) Vreken, <i>et al.</i> (1997)(33) Van Kuilenburg, <i>et al.</i> (1999)(36)	Clinical, Ex vivo	<b>weak</b>
	Toxicity	The T allele was observed in an individual without fluoropyrimidine toxicity	<b>Supports Statement:</b> Kristensen, <i>et al.</i> (2010)(65)	Clinical	<b>weak</b>
<p>*9B</p> <p>rs1801267 + rs1801265</p> <p>c.2657G&gt;A + c.85T&gt;C</p> <p>p.R886H + p.C29R</p>	Activity	*9B/*9B/the *9B allele was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Vreken, <i>et al.</i> (1997)(110) Vreken, <i>et al.</i> (1997)(33) Van Kuilenburg, <i>et al.</i> (1999)(36)	Clinical, Ex vivo	<b>weak</b>
		Within cell lines, the A allele of the rs1801267 variant (part of *9B) is not associated with altered DPD activity as compared to the G allele	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
	Toxicity	The A allele of the rs1801267 variant (part of *9B) is not associated with risk of fluoropyrimidine toxicity as compared to the G allele	Boige, <i>et al.</i> (2016)(69)	Clinical	<b>weak</b>

*10  rs1801268  c.2983G>T  p.V995F	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The TT genotype was observed in an individual with decreased DPD activity	<b>Supports Statement:</b> Van Kuilenburg, <i>et al.</i> (1999) (36)	Clinical	<b>weak</b>
*11  rs72549306  c.1003G>T  p.V335L	Activity	The T allele was observed in individuals with decreased DPD activity	<b>Supports Statement:</b> Kouwaki, <i>et al.</i> (1998)(111)	Ex vivo	<b>weak</b>
		The T allele was observed to result in decreased DPD activity in <i>E.coli</i> lysates	<b>Supports Statement:</b> Kouwaki, <i>et al.</i> (1998)(111)	In vitro	<b>weak</b>
		Within cell lines, the T allele is not associated with altered DPD activity as compared to the G allele	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
	Toxicity	The T allele was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Kouwaki, <i>et al.</i> (1998)(111)	Clinical	<b>weak</b>
*12	Activity	The *12 allele was observed in individuals with decreased DPD activity		Ex vivo	<b>weak</b>



rs80081766 + rs78060119  c.62G>A + c.1156G>T  p.R21Q + p.E386X		The T allele of the rs78060119 variant (part of *12) was observed to result in undetectable DPD activity in <i>E.coli</i> lysates	<b>Supports Statement:</b> Kouwaki, <i>et al.</i> (1998)(111)	In vitro	<b>weak</b>
		The A allele of the rs80081766 variant (part of *12) was not observed to result in altered DPD activity in <i>E.coli</i> lysates	Kouwaki, <i>et al.</i> (1998)(111)	In vitro	<b>weak</b>
		Within cell lines, the T allele of the rs78060119 variant (part of *12) is associated with decreased DPD activity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		Within cell lines, the A allele of the rs80081766 variant (part of *12) is not associated with altered DPD activity as compared to the G allele	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
	Toxicity	The *12 allele was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Kouwaki, <i>et al.</i> (1998)(111)	Clinical	<b>weak</b>

		The GG genotype of the rs78060119 variant (part of *12) is not associated with risk or severity of fluoropyrimidine toxicity as compared to the TT or GT genotypes	Zhao, <i>et al.</i> (2016)(52)	Clinical	<b>weak</b>
rs111858276 c.1484A>G p.D495G	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs112766203 c.2279C>T p.T760I	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs114096998 c.3067C>A p.P1023T	Activity	Within cell lines, the A allele is associated with increased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs115632870 c.151-69G>A	Activity	The AG genotype is associated with decreased DPD activity as compared to the GG genotype	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>weak</b>
rs12022243 1906-14763G>A	Toxicity	The A allele is associated with increased severity of fluoropyrimidine toxicity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Rosmarin, <i>et al.</i> (2015)(9)	Clinical	<b>weak</b>
rs12132152 g.97523004G>A <sup>e</sup>	Toxicity	The A allele is associated with increased severity of fluoropyrimidine toxicity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Rosmarin, <i>et al.</i> (2015)(9)	Clinical	<b>weak</b>
rs76387818 g.97539400G>A <sup>e</sup>	Toxicity	The A allele is associated with increased severity of fluoropyrimidine toxicity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Rosmarin, <i>et al.</i> (2015)(9)	Clinical	<b>weak</b>

rs7548189 c.1906-19696G>T	Toxicity	The T allele is associated with increased severity of fluoropyrimidine toxicity as compared to the G allele	<b>Supports Statement: <i>Same Direction of Association:</i></b> Rosmarin, <i>et al.</i> (2015)(9)	Clinical	<b>weak</b>
rs137999090 c.2021G>A p.G674D	Activity	Within cell lines, the A allele is associated with decreased DPD activity as compared to the G allele	<b>Supports Statement: <i>Statistically Significant:</i></b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs138616379 c.1775G>A p.R592Q	Activity	Within cell lines, the A allele is associated with decreased DPD activity as compared to the G allele	<b>Supports Statement: <i>Statistically Significant:</i></b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs141044036 c.2872A>G p.K958E	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the A allele	<b>Supports Statement: <i>Statistically Significant:</i></b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs143154602 c.1057C>T p.R353C	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs143986398 c.274C>G p.P92A	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs145773863 c.1777G>A p.G593R	Activity	Within cell lines, the A allele is associated with decreased DPD activity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs146356975 c.868A>G p.K290E	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs147601618 c.1796T>C p.M599T	Toxicity	The CT genotype was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Ofverholm, <i>et al.</i> (2010)(112)	Clinical	<b>weak</b>
	Activity	Within cell lines, the C allele is not associated with DPD activity as compared to the T allele	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs183105782 c.910T>C p.Y304H	Activity	Within cell lines, the C allele is associated with decreased DPD activity as compared to the T allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs183385770 c.1024G>A p.D342N	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs186169810 c.1314T>G p.F438L	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the T allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs188052243 c.2678A>G p.N893S	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs190577302 c.1054C>G p.L352V	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs200687447 c.2482G>A p.E828K	Activity	Within cell lines, the A allele is associated with increased DPD activity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs367619008 187A>G p.K63E	Toxicity	The AG genotype was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Kleibl, <i>et al.</i> (2009)(64)	Clinical	<b>weak</b>
rs376073289 c.623G>A p.R208Q	Toxicity	The AG genotype was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Thomas, <i>et al.</i> (2016)(48)	Clinical	<b>weak</b>
		The A allele is not associated with risk of fluoropyrimidine toxicity	Schwab, <i>et al.</i> (2008)(61)	Clinical	<b>weak</b>
rs45589337 c.775A>G p.K259E	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the A allele	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>



		The G allele is not associated with altered DPD activity as compared to the AA genotype	Sistonen, <i>et al.</i> (2014)(30)	Clinical	<b>weak</b>
		The AG genotype was observed in an individual without altered DPD activity	<b>Supports Statement:</b> Gross, <i>et al.</i> (2003)(94)	Ex vivo	<b>weak</b>
	Toxicity	The AG genotype was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Gross, <i>et al.</i> (2003)(94)	Clinical	<b>weak</b>
		The AG genotype/the G allele is not associated with risk or severity of fluoropyrimidine toxicity as compared to the AA genotype	Schwab, <i>et al.</i> (2008)(61) Rosmarin, <i>et al.</i> (2015)(9) Froehlich, <i>et al.</i> (2015)(3)	Clinical	<b>weak</b>

rs55674432 c.2639G>T p.G880V	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the G allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs56293913 c.1129-15T>C	Activity	The CC + CT genotypes are not associated with altered DPD activity as compared to the TT genotype	Offer, <i>et al.</i> (2013)(5) Sistonen, <i>et al.</i> (2014)(30)	Clinical, Ex vivo	<b>weak</b>
	Toxicity	The CC + CT genotypes are associated with increased severity of fluoropyrimidine toxicity as compared to the TT genotype	<b>Supports Statement: Statistically Significant:</b> Gross, <i>et al.</i> (2008)(77) <b>Same Direction of Association:</b> Amstutz, <i>et al.</i> (2009)(2) Deenen, <i>et al.</i> (2011)(50)	Clinical	<b>weak</b>
rs568132506 c.257C>T p.P86L	Activity	The CT + TT genotypes were observed in individuals with decreased DPD activity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2002)(40) Thomas, <i>et al.</i> (2016)(48)	Clinical, Ex vivo	<b>weak</b>

	Toxicity	The CT genotype was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Thomas, <i>et al.</i> (2016)(48)	Clinical	<b>weak</b>
rs59086055 c.1774C>T p.R592W	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs60139309 c.2582A>G p.K861R	Activity	Within cell lines, the G allele is associated with increased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs61757362 c.2948C>T p.T983I	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs72547601 c.2933A>G p.H978R	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The GG genotype was observed in an individual with decreased DPD activity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2002)(40)	Ex vivo	<b>weak</b>
rs72549304 c.1475C>T p.S492L	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The TT genotype was observed in an individual with decreased DPD activity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2002)(40)	Ex vivo	<b>weak</b>
rs72549305 c.1108A>G p.I370V	Activity	Within cell lines, the G allele is not associated with DPD activity as compared to the A allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The GG genotype was observed in an individual with decreased DPD activity	<b>Supports Statement:</b> van Kuilenburg, <i>et al.</i> (2002)(40)	Ex vivo	<b>weak</b>

rs72549307 c.632A>G p.Y211C	Activity	Within cell lines, the G allele is associated with decreased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The GG genotype was observed in an individual with decreased DPD activity	<b>Supports Statement:</b> (van Kuilenburg, <i>et al.</i> (2002)(40)	Ex vivo	<b>weak</b>
	Toxicity	The G allele is not associated with risk or severity of fluoropyrimidine toxicity	Froehlich, <i>et al.</i> (2015)(3)	Clinical	<b>weak</b>
rs72549308 c.601A>C p.S201R	Activity	Within cell lines, the C allele is associated with decreased DPD activity as compared to the A allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs72549310 c.61C>T p.R21X	Activity	Within cell lines, the T allele is associated with decreased DPD activity as compared to the C allele	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs72728438 c.1974+75A>G	Activity	The AG genotype is associated with decreased DPD activity as compared to the AA genotype	<b>Supports Statement: Statistically Significant:</b> Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>weak</b>

rs777425216 c.1651G>A p.A551T	Toxicity	The AG genotype was observed in an individual with fluoropyrimidine toxicity	<b>Supports Statement:</b> Rosmarin, <i>et al.</i> (2015)(9)	Clinical	<b>weak</b>
rs150036960 c.46C>G p.L16V	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs150385342 c.313G>A p.A105T	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs141462178 c.343A>G p.M115V	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the A allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs200562975 c.451A>G p.N151D	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the A allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs139834141 c.498G>A p.M166I	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs150437414 c.929T>C p.L310S	Activity	Within cell lines, the C allele is not associated with altered DPD activity as compared to the T allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs145112791 c.934C>T p.L312F	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs201018345 c.967G>A p.A323T	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(574)	In vitro	<b>weak</b>
rs143815742 c.1181G>T p.R394L	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs140602333 c.1180C>T p.R394W	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs200064537 c.1260T>A p.N420K	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the T allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs764666241 c.1278G>T p.M426I	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs142512579 c.1294G>A p.D432N	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs72975710 c.1349C>T p.A450V	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>



rs144395748 c.1358C>G p.P453R	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
		The GG genotype is not associated with altered DPD activity as compared to the CG genotype.	Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>weak</b>
rs199549923 c.1403C>A p.T468N	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs138391898 c.1519G>A p.V507I	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs148994843 c.1543G>A p.V515I	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs190951787 c.1577C>G p.T526S	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs142619737 c.1615G>A p.G539R	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs201615754 c.1682G>T p.R561L	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs3918289 c.1905C>G p.N635K	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs55971861 c.1906A>C p.I636L	Activity	Within cell lines, the C allele is not associated with altered DPD activity as compared to the A allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs138545885 c.1990G>T p.A664S	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs145548112 c.2161G>A p.A721T	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs146529561 c.2186C>T p.A729V	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs60511679 c.2195T>G p.V732G	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the T allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs56005131 c.2303C>A p.T768K	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs199634007 c.2336C>A p.T779N	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs201035051 c.2623A>C p.K875Q	Activity	Within cell lines, the C allele is not associated with altered DPD activity as compared to the A allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs147545709 c.2656C>T p.R886C	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs145529148 c.2915A>G p.Q972R	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the A allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs72547602 c.2921A>T p.D974V	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the A allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>

rs139459586 c.2978T>G p.L993R	Activity	Within cell lines, the G allele is not associated with altered DPD activity as compared to the T allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs202144771 c.2977C>T p.L993F	Activity	Within cell lines, the T allele is not associated with altered DPD activity as compared to the C allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs140114515 c.3049G>A p.V1017I	Activity	Within cell lines, the A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs148799944 c.3061G>C p.V1021L	Activity	Within cell lines, the C allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2014)(86)	In vitro	<b>weak</b>
rs6670886 c.525G>A p.S175S	Activity	The A allele is not associated with altered DPD activity as compared to the G allele.	Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>weak</b>

rs3790387 c.763-118A>G	Activity	The AG + GG genotypes are not associated with altered DPD activity as compared to the AA genotype.	Offer, <i>et al.</i> (2013)(5) Kuilenburg, <i>et al.</i> (2016)(28)	Ex vivo	<b>weak</b>
		The G allele is not associated with altered DPD activity as compared to the A allele.	Sistonen, <i>et al.</i> (2014)(30)	Clinical	<b>weak</b>
rs112550271 c.850+41T>C	Activity	The CT genotype is not associated with altered DPD activity as compared to the TT genotype.	Offer, <i>et al.</i> (2013)(5) Kuilenburg, <i>et al.</i> (2016)(28)	Ex vivo	<b>weak</b>
rs2811202 c.958+134T>G	Activity	The GG + GT genotypes are not associated with altered DPD activity as compared to the TT genotype.	Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>weak</b>
rs61789183 c.1340-106T>A	Activity	The AA + AT genotypes are not associated with altered DPD activity as compared to the TT genotype.	Offer, <i>et al.</i> (2013)(5) Kuilenburg, <i>et al.</i> (2016)(28)	Ex vivo	<b>weak</b>
rs57918000 c.1371C>T p.N457N	Activity	The CT genotype is not associated with altered DPD activity as compared to the CC genotype.	Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>weak</b>

rs2786783 c.1740+39C>T	Activity	The CT + TT genotypes are not associated with altered DPD activity as compared to the CC genotype.	Offer, <i>et al.</i> (2013)(5) Sistonen, <i>et al.</i> (2014)(30)	Ex vivo, Clinical	<b>weak</b>
rs2811178 1740+40A>G	Activity	The AG + GG genotypes are not associated with altered DPD activity as compared to the AA genotype.	Offer, <i>et al.</i> (2013)(5) Sistonen, <i>et al.</i> (2014)(30)	Ex vivo, Clinical	<b>weak</b>
rs12137711 c.2300-39G>A	Activity	The AG genotype is not associated with altered DPD activity as compared to the GG genotype.	Offer, <i>et al.</i> (2013)(5)	Ex vivo	<b>weak</b>
rs41309171 c.234-123G>C	Activity	The CG genotype is not associated with altered DPD activity as compared to the GG genotype.	Sistonen, <i>et al.</i> (2014)(30)	Clinical	<b>weak</b>
rs138924556 c.850+91C>T	Activity	The CT genotype is not associated with altered DPD activity as compared to the CC genotype.	Sistonen, <i>et al.</i> (2014)(30)	Clinical	<b>weak</b>

rs368600943 c.1129-28G>T	Activity	The GT genotype is not associated with altered DPD activity as compared to the GG genotype.	Sistonen, <i>et al.</i> (2014)(30)	Clinical	<b>weak</b>
-----------------------------	----------	---	------------------------------------	----------	-------------

<sup>a</sup>Nucleotide changes according to reference sequence NM\_000110.3 unless otherwise specified

<sup>b</sup>Protein changes according to reference sequence NP\_000101.2

<sup>c</sup>Rating Scheme for Quality of Evidence as per (16)

<sup>d</sup>Some of the small case series, although not strong individually, collectively do support a strong recommendation.

<sup>e</sup>Nucleotide changes according to NC\_000001.10

<sup>f</sup>Likely HapB3 causal variant. Proxy SNPs are c.1236G>A (rs56038477, E412E), c.483+18G>A (rs56276561) and c.959-51T>G (rs115349832). c.680+139G>A (rs6668296) is not exclusive to HapB3 and therefore not a suitable proxy.



## References

- (1) Whirl-Carrillo, M. *et al.* Pharmacogenomics knowledge for personalized medicine. *Clinical pharmacology and therapeutics* **92**, 414-7 (2012).
- (2) Amstutz, U., Farese, S., Aebi, S. & Largiader, C.R. Dihydropyrimidine dehydrogenase gene variation and severe 5-fluorouracil toxicity: a haplotype assessment. *Pharmacogenomics* **10**, 931-44 (2009).
- (3) Froehlich, T.K., Amstutz, U., Aebi, S., Joerger, M. & Largiader, C.R. Clinical importance of risk variants in the dihydropyrimidine dehydrogenase gene for the prediction of early-onset fluoropyrimidine toxicity. *Int J Cancer* **136**, 730-9 (2015).
- (4) CPIC. CPIC® Guideline for Fluoropyrimidines and DPYD.  
<<https://cpicpgx.org/guidelines/guideline-for-fluoropyrimidines-and-dpyd/>>.
- (5) Offer, S.M., Lee, A.M., Mattison, L.K., Fossum, C., Wegner, N.J. & Diasio, R.B. A DPYD Variant (Y186C) in Individuals of African Ancestry Is Associated With Reduced DPD Enzyme Activity. *Clinical pharmacology and therapeutics* **94**, 158-66 (2013).
- (6) Saif, M.W., Lee, A.M., Offer, S.M., McConnell, K., Relias, V. & Diasio, R.B. A DPYD variant (Y186C) specific to individuals of African descent in a patient with life-threatening 5-FU toxic effects: potential for an individualized medicine approach. *Mayo Clin Proc* **89**, 131-6 (2014).
- (7) Thorn, C.F., Marsh, S., Carrillo, M.W., McLeod, H.L., Klein, T.E. & Altman, R.B. PharmGKB summary: fluoropyrimidine pathways. *Pharmacogenet Genomics* **21**, 237-42 (2011).
- (8) Schwab, M. *et al.* Role of genetic and nongenetic factors for fluorouracil treatment-related severe toxicity: a prospective clinical trial by the German 5-FU Toxicity Study Group. *J Clin Oncol* **26**, 2131-8 (2008).
- (9) Rosmarin, D. *et al.* A candidate gene study of capecitabine-related toxicity in colorectal cancer identifies new toxicity variants at DPYD and a putative role for ENOSF1 rather than TYMS. *Gut* **64**, 111-20 (2015).
- (10) Jennings, B.A., Kwok, C.S., Willis, G., Matthews, V., Wawruch, P. & Loke, Y.K. Functional polymorphisms of folate metabolism and response to chemotherapy for colorectal cancer, a systematic review and meta-analysis. *Pharmacogenet Genomics* **22**, 290-304 (2012).
- (11) Magdy, T. *et al.* ABCC11/MRP8 polymorphisms affect 5-fluorouracil-induced severe toxicity and hepatic expression. *Pharmacogenomics* **14**, 1433-48 (2013).
- (12) Hamzic, S. *et al.* The impact of ABCC11 polymorphisms on the risk of early-onset fluoropyrimidine toxicity. *The pharmacogenomics journal*, (2016).
- (13) Loganayagam, A. *et al.* Pharmacogenetic variants in the DPYD, TYMS, CDA and MTHFR genes are clinically significant predictors of fluoropyrimidine toxicity. *British journal of cancer* **108**, 2505-15 (2013).

- (14) Hamzic, S. *et al.* Novel Genetic Variants in Carboxylesterase 1 Predict Severe Early-Onset Capecitabine-Related Toxicity. *Clinical pharmacology and therapeutics*, (2017).
- (15) Fernandez-Rozadilla, C. *et al.* Pharmacogenomics in colorectal cancer: a genome-wide association study to predict toxicity after 5-fluorouracil or FOLFOX administration. *The pharmacogenomics journal* **13**, 209-17 (2013).
- (16) Valdes, R., Payne, D.A. & Linder, M.W. Laboratory analysis and application of pharmacogenetics to clinical practice. (Washington, DC, NACB, 2010).
- (17) Deenen, M.J. *et al.* Upfront Genotyping of DPYD\*2A to Individualize Fluoropyrimidine Therapy: A Safety and Cost Analysis. *J Clin Oncol* **34**, 227-34 (2016).
- (18) Shuldiner, A.R. *et al.* The Pharmacogenomics Research Network Translational Pharmacogenetics Program: overcoming challenges of real-world implementation. *Clinical pharmacology and therapeutics* **94**, 207-10 (2013).
- (19) Wilke, R.A. *et al.* The emerging role of electronic medical records in pharmacogenomics. *Clinical pharmacology and therapeutics* **89**, 379-86 (2011).
- (20) Peterson, J.F. *et al.* Electronic health record design and implementation for pharmacogenomics: a local perspective. *Genetics in medicine : official journal of the American College of Medical Genetics* **15**, 833-41 (2013).
- (21) Gottesman, O. *et al.* The Electronic Medical Records and Genomics (eMERGE) Network: past, present, and future. *Genetics in medicine : official journal of the American College of Medical Genetics* **15**, 761-71 (2013).
- (22) Kullo, I.J., Jarvik, G.P., Manolio, T.A., Williams, M.S. & Roden, D.M. Leveraging the electronic health record to implement genomic medicine. *Genetics in medicine : official journal of the American College of Medical Genetics* **15**, 270-1 (2013).
- (23) Hicks, J.K. *et al.* A clinician-driven automated system for integration of pharmacogenetic interpretations into an electronic medical record. *Clinical pharmacology and therapeutics* **92**, 563-6 (2012).
- (24) Hoffman, J.M. *et al.* Developing knowledge resources to support precision medicine: principles from the Clinical Pharmacogenetics Implementation Consortium (CPIC). *J Am Med Inform Assoc* **23**, 796-801 (2016).
- (25) Pulley, J.M. *et al.* Operational Implementation of Prospective Genotyping for Personalized Medicine: The Design of the Vanderbilt PREDICT Project. *Clinical pharmacology and therapeutics* **92**, 87-95 (2012).
- (26) PGx Gene-specific Information Tables for CYP2C19.  
<<https://www.pharmgkb.org/page/cyp2c19RefMaterials>>. Accessed September 16 2016.
- (27) Wei, X., McLeod, H.L., McMurrough, J., Gonzalez, F.J. & Fernandez-Salguero, P. Molecular basis of the human dihydropyrimidine dehydrogenase deficiency and 5-fluorouracil toxicity. *The Journal of clinical investigation* **98**, 610-5 (1996).
- (28) Kuilenburg, A.B. *et al.* Phenotypic and clinical implications of variants in the dihydropyrimidine dehydrogenase gene. *Biochim Biophys Acta* **1862**, 754-62 (2016).

- (29) Nie, Q. *et al.* Quantitative contribution of rs75017182 to dihydropyrimidine dehydrogenase mRNA splicing and enzyme activity. *Clinical pharmacology and therapeutics*, (2017).
- (30) Sistonen, J. *et al.* Predicting 5-fluorouracil toxicity: DPD genotype and 5,6-dihydrouracil:uracil ratio. *Pharmacogenomics* **15**, 1653-66 (2014).
- (31) Offer, S.M., Wegner, N.J., Fossum, C., Wang, K. & Diasio, R.B. Phenotypic Profiling of DPYD Variations Relevant to 5-Fluorouracil Sensitivity Using Real-time Cellular Analysis and In Vitro Measurement of Enzyme Activity. *Cancer Res* **73**, 1958-68 (2013).
- (32) Holopainen, I. *et al.* Partial epilepsy in a girl with a symptom-free sister: first two Finnish patients with dihydropyrimidine dehydrogenase deficiency. *J Inherit Metab Dis* **20**, 719-20 (1997).
- (33) Vreken, P., Van Kuilenburg, A.B., Meinsma, R. & van Gennip, A.H. Dihydropyrimidine dehydrogenase (DPD) deficiency: identification and expression of missense mutations C29R, R886H and R235W. *Human genetics* **101**, 333-8 (1997).
- (34) Van Kuilenburg, A.B. *et al.* Heterozygosity for a point mutation in an invariant splice donor site of dihydropyrimidine dehydrogenase and severe 5-fluorouracil related toxicity. *Eur J Cancer* **33**, 2258-64 (1997).
- (35) Ridge, S.A. *et al.* Dihydropyrimidine dehydrogenase pharmacogenetics in patients with colorectal cancer. *British journal of cancer* **77**, 497-500 (1998).
- (36) Van Kuilenburg, A.B. *et al.* Genotype and phenotype in patients with dihydropyrimidine dehydrogenase deficiency. *Human genetics* **104**, 1-9 (1999).
- (37) van Kuilenburg, A.B. *et al.* Lethal outcome of a patient with a complete dihydropyrimidine dehydrogenase (DPD) deficiency after administration of 5-fluorouracil: frequency of the common IVS14+1G>A mutation causing DPD deficiency. *Clinical cancer research : an official journal of the American Association for Cancer Research* **7**, 1149-53 (2001).
- (38) Johnson, M.R., Wang, K. & Diasio, R.B. Profound dihydropyrimidine dehydrogenase deficiency resulting from a novel compound heterozygote genotype. *Clinical cancer research : an official journal of the American Association for Cancer Research* **8**, 768-74 (2002).
- (39) Morita, J. *et al.* A novel single nucleotide polymorphism (SNP) of the CYP2C19 gene in a Japanese subject with lowered capacity of mephobarbital 4'-hydroxylation. *Drug metabolism and pharmacokinetics* **19**, 236-8 (2004).
- (40) van Kuilenburg, A.B. *et al.* Novel disease-causing mutations in the dihydropyrimidine dehydrogenase gene interpreted by analysis of the three-dimensional protein structure. *Biochem J* **364**, 157-63 (2002).
- (41) Van Kuilenburg, A.B., Meinsma, R., Zoetekouw, L. & Van Gennip, A.H. Increased risk of grade IV neutropenia after administration of 5-fluorouracil due to a dihydropyrimidine dehydrogenase deficiency: high prevalence of the IVS14+1g>a mutation. *Int J Cancer* **101**, 253-8 (2002).
- (42) Al-Sanna'a, N.A., Van Kuilenburg, A.B., Atrak, T.M., Abdul-Jabbar, M.A. & Van Gennip, A.H. Dihydropyrimidine dehydrogenase deficiency presenting at birth. *J Inherit Metab Dis* **28**, 793-6 (2005).

- (43) Ezzeldin, H.H., Lee, A.M., Mattison, L.K. & Diasio, R.B. Methylation of the DPYD promoter: an alternative mechanism for dihydropyrimidine dehydrogenase deficiency in cancer patients. *Clinical cancer research : an official journal of the American Association for Cancer Research* **11**, 8699-705 (2005).
- (44) Largillier, R. *et al.* Pharmacogenetics of capecitabine in advanced breast cancer patients. *Clinical cancer research : an official journal of the American Association for Cancer Research* **12**, 5496-502 (2006).
- (45) Magne, N. *et al.* Dihydropyrimidine dehydrogenase activity and the IVS14+1G>A mutation in patients developing 5FU-related toxicity. *Br J Clin Pharmacol* **64**, 237-40 (2007).
- (46) Loganayagam, A., Arenas-Hernandez, M., Fairbanks, L., Ross, P., Sanderson, J.D. & Marinaki, A.M. The contribution of deleterious DPYD gene sequence variants to fluoropyrimidine toxicity in British cancer patients. *Cancer chemotherapy and pharmacology* **65**, 403-6 (2010).
- (47) van Kuilenburg, A.B. *et al.* Intragenic deletions and a deep intronic mutation affecting pre-mRNA splicing in the dihydropyrimidine dehydrogenase gene as novel mechanisms causing 5-fluorouracil toxicity. *Human genetics* **128**, 529-38 (2010).
- (48) Thomas, F. *et al.* Genotyping of a family with a novel deleterious DPYD mutation supports the pretherapeutic screening of DPD deficiency with dihydrouracil/uracil ratio. *Clinical pharmacology and therapeutics* **99**, 235-42 (2016).
- (49) Collie-Duguid, E.S., Etienne, M.C., Milano, G. & McLeod, H.L. Known variant DPYD alleles do not explain DPD deficiency in cancer patients. *Pharmacogenetics* **10**, 217-23 (2000).
- (50) Deenen, M.J. *et al.* Relationship between single nucleotide polymorphisms and haplotypes in DPYD and toxicity and efficacy of capecitabine in advanced colorectal cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research* **17**, 3455-68 (2011).
- (51) Joerger, M. *et al.* Germline TYMS genotype is highly predictive in patients with metastatic gastrointestinal malignancies receiving capecitabine-based chemotherapy. *Cancer chemotherapy and pharmacology* **75**, 763-72 (2015).
- (52) Zhao, X.Q. *et al.* DPYD gene polymorphisms are associated with risk and chemotherapy prognosis in pediatric patients with acute lymphoblastic leukemia. *Tumour Biol* **37**, 10393-402 (2016).
- (53) McLeod, H.L. *et al.* Pharmacogenetic predictors of adverse events and response to chemotherapy in metastatic colorectal cancer: results from North American Gastrointestinal Intergroup Trial N9741. *J Clin Oncol* **28**, 3227-33 (2010).
- (54) Cai, X. *et al.* The role of IVS14+1 G > A genotype detection in the dihydropyrimidine dehydrogenase gene and pharmacokinetic monitoring of 5-fluorouracil in the individualized adjustment of 5-fluorouracil for patients with local advanced and metastatic colorectal cancer: a preliminary report. *Eur Rev Med Pharmacol Sci* **18**, 1247-58 (2014).

- (55) Boisdron-Celle, M. *et al.* 5-Fluorouracil-related severe toxicity: a comparison of different methods for the pretherapeutic detection of dihydropyrimidine dehydrogenase deficiency. *Cancer letters* **249**, 271-82 (2007).
- (56) van Kuilenburg, A.B. *et al.* Pharmacokinetics of 5-fluorouracil in patients heterozygous for the IVS14+1G > A mutation in the dihydropyrimidine dehydrogenase gene. *Nucleosides Nucleotides Nucleic Acids* **27**, 692-8 (2008).
- (57) Gentile, G. *et al.* Genotype-phenotype correlations in 5-fluorouracil metabolism: a candidate DPYD haplotype to improve toxicity prediction. *The pharmacogenomics journal* **16**, 320-5 (2016).
- (58) Maring, J.G. *et al.* Reduced 5-FU clearance in a patient with low DPD activity due to heterozygosity for a mutant allele of the DPYD gene. *British journal of cancer* **86**, 1028-33 (2002).
- (59) Van Kuilenburg, A.B., Meinsma, R., Zoetekouw, L. & Van Gennip, A.H. High prevalence of the IVS14 + 1G>A mutation in the dihydropyrimidine dehydrogenase gene of patients with severe 5-fluorouracil-associated toxicity. *Pharmacogenetics* **12**, 555-8 (2002).
- (60) Salgueiro, N. *et al.* Mutations in exon 14 of dihydropyrimidine dehydrogenase and 5-Fluorouracil toxicity in Portuguese colorectal cancer patients. *Genetics in medicine : official journal of the American College of Medical Genetics* **6**, 102-7 (2004).
- (61) Schwab, M. *et al.* Role of genetic and nongenetic factors for fluorouracil treatment-related severe toxicity: a prospective clinical trial by the German 5-FU Toxicity Study Group. *J Clin Oncol* **26**, 2131-8 (2008).
- (62) Lee, A.M. *et al.* DPYD variants as predictors of 5-fluorouracil toxicity in adjuvant colon cancer treatment (NCCTG N0147). *J Natl Cancer Inst* **106**, (2014).
- (63) Toffoli, G. *et al.* Clinical validity of a DPYD-based pharmacogenetic test to predict severe toxicity to fluoropyrimidines. *Int J Cancer* **137**, 2971-80 (2015).
- (64) Kleibl, Z. *et al.* Influence of dihydropyrimidine dehydrogenase gene (DPYD) coding sequence variants on the development of fluoropyrimidine-related toxicity in patients with high-grade toxicity and patients with excellent tolerance of fluoropyrimidine-based chemotherapy. *Neoplasma* **56**, 303-16 (2009).
- (65) Kristensen, M.H., Pedersen, P.L., Melsen, G.V., Ellehauge, J. & Mejer, J. Variants in the dihydropyrimidine dehydrogenase, methylenetetrahydrofolate reductase and thymidylate synthase genes predict early toxicity of 5-fluorouracil in colorectal cancer patients. *J Int Med Res* **38**, 870-83 (2010).
- (66) Rosmarin, D. *et al.* Genetic markers of toxicity from capecitabine and other fluorouracil-based regimens: investigation in the QUASAR2 study, systematic review, and meta-analysis. *J Clin Oncol* **32**, 1031-9 (2014).
- (67) Braun, M.S. *et al.* Association of molecular markers with toxicity outcomes in a randomized trial of chemotherapy for advanced colorectal cancer: the FOCUS trial. *J Clin Oncol* **27**, 5519-28 (2009).

- (68) Dhawan, D., Panchal, H., Shukla, S. & Padh, H. Genetic variability & chemotoxicity of 5-fluorouracil & cisplatin in head & neck cancer patients: a preliminary study. *Indian J Med Res* **137**, 125-9 (2013).
- (69) Boige, V. *et al.* DPYD Genotyping to Predict Adverse Events Following Treatment With Fluorouracil-Based Adjuvant Chemotherapy in Patients With Stage III Colon Cancer: A Secondary Analysis of the PETACC-8 Randomized Clinical Trial. *JAMA Oncol*, (2016).
- (70) van Kuilenburg, A.B. *et al.* Clinical implications of dihydropyrimidine dehydrogenase (DPD) deficiency in patients with severe 5-fluorouracil-associated toxicity: identification of new mutations in the DPD gene. *Clinical cancer research : an official journal of the American Association for Cancer Research* **6**, 4705-12 (2000).
- (71) Raida, M. *et al.* Prevalence of a common point mutation in the dihydropyrimidine dehydrogenase (DPD) gene within the 5'-splice donor site of intron 14 in patients with severe 5-fluorouracil (5-FU)- related toxicity compared with controls. *Clinical cancer research : an official journal of the American Association for Cancer Research* **7**, 2832-9 (2001).
- (72) Steiner, M. *et al.* 5-Fluorouracil/irinotecan induced lethal toxicity as a result of a combined pharmacogenetic syndrome: report of a case. *J Clin Pathol* **58**, 553-5 (2005).
- (73) Morel, A. *et al.* Clinical relevance of different dihydropyrimidine dehydrogenase gene single nucleotide polymorphisms on 5-fluorouracil tolerance. *Molecular cancer therapeutics* **5**, 2895-904 (2006).
- (74) Saif, M.W., Ezzeldin, H., Vance, K., Sellers, S. & Diasio, R.B. DPYD\*2A mutation: the most common mutation associated with DPD deficiency. *Cancer chemotherapy and pharmacology* **60**, 503-7 (2007).
- (75) Salgado, J., Zabalegui, N., Gil, C., Monreal, I., Rodriguez, J. & Garcia-Foncillas, J. Polymorphisms in the thymidylate synthase and dihydropyrimidine dehydrogenase genes predict response and toxicity to capecitabine-raltitrexed in colorectal cancer. *Oncol Rep* **17**, 325-8 (2007).
- (76) Sulzyc-Bielicka, V. *et al.* 5-Fluorouracil toxicity-attributable IVS14 + 1G > A mutation of the dihydropyrimidine dehydrogenase gene in Polish colorectal cancer patients. *Pharmacol Rep* **60**, 238-42 (2008).
- (77) Gross, E. *et al.* Strong association of a common dihydropyrimidine dehydrogenase gene polymorphism with fluoropyrimidine-related toxicity in cancer patients. *PLoS One* **3**, e4003 (2008).
- (78) Boige, V. *et al.* Pharmacogenetic assessment of toxicity and outcome in patients with metastatic colorectal cancer treated with LV5FU2, FOLFOX, and FOLFIRI: FFCD 2000-05. *J Clin Oncol* **28**, 2556-64 (2010).
- (79) Ceric, T. *et al.* Investigation of IVS14 + 1G > A polymorphism of DPYD gene in a group of Bosnian patients treated with 5-Fluorouracil and capecitabine. *Bosn J Basic Med Sci* **10**, 133-9 (2010).
- (80) Savva-Bordalo, J. *et al.* Promoter methylation and large intragenic rearrangements of DPYD are not implicated in severe toxicity to 5-fluorouracil-based chemotherapy in gastrointestinal cancer patients. *BMC cancer* **10**, 470 (2010).

- (81) Suarez Martinez-Falero, B. & Gillmore, R. A rare cause of susceptibility to neutropenic sepsis in a patient with metastatic pancreas cancer. *BMJ Case Rep* **2014**, (2014).
- (82) Roberto, M. *et al.* Evaluation of 5-fluorouracil degradation rate and Pharmacogenetic profiling to predict toxicity following adjuvant Capecitabine. *Eur J Clin Pharmacol* **73**, 157-64 (2017).
- (83) Cellier, P. *et al.* Phase II study of preoperative radiation plus concurrent daily tegafur-uracil (UFT) with leucovorin for locally advanced rectal cancer. *BMC cancer* **11**, 98 (2011).
- (84) Dhelens, C. *et al.* Lethal 5-fluorouracil toxicity in a colorectal patient with severe dihydropyrimidine dehydrogenase (DPD) deficiency. *Int J Colorectal Dis* **31**, 699-701 (2016).
- (85) Seck, K. *et al.* Analysis of the DPYD gene implicated in 5-fluorouracil catabolism in a cohort of Caucasian individuals. *Clinical cancer research : an official journal of the American Association for Cancer Research* **11**, 5886-92 (2005).
- (86) Offer, S.M., Fossum, C.C., Wegner, N.J., Stuflesser, A.J., Butterfield, G.L. & Diasio, R.B. Comparative functional analysis of DPYD variants of potential clinical relevance to dihydropyrimidine dehydrogenase activity. *Cancer Res* **74**, 2545-54 (2014).
- (87) Obi, E.E., McDonald, A. & Kemp, E. A bilateral cicatricial ectropion and bilateral upper lid shortening caused by 5-fluorouracil toxicity in a patient with dihydropyrimidine dehydrogenase deficiency. *Cutan Ocul Toxicol* **30**, 157-9 (2011).
- (88) Meulendijks, D. *et al.* Patients homozygous for DPYD c.1129-5923C>G/haplotype B3 have partial DPD deficiency and require a dose reduction when treated with fluoropyrimidines. *Cancer chemotherapy and pharmacology* **78**, 875-80 (2016).
- (89) Lee, A.M. *et al.* Association between DPYD c.1129-5923 C>G/hapB3 and severe toxicity to 5-fluorouracil-based chemotherapy in stage III colon cancer patients: NCCTG N0147 (Alliance). *Pharmacogenet Genomics* **26**, 133-7 (2016).
- (90) Falvella, F.S. *et al.* DPD and UGT1A1 deficiency in colorectal cancer patients receiving triplet chemotherapy with fluoropyrimidines, oxaliplatin and irinotecan. *Br J Clin Pharmacol* **80**, 581-8 (2015).
- (91) Saif, M.W. Dihydropyrimidine dehydrogenase gene (DPYD) polymorphism among Caucasian and non-Caucasian patients with 5-FU- and capecitabine-related toxicity using full sequencing of DPYD. *Cancer Genomics Proteomics* **10**, 89-92 (2013).
- (92) Capitain, O., Boisdron-Celle, M., Poirier, A.L., Abadie-Lacourtoisie, S., Morel, A. & Gamelin, E. The influence of fluorouracil outcome parameters on tolerance and efficacy in patients with advanced colorectal cancer. *The pharmacogenomics journal* **8**, 256-67 (2008).
- (93) Jennings, B.A. *et al.* Evaluating predictive pharmacogenetic signatures of adverse events in colorectal cancer patients treated with fluoropyrimidines. *PLoS One* **8**, e78053 (2013).

- (94) Gross, E. *et al.* Detailed analysis of five mutations in dihydropyrimidine dehydrogenase detected in cancer patients with 5-fluorouracil-related side effects. *Human mutation* **22**, 498 (2003).
- (95) Lazar, A., Mau-Holzmann, U.A., Kolb, H., Reichenmiller, H.E., Riess, O. & Schomig, E. Multiple organ failure due to 5-fluorouracil chemotherapy in a patient with a rare dihydropyrimidine dehydrogenase gene variant. *Onkologie* **27**, 559-62 (2004).
- (96) He, Y.F. *et al.* Analysis of the DPYD gene implicated in 5-fluorouracil catabolism in Chinese cancer patients. *J Clin Pharm Ther* **33**, 307-14 (2008).
- (97) Zhang, H., Li, Y.M., Zhang, H. & Jin, X. DPYD\*5 gene mutation contributes to the reduced DPYD enzyme activity and chemotherapeutic toxicity of 5-FU: results from genotyping study on 75 gastric carcinoma and colon carcinoma patients. *Med Oncol* **24**, 251-8 (2007).
- (98) Teh, L.K. *et al.* Potential of dihydropyrimidine dehydrogenase genotypes in personalizing 5-fluorouracil therapy among colorectal cancer patients. *Ther Drug Monit* **35**, 624-30 (2013).
- (99) Rudek, M.A. *et al.* Fixed-dose capecitabine is feasible: results from a pharmacokinetic and pharmacogenetic study in metastatic breast cancer. *Breast Cancer Res Treat* **139**, 135-43 (2013).
- (100) Grau, J.J. *et al.* Dihydropyrimidine dehydrogenases and cytidine-deaminase gene polymorphisms as outcome predictors in resected gastric cancer patients treated with fluoropyrimidine adjuvant chemotherapy. *J Surg Oncol* **98**, 130-4 (2008).
- (101) Zhang, X.P. *et al.* Polymorphisms of dihydropyrimidine dehydrogenase gene and clinical outcomes of gastric cancer patients treated with fluorouracil-based adjuvant chemotherapy in Chinese population. *Chin Med J (Engl)* **125**, 741-6 (2012).
- (102) Farina-Sarasqueta, A., van Lijnschoten, G., Rutten, H.J. & van den Brule, A.J. Value of gene polymorphisms as markers of 5-FU therapy response in stage III colon carcinoma: a pilot study. *Cancer chemotherapy and pharmacology* **66**, 1167-71 (2010).
- (103) Cho, H.J., Park, Y.S., Kang, W.K., Kim, J.W. & Lee, S.Y. Thymidylate synthase (TYMS) and dihydropyrimidine dehydrogenase (DPYD) polymorphisms in the Korean population for prediction of 5-fluorouracil-associated toxicity. *Ther Drug Monit* **29**, 190-6 (2007).
- (104) Kim, S.R., Park, C.H., Park, S., Park, J.O., Lee, J. & Lee, S.Y. Genetic polymorphisms associated with 5-Fluorouracil-induced neurotoxicity. *Chemotherapy* **56**, 313-7 (2010).
- (105) Zaanani, A., Dumont, L.M., Lorient, M.A., Taieb, J. & Narjoz, C. A case of 5-FU-related severe toxicity associated with the p.Y186C DPYD variant. *Clinical pharmacology and therapeutics* **95**, 136 (2014).
- (106) Del Re, M. *et al.* Discovery of novel mutations in the dihydropyrimidine dehydrogenase gene associated with toxicity of fluoropyrimidines and viewpoint on preemptive pharmacogenetic screening in patients. *EPMA J* **6**, 17 (2015).



- (107) Van Kuilenburg, A.B. *et al.* Clinical and biochemical abnormalities in a patient with dihydropyrimidine dehydrogenase deficiency due to homozygosity for the C29R mutation. *J Inherit Metab Dis* **22**, 191-2 (1999).
- (108) Baskin, Y., Amirfallah, A., Unal, O.U., Calibasi, G. & Oztop, I. Dihydropyrimidine dehydrogenase 85T>C mutation is associated with ocular toxicity of 5-fluorouracil: a case report. *Am J Ther* **22**, e36-9 (2015).
- (109) Offer, S.M. & Diasio, R.B. Response to "A case of 5-FU-related severe toxicity associated with the P.Y186C DPYD variant". *Clinical pharmacology and therapeutics* **95**, 137 (2014).
- (110) Vreken, P., Van Kuilenburg, A.B., Meinsma, R. & van Gennip, A.H. Identification of novel point mutations in the dihydropyrimidine dehydrogenase gene. *J Inherit Metab Dis* **20**, 335-8 (1997).
- (111) Kouwaki, M. *et al.* Identification of novel mutations in the dihydropyrimidine dehydrogenase gene in a Japanese patient with 5-fluorouracil toxicity. *Clinical cancer research : an official journal of the American Association for Cancer Research* **4**, 2999-3004 (1998).
- (112) Ofverholm, A., Arkblad, E., Skrtic, S., Albertsson, P., Shubbar, E. & Enerback, C. Two cases of 5-fluorouracil toxicity linked with gene variants in the DPYD gene. *Clin Biochem* **43**, 331-4 (2010).