




HUST

ĐẠI HỌC BÁCH KHOA HÀ NỘI
HANOI UNIVERSITY OF SCIENCE AND TECHNOLOGY

ONE LOVE. ONE FUTURE.



Discrete Mathematics

PART 1

COMBINATORIAL THEORY

(Lý thuyết tổ hợp)

PART 2

GRAPH THEORY

(Lý thuyết đồ thị)

Content of Part 2

Chapter 1. Fundamental concepts

Chapter 2. Graph representation

Chapter 3. Graph Traversal

Chapter 4. Tree and Spanning tree

Chapter 5. Shortest path problem

Chapter 6. Maximum flow problem




Chapter 1. Fundamental concepts

1. Graph in practice

2. Graph types
3. Degree of vertex
4. Subgraph
5. Isomorphism of Graphs
6. Path and cycle
7. Connectedness
8. Special graphs
9. Graph Coloring problem

What is graph?


- In Maths:
Drawing or diagram represent data by using coordinate system



Not what we want to mention



- In discrete mathematics:
This is discrete structure with highly intuitive, very useful for expressing relationships.




Applications of graph in practice



Has the potential to be applied in many fields:

- Internet
- Traffic network
- Electrical network
- Water supply network
- Scheduling
- Flow optimization, circuit design
- DNA gene analysis
- Computer games
- Object oriented design
-

Relationship between subjects




Represent maze



Vertex = room


Edge = doorways or hallways



Vertex = source, switch, resistor, ...

Edge = connecting wire

Information Transmission in a Computer Network



Vertex = computer

Edge = communication speed

Traffic Flow on Highways




Vertex = city
Edge = the amount of traffic on highways connecting cities

Subway system



Street map



Chapter 1. Fundamental concepts

1. Graph in practice

2. Graph types

3. Degree of vertex

4. Subgraph

5. Isomorphism of Graphs

6. Path and cycle

7. Connectedness


8. Special graphs

9. Graph Coloring problem

Undirected Graphs

Definition. Undirected simple (**multi**) graph $G = (V, E)$ consists of 2 sets:

- Vertex set V is a finite set, each element is called **vertex**
- Edge set E is the set (**family**) of unordered pairs (u, v) , where $u, v \in V, u \neq v$




Undirected Multigraph

Undirected (simple) graph

(a, b) : multiple edges (parallel edges)

Undirected Simple Graph

- **Example:** Simple graph $G_1 = (V_1, E_1)$, where
 $V_1 = \{a, b, c, d, e, f, g, h\}$,
 $E_1 = \{(a,b), (b,c), (c,d), (a,d), (d,e), (a,e), (d,b), (f,g)\}$.




Graph G_1

Undirected MultiGraph

- **Example:** Multigraph $G_2 = (V_2, E_2)$, where

$$V_2 = \{a, b, c, d, e, f, g, h\},$$

$$E_2 = \{(a, b), (b, c), (b, c), (c, d), (a, d), (d, e), (a, e), (a, e), (a, e), (d, b), (f, g)\}.$$




Graph G_2

Directed Graph

Definition. Directed simple (**multi**) graph $G = (V, E)$ consists of 2 sets:

- Vertex set V is finite element, each element is called vertex
- Edge set E is set (**family**) of ordered pairs
 (u, v) , where $u, v \in V, u \neq v$



Directed multigraph


(Simple) Directed graph ~ Digraph

Simple digraph

Example: Simple digraph $G_3 = (V_3, E_3)$, where

$V_3 = \{a, b, c, d, e, f, g, h\}$,

$E_3 = \{(a, b), (b, c), (c, b), (d, c), (a, d), (b, d), (a, e), (d, e), (e, a), (f, g), (g, f)\}$




Graph G_3

Directed MultiGraph

Example: Directed multigraph $G_4 = (V_4, E_4)$, where

$$V_4 = \{a, b, c, d, e, f, g, h\},$$

$$E_4 = \{(a, b), (b, c), (c, b), (d, c), (a, d), (b, d), (a, e), (e, a), (d, e), (e, a), (f, g), (g, f)\}$$




Graph G_4


Graph Terminology

We have graph terminology related to relationship between vertices and edges:

- *Adjacency, connect, degree, start, end, indegree, outdegree,...*



Undirected edge $e=(u,v)$




Directed edge $e=(u,v)$

Adjacency

Given G the undirected graph with edge set E . Assume $e \in E$ is pair (u, v) .


Then we say:

- u, v are *adjacent / neighbors / connected*
- Edge e *connects* u and v .
- Vertices u and v are *endpoints* (đầu mút) of e .



Adjacency in directed graph

- G is directed graph (either simple or multiple) and assume $e = (u,v)$ is an edge of G . We say:
 - u and v are adjacent, u is adjacent to v , v is adjacent from u
 - e goes out of u , e goes into v .
 - e connects u with v , e goes from u to v
 - initial vertex of e is u
 - terminal vertex of e is v



Chapter 1. Fundamental concepts

1. Graph in practice
2. Graph types
- 3. Degree of vertex**
4. Subgraph
5. Isomorphism of Graphs
6. Path and cycle
7. Connectedness
8. Special graphs
9. Graph Coloring problem

Degree of a vertex in undirected graph

Assume G is undirected graph, $v \in V$ is a vertex.

- Degree of vertex v , $\deg(v)$, the number of edges incident on a vertex.
- Vertex with degree 0 is called isolated.
- Vertex with degree 1 is called pendant.
- Symbol often used:


$$\delta(G) = \min \{\deg(v): v \in V\},$$

$$\Delta(G) = \max \{\deg(v): v \in V\}.$$

Example

Edge(a, b) is **incident** with
two vertices a and b

b is adjacent to c and c is adjacent to b



$\deg(d) = 3$

$\deg(g) = 1$
g is pendant

$\delta(G) = \min \{\deg(v) : v \in V\} = 0,$
 $\Delta(G) = \max \{\deg(v) : v \in V\} = 3.$

Theorem: Undirected graph

Theorem 1 (The Handshaking theorem):

The undirected Graph $G = (V, E)$:

$$\sum_{v \in V} \deg(v) = 2|E|$$

Theorem 2:


An undirected graph has even number of vertices with odd degree

Degree of a vertex in directed graph

Given G directed graph, v is a vertex of G :

- *In-degree* of v , $\deg^-(v)$, number of edges for which v is terminal vertex (goes into v).
- *Out-degree* of v , $\deg^+(v)$, number of edges for which v is initial vertex (goes out of v).
- *Degree* of v , $\deg(v) = \deg^-(v) + \deg^+(v)$

Example



Theorem: Directed graph

Theorem 3 The directed Graph (either simple or multiple) $G = (V, E)$:

$$\sum_{v \in V} \deg^{-}(v) = \sum_{v \in V} \deg^{+}(v) = \frac{1}{2} \sum_{v \in V} \deg(v) = |E|$$

Chapter 1. Fundamental concepts


1. Graph in practice
2. Graph types
3. Degree of vertex
- 4. Subgraph**
5. Isomorphism of Graphs
6. Path and cycle
7. Connectedness
8. Special graphs
9. Graph Coloring problem

4. Subgraph

A **subgraph** of a graph $G = (V, E)$ is a graph $H = (V', E')$ where V' is a subset of V and E' is a subset of E .

Denoted by: $H \subseteq G$

Example: $V = \{u, v, w\}$, $E = \{\{u, v\}, \{v, w\}, \{w, u\}\}$



Spanning Subgraph

Definition.

Subgraph $H \subseteq G$ is called spanning subgraph of G if vertex set of H is vertex set of G : $V(H) = V(G)$.

Definition.


We write $H = G + \{(u,v), (u,w)\}$ to mean

$E(H) = E(G) \cup \{(u,v), (u,w)\}$, where $(u,v), (u,w) \notin E(G)$.

The union of graphs

The **union** of two simple graphs $G_1=(V_1, E_1)$ and $G_2=(V_2, E_2)$ is the simple graph $G_1 \cup G_2=(V_1 \cup V_2, E_1 \cup E_2)$.



Example:



The union of graphs

The **union** of two simple graphs $G_1=(V_1, E_1)$ and $G_2=(V_2, E_2)$ is the simple graph $G_1 \cup G_2=(V_1 \cup V_2, E_1 \cup E_2)$.



Example:



Chapter 1. Fundamental concepts

1. Graph in practice
2. Graph types
3. Degree of vertex
4. Subgraph
- 5. Isomorphism of Graphs**
6. Path and cycle
7. Connectedness
8. Special graphs
9. Graph Coloring problem

Isomorphism of Graphs




G is isomorphic to H

Definition. The simple graphs $G_1=(V_1,E_1)$ and $G_2=(V_2,E_2)$ are **isomorphic** if there is an one-to-one and onto function f from V_1 to V_2 with the property that a and b is adjacent in G_1 iff $f(a)$ and $f(b)$ is adjacent in G_2 , $\forall a,b \in V_1$

f is called an isomorphism.


Application Example:

In chemistry, to find if two compounds have the same structure




Example

Show that G and H are isomorphic.



G



H

Solution.

The function f with $f(u_1) = v_1$, $f(u_2) = v_4$, $f(u_3) = v_3$, and $f(u_4) = v_2$ is a one-to-one correspondence between $V(G)$ and $V(H)$.

Isomorphism graphs there will be:

- (1) The same number of vertices
- (2) The same number of edges
- (3) The same number of degree

Chapter 1. Fundamental concepts

1. Graph in practice
2. Graph types
3. Degree of vertex
4. Subgraph
5. Isomorphism of Graphs
- 6. Path and cycle**
7. Connectedness
8. Special graphs
9. Graph Coloring problem

Definition for Directed Graphs


A **Path** of length $n > 0$ from u to v in G is a sequence of n edges $e_1, e_2, e_3, \dots, e_n$ of G such that $f(e_1) = (x_0, x_1), f(e_2) = (x_1, x_2), \dots, f(e_n) = (x_{n-1}, x_n)$, where $x_0 = u$ and $x_n = v$.

A path is said to pass through $x_0, x_1, \dots, x_n$ or traverse $e_1, e_2, e_3, \dots, e_n$


- A path is called **elementary** if all the edges are distinct.
- A path is called **simple** if all the vertices are distinct.
- A path is **closed** if $v_0 = v_n$.
- A closed elementary path is called a **cycle**. A cycle is called simple if all the vertices are distinct (except $v_0 = v_n$).

A path is called *elementary* if all the edges are distinct.
A path is called *simple* if all the vertices are distinct.

Path: 5, 2, 3, 4 [OR: {5, 2}, {2, 3}, {3, 4}]
OR: c,b,e]
all the vertices are distinct → simple path




Path (directed) 1, 2, 5, 3, 4
[OR: (1, 2), (2, 5), (5, 3), (3, 4)]
OR: a, c, d, e]
• is the simple path



Path

- $P_1 = (1, 2, 3)$ is the simple path
- $P_2 = (4, 5, 2, 6, 5, 1)$ is the path but not the simple path




Cycle

- A closed elementary path is called a *cycle*.
 - A path is called *elementary* if all the edges are distinct.
 - A path is *closed* if $v_0 = v_n$.
- A cycle is called simple if all the vertices are distinct (except $v_0 = v_n$).


Simple cycle: (1, 2, 3, 1)

Simple cycle: (1, 2, 3, 4, 1)



Cycle on directed graph

- $C_1=(V,X,Y,W,U,V)$ is simple cycle
- $C_2=(U,W,X,Y,W,V,U)$ is cycle but not simple cycle




Chapter 1. Fundamental concepts

1. Graph in practice
2. Graph types
3. Degree of vertex
4. Subgraph
5. Isomorphism of Graphs
6. Path and cycle
- 7. Connectedness**
8. Special graphs
9. Graph Coloring problem

Undirected Graph

An undirected graph is **connected** if there exists a simple path between every pair of vertices


Example: $G(V, E)$ is connected since for $V = \{v_1, v_2, v_3, v_4, v_5\}$, there exists a path between $\{v_i, v_j\}$, $1 \leq i, j \leq 5$



Connectedness

Articulation Point (Cut vertex): removal of a vertex produces a subgraph with more connected components than in the original graph. The removal of a cut vertex from a connected graph produces a graph that is not connected



Bridge: An edge whose removal produces a subgraph with more connected components than in the original graph.



Connectedness

- If a graph is not connected then it splits up into a number of connected subgraphs, called its *connected components*.
- The connected components of G can be defined as its *maximal connected subgraphs*. This means that G_1 is a connected component of G if:
 - G_1 is a connected subgraph of G
 - G_1 is not itself a proper subgraph of any other *connected* subgraph of G . This second condition is what we mean by the term maximal; it says that if H is a connected subgraph such that $G_1 \subseteq H$, then $G_1 = H$.

Example: Graph G has 3 connected components: G_1 , G_2 , G_3




Connectedness

Directed Graph

- A directed graph is **strongly connected** if there is a path from u to v and from v to u whenever u and v are vertices in the graph
- A directed graph is **weakly connected** if its corresponding undirected graph is connected.

A strongly connected Graph can be weakly connected but the vice-versa is not true (why?)




Weakly connected graph


Directed Graph

- **Strongly connected Components:** subgraphs of a Graph G that are strongly connected

Example: $G1$ is the strongly connected component in G



G



$G1$

Chapter 1. Fundamental concepts

1. Graph in practice
2. Graph types
3. Degree of vertex
4. Subgraph
5. Isomorphism of Graphs
6. Path and cycle
7. Connectedness
- 8. Special graphs**
9. Graph Coloring problem


Some special graphs

1. Null graph
2. Complete graphs K_n
3. Cycles C_n
4. Wheels W_n
5. n -Cubes Q_n
6. Bipartite graphs
7. Complete bipartite graphs $K_{m,n}$
8. r -regular graph
9. Planar graph
10. Euler graph and Hamilton graph

Planar Graphs

- A graph is called *planar* if it can be drawn in the plane without any edges crossing.
 - A crossing of edges is the intersection of the lines or arcs representing them at a point other than their common endpoint.
 - Such a drawing is called a *planar representation* of the graph.



Example: K_4 is planar graph?



Yes, K_4 is planar graph

Example: Three house and three utilities problem

- Is it possible to join the three houses to the three utilities in such a way that none of the connections cross?





$K_{3,3}$

→ Phrased another way, this question is equivalent to: Given the complete bipartite graph $K_{3,3}$, can $K_{3,3}$ be drawn in the plane so that no two of its edges cross? ($K_{3,3}$ is planar graph?)

$K_{3,3}$ and K_5 are not planar graphs

- $K_{3,3}$ and K_5 are not planar graphs.




- We can prove that a particular graph is planar by showing how it can be drawn without any crossings.
- However, not all graphs are planar.
- It may be difficult to show that a graph is nonplanar. We would have to show that there is *no* way to draw the graph without any edges crossing.


Regions

- Euler showed that all planar representations of a graph split the plane into the same number of *regions*, including an unbounded region.

Example:




G_1 has 3 regions.



G_2 has 4 regions.




How to proof $K_{3,3}$ is not planar graph ?

- In any planar representation of $K_{3,3}$, vertex v_1 must be connected to both v_4 and v_5 , and v_2 also must be connected to both v_4 and v_5 .



How to proof $K_{3,3}$ is not planar graph ?

- The four edges $\{v_1, v_4\}, \{v_4, v_2\}, \{v_2, v_5\}, \{v_5, v_1\}$ form a closed curve that splits the plane into two regions, R_1 and R_2 .





- Next, we note that v_3 must be in either R_1 or R_2 .

Case 1: Assume v_3 is in R_2 . Then the edges $\{v_3, v_4\}$ and $\{v_4, v_5\}$ separate R_2 into two subregions, R_{21} and R_{22} .



How to proof $K_{3,3}$ is not planar graph ?

- Now there is no way to place vertex v_6 without forcing a crossing:
 - If v_6 is in R_1 then $\{v_6, v_3\}$ must cross an edge
 - If v_6 is in R_{21} then $\{v_6, v_2\}$ must cross an edge
 - If v_6 is in R_{22} then $\{v_6, v_1\}$ must cross an edge



How to proof $K_{3,3}$ is not planar graph ?

- The four edges $\{v_1, v_4\}, \{v_4, v_2\}, \{v_2, v_5\}, \{v_5, v_1\}$ form a closed curve that splits the plane into two regions, R_1 and R_2 .





- Next, we note that v_3 must be in either R_1 or R_2 .

Case 2: Assume v_3 is in R_1 . Then the edges $\{v_3, v_4\}$ and $\{v_4, v_5\}$ separate R_1 into two subregions, R_{11} and R_{12} .


How to proof $K_{3,3}$ is not planar graph ?

- Now there is no way to place vertex v_6 without forcing a crossing:
 - If v_6 is in R_2 then $\{v_6, v_3\}$ must cross an edge
 - If v_6 is in R_{11} then $\{v_6, v_2\}$ must cross an edge
 - If v_6 is in R_{12} then $\{v_6, v_1\}$ must cross an edge



How to proof $K_{3,3}$ is not planar graph ?

- Consequently, the graph $K_{3,3}$ must be nonplanar.




$K_{3,3}$

- Euler devised a formula for expressing the relationship between the number of vertices, edges, and regions of a planar graph.
- These *may* help us determine if a graph can be planar or not.

Euler's Formula

- Let G be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.



of edges, $e = 6$

of vertices, $v = 4$

of regions, $r = e - v + 2 = 4$


Euler's Formula (Cont.)

- **Corollary 1:** If G is a connected planar simple graph with e edges and v vertices where $v \geq 3$, then $e \leq 3v - 6$.

Is K_5 planar?

- K_5 has 5 vertices and 10 edges.
- We see that $v \geq 3$.
- So, if K_5 is planar, it must be true that $e \leq 3v - 6$.
- $3v - 6 = 3 \cdot 5 - 6 = 15 - 6 = 9$.
- So e must be ≤ 9 .
- But $e = 10$.
- So, K_5 is nonplanar.

K_5



Euler's Formula (Cont.)

- **Corollary 2:** If G is a connected planar simple graph, then G must have a vertex of degree not exceeding 5.

If G has one or two vertices, it is true;
thus, we assume that G has at least three vertices.

If the degree of each vertex were at least 6, then by Handshaking Theorem,
 $2e \geq 6v$, i.e., $e \geq 3v$,

but this contradicts the inequality from
Corollary 1: $e \leq 3v - 6$.


$$2e = \sum_{v \in V} \deg(v)$$

Euler's Formula (Cont.)

- **Corollary 3:** If a connected planar simple graph has e edges and v vertices with $v \geq 3$ and no circuits of length 3, then $e \leq 2v - 4$.

Is $K_{3,3}$ planar?

- $K_{3,3}$ has 6 vertices and 9 edges.
- Obviously, $v \geq 3$ and there are no circuits of length 3.
- If $K_{3,3}$ were planar, then $e \leq 2v - 4$ would have to be true.
- $2v - 4 = 2 \cdot 6 - 4 = 8$
- So e must be ≤ 8 .
- But $e = 9$.
- So $K_{3,3}$ is nonplanar.




Some special graphs

1. Null graph
2. Complete graphs K_n
3. Cycles C_n
4. Wheels W_n
5. n -Cubes Q_n
6. Bipartite graphs
7. Complete bipartite graphs $K_{m,n}$
8. r -regular graph
9. Planar graph
- 10. Euler graph and Hamilton graph**

Seven Bridges of Königsberg

- The town of Königsberg, Russia was set on both sides (A and C) of the Pregel river, and included two large islands (Kneiphof and Lomse, denoted as B and D respectively) which were connected to each other by 7 bridges. The residents of Königsberg wondered if it was possible to take a walking tour of the town that crossed each of the seven bridges exactly once. Is it possible to start at some node and take a walk that uses each edge exactly once, and ends at the starting node?
- In 1736, Euler proved that the problem has no solution.





Leonhard Euler

1707-1783


Seven Bridges of Königsberg

- To prove it, Euler reformulated the problem in graph terms:
 - Each land mass ~ a vertex
 - Each bridge ~ an edge



Is there a way to go through all 7 bridges, each exactly once, and then return to the starting position?

Whether or not there exists a cycle on a graph G that traverses through every edge of G exactly once.




Euler graph

- Definition
- Recognize Euler graph

Euler graph


- **Euler path** (*Eulerian trail*, *Euler walk*) in graph is a path that traverses through every edge exactly once.
- **Euler cycle** (*Eulerian circuit*, *Euler tour*) in graph is a Euler path begins and ends at the same vertex (is a cycle that traverses through every edge exactly once).
- Graph consisting of Euler cycle is called as **Euler graph**.
- Graph consisting of Euler path is called as **Half Euler graph**.
- Apparently, all Euler graphs are also half-Euler graphs.

Example



Half Euler graph

Euler path: a, c, d, b, e, d, a, b



Euler graph

Euler cycle: a, c, d, e, b, d, f, b, a

Euler path in graph is a path that traverses through every edge exactly once.

Euler cycle in graph is a cycle that traverses through every edge exactly once.

Half Euler graph: graph consists of Euler path

Euler graph: graph consists of Euler cycle

Euler's 1st theorem

- If a graph has any vertices of odd degree, then it can not have any Euler cycle.
- If a graph is connected and every vertex has an even degree, then it has at least one Euler cycle.

Proof:

- If a node has an odd degree, and the cycle starts at this node, then it must end elsewhere. This is because after we leave the node the first time the node has even degree, and every time we return to the node we must leave it. (On the paired arc.)
- If a node has an odd degree, and the cycle begins else where, then it must end at the node. This is a contradiction, since a cycle must end where it began.

Euler's 1st theorem

If a graph has all even degree nodes, then an Euler Circuit exists.

Algorithm:

- Step One: Randomly move from node to node, until stuck. Since all nodes had even degree, the circuit must have stopped at its starting point. (It is a circuit.)
- Step Two: If any of the arcs have not been included in our circuit, find an arc that touches our partial circuit, and add in a new circuit.
 - Each time we add a new circuit, we have included more nodes.
 - Since there are only a finite number of nodes, eventually the whole graph is included.


Euler's 2nd theorem

- If a graph has more than two vertices of odd degree, then it cannot have an Euler path.
- If a graph is connected and has exactly two vertices of odd degree, then it has at least one Euler path. Any such path must start at one of the odd degree vertices and must end at the other odd degree vertex.

Euler's theorems

If a graph is connected and if the number of odd degree vertices

- = 0, then Euler cycle (Theorem 1)
- = 2, then Euler path (Theorem 2)
 - This Euler path must start at one of the odd degree vertices and must end at the other odd degree vertex.



Is there a way to go through all 7 bridges, each exactly once, and then return to the starting position?




Whether or not there exists a cycle on a graph G that traverses through every edge of G exactly once.

Answer: *There exists vertex of odd degree → don't have Euler cycle*


Whether or not there exists Euler path in G ?

Answer: *There are 3 vertices of degree 3, one vertex of degree 5 → don't have Euler path*

Example



Half Euler graph



Euler graph

Hamilton graph


- Definition
- Recognize Hamilton graph

Hamilton graph

- **Hamilton path** in graph is a path that traverses every vertex exactly once.
- **Hamilton cycle** in graph is a cycle that traverses every vertex exactly once.
- Graph consisting of Hamilton cycle is called as **Hamilton graph**.
- Graph consisting of Hamilton path is called as **Half Hamilton graph**.
- Apparently, all Hamilton graphs are also half-Hamilton graphs.

Example: Hamilton graph

- Is graph consisting of Hamilton cycle: traverse every vertex exactly once.




Applications of Hamilton Cycles

- The famous **traveling salesman problem** or **TSP**:


The **travelling salesman problem (TSP)** asks the following question: "Given a list of cities and the distances between each pair of cities, what is the shortest possible route that visits each city exactly once and returns to the origin city?"

An equivalent formulation in terms of graph theory is: Find the Hamiltonian cycle with the least weight in a weighted graph.



Hamilton graph

- Graph has 2 vertices of degree 1 $\Rightarrow$ not Hamilton graph




- The above graph is Half Hamilton graph

Half Hamilton graph

- Vertices of degree 1 must be either the starting or ending position of the Hamilton path.

Graph has 3 vertices of degree 1
 $\Rightarrow$ Not Halft Hamilton graph



Theorem about existence of Hamilton path

- **Theorem Dirac:** If G is simple connected graph with $n \geq 3$ vertices, and $\forall v \deg(v) \geq n/2$, then G has Hamilton cycle.
- **Theorem Ore:** If G is simple connected graph with $n \geq 3$ vertices, and $\deg(u) + \deg(v) \geq n$ for all vertices pair u, v not adjacent, then G has Hamilton cycle.



Paul Adrien Maurice Dirac
1902 - 1984
(USA)





Oystein Ore
1899 - 1968
(Norway)

Chapter 1. Fundamental concepts

1. Graph in practice
2. Graph types
3. Degree of vertex
4. Subgraph
5. Isomorphism of Graphs
6. Path and cycle
7. Connectedness
8. Special graphs
- 9. Graph Coloring problem**

What is Coloring?

- Graph Coloring Problem is an assignment of colors to all vertices of a given graph such that two adjacent vertices get different colors.
- **Objective:** use **minimum** number of colors.



3-colorable

A graphic on the left side of the slide. It features a dark blue background with a large, stylized circular pattern of red dots. The dots are arranged in concentric, slightly irregular rings, creating a textured, halftone-like effect. In the center of this pattern, the word "HUST" is written in a bold, white, sans-serif font.

HUST

THANK YOU !