

Non-equilibrium transport in quantum chains under quasiperiodic potentials

Sourav Suresh, Dr. Devendra Singh Bhakuni, Prof. Auditya Sharma and Prof. Sanjeev Kumar
Department of Physical Sciences, Indian Institute of Science Education and Research, Mohali

Motivation

Non-trivial transport properties, localisation transitions and fractal properties appearing in one-dimensional quasiperiodic potentials have kept physicists (and mathematicians as well) interested for quite some time. With novel experimental tools and present-day technology allowing us to have extreme control over quantum systems, this interest has resurfaced. In this thesis, we numerically study an interpolating model between the two most paradigmatic one-dimensional quasiperiodic models, the Aubry-André-Harper (AAH) model and the Fibonacci model and study the non-equilibrium transport in an open and closed system setup.

Aubry-André-Harper Model





The Hamiltonian is given by the tight-binding Hamiltonian with on site potential,

$$\mathcal{H} = -J \sum_{i=1}^{L-1} (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) + \sum_{i=1}^L V_i c_i^\dagger c_i$$

$$V_i = 2\lambda \cos(2\pi b i + \theta), \quad i = 1, \dots, L$$

- λ is the potential strength, b is an irrational constant which we set to the golden ratio ϕ and θ is an arbitrary global phase.
- When b is irrational the potential never repeats itself on the lattice.
- It undergoes a localisation transition at $\lambda = 1.0$ (which is a multi-fractal state).
- The Inverse Participation Ratio (IPR) is one of the measures used to quantify localisation of states

$$\text{IPR}(\psi) = \sum_{n=1}^L |\psi_n|^4$$



Unitary Evolution

A particle (delta wavefunction) was initialised at the central site of the chain and was let to evolve unitarily. The animations of the unitary evolution of a gaussian wavefunction and a delta wavefunction are given below.







Animation: Gaussian



Animation: Delta


We use the Root Mean Squared Displacement (RMSD) from the central site to measure the spread.

$$\sqrt{(\Delta x)^2} = \sqrt{\sum_{n=1}^L (n - n_0)^2 |\psi_n|^2} \quad \sqrt{(\Delta x)^2} \sim t^\alpha$$



Based on the scaling, we get different transport regimes.


Transport regime	Transport coefficient
Localized	$\alpha = 0$
Ballistic	$\alpha = 1$
Diffusive	$\alpha = \frac{1}{2}$
Subdiffusive	$\alpha < \frac{1}{2}$



The interpolation is done in this region where the scaling is linear.

The transport coefficient was plotted against β for various λ

Boundary-Driven Chain



A current (J) flows through the boundary-driven chain as the spin baths give a bias.

$$J \sim \frac{1}{L^\nu}$$

Transport regime	Transport coefficient
Ballistic	$\nu = 0$
Superdiffusive	$0 < \nu < 1$
Diffusive	$\nu = 1$
Subdiffusive	$\nu > 1$
Localized	$\nu = \infty$



In many cases, α can be compared with ν as follows,

$$\alpha = \frac{1}{1 + \nu}$$

J scales with L as below and depending on ν we get transport regimes.

Using the covariance formalism in the Lindblad master equation with appropriate dissipators (we use the Local Master Equations (LMEs) approach), we get the Lyapunov Equation $WC + CW^\dagger = F$

with $\Gamma = \text{diag}(\gamma_1, \gamma_2, \dots, \gamma_L)$ and $F = \text{diag}(\gamma_1 f_1, \gamma_2 f_2, \dots, \gamma_L f_L)$ and $W = iH + \frac{\Gamma}{2}$



The largest five systems sizes are used for the linear regression.

Interpolating Aubry-André Fibonacci Model

The Hamiltonian is given by the tight-binding Hamiltonian with on site potential,


$$V_j(\beta) = -\frac{\tanh[\beta \cos(2\pi b j + \phi) - \beta \cos(\pi b)]}{\tanh \beta}$$

$\beta \rightarrow 0$ is the AAH limit
 $\beta \rightarrow \infty$ is the Fibonacci limit


The QR code on the right shows the animation of the potential going from the AAH limit to the Fibonacci limit.




Animation: IAAF Potential



Source: Fig 1.b DOI:10.1038/s41567-020-0908-7



Source: Fig 2.b arXiv:2106.13841v2



Source: Fig 1.a arXiv:2106.13841v2

Acknowledgement

I am grateful to Dr. D S Bhakuni for always being there when needed. He made sure that I never felt lost. The support and motivation from my guides Prof. A Sharma and Prof. S Kumar, are priceless. I am grateful to all of my CMT groupmates for their support. I would like to thank Prof. Arti Garg from Saha Institute of Nuclear Physics for the discussion we had.

References

- [1] Z. Rieder, J. L. Lebowitz, and E. Lieb, "Properties of a harmonic crystal in a stationary nonequilibrium state," Journal of Mathematical Physics, vol. 8, no. 5
- [2] Transport in boundary-driven quantum chains with quasiperiodic potentials – Thesis by Artur Machado Lacerda
- [3] Many-body localization in the interpolating Aubry-André-Fibonacci model – Štrkalj et al.
- [4] Many-body localization in a quasiperiodic Fibonacci chain SciPost Phys. 6, 050 (2019) – Nicolas Macé, Nicolas Laflorencie and Fabien Alet
- [5] Emergence of criticality through a cascade of delocalization transitions in quasiperiodic chains