



# Protein Residue Contact Network Analysis to understand function of Single Residue Mutations



**Dhanasekhar V.K, Dr. Shashi Bhushan Pandit, Prof. Somdatta Sinha**  
Department of Biological Science, IISER Mohali

## Introduction

- Insights into mechanistic details of protein function are crucial for protein design, rational drug discovery and understanding evolution of proteins.
- In general, Single Residue Mutations (SRM) or single site mutations have contributed immensely in identifying important residues and deciphering their roles in protein function.
- Recently, single site variation data from human 1000 genome sequencing projects and Deep Mutational Scanning (DMS) methods have generated plethora of SRMs, however, their effect on protein structure/function remains elusive.
- Previously, we have used Protein Contact Networks (PCN) [1] approach to understand functional defects in SRMs. We observed that global or local network properties between Wild type (WT) and mutant (MT) proteins are similar. There were some protein WT-MT pairs, which showed changes in global/local network properties.
- In continuation of the efforts to understand affect of SRM on protein structure/function, we make following two main objectives:
  - ✓ To study a subset of WT-MT protein pairs that have large change in amino acid properties.
  - ✓ To associate SRM to functional phenotypic effect we developed a score Amino acid Physiochemical Score Matrix (APSM).


## Dataset and Methodology

**Dataset A:** In-house dataset consists of a 3600 wild-type mutant pairs having single residue mutations used for the network and physico-chemical analysis.

**Dataset B:** Non-redundant single domain created from pdb database (PISCES) [2] having resolution  $\leq 2.5$  Å.

### Constuction of Protein residue Contact Network

- Nodes:** C- $\alpha$  residues
- Edges:** An edge between two nodes are drawn, if any two C $\alpha$  atoms is within Euclidian distance  $\leq 7$  Å [3]
- Degree (k):** Number of of connections of a node.
- Shortest path ( $L_{ij}$ ):** Number of links that must be traversed by the shortest route, from one to node to another.
- Clustering coefficient (CC)**  $C_i = 2 * n / (k_i(k_i - 1))$ ,
- Betweenness centrality:** measures the extent to which a vertex lies on paths between other vertices.  
 $g(v) = \sum_{s \neq v, t \neq v} \frac{\sigma_{st}(v)}{\sigma_{st}}$  where,  $\sigma_{st}(v)$  total number shortest path that passes through v  
 $\sigma_{st}$  is total number shortest path between nodes s and t



### Classification of contacts based on the linear residue separation

To study distribution of contacts for each residue in single domain, we classified contacts based on linear separation [4] between contacting residues as:

**Shorter Range:** less than 6 residue apart

**Medium range:** 12-23 residue apart


**Short range:** 6 – 11 residue apart

**Long range :** > 23 residue apart

## Results and Discussion


### A. PCN Analysis of Selected WT-MT pairs

We selected 15 WT-MT pairs (having no significant structural changes) to construct their protein contact networks and analysed several network parameters to find residue contact perturbation due to mutation. Below is an representative example.



### B. Distribution of contact types per residue



To understand whether short to long range contacts dominated in PCN, we classified types of contacts and examined their distribution in single domain proteins



Most fraction of contacts per residues are medium/long range.


### C. Construction of Physicochemical score matrix (APSM)

To quantitatively measure physicochemical change in SRMs, we developed an Amino acid Physicochemical Score matrix (APSM) [5], which can provides an interpretable metric.



Occurrence of mutants in buried/exposed residues in WT-MT pairs and colored with APSM score


## Case study of APSM score and phenotypic function




Distribution of average APSM score among all mutations in dataset A and score depedence on residue burial status

### D. Coorespondence between APSM and phenotypic function

DMS studies on Gentamicin 3-N-acetyltransferase provided series of 2004 SRMs with their phenotypic effect at 37°C in *E. coli*. We score these single residue mutations using APSM score and obtain correspondence with phenotypic fitness provided in the experimental studies[6]



### a. Association between APSM score and protein function b. Residue level distribution of APSM and fitness



## Conclusions

- In general, protein contact networks have limited ability to decipher affect of SRMs on protein structure of function. However, it can provide insights if the residue in the first/second shell shows perturbation on mutation.
- APSM score by itself is not able to discern effects of SRM on phenotypic function. We need to explore its combination with other features to predict phenotypic function.

## References

- (PDF) PROTEIN STRUCTURE: INSIGHTS FROM GRAPH THEORY (researchgate.net), <http://14.139.227.205:8080/jspui/handle/123456789/1971>
- <http://dunbrack.fccc.edu/lab/piscs>
- <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4894841/#:~:text=Short%2DRange%2Dcontacts%2Dare%2D0thosc,hy%2D0at%20least%2024%2Dresidues>
- <https://www.imt.org/IMG/Teducation/Aide-memoire/LK/aminoacids/IMG/Telassics.html#:~:text=There%2Dare%205%20MG%2DVolume,%2C%2D0C%2C%2D0P%2C%2D0T>
- <https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1007419#sec016>