


Introduction and Motivation



- Collagen is the most abundant structural protein in animals
- Famous for its strained periodic pattern with periodicity of 67 nm
- How individual molecules assemble into this pattern is not fully understood
- Aim : To develop an in silico model that mimics the behavior of collagen and understand the factors affecting periodicity

Molecule Models




Fig 1: Percentage length vs potential energy for 5 layers

Artificial molecule model






Fig 3: Mass signal and Fourier spectra of given fibril

Acknowledgements

- Thanks to OeAD and the ISTernship program for giving me the opportunity and funding my expenses during my stay, Kevin Klein and Prof. Anđela for very fruitful discussions and members of the Šarić lab for the constant support
- Thanks to committee members Prof. Tripta Bhatia, Prof. Abhishek Chaudhari and Prof. Kavita Dorai for constructive feedback

Simulation Protocol



Hydrophobic Interactions

$$E_{LJ} = \begin{cases} 4\epsilon_{LJ} \left[\left(\frac{\sigma}{x} \right)^{12} - \left(\frac{\sigma}{x} \right)^6 \right] + E_{LJ}^{\text{shift}} & \text{for } x \leq x_{\text{cutoff}} \\ 0 & \text{for } x > x_{\text{cutoff}} \end{cases}$$

Electrostatic Interactions

$$E_{CD} = \begin{cases} C \frac{q_i q_j}{\epsilon_{CD} x} e^{-kx} + E_{CD}^{\text{shift}} & \text{for } x \leq x_{\text{cutoff}} \\ 0 & \text{for } x > x_{\text{cutoff}} \end{cases}$$

- Molecules interact via two types of interactions
- Implementation of simulations is done in LAMMPS using modified Velocity Verlet algorithm

Observations and Results




Fig 2: Mass signal and Fourier spectrum of collagen fibril at the end of the simulation

- The periodicity emerges for some particular charge ratios and is affected by how the charge is distributed

Inter-Chain Comparison




Fig 4: Percentage similarity scores using Needleman Wunch algorithm for type I and type II collagen

- Multiple collagen sequence types show same value of periodicity
- Each D regions is 234 amino acids long
- Comparative analysis of type I and type II collagen suggests that there is more interchain similarity than intrachain similarity

References



- Hodge, A.J. and Petrushka, J.A. (1963) In: Ramachandran, G.N., Ed., Aspects of Protein Structure, Academic Press, New York, 289-300
- Image credits : [1] Schwarcz et. al. (2017). The Ultrastructure of Bone and Its Relevance to Mechanical Properties ; [2] Salvatore Luca et. al. Mimicking the Hierarchical Organization of Natural Collagen ; [3] Chapman et. al. Electron microscopy reviews 1990