

# Holographic Entanglement Entropy at Spatial Infinity

Abir Ghosh <sup>1</sup> Chethan Krishnan <sup>2</sup>

<sup>1</sup>Indian Institute of Science Education and Research Mohali

<sup>2</sup>Indian Institute of Science Bengaluru

## Abstract

We study the calculation of Entanglement entropy in AdS/CFT using RT surfaces and introduce a natural analogue of a "spherical subregion" for the spatial infinity (spi) of asymptotically flat spacetime. The holographic data of such a spi-subregion can be encoded via the asymptotic behavior of extremal surfaces on a bulk spatial slice or a pair of points on  $\mathcal{I}^+$  and  $\mathcal{I}^-$ . We define a holographic entropy associated with a spi-subregion as the area of an extremal Ryu-Takayanagi surface in the bulk, which is the waist of an Asymptotic Causal Diamond (ACD).

## Holographic Principle

The holographic principle was inspired by black hole entropy equation,  $S = A_{BH}/4G_N$  where,  $A_{BH}$  is the surface area of the black hole. The holographic principle suggests that a D-dimensional theory of quantum gravity can be equally represented as a (D-1) dimensional QFT written on the boundary of the space.

## Statement of AdS/CFT Correspondence

Any conformal field theory on  $\mathbb{R} \times \mathbb{S}^{d-1}$  is equivalent to a theory of quantum gravity in asymptotically  $AdS_{d+1} \times M$  spacetime where  $M$  is some compact manifold.

## Geometry of Anti-de Sitter space

AdS is a type of space-time geometry with constant negative curvature. It can be spanned by different coordinate systems. The coordinate systems are,

### Global coordinates

In these coordinates the metric is given by eq (1)

$$ds^2 = -\left(1 + \frac{r^2}{L^2}\right) dt^2 + \frac{dr^2}{(1 + r^2)} + r^2 d\Omega_{d-1}^2 \quad (1)$$

### Poincare Coordinates

In these coordinates the metric is given by eq (2)

$$ds^2 = \frac{L^2}{z^2} (dz^2 + (dx^2 - dx_0^2)) \quad (2)$$




Figure 1. Poincare coordinates embedded in compactified global coordinate. In fig (a) the yellow surfaces are constant  $z$  surfaces and the yellow region in fig (b) is the space spanned by poincare coordinates

## Ryu Takayanagi Surfaces

The RT prescription provides us a way to calculate the entanglement of sub regions on the boundary CFT. The prescription is as follows, we start with a boundary sub region (say A) on a constant time slice and then draw a co-dimension 2 surface (say  $\gamma_A$ ) anchored to the boundary of the sub-region such that,

- $\gamma_A$  has the same boundary as A.
- $\gamma_A$  is homologous to A.
- $\gamma_A$  extremizes the area. If there are multiple extremal surfaces,  $\gamma_A$  is the one with the least area.

Then the area of this extremal surface is proportional to the entropy of the sub region.




Figure 2. (a)  $AdS_3$  space and  $CFT_2$  living on its boundary and (b) a geodesics  $\gamma_A$  as a holographic screen

For empty AdS the entropy equation we get is,

$$S_A = \frac{\text{Area of } \gamma_A}{4G} = \frac{1}{4G} \log \left( \frac{L}{\pi a} \sinh \left( \frac{\pi l}{L} \right) \right) \quad (3)$$

where,  $L$  is the AdS-length scale,  $a$  is the UV-parameter and  $l$  is the size of the boundary sub-region.

## Bulk with black holes




Figure 3. (a) Minimal surfaces  $\gamma_A$  for various sizes of A. (b)  $\gamma_A$  and  $\gamma_B$  wrap the different parts of the horizon.

## Flat Space Holography

Flat space also has a compactified representation with the metric,

$$ds^2 = \Omega^2 (-dt'^2 + dr'^2 + r'^2 d\phi^2) \quad (4)$$

with the transformations,

$$t + r = \tan \left( \frac{t' + r'}{2} \right) \quad t - r = \tan \left( \frac{t' - r'}{2} \right) \quad (5)$$




Figure 4. Penrose diagram

## Asymptotic Causal Diamond

**Def:** An Asymptotic Causal Diamond,  $\mathfrak{C}(p, q)$ , is defined as the intersection of the past light cone of a point  $p$  at future null infinity  $\mathcal{I}^+$  and the future light cone of a point  $q$  at past null infinity  $\mathcal{I}^-$ . That is

$$\mathfrak{C}(p, q) = \mathcal{I}^-(p) \cap \mathcal{I}^+(q), \text{ where } p \in \mathcal{I}^+, q \in \mathcal{I}^-.$$




Figure 5. Illustration of a symmetric ACD inside a penrose diagram in 2+1 dimensions.

The waist of the ACD is given by,

$$r \cos(\phi - \phi_0) = -\tan q_0 \quad (7)$$

In conformal coordinates the eq (7) becomes eq(8)

$$\tan(r'/2) \cos(\theta - \phi) = -\tan q_0 \quad (8)$$

The entropy is then given as the area of the waist,

$$A = \frac{\pi^{(d-1)/2}}{\Gamma(\frac{d+1}{2})} (\Lambda^2 - \tan^2 q_0)^{(d-1)/2} \quad (9)$$




Figure 6. Plot of the waist ( $t = 0$  slice) for different positions of the vertex

## Bulk with black hole

The action functional for a co-dimension 2 surface in a schwarzchild metric,

$$A = \frac{\pi^{(d-1)/2}}{\Gamma(\frac{d+1}{2})} \int r^{d-2} \sin^{d-2}(\theta) \sqrt{\frac{r'^2}{1 - (\frac{r_s}{r})^{d-2}} + r^2} d\theta \quad (10)$$




Figure 7. Extremal surfaces in (7a) 3+1 and (7b) 4+1 dimensions with a blackhole in the bulk

## Conclusion and discussion

- In flat space the analogue of boundary data (sub-region) seems to be the angle at which the waist approaches the boundary
- The behaviour of extremal surface is similar in flat and AdS case. It suggests that the area could be associated to "boundary sub-region" defined by the asymptotic angle.
- Regulating the divergences in flat space case isn't as simple as in AdS case and is still work in progress.

## Acknowledgement

I would like to take a moment to thank my family and friends for their unwavering support and encouragement throughout my thesis project. I would also like to express my gratitude to my advisor and the members of my thesis committee for their guidance, feedback, and expertise. Your input and insights have been instrumental in shaping this work.

## References

- [1] Daniel Harlow. TASI Lectures on the Emergence of Bulk Physics in AdS/CFT. PoS, TASI2017:002, 2018.
- [2] Chethan Krishnan. Bulk locality and asymptotic causal diamonds. SciPost Physics, 7(4), oct 2019.
- [3] Shinsei Ryu and Tadashi Takayanagi. Aspects of Holographic Entanglement Entropy. JHEP, 08:045, 2006.