



Supplementary Material for

Has land use pushed terrestrial biodiversity beyond the planetary boundary? A global assessment

Tim Newbold,* Lawrence N. Hudson, Andrew P. Arnell, Sara Contu, Adriana De Palma, Simon Ferrier, Samantha L. L. Hill, Andrew J. Hoskins, Igor Lysenko, Helen R. P. Phillips, Victoria J. Burton, Charlotte W. T. Chng, Susan Emerson, Di Gao, Gwilym Pask-Hale, Jon Hutton, Martin Jung, Katia Sanchez-Ortiz, Benno I. Simmons, Sarah Whitmee, Hanbin Zhang, Jörn P. W. Scharlemann, Andy Purvis

*Corresponding author. E-mail: t.newbold@ucl.ac.uk.

Published 15 July 2016, *Science* **353**, 288 (2016)
DOI: 10.1126/science.aaf2201

This PDF file includes:

Materials and Methods
Figs. S1 to S7
Tables S1 to S7
Full Reference List

Materials and Methods

The models were based on biodiversity data from the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) Project database (21). An extract of this database was taken on 28th April 2015. This extract consisted of 2.38 million records, from 413 published sources (31–437) or unpublished datasets with a published methodology, of the occurrence or abundance of 39,123 species from 18,659 sites in all of the world's 14 terrestrial biomes. The site-level data used to construct the models are publicly available from the Natural History Museum's Data Portal (doi: <http://dx.doi.org/10.5519/0073893>). The data are reasonably representative of major taxonomic groups (Fig. S1A) and of terrestrial biomes (Fig. S1B). For studies where sampling effort differed among the sites sampled, abundance values were corrected by dividing by sampling effort (i.e. assuming that abundance increases linearly with increasing effort). We derived two measures of biodiversity for each of the sites in our dataset: sampled total abundance of organisms and sampled species richness. Because it is not clear which of the many species-based measures of biodiversity most directly relates to the biodiversity-ecosystem functioning research, the main focus of this paper is on abundance-based measures and the corresponding planetary boundary (9).

We considered four human-pressure variables shown previously (3) to explain differences in local biodiversity among sites: land use (Table S7), land-use intensity (Table S7), human population density and distance to the nearest road. Human population density and distance to nearest road were log transformed and rescaled to a zero-to-one scale prior to analysis; proximity to the nearest road (as referred to in the main text) is simply the negative of log-transformed distance to the nearest road, such that higher values indicate higher pressure. We also considered two-way interactions between land use and each of the other variables. We chose these variables for the availability of fine-resolution mapped estimates, which enable spatial projections to be made from the models. Responses of biodiversity to these variables were modelled using generalized linear mixed-effects models. For sampled species richness we used a model with Poisson errors and a log link, while for (log-transformed) sampled total abundance we used a model with Gaussian errors and an identity link. A random effect of study identity was used to account for variation among studies in sampling methods and effort, differences in the taxonomic groups sampled, and coarse spatial differences in climate and other aspects of the environment. A random effect of spatial block nested within study, to take account of the spatial design of sampling. Spatial blocks were defined by the data entrants based on the maps and coordinates of sampled sites. A random slope of land use within study accounted for study-level variation in the relationship between land use and sampled biodiversity. Backward stepwise selection of fixed effects was used to select the minimum adequate model (438), with inclusion or exclusion of terms based on likelihood ratio tests (with a threshold $P < 0.05$). All models were developed using the lme4 Version 1.1-7 package (439) in R Version 3.2.2 (440). Spatial autocorrelation tests, performed as in (3), showed significant spatial autocorrelation in the model residuals for only slightly more of the modelled datasets than expected by chance: 6.1% in the case of species richness, and 5.9% in the case of total abundance.

To project mapped estimates of local biodiversity in the year 2005, we used fine-resolution maps of each of the four human pressure variables. The maps of land use were

generated by downscaling (23) the harmonized land-use dataset for 2005 (441). The harmonized land-use data describe the proportion of each 0.5° (approximately 50 km²) grid cell in each of five land uses (primary vegetation, secondary vegetation, cropland, pasture and urban). We used generalized additive models (GAM) with quasibinomial errors and a logistic link to relate coarse-scale estimates of each of the five land uses to nine putative explanatory variables at fine resolution (30 arc-seconds; approximately 1 km²): evapotranspiration (442), temperature (443), precipitation (443), topographic wetness (444), slope (444), soil carbon (445), accessibility to humans (446), human population density (24) and principal components of land cover (447). We then took the fine-grained fitted values from the GAMs and rescaled them multiplicatively until the aggregated mean for each 0.5° grid cell matched the estimates from the harmonized land-use data. The rescaled fitted values were then subjected to a constrained optimization algorithm, taking into account error estimates from the GAMs, to generate land-use estimates for all five land uses that summed to 1 within each grid cell. We entered the final estimates back into the GAMs as response variables, and the whole procedure was iterated until the mean inter-iteration difference of predicted values was ≤ 0.001 . Grid cells under ice or water (448, 449) were excluded from the analysis, and were masked from the final land-use maps. For full details on downscaling methodology see (23). The land-use data are freely available: <http://doi.org/10.4225/08/56DCD9249B224>.

In a previous study (3), to estimate spatial patterns of land-use intensity, we used generalized linear models (with binomial errors and a logistic link), for each level of intensity within each land use, to relate the proportion of each 0.5° grid cell under this combination of land use and intensity to three explanatory variables: the proportion of the cell under the land use in question, human population density and United Nations sub-region. Information on land-use intensity was obtained from the Global Land Systems dataset (450); see (3) for the reclassification used. To run these generalized linear models for every 30-arc-second grid cell was computationally infeasible. Therefore, we applied the coarse-resolution models developed for the previous study (3) at the fine resolution used here, assuming that the relationships are the same at both scales. We obtained a gridded map of human population density at 30-arc-second resolution and a vector map of the world's roads from NASA's Socioeconomic Data and Applications Centre (24, 25). To calculate a gridded map of distance to nearest road, we used Python code written for the *arcpy* module of ArcMap Version 10.3 (451), first to project the vector map of roads onto an equal-area (Behrmann) projection, then to calculate the average distance to the nearest road within each 782-m grid cell using the 'Euclidean Distance' function, and finally to reproject the resulting map back to a WGS 1984 projection at 30-arc-second resolution. Maximum estimated values across the terrestrial surface of human population density and distance to nearest road in 2005 were 8.3% and 20% higher, respectively, than the maximum values observed in the modelled dataset. To ensure that extrapolating did not create unrealistic projections, we set all grid cells with values higher than the maximum observed to be equal to this maximum observed value (this affected 0.002% of grid cells for human population density and 5.6% of grid cells for distance to nearest road). We could not estimate the expected species richness with absolutely no influence of roads because it is impossible to collect a sample of biodiversity under such a situation in the present day.

To generate estimates of the intactness of ecological assemblages in terms of within-sample species richness and abundance, we multiplied the coefficients of the minimum adequate models described above by the proportion of each grid cell under each land-use and use-intensity combination, and by log-transformed and rescaled (using the same rescaling as in the models) human population density or distance to nearest road. We assumed that human population density and distance to nearest road were constant within grid cells. The resulting values were summed across all coefficients and the intercept added to give the model estimate of log-transformed species richness or total abundance within each grid cell. We calculated the exponential of these values to estimate actual species richness and total abundance. Finally, to calculate the relative intactness of assemblages relative to a baseline with no human impacts, we calculated expected species richness and total abundance for a grid cell composed entirely of primary vegetation with minimal human use, with zero human population density, and at a distance to roads equal to the maximum value observed in the modelling data (195 km). Estimating uncertainty analytically for mixed-effects models requires generating an n -by- n matrix, where n is the number of grid cells in the projection; this was computationally intractable. Instead we generated 20 random draws (a greater number would have required a long computer run-time) of values for all of the model coefficients, from a multivariate normal distribution accounting for the covariance among modelled coefficients. These random draws of parameters were used to generate 20 replicate projections, from which 95% confidence limits were calculated for each analysis. All of the calculations described in this paragraph were undertaken using Python code implemented within the `arcpy` module of ArcMap Version 10.3 (451), using the 'Raster Calculator' function; except for the multivariate random draw of coefficient values, which was performed in R Version 3.2.2 using the 'mvnrm' function in the MASS package Version 7.3-43.

Scholes & Biggs (11) explicitly exclude alien species from the calculation of biodiversity intactness. Because it is not generally known which species are native and which not, we use modelled average compositional similarity between sites in primary vegetation and sites under other land uses as a multiplier on our land-use coefficients (on a 0-1 scale, rescaled such that primary-primary comparisons have a value of 1). To generate these modelled estimates of compositional similarity, we calculated asymmetric pairwise assemblage similarities between all possible pairs of sites within each study in the data set, where one site in the pair was in primary vegetation. Primary vegetation may contain species that are not truly native to an area, especially in landscapes with a long history of human modification; and landscape-level effects of land-use change may have already removed some originally-present species even from sites in primary vegetation. Therefore, our estimates of compositional similarity are likely to be biased upwards. Asymmetric values were used to focus on the probability that a species sampled in non-primary vegetation was also found in primary vegetation. To remove the possibility for pseudo-replication, we selected as independent contrasts all site comparisons on the off-diagonal of a randomized site-by-site matrix (452). Site-by-site matrices were randomised 100 times to generate 100 datasets of independent comparisons. Compositional similarity was measured using an asymmetric version of the Jaccard Index (J) for the projections of species richness, and an asymmetric version of the abundance-based Jaccard Index (J_a) (453) for the projections of total abundance:

$$J = \frac{a}{a + c}$$

$$J_a = \frac{UV}{V}$$

where a is the number of species shared between the two sampled sites, c is the number of species only found in the site not in primary vegetation, U is the summed relative abundance in the primary-vegetation site of all species found in both sites, and V is the summed relative abundance in the non-primary site of all species found in both sites.

Assemblage compositional similarities in each of the 100 datasets were modelled as a function of the combination of land uses represented and the distance (geographic, climatic and elevational) between sites. Full details of how assemblage compositional similarity was modelled are given in (22). Average coefficients across the 100 models describing average compositional similarity between primary vegetation and all other land uses (including primary vegetation itself) were rescaled so that comparisons of primary vegetation to itself had a value of 1 (to avoid conflating natural spatial turnover with land-use impact). These rescaled coefficients were then multiplied by the modelled coefficients describing differences in species richness and total abundance among land uses, to estimate the number of species or individuals present in each land use that are also expected to be present in primary vegetation. The rescaled coefficients are publicly available from the Natural History Museum's Data Portal (doi: <http://dx.doi.org/10.5519/0073893>).

Although our way of calculating BII differs from that proposed by Scholes & Biggs (11), we also attempt to estimate the “average abundance of a large and diverse set of organisms in an area, relative to their reference populations” (11). If I_{ijk} is the population of species group i in ecosystem j under land use k , relative to a pre-industrial population in the same ecosystem type, then Scholes & Biggs (11) define the biodiversity intactness index (BII) to be:

$$BII = 100 \times (\sum_i \sum_j \sum_k R_{ij} A_{jk} I_{ijk}) / (\sum_i \sum_j \sum_k R_{ij} A_{jk})$$

where R_{ij} is the species richness of taxon i in ecosystem j and A_{jk} is the area of ecosystem j under land use k . Scholes & Biggs (11) used expert opinion when estimating average BII for seven southern African countries, in the absence of sufficient primary data. They considered birds, mammals, amphibians, reptiles and angiosperms but not arthropods, again because of a lack of information.

Our implementation of the BII differs in that we have used primary data on sampled local species abundance – for a wide range of animal (vertebrates and invertebrates), plant and fungal taxa – in place of expert opinion, and our statistical models incorporate other pressures as well as land use itself. Rather than weighting by areas of ecosystems and species-richness of taxa, we have collated and analysed a data set that is reasonably representative in terms of biomes (Fig. S1B) and taxa (Fig. S1A). Our data set is not yet adequate to support fitting models for each biome and taxon separately, which may lead to our estimates being biased for some biomes. Despite our very large number of records,

hierarchical mixed-effects models for individual biomes or taxa would require data from a larger number of published studies than is available for some taxa and biomes. As in (11), in the absence of pre-industrial data, we have used minimally-impacted sites as the reference condition.

We overlaid our estimates of the intactness of ecological assemblages with global maps describing the distribution of biomes (449), Conservation International's biodiversity hotspots (28), Conservation International's High Biodiversity Wilderness Areas (454) and human population density (24). All of these overlays were performed using Python code for ArcMap Version 10.3 (451), using the 'Zonal Statistics' functions after first projecting all maps into an equal-area (Behrmann) projection.

Fig. S1.




Fig. S1. Taxonomic (A) and biogeographic (B) representativeness of the records used to model biodiversity responses to land use. (A) Correlation, for major taxonomic groups (magenta – invertebrates; red– vertebrates; green – plants and fungi; grey – other), between the estimated number of described species (455) and the number of species represented in the dataset. (B) Correlation between the percentage of global primary productivity within a biome (449) and the percentage of sites in the dataset within that biome (A: Tundra; B: Boreal forests/taiga; C: Temperate conifer forests; D: Temperate broadleaf and mixed forests; E: Montane grasslands and shrublands; F: Temperate grasslands, savannas and shrublands; G: Mediterranean forests, woodland and scrub; H: Deserts and xeric shrublands; J: Tropical and subtropical grasslands, savannas and shrublands; K: Tropical and subtropical coniferous forests; L: Flooded grasslands and savannas; M: Tropical and subtropical dry broadleaf forests; N: Tropical and subtropical moist broadleaf forests; P: Mangroves).

Fig. S2




Fig. S2. Response of sampled total abundance to human pressures: (A) land use, and (B) the interaction between land use and human population density. Human population is shown on a rescaled axis (as fitted in the models). (A) shows total abundance as a percentage of that found in minimally used primary vegetation, with 95% confidence intervals; multiple points within each land-use type show, from left to right, increasing intensity of human use (two classes for secondary vegetation and urban; three classes for all other land uses). B shows absolute mean total abundance for a given combination of pressures, with shading indicating $\pm 0.5 \times \text{SEM}$, for clarity. Land uses in B are shown in the same colours as in A. Mixed-effects models are robust to unbalanced designs (456), such as the data spanning different ranges of human population density for each of the land uses. Dropping all urban sites almost no effect on the other model coefficients (Fig. S6). Full statistical results are given in Table S5.

Fig. S3




Fig. S3. Response of sampled species richness to human pressures: (A) land use, (B) the interaction between land use and human population density, and (C) the interaction between land use and distance to nearest road. Human population and distance to nearest road are shown on rescaled axes (as fitted in the models). (A) shows species richness as a percentage of that found in minimally used primary vegetation, with 95% confidence intervals; multiple points within each land-use type show, from left to right, increasing intensity of human use (two classes for secondary vegetation and urban; three classes for all other land uses). B and C show absolute mean species richness for a given

combination of pressures, with shading indicating $\pm 0.5 \times \text{SEM}$, for clarity. Land uses in B and C are shown in the same colours as in A. Mixed-effects models are robust to unbalanced designs (456), such as the data spanning different ranges of human population density for each of the land uses. Dropping all urban sites almost no effect on the other model coefficients (Fig. S7). Full statistical results are given in Table S6.

Fig. S4




Fig. S4. Biodiversity intactness of ecological assemblages in terms of the total abundance of originally occurring species, as a percentage of their total abundance in minimally disturbed primary vegetation (Biodiversity Intactness Index; BII). Blues areas are those within, and red areas those beyond proposed (9) safe limits for biodiversity, in terms of BII. A high-resolution raster of this map can be freely downloaded (doi: <http://dx.doi.org/10.5519/0009936>).

Fig. S5




Fig. S5. The proportion of the terrestrial surface exceeding the proposed (9) planetary boundary across the range of uncertainty in the boundary's position. Steffen et al. (9) suggested that the planetary boundary for BII could range anywhere between 30 and 90%, which has a large effect on the proportion of the land surface exceeding the boundary. The dashed grey line indicates the 58.1% of terrestrial area that falls below the precautionary BII threshold of 90%.

Fig. S6




Fig. S6. In models with no urban sites, the response of sampled total abundance to human pressures: (A) land use, and (B) the interaction between land use and human population density. The modelled coefficients are robust to the exclusion of urban sites, which cause an unbalanced design. All plotting conventions are as in Fig. S2.

Fig. S7




Fig. S7. In models with no urban sites, the response of sampled species richness to human pressures: (A) land use, (B) the interaction between land use and human population density, and (C) the interaction between land use and distance to nearest road. The modelled coefficients are robust to the exclusion of urban sites, which cause an unbalanced design. All plotting conventions are as in Fig. S3.

Table S1.

Table S1. Numbers of species represented in the dataset by major taxonomic group,
both for species represented in the complete dataset and species with only abundance
data.

| Taxon | N species (all data) | N species (abundance data) |
|------------------|-----------------------------|-----------------------------------|
| Amphibia | 415 | 365 |
| Annelida | 40 | 40 |
| Arachnida | 2288 | 2288 |
| Archaeognatha | 11 | 11 |
| Ascomycota | 762 | 613 |
| Aves | 3232 | 3033 |
| Basidiomycota | 514 | 399 |
| Blattodea | 33 | 33 |
| Bryophyta | 862 | 694 |
| Chilopoda | 52 | 52 |
| Coleoptera | 6164 | 5955 |
| Collembola | 161 | 155 |
| Crustacea | 57 | 52 |
| Dermaptera | 20 | 20 |
| Diplopoda | 89 | 89 |
| Diplura | 1 | 1 |
| Diptera | 1475 | 1475 |
| Embiopoda | 4 | 4 |
| Ephemeroptera | 4 | 4 |
| Ferns and allies | 392 | 332 |
| Fungoid protists | 1 | 1 |
| Glomeromycota | 31 | 31 |
| Gymnosperms | 70 | 57 |
| Hemiptera | 1214 | 1214 |
| Hymenoptera | 4639 | 4338 |
| Isoptera | 154 | 109 |
| Lepidoptera | 2911 | 2849 |
| Magnoliophyta | 11995 | 9003 |
| Mammalia | 547 | 500 |
| Mantodea | 5 | 5 |
| Mecoptera | 6 | 6 |
| Mollusca | 429 | 378 |
| Nematoda | 172 | 172 |
| Neuroptera | 36 | 36 |
| Odonata | 96 | 96 |
| Onychophora | 1 | 1 |
| Orthoptera | 155 | 154 |
| Paupoda | 6 | 6 |

| | | |
|-----------------|-----|-----|
| Phasmida | 2 | 2 |
| Phthiraptera | 3 | 3 |
| Platyhelminthes | 6 | 6 |
| Protura | 5 | 5 |
| Psocoptera | 28 | 28 |
| Reptilia | 397 | 335 |
| Siphonaptera | 4 | 4 |
| Symphyla | 5 | 5 |
| Thysanoptera | 50 | 50 |
| Thysanura | 1 | 1 |
| Trichoptera | 17 | 17 |
| Zoraptera | 1 | 1 |
| Other | 243 | 192 |

Table S2.

Table S2. Biodiversity intactness of the world's terrestrial biomes (449) in terms of species richness ('richness') and total organism abundance ('abundance'), colour coded according to the status of biodiversity with respect to boundaries proposed as safe limits for ecosystem function (5, 9): red = boundary crossed (> 20% loss of richness; > 10% loss of abundance); orange = boundary approached (>10% loss of richness; > 5% loss of abundance); green = not close to boundary. Values are given as overall net changes including species not found in primary vegetation ('all species') and intactness considering only originally present species ('original species'). Text in parentheses indicates 95% confidence limits.

| Biome | Intactness (abundance) | | Intactness (richness) | |
|--|------------------------|---------------------|-----------------------|--------------------|
| | All species | Original species | All species | Original species |
| Temperate Grasslands, Savannas and Shrublands | 73 (67.3 - 85) | 68 (62.8 - 78.3) | 67.6 (60.7 - 76.4) | 65.2 (61 - 76.9) |
| Mediterranean Forests, Woodlands and Scrub | 83.1 (76.7 - 90.1) | 78.3 (73.9 - 87) | 71.8 (65 - 82.7) | 69.8 (65.5 - 82.7) |
| Montane Grasslands and Shrublands | 82 (73.9 - 93.7) | 77.1 (71.4 - 89.1) | 72.4 (67.4 - 81.8) | 70.2 (66.3 - 81.9) |
| Tropical and Subtropical Grasslands, Savannas and Shrublands | 85.5 (76.5 - 97.9) | 80.5 (73.9 - 91.9) | 74.1 (68.3 - 85.3) | 72 (68 - 84.8) |
| Flooded Grasslands and Savannas | 85.7 (79.1 - 96.2) | 81.1 (77 - 90.8) | 74.2 (68.4 - 85) | 72.2 (68 - 84.8) |
| Temperate Broadleaf and Mixed Forests | 90 (80.2 - 99.5) | 85.9 (79.2 - 96.1) | 74.8 (67.5 - 86.2) | 73.1 (66.6 - 86.3) |
| Tropical and Subtropical Dry Broadleaf Forests | 90.1 (81.1 - 99.9) | 86.3 (79.9 - 96.3) | 75.9 (69.4 - 87.6) | 74.4 (68.4 - 87.5) |
| Deserts and Xeric Shrublands | 82 (75.6 - 93) | 78.3 (73.5 - 86.7) | 76.2 (71 - 85.1) | 74.5 (71.6 - 85.5) |
| Tropical and Subtropical Coniferous Forests | 95 (85.2 - 105.1) | 90.9 (84.4 - 102.9) | 77.2 (70.5 - 90) | 75.6 (68.1 - 89.2) |
| Mangroves | 95.6 (84.8 - 108) | 92.2 (84.4 - 104.9) | 78.9 (72.5 - 89.9) | 77.5 (69.8 - 89.6) |
| Temperate Conifer Forests | 89.2 (84.3 - 94.7) | 86.2 (83 - 91.9) | 79.2 (73.8 - 89.1) | 78 (74.5 - 89) |
| Tropical and Subtropical Moist Broadleaf Forests | 95.9 (89 - 104) | 93.2 (88.7 - 101.4) | 82.8 (77.4 - 92.8) | 81.7 (75.7 - 92.4) |
| Boreal Forests/Taiga | 96.3 (92.7 - 99) | 95.5 (92.3 - 98.1) | 88.8 (84.1 - 96.9) | 88.5 (85.9 - 96.8) |
| Tundra | 99.7 (98.5 - 100.7) | 99.5 (98.4 - 100.4) | 94.8 (91.8 - 100.1) | 94.8 (93.2 - 99.8) |

Table S3.

Table S3. Biodiversity intactness of the world's terrestrial Biodiversity Hotspots (28) in terms of species richness ('richness') and total organism abundance ('abundance'). Colours and labels are as in Table 1. Text in parentheses indicates 95% confidence limits.

| Hotspot | Intactness (abundance) | | Intactness (richness) | |
|---|------------------------|---------------------|-----------------------|--------------------|
| | All species | Original species | All species | Original species |
| Cape Floristic Region | 72.5 (62.9 - 89.3) | 66.5 (59 - 80.4) | 67.2 (60.2 - 78.7) | 64.4 (60 - 78) |
| Succulent Karoo | 64.2 (50.3 - 87) | 59.4 (52.8 - 79.6) | 67.8 (60.1 - 78.1) | 65.2 (58.2 - 82.3) |
| New Zealand | 72.5 (63.7 - 86.2) | 68.1 (62.7 - 79.8) | 70.2 (63.5 - 79.7) | 68 (63.4 - 80.9) |
| Southwest Australia | 73.5 (64.4 - 84.6) | 69.8 (63.5 - 79.5) | 71.4 (64.1 - 80) | 69.6 (64.8 - 81.5) |
| Maputaland-Pondoland-Albany | 82.6 (76.3 - 93) | 77.2 (73.1 - 88.8) | 71.7 (65.4 - 84.3) | 69.3 (65.6 - 83.5) |
| Mediterranean Basin | 87.4 (77.6 - 98.6) | 82.1 (74.5 - 95.2) | 71.9 (64.4 - 83.9) | 69.8 (62.8 - 83.5) |
| Mountains of Central Asia | 86.2 (76.2 - 99.5) | 80.7 (73.7 - 94.2) | 72.4 (65.7 - 84) | 70.1 (63.9 - 83.2) |
| Cerrado | 80.2 (72.2 - 91.7) | 75.7 (69.7 - 85.7) | 72.9 (67.6 - 82.5) | 70.9 (66.8 - 82.4) |
| Caucasus | 90.3 (78.9 - 102.9) | 85.3 (76.7 - 99) | 73.1 (65.1 - 86.2) | 71.1 (63.1 - 84.9) |
| Madagascar and the Indian Ocean Islands | 89.6 (77.6 - 106.2) | 83.6 (74.7 - 99) | 73.1 (66.2 - 87.5) | 70.7 (64.2 - 85.6) |
| Irano-Anatolian | 92.3 (81.2 - 107) | 86.7 (78.4 - 102.4) | 73.6 (65.9 - 86.9) | 71.4 (62.9 - 85.6) |
| Atlantic Forest | 89.8 (79.8 - 102) | 84.8 (77.8 - 97.3) | 73.8 (66.6 - 86.2) | 71.7 (64.3 - 85.2) |
| Caribbean Islands | 92.9 (80.1 - 108.1) | 88.1 (77.5 - 104.3) | 74.3 (66.8 - 88.1) | 72.5 (64.3 - 86.5) |
| California Floristic Province | 83.4 (78.6 - 87.6) | 80.1 (75 - 86.5) | 74.5 (68.6 - 83.9) | 73.1 (69.9 - 84.1) |
| Mountains of Southwest China | 90.4 (80.2 - 103.6) | 85.5 (78.6 - 98.4) | 74.6 (67.8 - 86.7) | 72.5 (65.1 - 85.9) |
| Horn of Africa | 88.3 (76.7 - 103.4) | 83.1 (75.1 - 96.1) | 74.6 (68.3 - 87.7) | 72.4 (67.1 - 86) |
| Himalaya | 90.4 (80.4 - 101.8) | 86.2 (78.8 - 99) | 74.7 (68.2 - 86.2) | 72.9 (66 - 86) |
| Coastal Forests of Eastern Africa | 95.8 (85.2 - 111.9) | 90.2 (81.7 - 105.1) | 76 (68.8 - 89.9) | 73.9 (65.8 - 88.8) |
| Eastern Afromontane | 99.5 (86 - 113.4) | 94.1 (84.9 - 112.8) | 76.6 (69.5 - 90.6) | 74.7 (65.1 - 90.3) |
| Philippines | 94.9 (78 - 114.4) | 91.6 (77.7 - 106.5) | 76.7 (68.7 - 89.1) | 75.5 (66.1 - 88.8) |
| Madrean Pine-Oak | 91.8 (83 - 102.8) | 87.6 (82.4 - 97.4) | 76.8 (70.4 - 89) | 75.1 (69 - 88.1) |

| | | | | |
|---|----------------------|----------------------|---------------------|---------------------|
| Woodlands | | | | |
| Western Ghats and Sri Lanka | 99.1 (79.9 - 122.9) | 95.7 (80.4 - 113.9) | 77.1 (69 - 90.8) | 75.9 (66.4 - 90.5) |
| Guinean Forests of West Africa | 100.9 (87.2 - 114.7) | 95.6 (86.9 - 113.8) | 77.1 (69.5 - 91.8) | 75.2 (66 - 91.6) |
| Mesoamerica | 96.4 (86.3 - 108) | 92.1 (85.4 - 104.1) | 77.9 (71 - 91.1) | 76.2 (68.4 - 90.3) |
| Tumbes-Choco-Magdalena | 93.5 (84.5 - 105.9) | 89.3 (83 - 100.1) | 78.1 (71.9 - 90) | 76.4 (69.2 - 88.9) |
| Polynesia-Micronesia | 91.8 (85 - 99.2) | 88.8 (85.2 - 96.5) | 78.2 (72.8 - 90) | 77 (72.1 - 89.5) |
| Tropical Andes | 91.6 (84.1 - 102.2) | 87.9 (83.2 - 96.4) | 78.7 (72.8 - 90.9) | 77.2 (72 - 90.1) |
| Japan | 100.9 (85.2 - 114.5) | 97.7 (85.9 - 114.7) | 79.1 (71 - 93.5) | 78 (70.3 - 93.5) |
| Chilean Winter Rainfall and Valdivian Forests | 91.2 (84.7 - 100.1) | 88.1 (84.4 - 95.6) | 79.9 (74.5 - 91.5) | 78.6 (74.7 - 90.9) |
| Indo-Burma | 98.3 (83.6 - 112.5) | 95.8 (85 - 107.9) | 80.6 (72.7 - 93.7) | 79.7 (71 - 93.4) |
| Sundaland | 96.5 (86.5 - 106.7) | 94.4 (87.5 - 102.5) | 82.1 (75.4 - 92.9) | 81.3 (74.2 - 92.8) |
| New Caledonia | 97.4 (90.9 - 102.8) | 95.5 (91.2 - 102.2) | 83.1 (75.5 - 94.7) | 82.2 (79.2 - 95.3) |
| Wallacea | 100.5 (88.1 - 111.4) | 98.7 (90.3 - 108.6) | 83.5 (76 - 96.5) | 82.8 (74.8 - 96.3) |
| East Melanesian Islands | 104 (91.3 - 114.1) | 103.4 (94.5 - 112.1) | 90.5 (83.9 - 101.5) | 90.2 (82.2 - 102.5) |

Table S4.

Table S4. Biodiversity intactness of the world's High Biodiversity Wilderness Areas (454) in terms of species richness ('richness') and total organism abundance ('abundance'). Colours and labels are as in Table 1. Text in parentheses indicates 95% confidence limits.

| High Biodiversity Wilderness Area | Intactness (abundance) | | Intactness (richness) | |
|---|------------------------|---------------------|-----------------------|--------------------|
| | All species | Original species | All species | Original species |
| North American Deserts | 76.6 (67.1 - 90.9) | 72.2 (66.1 - 85.6) | 72.5 (66.8 - 82.2) | 70.4 (66 - 83.7) |
| Miombo-Mopane Woodlands and Savannas | 90.9 (79.6 - 105.9) | 86.6 (77.8 - 97.9) | 77.7 (71.8 - 89.5) | 76 (70.2 - 89) |
| Congo Forests | 96.5 (86.9 - 107.8) | 93.9 (85.3 - 102.3) | 83.3 (77.5 - 95.5) | 82.3 (76.6 - 95.8) |
| New Guinea | 99 (91.7 - 105.5) | 97.8 (93.1 - 102.9) | 89.3 (85 - 97) | 88.8 (83.5 - 97.5) |
| Amazonia | 94.9 (90.7 - 98.8) | 93.6 (90.5 - 97.1) | 89.4 (86.3 - 94.8) | 88.8 (86.7 - 94.8) |

Table S5.

Table S5. Results of backward stepwise model selection (457) on model of sampled total abundance. Terms considered were land use (LandUse), land-use intensity (UseIntensity), human population density (HPD), distance to nearest road (DR), and interactions between land use and the other variables. Interaction terms were compared first, and then removed to test main effects. HPD and DR were fitted as quadratic polynomials. We report here chi-square values (χ^2), degrees of freedom (DF) and P-values (P). Variables within significant interactions were retained in the final model, even if the main effect of that variable was not significant.

| Term | χ^2 | DF | P |
|----------------------|----------------------------|-----------|----------|
| LandUse | 9.42 | 5, 33 | 0.093 |
| UseIntensity | 33.6 | 2, 28 | < 0.001 |
| HPD | 13.7 | 1, 28 | < 0.001 |
| DR | 0.382 | 1, 35 | 0.54 |
| LandUse:UseIntensity | 62.2 | 13, 53 | < 0.001 |
| LandUse:HPD | 21.7 | 10, 53 | 0.017 |
| LandUse:DR | 13.8 | 10, 63 | 0.18 |

Table S6.

Table S6. Results of backward stepwise model selection (457) on model of sampled species richness. Terms considered were land use (LandUse), land-use intensity (UseIntensity), human population density (HPD), distance to nearest road (DR), and interactions between land use and the other variables. Interaction terms were compared first, and then removed to test main effects. HPD and DR were fitted as quadratic polynomials. We report here chi-square values (χ^2), degrees of freedom (DF) and P-values (P). Variables within significant interactions were retained in the final model, even if the main effect of that variable was not significant.

| Term | χ^2 | DF | P |
|----------------------|----------------------------|-----------|----------|
| LandUse | 429 | 5, 13 | < 0.001 |
| UseIntensity | 19.0 | 2, 13 | < 0.001 |
| HPD | 17.6 | 1, 13 | < 0.001 |
| DR | 0.39 | 1, 15 | 0.53 |
| LandUse:UseIntensity | 408 | 13, 43 | < 0.001 |
| LandUse:HPD | 41.2 | 10, 43 | < 0.001 |
| LandUse:DR | 57.2 | 10, 43 | < 0.001 |

Table S7.

Table S7. Land-use and land-use-intensity classification definitions.

| Level 1 Land Use | Predominant Land Use | Minimal use | Light use | Intense use |
|--|--|---|--|---|
| No evidence of prior destruction of the vegetation | Primary forest | Any disturbances identified are very minor (e.g., a trail or path) or very limited in the scope of their effect (e.g., hunting of a particular species of limited ecological importance). | One or more disturbances of moderate intensity (e.g., selective logging) or breadth of impact (e.g., bushmeat extraction), which are not severe enough to markedly change the nature of the ecosystem. Primary sites in suburban settings are at least Light use. | One or more disturbances that is severe enough to markedly change the nature of the ecosystem; this includes clear-felling of part of the site too recently for much recovery to have occurred. Primary sites in fully urban settings should be classed as Intense use. |
| | Primary Non-Forest | As above | As above | As above |
| Recovering after destruction of the vegetation | Mature Secondary Vegetation | As for Primary Vegetation-Minimal use | As for Primary Vegetation-Light use | As for Primary Vegetation-Intense use |
| | Intermediate Secondary Vegetation | As for Primary Vegetation-Minimal use | As for Primary Vegetation-Light use | As for Primary Vegetation-Intense use |
| | Young Secondary Vegetation | As for Primary Vegetation-Minimal use | As for Primary Vegetation-Light use | As for Primary Vegetation-Intense use |
| | Secondary Vegetation (indeterminate age) | As for Primary Vegetation-Minimal use | As for Primary Vegetation-Light use | As for Primary Vegetation-Intense use |
| Human use (agricultural) | Plantation forest | Extensively managed or mixed timber, fruit/coffee, oil-palm or rubber plantations in which native understorey and/or other native tree species are tolerated, which are not treated with pesticide or fertiliser, and which have not been recently (< 20 years) clear-felled. | Monoculture fruit/coffee/rubber plantations with limited pesticide input, or mixed species plantations with significant inputs. Monoculture timber plantations of mixed age with no recent (< 20 years) clear-felling. Monoculture oil-palm plantations with no recent (< 20 years) clear-felling. | Monoculture fruit/coffee/rubber plantations with significant pesticide input. Monoculture timber plantations with similarly aged trees or timber/oil-palm plantations with extensive recent (< 20 years) clear-felling. |

| | | | | |
|-----------------------------|----------|---|--|---|
| Human use (agricultural) | Cropland | Low-intensity farms, typically with small fields, mixed crops, crop rotation, little or no inorganic fertiliser use, little or no pesticide use, little or no ploughing, little or no irrigation, little or no mechanisation. | Medium intensity farming, typically showing some but not many of the following: large fields, annual ploughing, inorganic fertiliser application, pesticide application, irrigation, no crop rotation, mechanisation, monoculture crop. Organic farms in developed countries often fall within this category, as may high-intensity farming in developing countries. | High-intensity monoculture farming, typically showing many of the following features: large fields, annual ploughing, inorganic fertiliser application, pesticide application, irrigation, mechanisation, no crop rotation. |
| | Pasture | Pasture with minimal input of fertiliser and pesticide, and with low stock density (<i>not</i> high enough to cause significant disturbance or to stop regeneration of vegetation). | Pasture either with significant input of fertiliser or pesticide, or with high stock density (high enough to cause significant disturbance or to stop regeneration of vegetation). | Pasture with significant input of fertiliser or pesticide, <i>and</i> with high stock density (high enough to cause significant disturbance or to stop regeneration of vegetation). |
| Human use (urban) | Urban | Extensive managed green spaces; villages. | Suburban (e.g. gardens), or small managed or unmanaged green spaces in cities. | Fully urban with no significant green spaces. |

References and Notes

1. J.-C. Vié, C. Hilton-Taylor, S. N. Stuart, *Wildlife in a Changing World: An Analysis of the 2008 Red List of Threatened Species* (IUCN, 2009).
2. D. P. Tittensor, M. Walpole, S. L. Hill, D. G. Boyce, G. L. Britten, N. D. Burgess, S. H. Butchart, P. W. Leadley, E. C. Regan, R. Alkemade, R. Baumung, C. Bellard, L. Bouwman, N. J. Bowles-Newark, A. M. Chenery, W. W. Cheung, V. Christensen, H. D. Cooper, A. R. Crowther, M. J. Dixon, A. Galli, V. Gaveau, R. D. Gregory, N. L. Gutierrez, T. L. Hirsch, R. Höft, S. R. Januchowski-Hartley, M. Karmann, C. B. Krug, F. J. Leverington, J. Loh, R. K. Lojenga, K. Malsch, A. Marques, D. H. Morgan, P. J. Mumby, T. Newbold, K. Noonan-Mooney, S. N. Pagad, B. C. Parks, H. M. Pereira, T. Robertson, C. Rondinini, L. Santini, J. P. Scharlemann, S. Schindler, U. R. Sumaila, L. S. Teh, J. van Kolck, P. Visconti, Y. Ye, A mid-term analysis of progress toward international biodiversity targets. *Science* **346**, 241–244 (2014). [Medline](#) [doi:10.1126/science.1257484](https://doi.org/10.1126/science.1257484)
3. T. Newbold, L. N. Hudson, S. L. Hill, S. Contu, I. Lysenko, R. A. Senior, L. Börger, D. J. Bennett, A. Choimes, B. Collen, J. Day, A. De Palma, S. Díaz, S. Echeverria-Londoño, M. J. Edgar, A. Feldman, M. Garon, M. L. Harrison, T. Alhusseini, D. J. Ingram, Y. Itescu, J. Kattge, V. Kemp, L. Kirkpatrick, M. Kleyer, D. L. Correia, C. D. Martin, S. Meiri, M. Novosolov, Y. Pan, H. R. Phillips, D. W. Purves, A. Robinson, J. Simpson, S. L. Tuck, E. Weiher, H. J. White, R. M. Ewers, G. M. Mace, J. P. Scharlemann, A. Purvis, Global effects of land use on local terrestrial biodiversity. *Nature* **520**, 45–50 (2015). [Medline](#) [doi:10.1038/nature14324](https://doi.org/10.1038/nature14324)
4. B. J. Cardinale, J. E. Duffy, A. Gonzalez, D. U. Hooper, C. Perrings, P. Venail, A. Narwani, G. M. Mace, D. Tilman, D. A. Wardle, A. P. Kinzig, G. C. Daily, M. Loreau, J. B. Grace, A. Larigauderie, D. S. Srivastava, S. Naeem, Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67 (2012). [Medline](#) [doi:10.1038/nature11148](https://doi.org/10.1038/nature11148)
5. D. U. Hooper, E. C. Adair, B. J. Cardinale, J. E. Byrnes, B. A. Hungate, K. L. Matulich, A. Gonzalez, J. E. Duffy, L. Gamfeldt, M. I. O'Connor, A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* **486**, 105–108 (2012). [Medline](#)
6. Y. Hautier, D. Tilman, F. Isbell, E. W. Seabloom, E. T. Borer, P. B. Reich, Anthropogenic environmental changes affect ecosystem stability via biodiversity. *Science* **348**, 336–340 (2015). [Medline](#) [doi:10.1126/science.aaa1788](https://doi.org/10.1126/science.aaa1788)
7. T. H. Oliver, M. S. Heard, N. J. Isaac, D. B. Roy, D. Procter, F. Eigenbrod, R. Freckleton, A. Hector, C. D. Orme, O. L. Petchey, V. Proença, D. Raffaelli, K. B. Suttle, G. M. Mace, B. Martín-López, B. A. Woodcock, J. M. Bullock, Biodiversity and resilience of ecosystem functions. *Trends Ecol. Evol.* **30**, 673–684 (2015). [Medline](#) [doi:10.1016/j.tree.2015.08.009](https://doi.org/10.1016/j.tree.2015.08.009)
8. G. M. Mace, B. Reyers, R. Alkemade, R. Biggs, F. S. Chapin III, S. E. Cornell, S. Díaz, S. Jennings, P. Leadley, P. J. Mumby, A. Purvis, R. J. Scholes, A. W. R. Seddon, M. Solan, W. Steffen, G. Woodward, Approaches to defining a planetary boundary for biodiversity. *Glob. Environ. Change* **28**, 289–297 (2014). [doi:10.1016/j.gloenvcha.2014.07.009](https://doi.org/10.1016/j.gloenvcha.2014.07.009)

9. W. Steffen, K. Richardson, J. Rockström, S. E. Cornell, I. Fetzer, E. M. Bennett, R. Biggs, S. R. Carpenter, W. de Vries, C. A. de Wit, C. Folke, D. Gerten, J. Heinke, G. M. Mace, L. M. Persson, V. Ramanathan, B. Reyers, S. Sörlin, Planetary boundaries: Guiding human development on a changing planet. *Science* **347**, 1259855 (2015). [Medline](#) [doi:10.1126/science.1259855](https://doi.org/10.1126/science.1259855)
10. S. Díaz, M. Cabido, Vive la différence: Plant functional diversity matters to ecosystem processes. *Trends Ecol. Evol.* **16**, 646–655 (2001). [doi:10.1016/S0169-5347\(01\)02283-2](https://doi.org/10.1016/S0169-5347(01)02283-2)
11. R. J. Scholes, R. Biggs, A biodiversity intactness index. *Nature* **434**, 45–49 (2005). [Medline](#) [doi:10.1038/nature03289](https://doi.org/10.1038/nature03289)
12. P. M. Vitousek, Biological invasions and ecosystem processes: Towards an integration of population biology and ecosystem studies. *Oikos* **57**, 7–13 (1990). [doi:10.2307/3565731](https://doi.org/10.2307/3565731)
13. J. M. Levine, M. Vilà, C. M. D’Antonio, J. S. Dukes, K. Grigulis, S. Lavorel, Mechanisms underlying the impacts of exotic plant invasions. *Proc. Biol. Sci.* **270**, 775–781 (2003). [Medline](#) [doi:10.1098/rspb.2003.2327](https://doi.org/10.1098/rspb.2003.2327)
14. L. Pejchar, H. A. Mooney, Invasive species, ecosystem services and human well-being. *Trends Ecol. Evol.* **24**, 497–504 (2009). [Medline](#) [doi:10.1016/j.tree.2009.03.016](https://doi.org/10.1016/j.tree.2009.03.016)
15. M. Vilà, J. L. Espinar, M. Hejda, P. E. Hulme, V. Jarošík, J. L. Maron, J. Pergl, U. Schaffner, Y. Sun, P. Pyšek, Ecological impacts of invasive alien plants: A meta-analysis of their effects on species, communities and ecosystems. *Ecol. Lett.* **14**, 702–708 (2011). [Medline](#) [doi:10.1111/j.1461-0248.2011.01628.x](https://doi.org/10.1111/j.1461-0248.2011.01628.x)
16. L. Gibson, T. M. Lee, L. P. Koh, B. W. Brook, T. A. Gardner, J. Barlow, C. A. Peres, C. J. Bradshaw, W. F. Laurance, T. E. Lovejoy, N. S. Sodhi, Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**, 378–381 (2011). [Medline](#) [doi:10.1038/nature10425](https://doi.org/10.1038/nature10425)
17. M. Vellend, L. Baeten, I. H. Myers-Smith, S. C. Elmendorf, R. Beauséjour, C. D. Brown, P. De Frenne, K. Verheyen, S. Wipf, Global meta-analysis reveals no net change in local-scale plant biodiversity over time. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 19456–19459 (2013). [Medline](#) [doi:10.1073/pnas.1312779110](https://doi.org/10.1073/pnas.1312779110)
18. C. D. Thomas, Local diversity stays about the same, regional diversity increases, and global diversity declines. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 19187–19188 (2013). [Medline](#) [doi:10.1073/pnas.1319304110](https://doi.org/10.1073/pnas.1319304110)
19. Materials and methods are available as supplementary materials on *Science* Online.
20. T. M. Brooks, R. A. Mittermeier, G. A. da Fonseca, J. Gerlach, M. Hoffmann, J. F. Lamoreux, C. G. Mittermeier, J. D. Pilgrim, A. S. Rodrigues, Global biodiversity conservation priorities. *Science* **313**, 58–61 (2006). [Medline](#) [doi:10.1126/science.1127609](https://doi.org/10.1126/science.1127609)
21. L. N. Hudson, T. Newbold, S. Contu, S. L. Hill, I. Lysenko, A. De Palma, H. R. Phillips, R. A. Senior, D. J. Bennett, H. Booth, A. Choimes, D. L. Correia, J. Day, S. Echeverría-Londoño, M. Garon, M. L. Harrison, D. J. Ingram, M. Jung, V. Kemp, L. Kirkpatrick, C. D. Martin, Y. Pan, H. J. White, J. Aben, S. Abrahamczyk, G. B. Adum, V. Aguilar-Barquero, M. A. Aizen, M. Ancrenaz, E. Arbeláez-Cortés, I. Armbrrecht, B. Azhar, A. B.

- Azpiroz, L. Baeten, A. Báldi, J. E. Banks, J. Barlow, P. Batáry, A. J. Bates, E. M. Bayne, P. Beja, Å. Berg, N. J. Berry, J. E. Bicknell, J. H. Bihn, K. Böhning-Gaese, T. Boekhout, C. Boutin, J. Bouyer, F. Q. Brearley, I. Brito, J. Brunet, G. Buczkowski, E. Buscardo, J. Cabra-García, M. Calviño-Cancela, S. A. Cameron, E. M. Cancelli, T. F. Carrijo, A. L. Carvalho, H. Castro, A. A. Castro-Luna, R. Cerda, A. Cerezo, M. Chauvat, F. M. Clarke, D. F. Cleary, S. P. Connop, B. D’Aniello, P. G. da Silva, B. Darvill, J. Dauber, A. Dejean, T. Diekötter, Y. Dominguez-Haydar, C. F. Dormann, B. Dumont, S. G. Dures, M. Dynesius, L. Edenius, Z. Elek, M. H. Entling, N. Farwig, T. M. Fayle, A. Felicioli, A. M. Felton, G. F. Ficetola, B. K. Filgueiras, S. J. Fonte, L. H. Fraser, D. Fukuda, D. Furlani, J. U. Ganzhorn, J. G. Garden, C. Gheler-Costa, P. Giordani, S. Giordano, M. S. Gottschalk, D. Goulson, A. D. Gove, J. Grogan, M. E. Hanley, T. Hanson, N. R. Hashim, J. E. Hawes, C. Hébert, A. J. Helden, J. A. Henden, L. Hernández, F. Herzog, D. Higuera-Díaz, B. Hilje, F. G. Horgan, R. Horváth, K. Hylander, P. Isaacs-Cubides, M. Ishitani, C. T. Jacobs, V. J. Jaramillo, B. Jauker, M. Jonsell, T. S. Jung, V. Kapoor, V. Kati, E. Katovai, M. Kessler, E. Knop, A. Kolb, Á. Kőrösi, T. Lachat, V. Lantschner, V. Le Féon, G. LeBuhn, J. P. Légaré, S. G. Letcher, N. A. Littlewood, C. A. López-Quintero, M. Louhaichi, G. L. Lövei, M. E. Lucas-Borja, V. H. Luja, K. Maeto, T. Magura, N. A. Mallari, E. Marin-Spiotta, E. J. Marshall, E. Martínez, M. M. Mayfield, G. Mikusinski, J. C. Milder, J. R. Miller, C. L. Morales, M. N. Muchane, M. Muchane, R. Naidoo, A. Nakamura, S. Naoe, G. Nates-Parra, D. A. Navarrete Gutierrez, E. L. Neuschulz, N. Noreika, O. Norfolk, J. A. Noriega, N. M. Nöske, N. O’Dea, W. Oduro, C. Ofori-Boateng, C. O. Oke, L. M. Osgathorpe, J. Paritsis, A. Parra-H, N. Pelegrin, C. A. Peres, A. S. Persson, T. Petanidou, B. Phalan, T. K. Philips, K. Poveda, E. F. Power, S. J. Presley, V. Proença, M. Quaranta, C. Quintero, N. A. Redpath-Downing, J. L. Reid, Y. T. Reis, D. B. Ribeiro, B. A. Richardson, M. J. Richardson, C. A. Robles, J. Römbke, L. P. Romero-Duque, L. Rosselli, S. J. Rossiter, T. H. Roulston, L. Rousseau, J. P. Sadler, S. Sáfián, R. A. Saldaña-Vázquez, U. Samnegård, C. Schüepp, O. Schweiger, J. L. Sedlock, G. Shahabuddin, D. Sheil, F. A. Silva, E. M. Slade, A. H. Smith-Pardo, N. S. Sodhi, E. J. Somarriba, R. A. Sosa, J. C. Stout, M. J. Struebig, Y. H. Sung, C. G. Threlfall, R. Tonietto, B. Tóthmérész, T. Tschardtke, E. C. Turner, J. M. Tylianakis, A. J. Vanbergen, K. Vassilev, H. A. Verboven, C. H. Vergara, P. M. Vergara, J. Verhulst, T. R. Walker, Y. Wang, J. I. Watling, K. Wells, C. D. Williams, M. R. Willig, J. C. Woinarski, J. H. Wolf, B. A. Woodcock, D. W. Yu, A. S. Zaitsev, B. Collen, R. M. Ewers, G. M. Mace, D. W. Purves, J. P. Scharlemann, A. Purvis, The PREDICTS database: A global database of how local terrestrial biodiversity responds to human impacts. *Ecol. Evol.* **4**, 4701–4735 (2014). [Medline](#)
22. T. Newbold, L. N. Hudson, S. L. L. Hill, S. Contu, C. L. Gray, J. P. W. Scharlemann, L. Borger, H. R. P. Phillips, D. Shell, I. Lysenko, A. Purvis, Global patterns of terrestrial assemblage turnover within and among land uses. *Ecography* 10.1111/ecog.01932 (2016).
23. A. J. Hoskins, A. Bush, J. Gilmore, T. Harwood, L. N. Hudson, C. Ware, K. J. Williams, S. Ferrier, Downscaling land-use data to provide global 30" estimates of five land-use classes. *Ecol. Evol.* 10.1002/ece3.2104 (2016).

24. CIESIN, IFPRI, CIAT, *Global Rural-Urban Mapping Project, Version 1 (GRUMPv1): Population Density Grid*. (NASA SEDAC, 2011); available at <http://sedac.ciesin.columbia.edu/data/dataset/grump-v1-population-density>.
25. CIESIN, ITOS, University of Georgia, *Global Roads Open Access Data Set, Version 1 (gROADSv1)* (NASA SEDAC, 2013); available at <http://dx.doi.org/10.7927/H4VD6WCT>.
26. R. A. Fuller, K. N. Irvine, P. Devine-Wright, P. H. Warren, K. J. Gaston, Psychological benefits of greenspace increase with biodiversity. *Biol. Lett.* **3**, 390–394 (2007). [Medline doi:10.1098/rsbl.2007.0149](#)
27. P. A. Sandifer, A. E. Sutton-Grier, B. P. Ward, Exploring connections among nature, biodiversity, ecosystem services, and human health and well-being: Opportunities to enhance health and biodiversity conservation. *Ecosyst. Serv.* **12**, 1–15 (2015). [doi:10.1016/j.ecoser.2014.12.007](#)
28. N. Myers, R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca, J. Kent, Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858 (2000). [Medline doi:10.1038/35002501](#)
29. United Nations, *UN General Assembly Resolution A/RES/70/1* (2015); available at www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E.
30. E. C. Ellis, J. O. Kaplan, D. Q. Fuller, S. Vavrus, K. Klein Goldewijk, P. H. Verburg, Used planet: A global history. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 7978–7985 (2013). [Medline doi:10.1073/pnas.1217241110](#)
31. J. Aben, M. Dorenbosch, S. K. Herzog, A. J. P. Smolders, G. Van Der Velde, Human disturbance affects a deciduous forest bird community in the Andean foothills of central Bolivia. *Bird Conserv. Int.* **18**, 363–380 (2008). [doi:10.1017/S0959270908007326](#)
32. G. B. Adum, M. P. Eichhorn, W. Oduro, C. Ofori-Boateng, M. O. Rödel, Two-stage recovery of amphibian assemblages following selective logging of tropical forests. *Conserv. Biol.* **27**, 354–363 (2013). [Medline doi:10.1111/cobi.12006](#)
33. V. Aguilar-Barquero, F. Jiménez-Hernández, Diversidad y distribución de palmas (Arecaceae) en tres fragmentos de bosque muy húmedo en Costa Rica. *Rev. Biol. Trop.* **57**, 83–92 (2009).
34. E. L. Alcala, C. Alcala, C. N. Dolino, Amphibians and reptiles in tropical rainforest fragments on Negros Island, the Philippines. *Environ. Conserv.* **31**, 254–261 (2004). [doi:10.1017/S0376892904001407](#)
35. O. E. Alcayaga, J. Pizarro-Araya, F. M. Alfaro, J. Cepeda-Pizarro, Spiders (Arachnida, Araneae) associated to agroecosystems in the Elqui Valley (Coquimbo Region, Chile). *Rev. Colomb. Entomol.* **39**, 150–154 (2013).
36. M. del M. Alguacil, E. Torrecillas, G. Hernández, A. Roldán, Changes in the diversity of soil arbuscular mycorrhizal fungi after cultivation for biofuel production in a Guantanamo (Cuba) tropical system. *PLOS ONE* **7**, e34887 (2012). [10.1371/journal.pone.0034887 Medline doi:10.1371/journal.pone.0034887](#)

37. A. Alignier, M. Deconchat, Patterns of forest vegetation responses to edge effect as revealed by a continuous approach. *Ann. For. Sci.* **70**, 601–609 (2013). [doi:10.1007/s13595-013-0301-0](https://doi.org/10.1007/s13595-013-0301-0)
38. A. N. Andersen, B. D. Hoffmann, Conservation value of low fire frequency in tropical savannas: Ants in monsoonal northern Australia. *Austral Ecol.* **36**, 497–503 (2011). [doi:10.1111/j.1442-9993.2010.02151.x](https://doi.org/10.1111/j.1442-9993.2010.02151.x)
39. E. Arbeláez-Cortés, H. A. Rodríguez-Correa, M. Restrepo-Chica, Mixed bird flocks: Patterns of activity and species composition in a region of the Central Andes of Colombia. *Rev. Mex. Biodivers.* **82**, 639–651 (2011).
40. I. Armbrrecht, I. Perfecto, E. Silverman, Limitation of nesting resources for ants in Colombian forests and coffee plantations. *Ecol. Entomol.* **31**, 403–410 (2006). [doi:10.1111/j.1365-2311.2006.00802.x](https://doi.org/10.1111/j.1365-2311.2006.00802.x)
41. J. Arroyo, J. C. Iturrondobeitia, C. Rad, S. Gonzalez-Carcedo, Oribatid mite (Acari) community structure in steppic habitats of Burgos Province, central northern Spain. *J. Nat. Hist.* **39**, 3453–3470 (2005). [doi:10.1080/00222930500240346](https://doi.org/10.1080/00222930500240346)
42. B. Azhar, D. B. Lindenmayer, J. Wood, J. Fischer, A. Manning, C. McElhinny, M. Zakaria, The influence of agricultural system, stand structural complexity and landscape context on foraging birds in oil palm landscapes. *Ibis* **155**, 297–312 (2013). [doi:10.1111/ibi.12025](https://doi.org/10.1111/ibi.12025)
43. N. M. Azman, N. S. Latip, S. A. Sah, M. A. Akil, N. J. Shafie, N. L. Khairuddin, Avian diversity and feeding guilds in a secondary forest, an oil palm plantation and a paddy field in riparian areas of the kerian river basin, perak, malaysia. *Trop. Life Sci. Res.* **22**, 45–64 (2011). [Medline](https://pubmed.ncbi.nlm.nih.gov/22111111/)
44. A. B. Azpiroz, J. G. Blake, Avian assemblages in altered and natural grasslands in the northern Campos of Uruguay. *Condor* **111**, 21–35 (2009). [doi:10.1525/cond.2009.080111](https://doi.org/10.1525/cond.2009.080111)
45. L. Baeten, D. Velghe, M. Vanhellemont, P. De Frenne, M. Hermy, K. Verheyen, Early trajectories of spontaneous vegetation recovery after intensive agricultural land use. *Restor. Ecol.* **18**, 379–386 (2010). [doi:10.1111/j.1526-100X.2009.00627.x](https://doi.org/10.1111/j.1526-100X.2009.00627.x)
46. L. Baeten, M. Hermy, S. Van Daele, K. Verheyen, Unexpected understorey community development after 30 years in ancient and post-agricultural forests. *J. Ecol.* **98**, 1447–1453 (2010). [doi:10.1111/j.1365-2745.2010.01711.x](https://doi.org/10.1111/j.1365-2745.2010.01711.x)
47. A. Báldi, P. Batáry, S. Erdős, Effects of grazing intensity on bird assemblages and populations of Hungarian grasslands. *Agric. Ecosyst. Environ.* **108**, 251–263 (2005). [doi:10.1016/j.agee.2005.02.006](https://doi.org/10.1016/j.agee.2005.02.006)
48. J. E. Banks, P. Sandvik, L. Keesecker, Beetle (Coleoptera) and spider (Araneae) diversity in a mosaic of farmland, edge, and tropical forest habitats in western Costa Rica. *Pan-Pac. Entomol.* **83**, 152–160 (2007). [doi:10.3956/0031-0603-83.2.152](https://doi.org/10.3956/0031-0603-83.2.152)
49. S. K. Baral, K. Katzensteiner, Diversity of vascular plant communities along a disturbance gradient in a central mid-hill community forest of Nepal. *Banko Janakari* **19**, 3–7 (2009). [doi:10.3126/banko.v19i1.2176](https://doi.org/10.3126/banko.v19i1.2176)

50. J. Barlow, T. A. Gardner, I. S. Araujo, T. C. Avila-Pires, A. B. Bonaldo, J. E. Costa, M. C. Esposito, L. V. Ferreira, J. Hawes, M. I. Hernandez, M. S. Hoogmoed, R. N. Leite, N. F. Lo-Man-Hung, J. R. Malcolm, M. B. Martins, L. A. Mestre, R. Miranda-Santos, A. L. Nunes-Gutjahr, W. L. Overal, L. Parry, S. L. Peters, M. A. Ribeiro-Junior, M. N. da Silva, C. da Silva Motta, C. A. Peres, Quantifying the biodiversity value of tropical primary, secondary, and plantation forests. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 18555–18560 (2007). [Medline](#) [doi:10.1073/pnas.0703333104](https://doi.org/10.1073/pnas.0703333104)
51. J. Barlow, L. A. M. Mestre, T. A. Gardner, C. A. Peres, The value of primary, secondary and plantation forests for Amazonian birds. *Biol. Conserv.* **136**, 212–231 (2007). [doi:10.1016/j.biocon.2006.11.021](https://doi.org/10.1016/j.biocon.2006.11.021)
52. J. Barlow, W. L. Overal, I. S. Araujo, T. A. Gardner, C. A. Peres, The value of primary, secondary and plantation forests for fruit-feeding butterflies in the Brazilian Amazon. *J. Appl. Ecol.* **44**, 1001–1012 (2007). [doi:10.1111/j.1365-2664.2007.01347.x](https://doi.org/10.1111/j.1365-2664.2007.01347.x)
53. B. I. P. Barratt, C. M. Ferguson, R. A. S. Logan, D. Barton, N. L. Bell, S. U. Sarathchandra, R. J. Townsend, Biodiversity of indigenous tussock grassland sites in Otago, Canterbury and the central North Island of New Zealand I. The macro-invertebrate fauna. *J. R. Soc. N. Z.* **35**, 287–301 (2005). [doi:10.1080/03014223.2005.9517785](https://doi.org/10.1080/03014223.2005.9517785)
54. B. I. P. Barratt, S. P. Worner, K. Affeld, C. M. Ferguson, D. M. Barton, N. L. Bell, R. J. Townsend, Biodiversity of indigenous tussock grassland sites in Otago, Canterbury and the Central North Island of New Zealand VI. Coleoptera biodiversity, community structure, exotic species invasion, and the effect of disturbance by agricultural development. *J. R. Soc. N. Z.* **42**, 217–239 (2012). [doi:10.1080/03036758.2011.559664](https://doi.org/10.1080/03036758.2011.559664)
55. L. Barrico, A. M. Azul, M. C. Morais, A. Pereira Coutinho, H. Freitas, P. Castro, Biodiversity in urban ecosystems: Plants and macromycetes as indicators for conservation planning in the city of Coimbra (Portugal). *Landsc. Urban Plan.* **106**, 88–102 (2012). [doi:10.1016/j.landurbplan.2012.02.011](https://doi.org/10.1016/j.landurbplan.2012.02.011)
56. P. Bartolommei, A. Mortelliti, F. Pezzo, L. Puglisi, Distribution of nocturnal birds (Strigiformes and Caprimulgidae) in relation to land-use types, extent and configuration in agricultural landscapes of Central Italy. *Rend. Lincei* **24**, 13–21 (2013). [doi:10.1007/s12210-012-0211-3](https://doi.org/10.1007/s12210-012-0211-3)
57. Y. Basset, O. Missa, A. Alonso, S. E. Miller, G. Curletti, M. De Meyer, C. Eardley, O. T. Lewis, M. W. Mansell, V. Novotny, T. Wagner, Changes in arthropod assemblages along a wide gradient of disturbance in Gabon. *Conserv. Biol.* **22**, 1552–1563 (2008). [Medline](#) [doi:10.1111/j.1523-1739.2008.01017.x](https://doi.org/10.1111/j.1523-1739.2008.01017.x)
58. A. J. Bates, J. P. Sadler, A. J. Fairbrass, S. J. Falk, J. D. Hale, T. J. Matthews, Changing bee and hoverfly pollinator assemblages along an urban-rural gradient. *PLOS ONE* **6**, e23459 (2011). [Medline](#) [doi:10.1371/journal.pone.0023459](https://doi.org/10.1371/journal.pone.0023459)
59. B. Baur, C. Cremene, G. Groza, L. Rakosy, A. A. Schileiko, A. Baur, P. Stoll, A. Erhardt, Effects of abandonment of subalpine hay meadows on plant and invertebrate diversity in Transylvania, Romania. *Biol. Conserv.* **132**, 261–273 (2006). [doi:10.1016/j.biocon.2006.04.018](https://doi.org/10.1016/j.biocon.2006.04.018)

60. S. Benedick, J. K. Hill, N. Mustaffa, V. K. Chey, M. Maryati, J. B. Searle, M. Schilthuizen, K. C. Hamer, Impacts of rain forest fragmentation on butterflies in northern Borneo: Species richness, turnover and the value of small fragments. *J. Appl. Ecol.* **43**, 967–977 (2006). [doi:10.1111/j.1365-2664.2006.01209.x](https://doi.org/10.1111/j.1365-2664.2006.01209.x)
61. A. Berg, K. Ahrne, E. Ockinger, R. Svensson, B. Soderstrom, Butterfly distribution and abundance is affected by variation in the Swedish forest-farmland landscape. *Biol. Conserv.* **144**, 2819–2831 (2011). [doi:10.1016/j.biocon.2011.07.035](https://doi.org/10.1016/j.biocon.2011.07.035)
62. H. Bernard, J. Fjeldsa, M. Mohamed, A case study on the effects of disturbance and conversion of tropical lowland rain forest on the non-volant small mammals in north Borneo: Management implications. *Mammal Study* **34**, 85–96 (2009). [doi:10.3106/041.034.0204](https://doi.org/10.3106/041.034.0204)
63. N. J. Berry, O. L. Phillips, S. L. Lewis, J. K. Hill, D. P. Edwards, N. B. Tawatao, N. Ahmad, D. Magintan, C. V. Khen, M. Maryati, R. C. Ong, K. C. Hamer, The high value of logged tropical forests: Lessons from northern Borneo. *Biodivers. Conserv.* **19**, 985–997 (2010). [doi:10.1007/s10531-010-9779-z](https://doi.org/10.1007/s10531-010-9779-z)
64. J. Bicknell, C. A. Peres, Vertebrate population responses to reduced-impact logging in a neotropical forest. *For. Ecol. Manage.* **259**, 2267–2275 (2010). [doi:10.1016/j.foreco.2010.02.027](https://doi.org/10.1016/j.foreco.2010.02.027)
65. J. H. Bihn, M. Verhaagh, M. Brändle, R. Brandl, Do secondary forests act as refuges for old growth forest animals? Recovery of ant diversity in the Atlantic forest of Brazil. *Biol. Conserv.* **141**, 733–743 (2008). [doi:10.1016/j.biocon.2007.12.028](https://doi.org/10.1016/j.biocon.2007.12.028)
66. R. Billeter, J. Liira, D. Bailey, R. Bugter, P. Arens, I. Augenstein, S. Aviron, J. Baudry, R. Bukacek, F. Burel, M. Cerny, G. De Blust, R. De Cock, T. Diekötter, H. Dietz, J. Dirksen, C. Dormann, W. Durka, M. Frenzel, R. Hamersky, F. Hendrickx, F. Herzog, S. Klotz, B. Koolstra, A. Lausch, D. Le Coeur, J. P. Maelfait, P. Opdam, M. Roubalova, A. Schermann, N. Schermann, T. Schmidt, O. Schweiger, M. J. M. Smulders, M. Speelmans, P. Simova, J. Verboom, W. K. R. E. Van Wingerden, M. Zobel, P. J. Edwards, Indicators for biodiversity in agricultural landscapes: A pan-European study. *J. Appl. Ecol.* **45**, 141–150 (2008). [doi:10.1111/j.1365-2664.2007.01393.x](https://doi.org/10.1111/j.1365-2664.2007.01393.x)
67. R. J. Blake, D. B. Westbury, B. A. Woodcock, P. Sutton, S. G. Potts, Enhancing habitat to help the plight of the bumblebee. *Pest Manag. Sci.* **67**, 377–379 (2011). [Medline doi:10.1002/ps.2136](https://doi.org/10.1002/ps.2136)
68. K. R. Blanche, J. A. Ludwig, S. A. Cunningham, Proximity to rainforest enhances pollination and fruit set in orchards. *J. Appl. Ecol.* **43**, 1182–1187 (2006). [doi:10.1111/j.1365-2664.2006.01230.x](https://doi.org/10.1111/j.1365-2664.2006.01230.x)
69. R. Blanche, S. A. Cunningham, Rain forest provides pollinating beetles for atemoya crops. *J. Econ. Entomol.* **98**, 1193–1201 (2005). [Medline doi:10.1603/0022-0493-98.4.1193](https://doi.org/10.1603/0022-0493-98.4.1193)
70. R. Bóçon, thesis, “Riqueza e Abundância de Aves em três Estágios Sucessionais da Floresta Ombrófila Densa Submontana, Antonina, Paraná,” Hungarian Academy of Sciences, 2010.
71. P. A. V. Borges, J. M. Lobo, E. B. Azevedo, C. S. Gaspar, C. Melo, L. V. Nunes; P. A. V. Borges et al, Invasibility and species richness of island endemic arthropods: A general

- model of endemic vs. exotic species. *J. Biogeogr.* **33**, 169–187 (2006).
[doi:10.1111/j.1365-2699.2005.01324.x](https://doi.org/10.1111/j.1365-2699.2005.01324.x)
72. S. H. Borges, Bird assemblages in secondary forests developing after slash-and-burn agriculture in the Brazilian Amazon. *J. Trop. Ecol.* **23**, 469–477 (2007).
[doi:10.1017/S0266467407004105](https://doi.org/10.1017/S0266467407004105)
73. M. M. Bos, I. Steffan-Dewenter, T. Tscharntke, The contribution of cacao agroforests to the conservation of lower canopy ant and beetle diversity in Indonesia. *Biodivers. Conserv.* **16**, 2429–2444 (2007). [doi:10.1007/s10531-007-9196-0](https://doi.org/10.1007/s10531-007-9196-0)
74. B. M. Bösing, D. H. Haarmeyer, J. Dengler, J. U. Ganzhorn, U. Schmiedel, Effects of livestock grazing and habitat characteristics on small mammal communities in the Knersvlakte, South Africa. *J. Arid Environ.* **104**, 124–131 (2014).
[doi:10.1016/j.jaridenv.2014.02.011](https://doi.org/10.1016/j.jaridenv.2014.02.011)
75. C. Boutin, P. A. Martin, A. Baril, Arthropod diversity as affected by agricultural management (organic and conventional farming), plant species, and landscape context. *Ecoscience* **16**, 492–501 (2009). [doi:10.2980/16-4-3250](https://doi.org/10.2980/16-4-3250)
76. J. Bouyer, Y. Sana, Y. Samandoulgou, J. Cesar, L. Guerrini, C. Kabore-Zoungrana, D. Dulieu, Identification of ecological indicators for monitoring ecosystem health in the trans-boundary W Regional park: A pilot study. *Biol. Conserv.* **138**, 73–88 (2007).
[doi:10.1016/j.biocon.2007.04.001](https://doi.org/10.1016/j.biocon.2007.04.001)
77. C. Bragagnolo, A. A. Nogueira, R. Pinto-da-Rocha, R. Pardini, Harvestmen in an Atlantic forest fragmented landscape: Evaluating assemblage response to habitat quality and quantity. *Biol. Conserv.* **139**, 389–400 (2007). [doi:10.1016/j.biocon.2007.07.008](https://doi.org/10.1016/j.biocon.2007.07.008)
78. J. S. Brandt, E. M. Wood, A. M. Pidgeon, L.-X. Han, Z. Fang, V. C. Radeloff, Sacred forests are keystone structures for forest bird conservation in southwest China's Himalayan Mountains. *Biol. Conserv.* **166**, 34–42 (2013). [doi:10.1016/j.biocon.2013.06.014](https://doi.org/10.1016/j.biocon.2013.06.014)
79. F. Q. Brearley, Below-ground secondary succession in tropical forests of Borneo. *J. Trop. Ecol.* **27**, 413–420 (2011). [doi:10.1017/S0266467411000149](https://doi.org/10.1017/S0266467411000149)
80. J. A. D. Breedts, N. Dreber, K. Kellner, Post-wildfire regeneration of rangeland productivity and functionality - observations across three semi-arid vegetation types in South Africa. *African J. Range Forage Sci.* **30**, 161–167 (2013). [doi:10.2989/10220119.2013.816367](https://doi.org/10.2989/10220119.2013.816367)
81. I. Brito, M. J. Goss, M. de Carvalho, O. Chatagnier, D. van Tuinen, Impact of tillage system on arbuscular mycorrhiza fungal communities in the soil under Mediterranean conditions. *Soil Tillage Res.* **121**, 63–67 (2012). [doi:10.1016/j.still.2012.01.012](https://doi.org/10.1016/j.still.2012.01.012)
82. J. Brunet, K. Valtinat, M. L. Mayr, A. Felton, M. Lindblad, H. H. Bruun, Understory succession in post-agricultural oak forests: Habitat fragmentation affects forest specialists and generalists differently. *For. Ecol. Manage.* **262**, 1863–1871 (2011).
[doi:10.1016/j.foreco.2011.08.007](https://doi.org/10.1016/j.foreco.2011.08.007)
83. G. Buczkowski, Extreme life history plasticity and the evolution of invasive characteristics in a native ant. *Biol. Invasions* **12**, 3343–3349 (2010). [doi:10.1007/s10530-010-9727-6](https://doi.org/10.1007/s10530-010-9727-6)

84. G. Buczkowski, D. S. Richmond, The effect of urbanization on ant abundance and diversity: A temporal examination of factors affecting biodiversity. *PLOS ONE* **7**, e41729 (2012). [Medline](#) [doi:10.1371/journal.pone.0041729](#)
85. C. M. Buddle, D. P. Shorthouse, Effects of experimental harvesting on spider (Araneae) assemblages in boreal deciduous forests. *Can. Entomol.* **140**, 437–452 (2008). [doi:10.4039/n07-LS01](#)
86. E. Buscardo, G. F. Smith, D. L. Kelly, H. Freitas, S. Iremonger, F. J. G. Mitchell, S. O'Donoghue, A.-M. McKee, The early effects of afforestation on biodiversity of grasslands in Ireland. *Biodivers. Conserv.* **17**, 1057–1072 (2008). [doi:10.1007/s10531-007-9275-2](#)
87. J. Cabra-García, C. Bermúdez-Rivas, A. M. Osorio, P. Chacón, Cross-taxon congruence of alpha and beta diversity among five leaf litter arthropod groups in Colombia. *Biodivers. Conserv.* **21**, 1493–1508 (2012). [doi:10.1007/s10531-012-0259-5](#)
88. N. C. Cáceres, R. P. Nápoli, J. Casella, W. Hannibal, Mammals in a fragmented savannah landscape in south-western Brazil. *J. Nat. Hist.* **44**, 491–512 (2010). [doi:10.1080/00222930903477768](#)
89. N. L. Cagle, Snake species distributions and temperate grasslands: A case study from the American tallgrass prairie. *Biol. Conserv.* **141**, 744–755 (2008). [doi:10.1016/j.biocon.2008.01.003](#)
90. M. Calviño-Cancela, M. Rubido-Bará, E. J. B. van Etten, Do eucalypt plantations provide habitat for native forest biodiversity? *For. Ecol. Manage.* **270**, 153–162 (2012). [doi:10.1016/j.foreco.2012.01.019](#)
91. S. A. Cameron, J. D. Lozier, J. P. Strange, J. B. Koch, N. Cordes, L. F. Solter, T. L. Griswold, Patterns of widespread decline in North American bumble bees. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 662–667 (2011). [Medline](#) [doi:10.1073/pnas.1014743108](#)
92. D. Carpenter, P. M. Hammond, E. Sherlock, A. Lidgett, K. Leigh, P. Eggleton, Biodiversity of soil macrofauna in the New Forest: A benchmark study across a national park landscape. *Biodivers. Conserv.* **21**, 3385–3410 (2012). [doi:10.1007/s10531-012-0369-0](#)
93. T. F. Carrijo, D. Brandao, D. E. de Oliveira, D. A. Costa, T. Santos, Effects of pasture implantation on the termite (Isoptera) fauna in the Central Brazilian Savanna (Cerrado). *J. Insect Conserv.* **13**, 575–581 (2009). [doi:10.1007/s10841-008-9205-y](#)
94. A. L. de Carvalho, E. J. L. Ferreira, J. M. T. Lima, A. L. de Carvalho, Floristic and structural comparisons among palm communities in primary and secondary forest fragments of the Raimundo Irineu Serra Environmental Protection Area–Rio Branco, Acre, Brazil. *Acta Amazon.* **40**, 657–666 (2010). [doi:10.1590/S0044-59672010000400004](#)
95. C. Cassano, J. Barlow, R. Pardini, Forest loss or management intensification? Identifying causes of mammal decline in cacao agroforests. *Biol. Conserv.* **169**, 14–22 (2014). [doi:10.1016/j.biocon.2013.10.006](#)
96. H. Castro, V. Lehsten, S. Lavorel, H. Freitas, Functional response traits in relation to land use change in the Montado. *Agric. Ecosyst. Environ.* **137**, 183–191 (2010). [doi:10.1016/j.agee.2010.02.002](#)

97. A. A. Castro-Luna, V. J. Sosa, G. Castillo-Campos, Bat diversity and abundance associated with the degree of secondary succession in a tropical forest mosaic in south-eastern Mexico. *Anim. Conserv.* **10**, 219–228 (2007). [doi:10.1111/j.1469-1795.2007.00097.x](https://doi.org/10.1111/j.1469-1795.2007.00097.x)
98. Center for International Forestry Research, *Multidisciplinary Landscape Assessment—Cameroon* (Center for International Forestry Research, 2013); available at www.cifor.org/mla/_ref/method/index.htm.
99. Center for International Forestry Research, *Multidisciplinary Landscape Assessment—Philippines* (Center for International Forestry Research, 2013); available at www.cifor.org/mla/_ref/method/index.htm.
100. A. Cerezo, M. C. Conde, S. L. Poggio, Pasture area and landscape heterogeneity are key determinants of bird diversity in intensively managed farmland. *Biodivers. Conserv.* **20**, 2649–2667 (2011). [doi:10.1007/s10531-011-0096-y](https://doi.org/10.1007/s10531-011-0096-y)
101. K. A. Chapman, P. B. Reich, Land use and habitat gradients determine bird community diversity and abundance in suburban, rural and reserve landscapes of Minnesota, USA. *Biol. Conserv.* **135**, 527–541 (2007). [doi:10.1016/j.biocon.2006.10.050](https://doi.org/10.1016/j.biocon.2006.10.050)
102. M. Chauvat, V. Wolters, J. Dauber, Response of collembolan communities to land-use change and grassland succession. *Ecography* **30**, 183–192 (2007). [doi:10.1111/j.0906-7590.2007.04888.x](https://doi.org/10.1111/j.0906-7590.2007.04888.x)
103. M. Christensen, J. Heilmann-Clausen, Forest biodiversity gradients and the human impact in Annapurna Conservation Area, Nepal. *Biodivers. Conserv.* **18**, 2205–2221 (2009). [doi:10.1007/s10531-009-9583-9](https://doi.org/10.1007/s10531-009-9583-9)
104. R. J. Clark, P. J. Gerard, J. M. Mellsop, Spider biodiversity and density following cultivation in pastures in the Waikato, New Zealand. *N. Z. J. Agric. Res.* **47**, 247–259 (2004). [doi:10.1080/00288233.2004.9513592](https://doi.org/10.1080/00288233.2004.9513592)
105. F. M. Clarke, L. V. Rostant, P. A. Racey, Life after logging: Post-logging recovery of a neotropical bat community. *J. Appl. Ecol.* **42**, 409–420 (2005). [doi:10.1111/j.1365-2664.2005.01024.x](https://doi.org/10.1111/j.1365-2664.2005.01024.x)
106. D. F. R. Cleary, A. Ø. Mooers, K. A. O. Eichhorn, J. van Tol, R. de Jong, S. B. J. Menken, Diversity and community composition of butterflies and odonates in an ENSO-induced fire affected habitat mosaic: A case study from East Kalimantan, Indonesia. *Oikos* **105**, 426–448 (2004). [doi:10.1111/j.0030-1299.2004.12219.x](https://doi.org/10.1111/j.0030-1299.2004.12219.x)
107. D. F. R. Cleary, A. O. Mooers, Burning and logging differentially affect endemic vs. widely distributed butterfly species in Borneo. *Divers. Distrib.* **12**, 409–416 (2006). [doi:10.1111/j.1366-9516.2006.00256.x](https://doi.org/10.1111/j.1366-9516.2006.00256.x)
108. K. L. Cockle, M. L. Leonard, A. A. Bodrati, Presence and abundance of birds in an Atlantic forest reserve and adjacent plantation of shade-grown yerba mate, in Paraguay. *Biodivers. Conserv.* **14**, 3265–3288 (2005). [doi:10.1007/s10531-004-0446-0](https://doi.org/10.1007/s10531-004-0446-0)
109. S. Connop, T. Hill, J. Steer, P. Shaw, Microsatellite analysis reveals the spatial dynamics of *Bombus humilis* and *Bombus sylvarum*. *Insect Conserv. Divers.* **4**, 212–221 (2011). [doi:10.1111/j.1752-4598.2010.00116.x](https://doi.org/10.1111/j.1752-4598.2010.00116.x)

110. M. D. Craig, A. H. Grigg, M. J. Garkaklis, R. J. Hobbs, C. D. Grant, P. A. Fleming, G. E. S. J. Hardy, Does habitat structure influence capture probabilities? A study of reptiles in a eucalypt forest. *Wildl. Res.* **36**, 509–515 (2009). [doi:10.1071/WR09014](https://doi.org/10.1071/WR09014)
111. M. D. Craig, A. H. Grigg, R. J. Hobbs, G. E. S. J. Hardy, Does coarse woody debris density and volume influence the terrestrial vertebrate community in restored bauxite mines? *For. Ecol. Manage.* **318**, 142–150 (2014). [doi:10.1016/j.foreco.2014.01.011](https://doi.org/10.1016/j.foreco.2014.01.011)
112. M. D. Craig, G. E. S. J. Hardy, J. B. Fontaine, M. J. Garkakalis, A. H. Grigg, C. D. Grant, P. A. Fleming, R. J. Hobbs, Identifying unidirectional and dynamic habitat filters to faunal recolonisation in restored mine-pits. *J. Appl. Ecol.* **49**, 919–928 (2012). [doi:10.1111/j.1365-2664.2012.02152.x](https://doi.org/10.1111/j.1365-2664.2012.02152.x)
113. S. A. Cunningham, N. A. Schellhorn, A. Marcora, M. Batley, Movement and phenology of bees in a subtropical Australian agricultural landscape. *Austral Ecol.* **38**, 456–464 (2013). [doi:10.1111/j.1442-9993.2012.02432.x](https://doi.org/10.1111/j.1442-9993.2012.02432.x)
114. P. G. da Silva, thesis, “Espécies de Scarabaeinae (Coleoptera: Scarabaeidae) de Fragmentos Florestais com Diferentes Níveis de Alteração em Santa Maria, Rio Grande do Sul,” Universidade Federal de Santa Maria, 2011).
115. B. D’Aniello, I. Stanislao, S. Bonelli, E. Balletto, Haying and grazing effects on the butterfly communities of two Mediterranean-area grasslands. *Biodivers. Conserv.* **20**, 1731–1744 (2011). [doi:10.1007/s10531-011-0058-4](https://doi.org/10.1007/s10531-011-0058-4)
116. E. Danquah, S. K. Oppong, M. E. Nutsuakor, Effect of protected area category on mammal abundance in Western Ghana. *J. Biodivers. Environ. Sci.* **2**, 50–57 (2012).
117. B. Darvill, M. E. Knight, D. Goulson, Use of genetic markers to quantify bumblebee foraging range and nest density. *Oikos* **107**, 471–478 (2004). [doi:10.1111/j.0030-1299.2004.13510.x](https://doi.org/10.1111/j.0030-1299.2004.13510.x)
118. A. L. V. Davis, T. K. Philips, Effect of deforestation on a Southwest Ghana dung beetle assemblage (Coleoptera : Scarabaeidae) at the periphery of Ankasa conservation area. *Environ. Entomol.* **34**, 1081–1088 (2005). [doi:10.1093/ee/34.5.1081](https://doi.org/10.1093/ee/34.5.1081)
119. J. Dawson *et al.*, Bird communities of the lower Waria Valley, Morobe Province, Papua New Guinea: A comparison between habitat types. *Trop. Conserv. Sci.* **4**, 317–348 (2011).
120. N. D’Cruze, S. Kumar, Effects of anthropogenic activities on lizard communities in northern Madagascar. *Anim. Conserv.* **14**, 542–552 (2011). [doi:10.1111/j.1469-1795.2011.00459.x](https://doi.org/10.1111/j.1469-1795.2011.00459.x)
121. C. de Sassi, O. T. Lewis, J. M. Tylianakis, Plant-mediated and nonadditive effects of two global change drivers on an insect herbivore community. *Ecology* **93**, 1892–1901 (2012). [Medline doi:10.1890/11-1839.1](https://doi.org/10.1890/11-1839.1)
122. V. M. de Souza, B. de Souza, E. F. Morato, Effect of the forest succession on the anurans (Amphibia: Anura) of the Reserve Catuaba and its periphery, Acre, southwestern Amazonia. *Rev. Bras. Zool.* **25**, 49–57 (2008).
123. B. de Thoisy, C. Richard-Hansen, B. Goguillon, P. Joubert, J. Obstancias, P. Winterton, S. Brosse, Rapid evaluation of threats to biodiversity: Human footprint score and large

- vertebrate species responses in French Guiana. *Biodivers. Conserv.* **19**, 1567–1584 (2010). [doi:10.1007/s10531-010-9787-z](https://doi.org/10.1007/s10531-010-9787-z)
124. J. H. C. Delabie, R. Céréghino, S. Groc, A. Dejean, M. Gibernau, B. Corbara, A. Dejean, Ants as biological indicators of Wayana Amerindian land use in French Guiana. *C. R. Biol.* **332**, 673–684 (2009). [Medline](https://pubmed.ncbi.nlm.nih.gov/19111111/) [doi:10.1016/j.crv.2009.01.006](https://doi.org/10.1016/j.crv.2009.01.006)
125. J. L. Devineau, A. Fournier, S. Nignan, “Ordinary biodiversity” in western Burkina Faso (West Africa): What vegetation do the state forests conserve? *Biodivers. Conserv.* **18**, 2075–2099 (2009). [doi:10.1007/s10531-008-9574-2](https://doi.org/10.1007/s10531-008-9574-2)
126. T. Diekötter, K. Walther-Hellwig, M. Conradi, M. Suter, R. Frankl, Effects of landscape elements on the distribution of the rare bumblebee species *Bombus muscorum* in an agricultural landscape. *Biodivers. Conserv.* **15**, 57–68 (2006). [doi:10.1007/s10531-004-2932-9](https://doi.org/10.1007/s10531-004-2932-9)
127. J. Dolia, M. S. Devy, N. A. Aravind, A. Kumar, Adult butterfly communities in coffee plantations around a protected area in the Western Ghats, India. *Anim. Conserv.* **11**, 26–34 (2008). [doi:10.1111/j.1469-1795.2007.00143.x](https://doi.org/10.1111/j.1469-1795.2007.00143.x)
128. E. Domínguez, N. Bahamonde, C. Muñoz-Escobar, Efectos de la extracción de turba sobre la composición y estructura de una turbera de *Sphagnum* explotada y abandonada hace 20 años, Chile. *An. Inst. Patagon.* **40**, 37–45 (2012). [doi:10.4067/S0718-686X2012000200003](https://doi.org/10.4067/S0718-686X2012000200003)
129. Y. Dominguez-Haydar, I. Armbrrecht, Response of ants and their seed removal in rehabilitation areas and forests at El Cerrejon coal mine in Colombia. *Restor. Ecol.* **19**, 178–184 (2010). [doi:10.1111/j.1526-100X.2010.00735.x](https://doi.org/10.1111/j.1526-100X.2010.00735.x)
130. B. Dumont, A. Farruggia, J.-P. Garel, P. Bachelard, E. Boitier, M. Frain, How does grazing intensity influence the diversity of plants and insects in a species-rich upland grassland on basalt soils? *Grass Forage Sci.* **64**, 92–105 (2009). [doi:10.1111/j.1365-2494.2008.00674.x](https://doi.org/10.1111/j.1365-2494.2008.00674.x)
131. S. G. Dures, G. S. Cumming, The confounding influence of homogenising invasive species in a globally endangered and largely urban biome: Does habitat quality dominate avian biodiversity? *Biol. Conserv.* **143**, 768–777 (2010). [doi:10.1016/j.biocon.2009.12.019](https://doi.org/10.1016/j.biocon.2009.12.019)
132. L. Edenius, G. Mikusiński, J. Bergh, Can repeated fertilizer applications to young Norway spruce enhance avian diversity in intensively managed forests? *Ambio* **40**, 521–527 (2011). [Medline](https://pubmed.ncbi.nlm.nih.gov/21111111/) [doi:10.1007/s13280-011-0137-5](https://doi.org/10.1007/s13280-011-0137-5)
133. F. Eigenbrod, S. J. Hecnar, L. Fahrig, Accessible habitat: An improved measure of the effects of habitat loss and roads on wildlife populations. *Landscape Ecol.* **23**, 159–168 (2008). [doi:10.1007/s10980-007-9174-7](https://doi.org/10.1007/s10980-007-9174-7)
134. Z. Elek, G. L. Lovei, Patterns in ground beetle (Coleoptera: Carabidae) assemblages along an urbanisation gradient in Denmark. *Acta Oecol.* **32**, 104–111 (2007). [doi:10.1016/j.actao.2007.03.008](https://doi.org/10.1016/j.actao.2007.03.008)
135. W. Endo, C. A. Peres, E. Salas, S. Mori, J.-L. Sanchez-Vega, G. H. Shepard, V. Pacheco, D. W. Yu, Game vertebrate densities in hunted and nonhunted forest sites in Manu National Park, Peru. *Biotropica* **42**, 251–261 (2010). [doi:10.1111/j.1744-7429.2009.00546.x](https://doi.org/10.1111/j.1744-7429.2009.00546.x)

136. R. M. Ewers, S. Thorpe, R. K. Didham, Synergistic interactions between edge and area effects in a heavily fragmented landscape. *Ecology* **88**, 96–106 (2007). [Medline](#) [doi:10.1890/0012-9658\(2007\)88\[96:SIBEAA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2007)88[96:SIBEAA]2.0.CO;2)
137. A. Faruk, D. Belabut, N. Ahmad, R. J. Knell, T. W. J. Garner, Effects of oil-palm plantations on diversity of tropical anurans. *Conserv. Biol.* **27**, 615–624 (2013). [Medline](#) [doi:10.1111/cobi.12062](https://doi.org/10.1111/cobi.12062)
138. N. Farwig, D. Bailey, E. Bochud, J. D. Herrmann, E. Kindler, N. Reusser, C. Schüepp, M. H. Schmidt-Entling, Isolation from forest reduces pollination, seed predation and insect scavenging in Swiss farmland. *Landscape Ecol.* **24**, 919–927 (2009). [doi:10.1007/s10980-009-9376-2](https://doi.org/10.1007/s10980-009-9376-2)
139. N. Farwig, N. Sajita, K. Böhning-Gaese, Conservation value of forest plantations for bird communities in western Kenya. *For. Ecol. Manage.* **255**, 3885–3892 (2008). [doi:10.1016/j.foreco.2008.03.042](https://doi.org/10.1016/j.foreco.2008.03.042)
140. T. M. Fayle, E. C. Turner, J. L. Snaddon, V. K. Chey, A. Y. C. Chung, P. Eggleton, W. A. Foster, Oil palm expansion into rain forest greatly reduces ant biodiversity in canopy, epiphytes and leaf-litter. *Basic Appl. Ecol.* **11**, 337–345 (2010). [doi:10.1016/j.baec.2009.12.009](https://doi.org/10.1016/j.baec.2009.12.009)
141. R. J. Fensham, J. M. Dwyer, T. J. Eyre, R. J. Fairfax, J. Wang, The effect of clearing on plant composition in mulga (*Acacia aneura*) dry forest, Australia. *Austral Ecol.* **37**, 183–192 (2012). [doi:10.1111/j.1442-9993.2011.02261.x](https://doi.org/10.1111/j.1442-9993.2011.02261.x)
142. H. Fermon, M. Waltert, R. I. Vane-Wright, M. Muhlenberg, Forest use and vertical stratification in fruit-feeding butterflies of Sulawesi, Indonesia: Impacts for conservation. *Biodivers. Conserv.* **14**, 333–350 (2005). [doi:10.1007/s10531-004-5354-9](https://doi.org/10.1007/s10531-004-5354-9)
143. I. C. Fernández, J. A. Simonetti, Small mammal assemblages in fragmented shrublands of urban areas of Central Chile. *Urban Ecosyst.* **16**, 377–387 (2013). [doi:10.1007/s11252-012-0272-1](https://doi.org/10.1007/s11252-012-0272-1)
144. C. Fiera, Preliminary data on the species diversity of Collembola (Hexapoda: Collembola) along an urban gradient in București. *Trav. Mus. Antipa* **51**, 363–367 (2008).
145. M. M. Fierro, L. Cruz-López, D. Sánchez, R. Villanueva-Gutiérrez, R. Vandame, Effect of biotic factors on the spatial distribution of stingless bees (Hymenoptera: Apidae, Meliponini) in fragmented neotropical habitats. *Neotrop. Entomol.* **41**, 95–104 (2012). [Medline](#) [doi:10.1007/s13744-011-0009-5](https://doi.org/10.1007/s13744-011-0009-5)
146. B. K. C. Filgueiras, L. Iannuzzi, I. R. Leal, Habitat fragmentation alters the structure of dung beetle communities in the Atlantic Forest. *Biol. Conserv.* **144**, 362–369 (2011). [doi:10.1016/j.biocon.2010.09.013](https://doi.org/10.1016/j.biocon.2010.09.013)
147. H. K. Fırıncioğlu, S. S. Seefeldt, B. Sahin, M. Vural, Assessment of grazing effect on sheep fescue (*Festuca valesiaca*) dominated steppe rangelands, in the semi-arid Central Anatolian region of Turkey. *J. Arid Environ.* **73**, 1149–1157 (2009). [doi:10.1016/j.jaridenv.2009.05.012](https://doi.org/10.1016/j.jaridenv.2009.05.012)

148. D. J. Flaspohler, C. P. Giardina, G. P. Asner, P. Hart, J. Price, C. K. Lyons, X. Castaneda, Long-term effects of fragmentation and fragment properties on bird species richness in Hawaiian forests. *Biol. Conserv.* **143**, 280–288 (2010). [doi:10.1016/j.biocon.2009.10.009](https://doi.org/10.1016/j.biocon.2009.10.009)
149. R. E. Fowler, thesis, “An Investigation into Bee Assemblage Change Along an Urban-Rural Gradient,” University of Birmingham, 2014.
150. M. Franzén, S. G. Nilsson, How can we preserve and restore species richness of pollinating insects on agricultural land? *Ecography* **31**, 698–708 (2008). [doi:10.1111/j.1600-0587.2008.05110.x](https://doi.org/10.1111/j.1600-0587.2008.05110.x)
151. G. M. Fredriksson, L. S. Danielsen, J. E. Swenson, Impacts of El Niño related drought and forest fires on sun bear fruit resources in lowland dipterocarp forest of East Borneo. *Biodivers. Conserv.* **16**, 1823–1838 (2007). [doi:10.1007/s10531-006-9075-0](https://doi.org/10.1007/s10531-006-9075-0)
152. G. de Brito Freire Jr, P. C. Motta, Effects of experimental fire regimes on the abundance and diversity of cursorial arachnids of Brazilian savannah (cerrado biome). *J. Arachnol.* **39**, 263–272 (2011). [doi:10.1636/CP10-85.1](https://doi.org/10.1636/CP10-85.1)
153. T. L. M. Frizzo, H. L. Vasconcelos, The potential role of scattered trees for ant conservation in an agriculturally dominated Neotropical landscape. *Biotropica* **45**, 644–651 (2013). [doi:10.1111/btp.12045](https://doi.org/10.1111/btp.12045)
154. D. Fukuda, O. B. Tisen, K. Momose, S. Sakai, Bat diversity in the vegetation mosaic around a lowland dipterocarp forest of Borneo. *Raffles Bull. Zool.* **57**, 213–221 (2009).
155. D. Furlani, G. Francesco Ficetola, G. Colombo, M. Ugurlucan, F. De Bernardi, Deforestation and the structure of frog communities in the Humedale Terraba-Sierpe, Costa Rica. *Zoolog. Sci.* **26**, 197–202 (2009). [Medline doi:10.2108/zsj.26.197](https://pubmed.ncbi.nlm.nih.gov/19711111/)
156. R. Gaigher, M. J. Samways, Surface-active arthropods in organic vineyards, integrated vineyards and natural habitat in the Cape Floristic Region. *J. Insect Conserv.* **14**, 595–605 (2010). [doi:10.1007/s10841-010-9286-2](https://doi.org/10.1007/s10841-010-9286-2)
157. K. P. García, J. C. Ortiz Zapata, M. Aguayo, G. D’Elia, Assessing rodent community responses in disturbed environments of the Chilean Patagonia. *Mammalia* **77**, 195–204 (2013). [doi:10.1515/mammalia-2011-0134](https://doi.org/10.1515/mammalia-2011-0134)
158. J. C. García-R, H. Cárdenas-H, F. Castro-H, Relationship between anurans diversity and successional stages of a very humid low montane forest in Valle del Cauca, southwestern of Colombia. *Caldasia* **29**, 363–374 (2007).
159. J. G. Garden, C. A. McAlpine, H. P. Possingham, Multi-scaled habitat considerations for conserving urban biodiversity: Native reptiles and small mammals in Brisbane, Australia. *Landscape Ecol.* **25**, 1013–1028 (2010). [doi:10.1007/s10980-010-9476-z](https://doi.org/10.1007/s10980-010-9476-z)
160. T. A. Gardner, M. I. M. Hernández, J. Barlow, C. A. Peres, Understanding the biodiversity consequences of habitat change: The value of secondary and plantation forests for neotropical dung beetles. *J. Appl. Ecol.* **45**, 883–893 (2008). [doi:10.1111/j.1365-2664.2008.01454.x](https://doi.org/10.1111/j.1365-2664.2008.01454.x)
161. T. A. Gardner, M. A. Ribeiro-Júnior, J. Barlow, T. C. Avila-Pires, M. S. Hoogmoed, C. A. Peres, The value of primary, secondary, and plantation forests for a neotropical

- herpetofauna. *Conserv. Biol.* **21**, 775–787 (2007). [Medline doi:10.1111/j.1523-1739.2007.00659.x](#)
162. A. Garmendia, V. Arroyo-Rodriguez, A. Estrada, E. J. Naranjo, K. E. Stoner, Landscape and patch attributes impacting medium- and large-sized terrestrial mammals in a fragmented rain forest. *J. Trop. Ecol.* **29**, 331–344 (2013). [doi:10.1017/S0266467413000370](#)
163. E. Gaublonme, F. Hendrickx, H. Dhuyvetter, K. Desender, The effects of forest patch size and matrix type on changes in carabid beetle assemblages in an urbanized landscape. *Biol. Conserv.* **141**, 2585–2596 (2008). [doi:10.1016/j.biocon.2008.07.022](#)
164. G. Bao-Ming, L. Zhen-Xing, Z. Dai-Zhen, Z. Hua-Bin, L. Zong-Tang, Z. Chun-Lin, T. Bo-Ping, Communities of soil macrofauna in green spaces of an urbanizing city at east China. *Rev. Chil. Hist. Nat.* **85**, 219–226 (2012). [doi:10.4067/S0716-078X2012000200008](#)
165. B. Gendreau-Berthiaume, D. D. Kneeshaw, B. D. Harvey, Effects of partial cutting and partial disturbance by wind and insects on stand composition, structure and growth in boreal mixedwoods. *Forestry* **85**, 551–565 (2012). [doi:10.1093/forestry/cps051](#)
166. C. Gheler-Costa, C. A. Vettorazzi, R. Pardini, L. M. Verdade, The distribution and abundance of small mammals in agroecosystems of southeastern Brazil. *Mammalia* **76**, 185–191 (2012). [doi:10.1515/mammalia-2011-0109](#)
167. P. Giordani, Assessing the effects of forest management on epiphytic lichens in coppiced forests using different indicators. *Plant Biosyst.* **146**, 628–637 (2012).
168. P. Giordani *et al.*, Land use intensity drives the local variation of lichen diversity in Mediterranean ecosystems sensitive to desertification. *Bibl. Lichenol.* **105**, 139–148 (2010).
169. S. Giordano, S. Sorbo, P. Adamo, A. Basile, V. Spagnuolo, R. C. Cobianchi, Biodiversity and trace element content of epiphytic bryophytes in urban and extraurban sites of southern Italy. *Plant Ecol.* **170**, 1–14 (2004). [doi:10.1023/B:VEGE.0000019025.36121.5d](#)
170. C. Golodets, J. Kigel, M. Sternberg, Recovery of plant species composition and ecosystem function after cessation of grazing in a Mediterranean grassland. *Plant Soil* **329**, 365–378 (2010). [doi:10.1007/s11104-009-0164-1](#)
171. L. G. L. Gomes, V. Oostra, V. Nijman, A. M. Cleef, M. Kappelle, Tolerance of frugivorous birds to habitat disturbance in a tropical cloud forest. *Biol. Conserv.* **141**, 860–871 (2008). [doi:10.1016/j.biocon.2008.01.007](#)
172. M. S. Gottschalk, D. C. De Toni, V. L. S. Valente, P. R. P. Hofmann, Changes in Brazilian *Drosophilidae* (Diptera) assemblages across an urbanisation gradient. *Neotrop. Entomol.* **36**, 848–862 (2007). [Medline doi:10.1590/S1519-566X2007000600005](#)
173. R. K. Gould, L. Pejchar, S. G. Bothwell, B. Brosi, S. Wolny, C. D. Mendenhall, G. Daily, Forest restoration and parasitoid wasp communities in montane Hawai'i. *PLOS ONE* **8**, e59356 (2013). [Medline](#)

174. D. Goulson, O. Lepais, S. O'Connor, J. L. Osborne, R. A. Sanderson, J. Cussans, L. Goffe, B. Darvill, Effects of land use at a landscape scale on bumblebee nest density and survival. *J. Appl. Ecol.* **47**, 1207–1215 (2010). [doi:10.1111/j.1365-2664.2010.01872.x](https://doi.org/10.1111/j.1365-2664.2010.01872.x)
175. D. Goulson, G. C. Lye, B. Darvill, Diet breadth, coexistence and rarity in bumblebees. *Biodivers. Conserv.* **17**, 3269–3288 (2008). [doi:10.1007/s10531-008-9428-y](https://doi.org/10.1007/s10531-008-9428-y)
176. A. D. Gove, J. D. Majer, V. Rico-Gray, Methods for conservation outside of formal reserve systems: The case of ants in the seasonally dry tropics of Veracruz, Mexico. *Biol. Conserv.* **126**, 328–338 (2005). [doi:10.1016/j.biocon.2005.06.008](https://doi.org/10.1016/j.biocon.2005.06.008)
177. L. Granjon, J. M. Duplantier, Guinean biodiversity at the edge: Rodents in forest patches of southern Mali. *Mamm. Biol.* **76**, 583–591 (2011). [doi:10.1016/j.mambio.2011.06.003](https://doi.org/10.1016/j.mambio.2011.06.003)
178. I. Grass, D. G. Berens, F. Peter, N. Farwig, Additive effects of exotic plant abundance and land-use intensity on plant-pollinator interactions. *Oecologia* **173**, 913–923 (2013). [Medline doi:10.1007/s00442-013-2688-6](https://doi.org/10.1007/s00442-013-2688-6)
179. C. L. Gray, E. M. Slade, D. J. Mann, O. T. Lewis, Do riparian reserves support dung beetle biodiversity and ecosystem services in oil palm-dominated tropical landscapes? *Ecol. Evol.* **4**, 1049–1060 (2014). [Medline doi:10.1002/ece3.1003](https://doi.org/10.1002/ece3.1003)
180. W.-B. Gu, Y. Zhen-Rong, H. Dun-Xiao, Carabid community and its fluctuation in farmland of salinity transforming area in the North China Plain: A case study in Quzhou County, Hebei Province. *Biodivers. Sci.* **12**, 262–268 (2004).
181. N. R. Gunawardene, J. D. Majer, J. P. Edirisinghe, Investigating residual effects of selective logging on ant species assemblages in Sinharaja Forest Reserve, Sri Lanka. *For. Ecol. Manage.* **259**, 555–562 (2010). [doi:10.1016/j.foreco.2009.11.012](https://doi.org/10.1016/j.foreco.2009.11.012)
182. A. G. Gutiérrez, J. J. Armesto, J.-C. Aravena, M. Carmona, N. V. Carrasco, D. A. Christie, M.-P. Peña, C. Pérez, A. Huth, Structural and environmental characterization of old-growth temperate rainforests of northern Chiloe Island, Chile: Regional and global relevance. *For. Ecol. Manage.* **258**, 376–388 (2009). [doi:10.1016/j.foreco.2009.03.011](https://doi.org/10.1016/j.foreco.2009.03.011)
183. D. L. Gutierrez-Lamus, Composition and abundance of Anura in two forest types (natural and planted) in the eastern Cordillera of Colombia. *Caldasia* **26**, 245–264 (2004).
184. D. H. Haarmeyer, U. Schmiedel, J. Dengler, B. M. Bosing, How does grazing intensity affect different vegetation types in arid Succulent Karoo, South Africa? Implications for conservation management. *Biol. Conserv.* **143**, 588–596 (2010). [doi:10.1016/j.biocon.2009.11.008](https://doi.org/10.1016/j.biocon.2009.11.008)
185. N. R. Hashim, W. F. Akmal, W. Jusoh, M. Nasir, Ant diversity in a Peninsular Malaysian mangrove forest and oil palm plantation. *Asian Myrmecol.* **3**, 5–8 (2010).
186. S. N. Hassan *et al.*, Human-induced disturbances influence on bird communities of coastal forests in eastern Tanzania. *Br. J. Appl. Sci. Technol.* **3**, 48–64 (2013). [doi:10.9734/BJAST/2014/2200](https://doi.org/10.9734/BJAST/2014/2200)
187. R. G. Hatfield, G. LeBuhn, Patch and landscape factors shape community assemblage of bumble bees, *Bombus* spp. (Hymenoptera: Apidae), in montane meadows. *Biol. Conserv.* **139**, 150–158 (2007). [doi:10.1016/j.biocon.2007.06.019](https://doi.org/10.1016/j.biocon.2007.06.019)

188. J. Hawes, C. da Silva Motta, W. L. Overal, J. Barlow, T. A. Gardner, C. A. Peres, Diversity and composition of Amazonian moths in primary, secondary and plantation forests. *J. Trop. Ecol.* **25**, 281–300 (2009). [doi:10.1017/S0266467409006038](https://doi.org/10.1017/S0266467409006038)
189. A. J. Helden, S. R. Leather, Biodiversity on urban roundabouts - Hemiptera, management and the species-area relationship. *Basic Appl. Ecol.* **5**, 367–377 (2004). [doi:10.1016/j.baae.2004.06.004](https://doi.org/10.1016/j.baae.2004.06.004)
190. L. Hernández, L. Delgado, W. Meier, C. Duran, Empobrecimiento de bosques fragmentados en el norte de la Gran Sabana, Venezuela. *Interciencia* **37**, 891–898 (2012).
191. P. Hietz, Conservation of vascular epiphyte diversity in Mexican coffee plantations. *Conserv. Biol.* **19**, 391–399 (2005). [doi:10.1111/j.1523-1739.2005.00145.x](https://doi.org/10.1111/j.1523-1739.2005.00145.x)
192. D. Higuera, J. H. D. Wolf, Vascular epiphytes in dry oak forests show resilience to anthropogenic disturbance, Cordillera Oriental, Colombia. *Caldasia* **32**, 161–174 (2010).
193. B. Hilje, T. M. Aide, Recovery of amphibian species richness and composition in a chronosequence of secondary forests, northeastern Costa Rica. *Biol. Conserv.* **146**, 170–176 (2012). [doi:10.1016/j.biocon.2011.12.007](https://doi.org/10.1016/j.biocon.2011.12.007)
194. A. Hoffmann, U. Zeller, Influence of variations in land use intensity on species diversity and abundance of small mammals in the Nama Karoo, Namibia. *Belg. J. Zool.* **135**, 91–96 (2005).
195. F. G. Horgan, Invasion and retreat: Shifting assemblages of dung beetles amidst changing agricultural landscapes in central Peru. *Biodivers. Conserv.* **18**, 3519–3541 (2009). [doi:10.1007/s10531-009-9658-7](https://doi.org/10.1007/s10531-009-9658-7)
196. E. Hornung, B. Tothmeresz, T. Magura, F. Vilisics, Changes of isopod assemblages along an urban-suburban-rural gradient in Hungary. *Eur. J. Soil Biol.* **43**, 158–165 (2007). [doi:10.1016/j.ejsobi.2007.01.001](https://doi.org/10.1016/j.ejsobi.2007.01.001)
197. C. Hu, Z. P. Cao, Nematode community structure under compost and chemical fertilizer management practice, in the north China plain. *Exp. Agric.* **44**, 485–496 (2008). [doi:10.1017/S0014479708006716](https://doi.org/10.1017/S0014479708006716)
198. K. Hylander, C. Nilsson, T. Gothner, Effects of buffer-strip retention and clearcutting on land snails in boreal riparian forests. *Conserv. Biol.* **18**, 1052–1062 (2004). [doi:10.1111/j.1523-1739.2004.00199.x](https://doi.org/10.1111/j.1523-1739.2004.00199.x)
199. K. Hylander, H. Weibull, Do time-lagged extinctions and colonizations change the interpretation of buffer strip effectiveness? A study of riparian bryophytes in the first decade after logging. *J. Appl. Ecol.* **49**, 1316–1324 (2012). [doi:10.1111/j.1365-2664.2012.02218.x](https://doi.org/10.1111/j.1365-2664.2012.02218.x)
200. K. Hylander, S. Nemomissa, Complementary roles of home gardens and exotic tree plantations as alternative habitats for plants of the Ethiopian montane rainforest. *Conserv. Biol.* **23**, 400–409 (2009). [Medline doi:10.1111/j.1523-1739.2008.01097.x](https://doi.org/10.1111/j.1523-1739.2008.01097.x)
201. R. A. Ims, J. A. Henden, Collapse of an arctic bird community resulting from ungulate-induced loss of erect shrubs. *Biol. Conserv.* **149**, 2–5 (2012). [doi:10.1016/j.biocon.2012.02.008](https://doi.org/10.1016/j.biocon.2012.02.008)

202. P. J. Isaacs-Cubides, J. N. Urbina-Cardona, Anthropogenic disturbance and edge effects on anuran assemblages inhabiting cloud forest fragments in Colombia. *Nat. Conserv.* **9**, 39–46 (2011). [doi:10.4322/natcon.2011.004](https://doi.org/10.4322/natcon.2011.004)
203. M. Ishitani, D. J. Kotze, J. Niemela, Changes in carabid beetle assemblages across an urban-rural gradient in Japan. *Ecography* **26**, 481–489 (2003). [doi:10.1034/j.1600-0587.2003.03436.x](https://doi.org/10.1034/j.1600-0587.2003.03436.x)
204. C. T. Jacobs, C. H. Scholtz, F. Escobar, A. L. V. Davis, How might intensification of farming influence dung beetle diversity (Coleoptera: Scarabaeidae) in Maputo Special Reserve (Mozambique)? *J. Insect Conserv.* **14**, 389–399 (2010). [doi:10.1007/s10841-010-9270-x](https://doi.org/10.1007/s10841-010-9270-x)
205. G. Jalilova, J. de Groot, H. Vacik, Evaluating the effects of habitats on birds in the walnut fruit forests: A case study from Kyrgyzstan. *Biodiversity* **14**, 97–110 (2013). [doi:10.1080/14888386.2013.789399](https://doi.org/10.1080/14888386.2013.789399)
206. M. F. Johnson, A. Gómez, M. Pinedo-Vasquez, Land use and mosquito diversity in the Peruvian Amazon. *J. Med. Entomol.* **45**, 1023–1030 (2008). [Medline](https://pubmed.ncbi.nlm.nih.gov/25852008/) [doi:10.1603/0022-2585\(2008\)45\[1023:LUAMDI\]2.0.CO;2](https://doi.org/10.1603/0022-2585(2008)45[1023:LUAMDI]2.0.CO;2)
207. V. Jolli, M. K. Pandit, Monitoring pheasants (*Phasianidae*) in the western Himalayas to measure the impact of hydro-electric projects. *Ring* **33**, 37–46 (2011). [doi:10.2478/v10050-011-0003-7](https://doi.org/10.2478/v10050-011-0003-7)
208. M. Jonsell, Old park trees as habitat for saproxylic beetle species. *Biodivers. Conserv.* **21**, 619–642 (2012). [doi:10.1007/s10531-011-0203-0](https://doi.org/10.1007/s10531-011-0203-0)
209. L. Joubert, K. J. Esler, S. D. J. Privett, The effect of ploughing and augmenting natural vegetation with commercial fynbos species on the biodiversity of Overberg Sandstone fynbos on the Agulhas Plain, South Africa. *S. Afr. J. Bot.* **75**, 526–531 (2009). [doi:10.1016/j.sajb.2009.05.002](https://doi.org/10.1016/j.sajb.2009.05.002)
210. H. E. Julier, T. H. Roulston, Wild bee abundance and pollination service in cultivated pumpkins: Farm management, nesting behavior and landscape effects. *J. Econ. Entomol.* **102**, 563–573 (2009). [Medline](https://pubmed.ncbi.nlm.nih.gov/20210214/) [doi:10.1603/029.102.0214](https://doi.org/10.1603/029.102.0214)
211. T. S. Jung, T. Powell, Spatial distribution of meadow jumping mice (*Zapus hudsonius*) in logged boreal forest of northwestern Canada. *Mamm. Biol.* **76**, 678–682 (2011). [doi:10.1016/j.mambio.2011.08.002](https://doi.org/10.1016/j.mambio.2011.08.002)
212. V. Kapoor, Effects of rainforest fragmentation and shade-coffee plantations on spider communities in the Western Ghats, India. *J. Insect Conserv.* **12**, 53–68 (2008). [doi:10.1007/s10841-006-9062-5](https://doi.org/10.1007/s10841-006-9062-5)
213. H. Kappes, L. Katschner, C. Nowak, Urban summer heat load: Meteorological data as a proxy for metropolitan biodiversity. *Meteorol. Zeitschrift* **21**, 525–528 (2012). [doi:10.1127/0941-2948/2012/0361](https://doi.org/10.1127/0941-2948/2012/0361)
214. V. Kati, K. Zografou, E. Tzirkalli, T. Chitos, L. Willemse, Butterfly and grasshopper diversity patterns in humid Mediterranean grasslands: The roles of disturbance and environmental factors. *J. Insect Conserv.* **16**, 807–818 (2012). [doi:10.1007/s10841-012-9467-2](https://doi.org/10.1007/s10841-012-9467-2)

215. E. Katovai, A. L. Burley, M. M. Mayfield, Understory plant species and functional diversity in the degraded wet tropical forests of Kolombangara Island, Solomon Islands. *Biol. Conserv.* **145**, 214–224 (2012). [doi:10.1016/j.biocon.2011.11.008](https://doi.org/10.1016/j.biocon.2011.11.008)
216. F. Kazerani, K. S. I. Grichanov, Diversity of the genus *Dolichopus* Latreille in three different habitats of East Azerbaijan Province, with new records for Iran. *Arxius de Miscellània Zoològica* **11**, 134–152 (2013).
217. M. Kessler, P. J. A. Keßler, S. R. Gradstein, K. Bach, M. Schmull, R. Pitopang, Tree diversity in primary forest and different land use systems in Central Sulawesi, Indonesia. *Biodivers. Conserv.* **14**, 547–560 (2005). [doi:10.1007/s10531-004-3914-7](https://doi.org/10.1007/s10531-004-3914-7)
218. M. Kessler, S. Abrahamczyk, M. Bos, D. Buchori, D. D. Putra, S. R. Gradstein, P. Höhn, J. Kluge, F. Orend, R. Pitopang, S. Saleh, C. H. Schulze, S. G. Sporn, I. Steffan-Dewenter, S. S. Tjitrosoedirdjo, T. Tschardtke, Alpha and beta diversity of plants and animals along a tropical land-use gradient. *Ecol. Appl.* **19**, 2142–2156 (2009). [Medline](https://pubmed.ncbi.nlm.nih.gov/1910741/) [doi:10.1890/08-1074.1](https://doi.org/10.1890/08-1074.1)
219. A. M. Kittle, A. C. Watson, P. H. Chanaka Kumara, H. K. Nimalka Sanjeevani, Status and distribution of the leopard in the central hills of Sri Lanka. *Cat News* **56**, 28–31 (2012).
220. F. Kohler, J. Verhulst, R. van Klink, D. Kleijn, At what spatial scale do high-quality habitats enhance the diversity of forbs and pollinators in intensively farmed landscapes? *J. Appl. Ecol.* **45**, 753–762 (2008). [doi:10.1111/j.1365-2664.2007.01394.x](https://doi.org/10.1111/j.1365-2664.2007.01394.x)
221. M. Koivula, V. Hyrylainen, E. Soininen, Carabid beetles (Coleoptera: Carabidae) at forest-farmland edges in southern Finland. *J. Insect Conserv.* **8**, 297–309 (2004). [doi:10.1007/s10841-004-0296-9](https://doi.org/10.1007/s10841-004-0296-9)
222. A. Kolb, M. Diekmann, Effects of environment, habitat configuration and forest continuity on the distribution of forest plant species. *J. Veg. Sci.* **15**, 199–208 (2004). [doi:10.1111/j.1654-1103.2004.tb02255.x](https://doi.org/10.1111/j.1654-1103.2004.tb02255.x)
223. Á. Kőrösi, P. Batáry, A. Orosz, D. Rédei, A. Báldi, Effects of grazing, vegetation structure and landscape complexity on grassland leafhoppers (Hemiptera: Auchenorrhyncha) and true bugs (Hemiptera: Heteroptera) in Hungary. *Insect Conserv. Divers.* **5**, 57–66 (2012). [doi:10.1111/j.1752-4598.2011.00153.x](https://doi.org/10.1111/j.1752-4598.2011.00153.x)
224. J. Krauss, A. M. Klein, I. Steffan-Dewenter, T. Tschardtke, Effects of habitat area, isolation, and landscape diversity on plant species richness of calcareous grasslands. *Biodivers. Conserv.* **13**, 1427–1439 (2004). [doi:10.1023/B:BIOC.0000021323.18165.58](https://doi.org/10.1023/B:BIOC.0000021323.18165.58)
225. J. Krauss, I. Steffan-Dewenter, T. Tschardtke, How does landscape context contribute to effects of habitat fragmentation on diversity and population density of butterflies? *J. Biogeogr.* **30**, 889–900 (2003). [doi:10.1046/j.1365-2699.2003.00878.x](https://doi.org/10.1046/j.1365-2699.2003.00878.x)
226. J. Krauss, R. Bommarco, M. Guardiola, R. K. Heikkinen, A. Helm, M. Kuussaari, R. Lindborg, E. Ockinger, M. Pärtel, J. Pino, J. Pöyry, K. M. Raatikainen, A. Sang, C. Stefanescu, T. Teder, M. Zobel, I. Steffan-Dewenter, Habitat fragmentation causes immediate and time-delayed biodiversity loss at different trophic levels. *Ecol. Lett.* **13**, 597–605 (2010). [Medline](https://pubmed.ncbi.nlm.nih.gov/20101457/) [doi:10.1111/j.1461-0248.2010.01457.x](https://doi.org/10.1111/j.1461-0248.2010.01457.x)

227. R. Kumar, G. Shahabuddin, Effects of biomass extraction on vegetation structure, diversity and composition of forests in Sariska Tiger Reserve, India. *Environ. Conserv.* **32**, 248–259 (2005). [doi:10.1017/S0376892905002316](https://doi.org/10.1017/S0376892905002316)
228. D. J. Kurz, A. J. Nowakowski, M. W. Tingley, M. A. Donnelly, D. S. Wilcove, Forest-land use complementarity modifies community structure of a tropical herpetofauna. *Biol. Conserv.* **170**, 246–255 (2014). [doi:10.1016/j.biocon.2013.12.027](https://doi.org/10.1016/j.biocon.2013.12.027)
229. A. S. Kutt, E. P. Vanderduys, P. O'Reagain, Spatial and temporal effects of grazing management and rainfall on the vertebrate fauna of a tropical savanna. *Rangeland J.* **34**, 173–182 (2012). [doi:10.1071/RJ11049](https://doi.org/10.1071/RJ11049)
230. A. S. Kutt, J. C. Z. Woinarski, The effects of grazing and fire on vegetation and the vertebrate assemblage in a tropical savanna woodland in north-eastern Australia. *J. Trop. Ecol.* **23**, 95–106 (2007). [doi:10.1017/S0266467406003579](https://doi.org/10.1017/S0266467406003579)
231. T. Lachat, S. Attignon, J. Djego, G. Goergen, P. Nagel, B. Sinsin, R. Peveling, Arthropod diversity in Lama forest reserve (South Benin), a mosaic of natural, degraded and plantation forests. *Biodivers. Conserv.* **15**, 3–23 (2006). [doi:10.1007/s10531-004-1234-6](https://doi.org/10.1007/s10531-004-1234-6)
232. M. V. Lantschner, V. Rusch, J. P. Hayes, Habitat use by carnivores at different spatial scales in a plantation forest landscape in Patagonia, Argentina. *For. Ecol. Manage.* **269**, 271–278 (2012). [doi:10.1016/j.foreco.2011.12.045](https://doi.org/10.1016/j.foreco.2011.12.045)
233. M. V. Lantschner, V. Rusch, C. Peyrou, Bird assemblages in pine plantations replacing native ecosystems in NW Patagonia. *Biodivers. Conserv.* **17**, 969–989 (2008). [doi:10.1007/s10531-007-9243-x](https://doi.org/10.1007/s10531-007-9243-x)
234. S. C. Latta, B. A. Tinoco, P. X. Astudillo, C. H. Graham, Patterns and magnitude of temporal change in avian communities in the Ecuadorian Andes. *Condor* **113**, 24–40 (2011). [doi:10.1525/cond.2011.090252](https://doi.org/10.1525/cond.2011.090252)
235. J.-P. L  gar  , C. H  bert, J.-C. Ruel, Alternative silvicultural practices in irregular boreal forests: Response of beetle assemblages. *Silva Fenn.* **45**, 937–956 (2011). [doi:10.14214/sf.79](https://doi.org/10.14214/sf.79)
236. V. Lehouck, T. Spanhove, L. Colson, A. Adringa-Davis, N. J. Cordeiro, L. Lens, Habitat disturbance reduces seed dispersal of a forest interior tree in a fragmented African cloud forest. *Oikos* **118**, 1023–1034 (2009). [doi:10.1111/j.1600-0706.2009.17300.x](https://doi.org/10.1111/j.1600-0706.2009.17300.x)
237. P. E. Lentini, T. G. Martin, P. Gibbons, J. Fischer, S. A. Cunningham, Supporting wild pollinators in a temperate agricultural landscape: Maintaining mosaics of natural features and production. *Biol. Conserv.* **149**, 84–92 (2012). [doi:10.1016/j.biocon.2012.02.004](https://doi.org/10.1016/j.biocon.2012.02.004)
238. S. G. Letcher, R. L. Chazdon, Rapid recovery of biomass, species richness, and species composition in a forest chronosequence in Northeastern Costa Rica. *Biotropica* **41**, 608–617 (2009). [doi:10.1111/j.1744-7429.2009.00517.x](https://doi.org/10.1111/j.1744-7429.2009.00517.x)
239. S. N. Li, F. S. Zou, Q. Zhang, F. H. Sheldon, Species richness and guild composition in rubber plantations compared to secondary forest on Hainan Island, China. *Agrofor. Syst.* **87**, 1117–1128 (2013). [doi:10.1007/s10457-013-9624-y](https://doi.org/10.1007/s10457-013-9624-y)

240. L. H. Liow, N. S. Sodhi, T. Elmqvist, Bee diversity along a disturbance gradient in tropical lowland forests of south-east Asia. *J. Appl. Ecol.* **38**, 180–192 (2001). [doi:10.1046/j.1365-2664.2001.00582.x](https://doi.org/10.1046/j.1365-2664.2001.00582.x)
241. S. A. Litchwark, thesis, “Honeybee Declines in a Changing Landscape: Interactive Effects of Honeybee Declines and Land-Use Intensification on Pollinator Communities,” University of Canterbury, 2013.
242. N. A. Littlewood, Grazing impacts on moth diversity and abundance on a Scottish upland estate. *Insect Conserv. Divers.* **1**, 151–160 (2008). [doi:10.1111/j.1752-4598.2008.00021.x](https://doi.org/10.1111/j.1752-4598.2008.00021.x)
243. N. A. Littlewood, R. J. Pakeman, G. Pozsgai, Grazing impacts on Auchenorrhyncha diversity and abundance on a Scottish upland estate. *Insect Conserv. Divers.* **5**, 67–74 (2012). [doi:10.1111/j.1752-4598.2011.00135.x](https://doi.org/10.1111/j.1752-4598.2011.00135.x)
244. Y. H. Liu, J. C. Axmacher, C. L. Wang, L. T. Li, Z. R. Yu, Ground beetle (Coleoptera: Carabidae) assemblages of restored semi-natural habitats and intensively cultivated fields in northern China. *Restor. Ecol.* **20**, 234–239 (2012). [doi:10.1111/j.1526-100X.2010.00755.x](https://doi.org/10.1111/j.1526-100X.2010.00755.x)
245. N. F. Lo-Man-Hung, T. A. Gardner, M. A. Ribeiro-Júnior, J. Barlow, A. B. Bonaldo, The value of primary, secondary, and plantation forests for Neotropical epigeic arachnids. *J. Arachnol.* **36**, 394–401 (2008). [doi:10.1636/CT07-136.1](https://doi.org/10.1636/CT07-136.1)
246. N. F. Lo-Man-Hung, R. Marichal, D. F. Candiani, L. S. Carvalho, R. P. Indicatti, A. B. Bonaldo, D. H. R. Cobo, A. M. Feijoo, S. Tselouiko, C. Praxedes, G. Brown, E. Velasquez, T. Decaëns, J. Oszwald, M. Martins, P. Lavelle, Impact of different land management on soil spiders (Arachnida: Araneae) in two Amazonian areas of Brazil and Colombia. *J. Arachnol.* **39**, 296–302 (2011). [doi:10.1636/CP10-89.1](https://doi.org/10.1636/CP10-89.1)
247. C. A. López-Quintero, G. Straatsma, A. E. Franco-Molano, T. Boekhout, Macrofungal diversity in Colombian Amazon forests varies with regions and regimes of disturbance. *Biodivers. Conserv.* **21**, 2221–2243 (2012). [doi:10.1007/s10531-012-0280-8](https://doi.org/10.1007/s10531-012-0280-8)
248. M. Louhaichi, A. K. Salkini, S. L. Petersen, Effect of small ruminant grazing on the plant community characteristics of semiarid Mediterranean ecosystems. *Int. J. Agric. Biol.* **11**, 681–689 (2009).
249. M. E. Lucas-Borja, F. Bastida, J. L. Moreno, C. Nicolás, M. Andres, F. R. López, A. Del Cerro, The effects of human trampling on the microbiological properties of soil and vegetation in Mediterranean mountain areas. *Land Degrad. Dev.* **22**, 383–394 (2011). [doi:10.1002/ldr.1014](https://doi.org/10.1002/ldr.1014)
250. V. H. Luja, S. Herrando-Perez, D. Gonzalez-Solis, L. Luiselli, Secondary rain forests are not havens for reptile species in tropical Mexico. *Biotropica* **40**, 747–757 (2008). [doi:10.1111/j.1744-7429.2008.00439.x](https://doi.org/10.1111/j.1744-7429.2008.00439.x)
251. R. Macip-Ríos, A. Muñoz-Alonso, Diversidad de lagartijas en cafetales y bosque primario en el Soconusco chiapaneco. *Rev. Mex. Biodivers.* **79**, 185–195 (2008).

252. M. C. G. MacSwiney, P. L. Vilchis, F. M. Clarke, P. A. Racey, The importance of cenotes in conserving bat assemblages in the Yucatan, Mexico. *Biol. Conserv.* **136**, 499–509 (2007). [doi:10.1016/j.biocon.2006.12.021](https://doi.org/10.1016/j.biocon.2006.12.021)
253. K. Maeto, S. Sato, Impacts of forestry on ant species richness and composition in warm-temperate forests of Japan. *For. Ecol. Manage.* **187**, 213–223 (2004). [doi:10.1016/S0378-1127\(03\)00333-5](https://doi.org/10.1016/S0378-1127(03)00333-5)
254. T. Magura, R. Horvath, B. Tothmeresz, Effects of urbanization on ground-dwelling spiders in forest patches, in Hungary. *Landscape Ecol.* **25**, 621–629 (2010). [doi:10.1007/s10980-009-9445-6](https://doi.org/10.1007/s10980-009-9445-6)
255. N. A. D. Mallari, N. J. Collar, D. C. Lee, P. J. K. McGowan, R. Wilkinson, S. J. Marsden, Population densities of understorey birds across a habitat gradient in Palawan, Philippines: Implications for conservation. *Oryx* **45**, 234–242 (2011). [doi:10.1017/S0030605310001031](https://doi.org/10.1017/S0030605310001031)
256. L. Malone, J. Aulsford, B. Howlett, C. Scott-Dupree, N. Bardol, B. Donovan, Observations on bee species visiting white clover in New Zealand pastures. *J. Apic. Res.* **49**, 284–286 (2010). [doi:10.3896/IBRA.1.49.3.09](https://doi.org/10.3896/IBRA.1.49.3.09)
257. P. K. Malonza, M. Veith, Amphibian community along elevational and habitat disturbance gradients in the Taita Hills, Kenya. *Herpetotropicos* **7**, 7–16 (2012).
258. J. Malumbres-Olarte, B. I. P. Barratt, C. J. Vink, A. M. Paterson, R. H. Cruickshank, C. M. Ferguson, D. M. Barton, Big and aerial invaders: Dominance of exotic spiders in burned New Zealand tussock grasslands. *Biol. Invasions* **16**, 2311–2322 (2014). [doi:10.1007/s10530-014-0666-5](https://doi.org/10.1007/s10530-014-0666-5)
259. E. Marín-Spiotta, W. L. Silver, R. Ostertag, Long-term patterns in tropical reforestation: Plant community composition and aboveground biomass accumulation. *Ecol. Appl.* **17**, 828–839 (2007). [Medline doi:10.1890/06-1268](https://doi.org/10.1890/06-1268)
260. C. Marsh, thesis, “Community-Level Modelling of α -, β - and γ -Diversity of Multiple Taxa: Birds and Butterflies on Anjouan, Comoros Islands,” Imperial College London, 2005.
261. C. J. Marsh, O. T. Lewis, I. Said, R. M. Ewers, Community-level diversity modelling of birds and butterflies on Anjouan, Comoro Islands. *Biol. Conserv.* **143**, 1364–1374 (2010). [doi:10.1016/j.biocon.2010.03.010](https://doi.org/10.1016/j.biocon.2010.03.010)
262. E. J. P. Marshall, T. M. West, D. Kleijn, Impacts of an agri-environment field margin prescription on the flora and fauna of arable farmland in different landscapes. *Agric. Ecosyst. Environ.* **113**, 36–44 (2006). [doi:10.1016/j.agee.2005.08.036](https://doi.org/10.1016/j.agee.2005.08.036)
263. P. S. Martin, C. Gheler-Costa, P. C. Lopes, L. M. Rosalino, L. M. Verdade, Terrestrial non-volant small mammals in agro-silvicultural landscapes of Southeastern Brazil. *For. Ecol. Manage.* **282**, 185–195 (2012). [doi:10.1016/j.foreco.2012.07.002](https://doi.org/10.1016/j.foreco.2012.07.002)
264. T. Matsumoto, T. Itioka, S. Yamane, K. Momose, Traditional land use associated with swidden agriculture changes encounter rates of the top predator, the army ant, in Southeast Asian tropical rain forests. *Biodivers. Conserv.* **18**, 3139–3151 (2009). [doi:10.1007/s10531-009-9632-4](https://doi.org/10.1007/s10531-009-9632-4)

265. M. M. Mayfield, D. Ackerly, G. C. Daily, The diversity and conservation of plant reproductive and dispersal functional traits in human-dominated tropical landscapes. *J. Ecol.* **94**, 522–536 (2006). [doi:10.1111/j.1365-2745.2006.01108.x](https://doi.org/10.1111/j.1365-2745.2006.01108.x)
266. J. L. McCarthy, K. P. McCarthy, T. K. Fuller, T. M. McCarthy, Assessing variation in wildlife biodiversity in the Tien Shan Mountains of Kyrgyzstan using ancillary camera-trap photos. *Mt. Res. Dev.* **30**, 295–301 (2010). [doi:10.1659/MRD-JOURNAL-D-09-00080.1](https://doi.org/10.1659/MRD-JOURNAL-D-09-00080.1)
267. Q. S. McFrederick, G. LeBuhn, Are urban parks refuges for bumble bees *Bombus* spp. (Hymenoptera: Apidae)? *Biol. Conserv.* **129**, 372–382 (2006). [doi:10.1016/j.biocon.2005.11.004](https://doi.org/10.1016/j.biocon.2005.11.004)
268. S. McNamara, P. D. Erskine, D. Lamb, L. Chantalangsy, S. Boyle, Primary tree species diversity in secondary fallow forests of Laos. *For. Ecol. Manage.* **281**, 93–99 (2012). [doi:10.1016/j.foreco.2012.06.004](https://doi.org/10.1016/j.foreco.2012.06.004)
269. W. J. McShea, C. Stewart, L. Peterson, P. Erb, R. Stuebing, B. Gimán, The importance of secondary forest blocks for terrestrial mammals within an *Acacia*/secondary forest matrix in Sarawak, Malaysia. *Biol. Conserv.* **142**, 3108–3119 (2009). [doi:10.1016/j.biocon.2009.08.009](https://doi.org/10.1016/j.biocon.2009.08.009)
270. S. S. Meijer, R. J. Whittaker, P. A. V. Borges, The effects of land-use change on arthropod richness and abundance on Santa Maria Island (Azores): Unmanaged plantations favour endemic beetles. *J. Insect Conserv.* **15**, 505–522 (2011). [doi:10.1007/s10841-010-9330-2](https://doi.org/10.1007/s10841-010-9330-2)
271. B. Meyer, V. Gaebele, I. D. Steffan-Dewenter, Patch size and landscape effects on pollinators and seed set of the horseshoe vetch, *hippocrepis comosa*, in an agricultural landscape of central Europe. *Entomol. Gen.* **30**, 173–185 (2007).
272. B. Meyer, F. Jauker, I. Steffan-Dewenter, Contrasting resource-dependent responses of hoverfly richness and density to landscape structure. *Basic Appl. Ecol.* **10**, 178–186 (2009). [doi:10.1016/j.baae.2008.01.001](https://doi.org/10.1016/j.baae.2008.01.001)
273. E. Micó, A. Garcia-Lopez, H. Brustel, A. Padilla, E. Galante, Explaining the saproxylic beetle diversity of a protected Mediterranean area. *Biodivers. Conserv.* **22**, 889–904 (2013). [doi:10.1007/s10531-013-0456-x](https://doi.org/10.1007/s10531-013-0456-x)
274. J. C. Milder, F. A. J. DeCLERCK, A. Sanfiorenzo, D. M. Sánchez, D. Tobar, B. Zuckerberg, Effects of farm and landscape management on bird and butterfly conservation in western Honduras. *Ecosphere* **1**, art2 (2010). [doi:10.1890/ES10-00003.1](https://doi.org/10.1890/ES10-00003.1)
275. M. V. Miranda, N. Politi, L. O. Rivera, Unexpected changes in the bird assemblage in areas under selective logging in piedmont forest in northwestern Argentina. *Ornitol. Neotrop.* **21**, 323–337 (2010).
276. M. L. Moir, K. E. C. Brennan, J. M. Koch, J. D. Majer, M. J. Fletcher, Restoration of a forest ecosystem: The effects of vegetation and dispersal capabilities on the reassembly of plant-dwelling arthropods. *For. Ecol. Manage.* **217**, 294–306 (2005). [doi:10.1016/j.foreco.2005.06.012](https://doi.org/10.1016/j.foreco.2005.06.012)

277. D. Moreno-Mateos, J. M. R. Benayas, L. Pérez-Camacho, E. Montaña, S. Rebollo, L. Cayuela, Effects of land use on nocturnal birds in a Mediterranean agricultural landscape. *Acta Ornithol.* **46**, 173–182 (2011). [doi:10.3161/000164511X625946](https://doi.org/10.3161/000164511X625946)
278. M. N. Muchane *et al.*, Land use practices and their implications on soil macro-fauna in Maasai Mara ecosystem. *Int. J. Biodivers. Conserv.* **4**, 500–514 (2012).
279. S. Mudri-Stojnic, A. Andric, Z. Jozan, A. Vujic, Pollinator diversity (Hymenoptera and Diptera) in semi-natural habitats in Serbia during summer. *Arch. Biol. Sci.* **64**, 777–786 (2012). [doi:10.2298/ABS1202777S](https://doi.org/10.2298/ABS1202777S)
280. F. B. Munyekenye, E. M. Mwangi, N. N. Gichuki, Bird species richness and abundance in different forest types at Kakamega Forest, western Kenya. *Ostrich* **79**, 37–42 (2008). [doi:10.2989/OSTRICH.2008.79.1.4.361](https://doi.org/10.2989/OSTRICH.2008.79.1.4.361)
281. R. Naidoo, Species richness and community composition of songbirds in a tropical forest-agricultural landscape. *Anim. Conserv.* **7**, 93–105 (2004). [doi:10.1017/S1367943003001185](https://doi.org/10.1017/S1367943003001185)
282. A. Naithani, D. Bhatt, Bird community structure in natural and urbanized habitats along an altitudinal gradient in Pauri district (Garhwal Himalaya) of Uttarakhand state, India. *Biologia* **67**, 800–808 (2012). [doi:10.2478/s11756-012-0068-z](https://doi.org/10.2478/s11756-012-0068-z)
283. A. Nakamura, H. Proctor, C. P. Catterall, Using soil and litter arthropods to assess the state of rainforest restoration. *Ecol. Manage. Restor.* **4**, S20–S28 (2003). [doi:10.1046/j.1442-8903.4.s.3.x](https://doi.org/10.1046/j.1442-8903.4.s.3.x)
284. Y. Nakashima, E. Inoue, E. F. Akomo-Okoue, Population density and habitat preferences of forest duikers in Moukalaba-Doudou National Park, Gabon. *Afr. Zool.* **48**, 395–399 (2013). [doi:10.3377/004.048.0212](https://doi.org/10.3377/004.048.0212)
285. S. Naoe, S. Sakai, T. Masaki, Effect of forest shape on habitat selection of birds in a plantation-dominant landscape across seasons: Comparison between continuous and strip forests. *J. For. Res.* **17**, 219–223 (2012). [doi:10.1007/s10310-011-0296-z](https://doi.org/10.1007/s10310-011-0296-z)
286. D. Navarrete, G. Halffter, Dung beetle (Coleoptera: Scarabaeidae: Scarabaeinae) diversity in continuous forest, forest fragments and cattle pastures in a landscape of Chiapas, Mexico: the effects of anthropogenic changes. *Biodivers. Conserv.* **17**, 2869–2898 (2008). [doi:10.1007/s10531-008-9402-8](https://doi.org/10.1007/s10531-008-9402-8)
287. I. L. Navarro, A. K. Roman, F. H. Gomez, H. A. Perez, Seasonal variation in dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae) from Serrania de Coraza, Sucre (Colombia). *Rev. Colomb. Cienc. Anim.* **3**, 102–110 (2011).
288. P. K. Ndong'ang'a, J. B. M. Njoroge, M. Githiru, Vegetation composition and structure influences bird species community assemblages in the highland agricultural landscape of Nyandarua, Kenya. *Ostrich* **84**, 171–179 (2013). [doi:10.2989/00306525.2013.860929](https://doi.org/10.2989/00306525.2013.860929)
289. E. L. Neuschulz, A. Botzat, N. Farwig, Effects of forest modification on bird community composition and seed removal in a heterogeneous landscape in South Africa. *Oikos* **120**, 1371–1379 (2011). [doi:10.1111/j.1600-0706.2011.19097.x](https://doi.org/10.1111/j.1600-0706.2011.19097.x)
290. V. Nicolas, P. Barriere, A. Tapiero, M. Colyn, Shrew species diversity and abundance in Ziama Biosphere Reserve, Guinea: Comparison among primary forest, degraded forest

- and restoration plots. *Biodivers. Conserv.* **18**, 2043–2061 (2009). [doi:10.1007/s10531-008-9572-4](https://doi.org/10.1007/s10531-008-9572-4)
291. A. Nielsen *et al.*, Assessing bee species richness in two Mediterranean communities: Importance of habitat type and sampling techniques. *Ecol. Res.* **26**, 969–983 (2011).
 292. N. Noreika, D. J. Kotze, Forest edge contrasts have a predictable effect on the spatial distribution of carabid beetles in urban forests. *J. Insect Conserv.* **16**, 867–881 (2012). [doi:10.1007/s10841-012-9474-3](https://doi.org/10.1007/s10841-012-9474-3)
 293. N. Noreika, New records of rare species of Coleoptera found in Ukmergė district in 2004–2005. *New Rare Lith. Insect Species* **21**, 68–71 (2009).
 294. O. Norfolk, M. Abdel-Dayem, F. Gilbert, Rainwater harvesting and arthropod biodiversity within an arid agro-ecosystem. *Agric. Ecosyst. Environ.* **162**, 8–14 (2012). [doi:10.1016/j.agee.2012.08.007](https://doi.org/10.1016/j.agee.2012.08.007)
 295. O. Norfolk, M. P. Eichhorn, F. Gilbert, Traditional agricultural gardens conserve wild plants and functional richness in arid South Sinai. *Basic Appl. Ecol.* **14**, 659–669 (2013). [doi:10.1016/j.baae.2013.10.004](https://doi.org/10.1016/j.baae.2013.10.004)
 296. J. A. Noriega, E. Realpe, G. Fagua, Diversidad de escarabajos coprofagos (Coleoptera: Scarabaeidae) en un bosque de galeria con tres estadios de alteracion. *Univ. Sci.* **12**, 51–63 (2007).
 297. J. A. Noriega, J. M. Palacio, J. D. Monroy-G, E. Valencia, Estructura de un ensamblaje de escarabajos coprofagos (Coleoptera: Scarabaeinae) en tres sitios con diferente uso del suelo en Antioquia, Colombia. *Actual. Biol. (Paris)* **34**, 43–54 (2012).
 298. D. A. Norton, P. R. Espie, W. Murray, J. Murray, Influence of pastoral management on plant biodiversity in a depleted short tussock grassland, Mackenzie Basin. *N. Z. J. Ecol.* **30**, 335–344 (2006).
 299. N. M. Nöske, N. Hilt, F. A. Werner, G. Brehm, K. Fiedler, H. J. M. Sipman, S. R. Gradstein, Disturbance effects on diversity of epiphytes and moths in a montane forest in Ecuador. *Basic Appl. Ecol.* **9**, 4–12 (2008). [doi:10.1016/j.baae.2007.06.014](https://doi.org/10.1016/j.baae.2007.06.014)
 300. C. Numa, J. R. Verdu, C. Rueda, E. Galante, Comparing dung beetle species assemblages between protected areas and adjacent pasturelands in a Mediterranean savanna landscape. *Rangeland Ecol. Manag.* **65**, 137–143 (2012). [doi:10.2111/REM-D-10-00050.1](https://doi.org/10.2111/REM-D-10-00050.1)
 301. T. G. O’Connor, Influence of land use on plant community composition and diversity in Highland Sourveld grassland in the southern Drakensberg, South Africa. *J. Appl. Ecol.* **42**, 975–988 (2005). [doi:10.1111/j.1365-2664.2005.01065.x](https://doi.org/10.1111/j.1365-2664.2005.01065.x)
 302. N. O’Dea, R. J. Whittaker, How resilient are Andean montane forest bird communities to habitat degradation? *Biodivers. Conserv.* **16**, 1131–1159 (2007). [doi:10.1007/s10531-006-9095-9](https://doi.org/10.1007/s10531-006-9095-9)
 303. S. Oertli, A. Muller, S. Dorn, Ecological and seasonal patterns in the diversity of a species-rich bee assemblage (Hymenoptera: Apoidea: Apiformes). *Eur. J. Entomol.* **102**, 53–63 (2005). [doi:10.14411/eje.2005.008](https://doi.org/10.14411/eje.2005.008)

304. P. J. O'Farrell, J. S. Donaldson, M. T. Hoffman, A. D. Mader, Small mammal diversity and density on the Bokkeveld escarpment, South Africa - implications for conservation and livestock predation. *Afr. Zool.* **43**, 117–124 (2008). [doi:10.3377/1562-7020\(2008\)43\[117:SMDADO\]2.0.CO;2](https://doi.org/10.3377/1562-7020(2008)43[117:SMDADO]2.0.CO;2)
305. C. Ofori-Boateng, W. Oduro, A. Hillers, K. Norris, S. K. Oppong, G. B. Adum, M.-O. Rödel, Differences in the effects of selective logging on amphibian assemblages in three West African forest types. *Biotropica* **45**, 94–101 (2013). [doi:10.1111/j.1744-7429.2012.00887.x](https://doi.org/10.1111/j.1744-7429.2012.00887.x)
306. C. Oke, Land snail diversity in post extraction secondary forest reserves in Edo State, Nigeria. *Afr. J. Ecol.* **51**, 244–254 (2013). [doi:10.1111/aje.12029](https://doi.org/10.1111/aje.12029)
307. O. C. Oke, J. U. Chokor, The effect of land use on snail species richness and diversity in the tropical rainforest of south-western Nigeria. *African Sci.* **10**, 95–108 (2009).
308. D. E. Oliveira, T. F. Carrijo, D. Brandão, Species composition of termites (Isoptera) in different Cerrado vegetation physiognomies. *Sociobiology* **60**, 190–197 (2013). [doi:10.13102/sociobiology.v60i2.190-197](https://doi.org/10.13102/sociobiology.v60i2.190-197)
309. L. M. Osgathorpe, K. Park, D. Goulson, The use of off-farm habitats by foraging bumblebees in agricultural landscapes: Implications for conservation management. *Apidologie (Celle)* **43**, 113–127 (2012). [doi:10.1007/s13592-011-0083-z](https://doi.org/10.1007/s13592-011-0083-z)
310. S. E. Otavo, A. Parrado-Rosselli, J. A. Noriega, [Scarabaeoidea superfamily (Insecta: Coleoptera) as a bioindicator element of anthropogenic disturbance in an amazon national park]. *Rev. Biol. Trop.* **61**, 735–752 (2013). [Medline](https://pubmed.ncbi.nlm.nih.gov/24511111/)
311. C. R. V. Otto, G. J. Roloff, Songbird response to green-tree retention prescriptions in clearcut forests. *For. Ecol. Manage.* **284**, 241–250 (2012). [doi:10.1016/j.foreco.2012.07.016](https://doi.org/10.1016/j.foreco.2012.07.016)
312. Ö. Özden, W. M. Ciesla, W. J. Fuller, D. J. Hodgson, Butterfly diversity in Mediterranean islands and in Pentadaktylos *Pinus brutia* forests of Cyprus. *Biodivers. Conserv.* **17**, 2821–2832 (2008). [doi:10.1007/s10531-008-9382-8](https://doi.org/10.1007/s10531-008-9382-8)
313. N. V. Page, Q. Qureshi, G. S. Rawat, C. G. Kushalappa, Plant diversity in sacred forest fragments of Western Ghats: A comparative study of four life forms. *Plant Ecol.* **206**, 237–250 (2010). [doi:10.1007/s11258-009-9638-8](https://doi.org/10.1007/s11258-009-9638-8)
314. S. Paradis, T. T. Work, Partial cutting does not maintain spider assemblages within the observed range of natural variability in Eastern Canadian black spruce forests. *For. Ecol. Manage.* **262**, 2079–2093 (2011). [doi:10.1016/j.foreco.2011.08.032](https://doi.org/10.1016/j.foreco.2011.08.032)
315. J. Paritsis, M. A. Aizen, Effects of exotic conifer plantations on the biodiversity of understory plants, epigeal beetles and birds in *Nothofagus dombeyi* forests. *For. Ecol. Manage.* **255**, 1575–1583 (2008). [doi:10.1016/j.foreco.2007.11.015](https://doi.org/10.1016/j.foreco.2007.11.015)
316. A. Parra-H, G. Nates-Parra, [Variation of the orchid bees community (Hymenoptera: Apidae) in three altered habitats of the Colombian “llano” piedmont]. *Rev. Biol. Trop.* **55**, 931–941 (2007). [Medline](https://pubmed.ncbi.nlm.nih.gov/18411111/)
317. L. Parry, J. Barlow, C. A. Peres, Hunting for sustainability in tropical secondary forests. *Conserv. Biol.* **23**, 1270–1280 (2009). [Medline](https://pubmed.ncbi.nlm.nih.gov/19211111/) [doi:10.1111/j.1523-1739.2009.01224.x](https://doi.org/10.1111/j.1523-1739.2009.01224.x)

318. N. Pelegrin, E. H. Bucher, Effects of habitat degradation on the lizard assemblage in the Arid Chaco, central Argentina. *J. Arid Environ.* **79**, 13–19 (2012). [doi:10.1016/j.jaridenv.2011.11.004](https://doi.org/10.1016/j.jaridenv.2011.11.004)
319. C. A. Peres, H. S. Nascimento, Impact of game hunting by the Kayapo of south-eastern Amazonia: Implications for wildlife conservation in tropical forest indigenous reserves. *Biodivers. Conserv.* **15**, 2627–2653 (2006). [doi:10.1007/s10531-005-5406-9](https://doi.org/10.1007/s10531-005-5406-9)
320. R. S. Pethiyagoda Jr., K. Manamendra-Arachchi, Endangered anurans in a novel forest in the highlands of Sri Lanka. *Wildl. Res.* **39**, 641–648 (2012). [doi:10.1071/WR12079](https://doi.org/10.1071/WR12079)
321. B. Phalan, M. Onial, A. Balmford, R. E. Green, Reconciling food production and biodiversity conservation: Land sharing and land sparing compared. *Science* **333**, 1289–1291 (2011). [Medline doi:10.1126/science.1208742](https://doi.org/10.1126/science.1208742)
322. J. Pincheira-Ulbrich, J. R. Rau, C. Smith-Ramirez, Vascular epiphytes and climbing plants diversity in an agroforestral landscape in southern Chile: A comparison among native forest fragments. *Bol. La Soc. Argentina Bot.* **47**, 411–426 (2012).
323. E. Pineda, G. Halffter, Species diversity and habitat fragmentation: Frogs in a tropical montane landscape in Mexico. *Biol. Conserv.* **117**, 499–508 (2004). [doi:10.1016/j.biocon.2003.08.009](https://doi.org/10.1016/j.biocon.2003.08.009)
324. S. L. Poggio, E. J. Chaneton, C. M. Ghersa, The arable plant diversity of intensively managed farmland: Effects of field position and crop type at local and landscape scales. *Agric. Ecosyst. Environ.* **166**, 55–64 (2013). [doi:10.1016/j.agee.2012.01.013](https://doi.org/10.1016/j.agee.2012.01.013)
325. N. Politi, M. Hunter Jr., L. Rivera, Assessing the effects of selective logging on birds in Neotropical piedmont and cloud montane forests. *Biodivers. Conserv.* **21**, 3131–3155 (2012). [doi:10.1007/s10531-012-0358-3](https://doi.org/10.1007/s10531-012-0358-3)
326. P. Pons, C. Wendenburg, The impact of fire and forest conversion into savanna on the bird communities of West Madagascan dry forests. *Anim. Conserv.* **8**, 183–193 (2005). [doi:10.1017/S1367943005001940](https://doi.org/10.1017/S1367943005001940)
327. K. Poveda, E. Martinez, M. F. Kersch-Becker, M. A. Bonilla, T. Tschardtke, Landscape simplification and altitude affect biodiversity, herbivory and Andean potato yield. *J. Appl. Ecol.* **49**, 513–522 (2012). [doi:10.1111/j.1365-2664.2012.02120.x](https://doi.org/10.1111/j.1365-2664.2012.02120.x)
328. E. F. Power, D. L. Kelly, J. C. Stout, Organic farming and landscape structure: Effects on insect-pollinated plant diversity in intensively managed grasslands. *PLOS ONE* **7**, e38073 (2012). [Medline doi:10.1371/journal.pone.0038073](https://doi.org/10.1371/journal.pone.0038073)
329. E. F. Power, J. C. Stout, Organic dairy farming: Impacts on insect-flower interaction networks and pollination. *J. Appl. Ecol.* **48**, 561–569 (2011). [doi:10.1111/j.1365-2664.2010.01949.x](https://doi.org/10.1111/j.1365-2664.2010.01949.x)
330. S. J. Presley, M. R. Willig, J. M. Wunderle Jr., L. N. Saldanha, Effects of reduced-impact logging and forest physiognomy on bat populations of lowland Amazonian forest. *J. Appl. Ecol.* **45**, 14–25 (2008). [doi:10.1111/j.1365-2664.2007.01373.x](https://doi.org/10.1111/j.1365-2664.2007.01373.x)
331. V. M. Proença, H. M. Pereira, J. Guilherme, L. Vicente, Plant and bird diversity in natural forests and in native and exotic plantations in NW Portugal. *Acta Oecol.* **36**, 219–226 (2010). [doi:10.1016/j.actao.2010.01.002](https://doi.org/10.1016/j.actao.2010.01.002)

332. M. Quaranta *et al.*, Wild bees in agroecosystems and semi-natural landscapes. 1997-2000 collection period in Italy. *Bull. Insectology* **57**, 11–62 (2004).
333. C. Quintero, C. L. Morales, M. A. Aizen, Effects of anthropogenic habitat disturbance on local pollinator diversity and species turnover across a precipitation gradient. *Biodivers. Conserv.* **19**, 257–274 (2010). [doi:10.1007/s10531-009-9720-5](https://doi.org/10.1007/s10531-009-9720-5)
334. R. Rader, I. Bartomeus, J. M. Tylianakis, E. Laliberté, The winners and losers of land use intensification: Pollinator community disassembly is non-random and alters functional diversity. *Divers. Distrib.* **20**, 908–917 (2014).
335. J. Ranganathan, K. M. A. Chan, G. C. Daily, Satellite detection of bird communities in tropical countryside. *Ecol. Appl.* **17**, 1499–1510 (2007). [Medline](https://pubmed.ncbi.nlm.nih.gov/16811111/) [doi:10.1890/06-0285.1](https://doi.org/10.1890/06-0285.1)
336. J. Ranganathan, R. J. R. Daniels, M. D. S. Chandran, P. R. Ehrlich, G. C. Daily, Sustaining biodiversity in ancient tropical countryside. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 17852–17854 (2008). [Medline](https://pubmed.ncbi.nlm.nih.gov/18311111/) [doi:10.1073/pnas.0808874105](https://doi.org/10.1073/pnas.0808874105)
337. F. Raub, H. Höfer, L. Scheuermann, R. Brandl, The conservation value of secondary forests in the southern Brazilian Mata Atlantica from a spider perspective. *J. Arachnol.* **42**, 52–73 (2014). [doi:10.1636/P13-47.1](https://doi.org/10.1636/P13-47.1)
338. N. Redpath, L. M. Osgathorpe, K. Park, D. Goulson, Crofting and bumblebee conservation: The impact of land management practices on bumblebee populations in northwest Scotland. *Biol. Conserv.* **143**, 492–500 (2010). [doi:10.1016/j.biocon.2009.11.019](https://doi.org/10.1016/j.biocon.2009.11.019)
339. J. L. Reid, J. B. C. Harris, R. A. Zahawi, Avian habitat preference in tropical forest restoration in southern Costa Rica. *Biotropica* **44**, 350–359 (2012). [doi:10.1111/j.1744-7429.2011.00814.x](https://doi.org/10.1111/j.1744-7429.2011.00814.x)
340. Y. T. Reis, E. M. Cancellato, Termite (Insecta, Isoptera) richness in primary and secondary Atlantic Forest in southeastern Bahia. *Iheringia Ser. Zool.* **97**, 229–234 (2007).
341. J. M. Rey-Benayas, I. Galvan, L. M. Carrascal, Differential effects of vegetation restoration in Mediterranean abandoned cropland by secondary succession and pine plantations on bird assemblages. *For. Ecol. Manage.* **260**, 87–95 (2010). [doi:10.1016/j.foreco.2010.04.004](https://doi.org/10.1016/j.foreco.2010.04.004)
342. C. Reynolds, C. T. Symes, Grassland bird response to vegetation structural heterogeneity and clearing of invasive bramble. *Afr. Zool.* **48**, 228–239 (2013). [doi:10.3377/004.048.0217](https://doi.org/10.3377/004.048.0217)
343. D. B. Ribeiro, A. V. L. Freitas, The effect of reduced-impact logging on fruit-feeding butterflies in Central Amazon, Brazil. *J. Insect Conserv.* **16**, 733–744 (2012). [doi:10.1007/s10841-012-9458-3](https://doi.org/10.1007/s10841-012-9458-3)
344. B. A. Richardson, M. J. Richardson, F. N. Soto-Adames, Separating the effects of forest type and elevation on the diversity of litter invertebrate communities in a humid tropical forest in Puerto Rico. *J. Anim. Ecol.* **74**, 926–936 (2005). [doi:10.1111/j.1365-2656.2005.00990.x](https://doi.org/10.1111/j.1365-2656.2005.00990.x)
345. C. A. Robles, C. C. Carmaran, S. E. Lopez, Screening of xylophagous fungi associated with *Platanus acerifolia* in urban landscapes: Biodiversity and potential biodeterioration. *Landsc. Urban Plan.* **100**, 129–135 (2011). [doi:10.1016/j.landurbplan.2010.12.003](https://doi.org/10.1016/j.landurbplan.2010.12.003)

346. M. M. Rodrigues, M. A. Uchôa, S. Ide, Dung beetles (Coleoptera: Scarabaeoidea) in three landscapes in Mato Grosso do Sul, Brazil. *Braz. J. Biol.* **73**, 211–220 (2013). [Medline](#) [doi:10.1590/S1519-69842013000100023](#)
347. J. Römbke, P. Schmidt, H. Höfer, The earthworm fauna of regenerating forests and anthropogenic habitats in the coastal region of Paraná. *Pesquisa Agropecu. Bras.* **44**, 1040–1049 (2009). [doi:10.1590/S0100-204X2009000800037](#)
348. L. P. Romero-Duque, V. J. Jaramillo, A. Perez-Jimenez, Structure and diversity of secondary tropical dry forests in Mexico, differing in their prior land-use history. *For. Ecol. Manage.* **253**, 38–47 (2007). [doi:10.1016/j.foreco.2007.07.002](#)
349. L. Rosselli, thesis, “Factores Ambientales Relacionados con la Presencia y Abundancia de las Aves de los Humedales de la Sabana de Bogotá,” Universidad Nacional de Colombia, 2011.
350. L. Rousseau, S. J. Fonte, O. Tellez, R. van der Hoek, P. Lavelle, Soil macrofauna as indicators of soil quality and land use impacts in smallholder agroecosystems of western Nicaragua. *Ecol. Indic.* **27**, 71–82 (2013). [doi:10.1016/j.ecolind.2012.11.020](#)
351. A. V. Rubio, J. A. Simonetti, Lizard assemblages in a fragmented landscape of central Chile. *Eur. J. Wildl. Res.* **57**, 195–199 (2011). [doi:10.1007/s10344-010-0434-5](#)
352. S. Sáfián, G. Csontos, D. Winkler, Butterfly community recovery in degraded rainforest habitats in the Upper Guinean Forest Zone (Kakum forest, Ghana). *J. Insect Conserv.* **15**, 351–359 (2011). [doi:10.1007/s10841-010-9343-x](#)
353. W. Sakchoowong, S. Nomura, K. Ogata, J. Chanpaisaeng, Diversity of pselaphine beetles (Coleoptera: Staphylinidae: Pselaphinae) in eastern Thailand. *Entomol. Sci.* **11**, 301–313 (2008). [doi:10.1111/j.1479-8298.2008.00281.x](#)
354. R. A. Saldaña-Vázquez, V. J. Sosa, J. R. Hernandez-Montero, F. Lopez-Barrera, Abundance responses of frugivorous bats (Stenodermatinae) to coffee cultivation and selective logging practices in mountainous central Veracruz, Mexico. *Biodivers. Conserv.* **19**, 2111–2124 (2010). [doi:10.1007/s10531-010-9829-6](#)
355. K. Sam, B. Koane, S. Jeppy, V. Novotny, Effect of forest fragmentation on bird species richness in Papua New Guinea. *J. Field Ornithol.* **85**, 152–167 (2014). [doi:10.1111/jfo.12057](#)
356. U. Samnegård, A. S. Persson, H. G. Smith, Gardens benefit bees and enhance pollination in intensively managed farmland. *Biol. Conserv.* **144**, 2602–2606 (2011). [doi:10.1016/j.biocon.2011.07.008](#)
357. J. Santana, M. Porto, L. Gordinho, L. Reino, P. Beja, Long-term responses of Mediterranean birds to forest fuel management. *J. Appl. Ecol.* **49**, 632–643 (2012).
358. J. Savage, T. A. Wheeler, A. M. A. Moores, A. G. Taillefer, Effects of habitat size, vegetation cover, and surrounding land use on Diptera diversity in temperate Nearctic bogs. *Wetlands* **31**, 125–134 (2011). [doi:10.1007/s13157-010-0133-8](#)
359. M. Schilthuizen, T. S. Liew, B. Bin Elahan, I. Lackman-Ancrenaz, Effects of karst forest degradation on pulmonate and prosobranch land snail communities in Sabah, Malaysian Borneo. *Conserv. Biol.* **19**, 949–954 (2005).

360. N. L. Schon, A. D. Mackay, M. A. Minor, G. W. Yeates, M. J. Hedley, Soil fauna in grazed New Zealand hill country pastures at two management intensities. *Appl. Soil Ecol.* **40**, 218–228 (2008). [doi:10.1016/j.apsoil.2008.04.007](https://doi.org/10.1016/j.apsoil.2008.04.007)
361. N. L. Schon, A. D. Mackay, G. W. Yeates, M. A. Minor, Separating the effects of defoliation and dairy cow treading pressure on the abundance and diversity of soil invertebrates in pastures. *Appl. Soil Ecol.* **46**, 209–221 (2010). [doi:10.1016/j.apsoil.2010.08.011](https://doi.org/10.1016/j.apsoil.2010.08.011)
362. C. Schüepp, J. D. Herrmann, F. Herzog, M. H. Schmidt-Entling, Differential effects of habitat isolation and landscape composition on wasps, bees, and their enemies. *Oecologia* **165**, 713–721 (2011). [Medline](https://pubmed.ncbi.nlm.nih.gov/21544420/) [doi:10.1007/s00442-010-1746-6](https://doi.org/10.1007/s00442-010-1746-6)
363. C. Schüepp, S. Rittiner, M. H. Entling, High bee and wasp diversity in a heterogeneous tropical farming system compared to protected forest. *PLOS ONE* **7**, e52109 (2012). [Medline](https://pubmed.ncbi.nlm.nih.gov/2252109/) [doi:10.1371/journal.pone.0052109](https://doi.org/10.1371/journal.pone.0052109)
364. K. Schumann, R. Wittig, A. Thiombiano, U. Becker, K. Hahn, Impact of land-use type and harvesting on population structure of a non-timber forest product-providing tree in a semi-arid savanna, West Africa. *Biol. Conserv.* **144**, 2369–2376 (2011). [doi:10.1016/j.biocon.2011.06.018](https://doi.org/10.1016/j.biocon.2011.06.018)
365. D. M. Scott, D. Brown, S. Mahood, B. Denton, A. Silburn, F. Rakotondraparany, The impacts of forest clearance on lizard, small mammal and bird communities in the arid spiny forest, southern Madagascar. *Biol. Conserv.* **127**, 72–87 (2006). [doi:10.1016/j.biocon.2005.07.014](https://doi.org/10.1016/j.biocon.2005.07.014)
366. K. A. Scott, S. A. Setterfield, M. M. Douglas, A. N. Andersen, Fire tolerance of perennial grass tussocks in a savanna woodland. *Austral Ecol.* **35**, 858–861 (2010). [doi:10.1111/j.1442-9993.2009.02091.x](https://doi.org/10.1111/j.1442-9993.2009.02091.x)
367. J. L. Sedlock, S. E. Weyandt, L. Cororan, M. Damerow, S.-H. Hwa, B. Pauli, Bat diversity in tropical forest and agro-pastoral habitats within a protected area in the Philippines. *Acta Chiropt.* **10**, 349–358 (2008). [doi:10.3161/150811008X414926](https://doi.org/10.3161/150811008X414926)
368. N. J. Shafie, S. A. M. Sah, N. S. A. Latip, N. M. Azman, N. L. Khairuddin, Diversity pattern of bats at two contrasting habitat types along Kerian River, Perak, Malaysia. *Trop. Life Sci. Res.* **22**, 13–22 (2011). [Medline](https://pubmed.ncbi.nlm.nih.gov/21544420/)
369. G. Shahabuddin, R. Kumar, Effects of extractive disturbance on bird assemblages, vegetation structure and floristics in tropical scrub forest, Sariska Tiger Reserve, India. *For. Ecol. Manage.* **246**, 175–185 (2007). [doi:10.1016/j.foreco.2007.03.061](https://doi.org/10.1016/j.foreco.2007.03.061)
370. G. Shannon, D. J. Druce, B. R. Page, H. C. Eckhardt, R. Grant, R. Slotow, The utilization of large savanna trees by elephant in southern Kruger National Park. *J. Trop. Ecol.* **24**, 281–289 (2008). [doi:10.1017/S0266467408004951](https://doi.org/10.1017/S0266467408004951)
371. D. Sheil *et al.*, Exploring Biological Diversity, Environment and Local People's Perspectives in Forest Landscapes: Methods for a Multidisciplinary Landscape Assessment," Center for International Forestry Research, Jakarta, Indonesia, 2002.

372. F. H. Sheldon, A. Styring, P. A. Hosner, Bird species richness in a Bornean exotic tree plantation: A long-term perspective. *Biol. Conserv.* **143**, 399–407 (2010). [doi:10.1016/j.biocon.2009.11.004](https://doi.org/10.1016/j.biocon.2009.11.004)
373. R. E. Shuler, T. H. Roulston, G. E. Farris, Farming practices influence wild pollinator populations on squash and pumpkin. *J. Econ. Entomol.* **98**, 790–795 (2005). [Medline](https://pubmed.ncbi.nlm.nih.gov/1603/0022-0493-98.3.790/) [doi:10.1603/0022-0493-98.3.790](https://doi.org/10.1603/0022-0493-98.3.790)
374. S. J. Siebert, Patterns of plant species richness of temperate and tropical grassland in South Africa. *Plant Ecol. Evol.* **144**, 249–254 (2011). [doi:10.5091/plecevo.2011.501](https://doi.org/10.5091/plecevo.2011.501)
375. F. A. B. Silva, C. M. Q. Costa, R. C. Moura, A. I. Farias, Study of the dung beetle (Coleoptera: Scarabaeidae) community at two sites: Atlantic forest and clear-cut, Pernambuco, Brazil. *Environ. Entomol.* **39**, 359–367 (2010). [Medline](https://pubmed.ncbi.nlm.nih.gov/1603/EN09180/) [doi:10.1603/EN09180](https://doi.org/10.1603/EN09180)
376. E. M. Slade, D. J. Mann, O. T. Lewis, Biodiversity and ecosystem function of tropical forest dung beetles under contrasting logging regimes. *Biol. Conserv.* **144**, 166–174 (2011). [doi:10.1016/j.biocon.2010.08.011](https://doi.org/10.1016/j.biocon.2010.08.011)
377. A. Smith-Pardo, V. H. Gonzalez, Diversidad de abejas (Hymenoptera: Apoidea) en estados sucesionales del bosque humedo tropical. *Acta Biológica Colomb.* **12**, 43–55 (2007).
378. N. S. Sodhi, D. S. Wilcove, T. M. Lee, C. H. Sekercioglu, R. Subaraj, H. Bernard, D. L. Yong, S. L. Lim, D. M. Prawiradilaga, B. W. Brook, Deforestation and avian extinction on tropical landbridge islands. *Conserv. Biol.* **24**, 1290–1298 (2010). [Medline](https://pubmed.ncbi.nlm.nih.gov/1111/j.1523-1739.2010.01495.x/) [doi:10.1111/j.1523-1739.2010.01495.x](https://doi.org/10.1111/j.1523-1739.2010.01495.x)
379. M. C. K. Soh, N. S. Sodhi, S. L. H. Lim, High sensitivity of montane bird communities to habitat disturbance in Peninsular Malaysia. *Biol. Conserv.* **129**, 149–166 (2006). [doi:10.1016/j.biocon.2005.10.030](https://doi.org/10.1016/j.biocon.2005.10.030)
380. R. A. Sosa, V. A. Benz, J. M. Galea, I. V. Poggio Herrero, Efecto del grado de disturbio sobre el ensamble de aves en la reserva provincial Parque Luro, La Pampa, Argentina. *Rev. la Asoc. Argentina Ecol. Paisajes* **1**, 101–110 (2010).
381. H. Sridhar, T. R. S. Raman, D. Mudappa, Mammal persistence and abundance in tropical rainforest remnants in the southern Western Ghats, India. *Curr. Sci.* **94**, 748–757 (2008).
382. P. C. Stouffer, E. I. Johnson, R. O. Bierregaard Jr., T. E. Lovejoy, Understory bird communities in Amazonian rainforest fragments: Species turnover through 25 years post-isolation in recovering landscapes. *PLOS ONE* **6**, e20543 (2011). [10.1371/journal.pone.0020543](https://doi.org/10.1371/journal.pone.0020543) [Medline](https://pubmed.ncbi.nlm.nih.gov/10.1371/journal.pone.0020543/) [doi:10.1371/journal.pone.0020543](https://doi.org/10.1371/journal.pone.0020543)
383. L. Ström, K. Hylander, M. Dynesius, Different long-term and short-term responses of land snails to clear-cutting of boreal stream-side forests. *Biol. Conserv.* **142**, 1580–1587 (2009). [doi:10.1016/j.biocon.2009.02.028](https://doi.org/10.1016/j.biocon.2009.02.028)
384. M. J. Struebig, T. Kingston, A. Zubaid, A. Mohd-Adnan, S. J. Rossiter, Conservation value of forest fragments to Palaeotropical bats. *Biol. Conserv.* **141**, 2112–2126 (2008). [doi:10.1016/j.biocon.2008.06.009](https://doi.org/10.1016/j.biocon.2008.06.009)

385. Z. M. Su, R. Z. Zhang, J. X. Qiu, Decline in the diversity of willow trunk-dwelling weevils (Coleoptera: Curculionoidea) as a result of urban expansion in Beijing, China. *J. Insect Conserv.* **15**, 367–377 (2011). [doi:10.1007/s10841-010-9310-6](https://doi.org/10.1007/s10841-010-9310-6)
386. M. Suarez-Rubio, J. R. Thomlinson, Landscape and patch-level factors influence bird communities in an urbanized tropical island. *Biol. Conserv.* **142**, 1311–1321 (2009). [doi:10.1016/j.biocon.2008.12.035](https://doi.org/10.1016/j.biocon.2008.12.035)
387. S. Sugiura, T. Tsuru, Y. Yamaura, H. Makihara, Small off-shore islands can serve as important refuges for endemic beetle conservation. *J. Insect Conserv.* **13**, 377–385 (2009). [doi:10.1007/s10841-008-9185-y](https://doi.org/10.1007/s10841-008-9185-y)
388. K. S. Summerville, Managing the forest for more than the trees: Effects of experimental timber harvest on forest Lepidoptera. *Ecol. Appl.* **21**, 806–816 (2011). [Medline](https://pubmed.ncbi.nlm.nih.gov/21511111/) [doi:10.1890/10-0715.1](https://doi.org/10.1890/10-0715.1)
389. K. S. Summerville, C. J. Conoan, R. M. Steichen, Species traits as predictors of lepidopteran composition in restored and remnant tallgrass prairies. *Ecol. Appl.* **16**, 891–900 (2006). [Medline](https://pubmed.ncbi.nlm.nih.gov/16481111/) [doi:10.1890/1051-0761\(2006\)016\[0891:STAPOL\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[0891:STAPOL]2.0.CO;2)
390. K. S. Summerville, T. O. Crist, Effects of timber harvest on forest Lepidoptera: Community, guild and species responses. *Ecol. Appl.* **12**, 820–835 (2002). [doi:10.1890/1051-0761\(2002\)012\[0820:EOTHOF\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2002)012[0820:EOTHOF]2.0.CO;2)
391. Y. H. Sung, N. E. Karraker, B. C. H. Hau, Terrestrial herpetofaunal assemblages in secondary forests and exotic *Lophostemon confertus* plantations in South China. *For. Ecol. Manage.* **270**, 71–77 (2012). [doi:10.1016/j.foreco.2012.01.011](https://doi.org/10.1016/j.foreco.2012.01.011)
392. C. G. Threlfall, B. Law, P. B. Banks, Sensitivity of insectivorous bats to urbanization: Implications for suburban conservation planning. *Biol. Conserv.* **146**, 41–52 (2012). [doi:10.1016/j.biocon.2011.11.026](https://doi.org/10.1016/j.biocon.2011.11.026)
393. R. Tonietto, J. Fant, J. Ascher, K. Ellis, D. Larkin, A comparison of bee communities of Chicago green roofs, parks and prairies. *Landsc. Urban Plan.* **103**, 102–108 (2011). [doi:10.1016/j.landurbplan.2011.07.004](https://doi.org/10.1016/j.landurbplan.2011.07.004)
394. I. Torre, V. Bros, X. Santos, Assessing the impact of reforestation on the diversity of Mediterranean terrestrial Gastropoda. *Biodivers. Conserv.* **23**, 2579–2589 (2014). [doi:10.1007/s10531-014-0740-4](https://doi.org/10.1007/s10531-014-0740-4)
395. E. C. Turner, W. A. Foster, The impact of forest conversion to oil palm on arthropod abundance and biomass in Sabah, Malaysia. *J. Trop. Ecol.* **25**, 23–30 (2009). [doi:10.1017/S0266467408005658](https://doi.org/10.1017/S0266467408005658)
396. J. M. Tylianakis, A.-M. Klein, T. Tscharntke, Spatiotemporal variation in the diversity of Hymenoptera across a tropical habitat gradient. *Ecology* **86**, 3296–3302 (2005). [doi:10.1890/05-0371](https://doi.org/10.1890/05-0371)
397. M. Uehara-Prado, Effects of land use on ant species composition and diaspore removal in exotic grasslands in the Brazilian Pantanal (Hymenoptera: Formicidae). *Sociobiology* **45**, 915–923 (2005).
398. M. Uehara-Prado, K. S. Brown, A. V. L. Freitas, Species richness, composition and abundance of fruit-feeding butterflies in the Brazilian Atlantic Forest: Comparison

- between a fragmented and a continuous landscape. *Glob. Ecol. Biogeogr.* **16**, 43–54 (2007). [doi:10.1111/j.1466-8238.2006.00267.x](https://doi.org/10.1111/j.1466-8238.2006.00267.x)
399. J. N. Urbina-Cardona, M. Olivares-Perez, V. H. Reynoso, Herpetofauna diversity and microenvironment correlates across a pasture-edge-interior ecotone in tropical rainforest fragments in the Los Tuxtlas Biosphere Reserve of Veracruz, Mexico. *Biol. Conserv.* **132**, 61–75 (2006). [doi:10.1016/j.biocon.2006.03.014](https://doi.org/10.1016/j.biocon.2006.03.014)
400. J. N. Urbina-Cardona, M. C. Londoño-Murcia, D. G. García-Ávila, Spatio-temporal dynamics of snake diversity in four habitats with different degrees of anthropogenic disturbance in the Gorgona Island National Natural Park in the Colombian Pacific. *Caldasia* **30**, 479–493 (2008).
401. D. Vallan, Effects of anthropogenic environmental changes on amphibian diversity in the rain forests of eastern Madagascar. *J. Trop. Ecol.* **18**, 725–742 (2002). [doi:10.1017/S026646740200247X](https://doi.org/10.1017/S026646740200247X)
402. A. J. Vanbergen, B. A. Woodcock, A. D. Watt, J. Niemela, Effect of land-use heterogeneity on carabid communities at the landscape scale. *Ecography* **28**, 3–16 (2005). [doi:10.1111/j.0906-7590.2005.03991.x](https://doi.org/10.1111/j.0906-7590.2005.03991.x)
403. H. L. Vasconcelos, R. Pacheco, R. C. Silva, P. B. Vasconcelos, C. T. Lopes, A. N. Costa, E. M. Bruna, Dynamics of the leaf-litter arthropod fauna following fire in a neotropical woodland savanna. *PLOS ONE* **4**, e7762 (2009). [Medline doi:10.1371/journal.pone.0007762](https://doi.org/10.1371/journal.pone.0007762)
404. K. Vassilev, H. Pedashenko, S. C. Nikolov, I. Apostolova, J. Dengler, Effect of land abandonment on the vegetation of upland semi-natural grasslands in the Western Balkan Mts., Bulgaria. *Plant Biosyst.* **145**, 654–665 (2011). [doi:10.1080/11263504.2011.601337](https://doi.org/10.1080/11263504.2011.601337)
405. D. P. Vázquez, D. Simberloff, Ecological specialization and susceptibility to disturbance: Conjectures and refutations. *Am. Nat.* **159**, 606–623 (2002). [Medline doi:10.1086/339991](https://doi.org/10.1086/339991)
406. H. A. F. Verboven, R. Brys, M. Hermy, Sex in the city: Reproductive success of *Digitalis purpurea* in a gradient from urban to rural sites. *Landsc. Urban Plan.* **106**, 158–164 (2012). [doi:10.1016/j.landurbplan.2012.02.015](https://doi.org/10.1016/j.landurbplan.2012.02.015)
407. M. J. Verdasca, A. S. Leitão, J. Santana, M. Porto, S. Dias, P. Beja, Forest fuel management as a conservation tool for early successional species under agricultural abandonment: The case of Mediterranean butterflies. *Biol. Conserv.* **146**, 14–23 (2012). [doi:10.1016/j.biocon.2011.10.031](https://doi.org/10.1016/j.biocon.2011.10.031)
408. J. R. Verdú, C. E. Moreno, G. Sánchez-Rojas, C. Numa, E. Galante, G. Halfpeter, Grazing promotes dung beetle diversity in the xeric landscape of a Mexican Biosphere Reserve. *Biol. Conserv.* **140**, 308–317 (2007). [doi:10.1016/j.biocon.2007.08.015](https://doi.org/10.1016/j.biocon.2007.08.015)
409. C. H. Vergara, E. I. Badano, Pollinator diversity increases fruit production in Mexican coffee plantations: The importance of rustic management systems. *Agric. Ecosyst. Environ.* **129**, 117–123 (2009). [doi:10.1016/j.agee.2008.08.001](https://doi.org/10.1016/j.agee.2008.08.001)
410. P. M. Vergara, J. A. Simonetti, Avian responses to fragmentation of the Maulino Forest in central Chile. *Oryx* **38**, 383–388 (2004). [doi:10.1017/S0030605304000742](https://doi.org/10.1017/S0030605304000742)

411. M. Virgilio, T. Backeljau, R. Emeleme, J. L. Juakali, M. De Meyer, A quantitative comparison of frugivorous tephritids (Diptera: Tephritidae) in tropical forests and rural areas of the Democratic Republic of Congo. *Bull. Entomol. Res.* **101**, 591–597 (2011). [Medline](#) [doi:10.1017/S0007485311000216](#)
412. L. Van Vu, C. Q. Vu, Diversity pattern of butterfly communities (Lepidoptera, Papilionoidea) in different habitat types in a tropical rain forest of Southern Vietnam. *ISRN Zool.* **2011**, 1–8 (2011).
413. E. M. Waite, G. Closs, Y. Van Heezik, C. Berry, K. Dickinson, Arboreal arthropod sampling methods for urban trees. *J. Insect Conserv.* **16**, 931–939 (2012). [doi:10.1007/s10841-012-9480-5](#)
414. E. Waite, G. P. Closs, Y. van Heezik, K. J. M. Dickinson, Resource availability and foraging of silvereyes (*Zosterops lateralis*) in urban trees. *Emu* **113**, 26–32 (2013). [doi:10.1071/MU11093](#)
415. S. Walker, D. J. Wilson, G. Norbury, A. Monks, A. J. Tanentzap, Complementarity of indigenous flora in shrublands and grasslands in a New Zealand dryland landscape. *N. Z. J. Ecol.* **38**, 230–241 (2014).
416. T. R. Walker, P. D. Crittenden, S. D. Young, T. Prystina, An assessment of pollution impacts due to the oil and gas industries in the Pechora Basin, north-eastern European Russia. *Ecol. Indic.* **6**, 369–387 (2006). [doi:10.1016/j.ecolind.2005.03.015](#)
417. Y. Wang, Y. Bao, M. Yu, G. Xu, P. Ding, Nestedness for different reasons: The distributions of birds, lizards and small mammals on islands of an inundated lake. *Divers. Distrib.* **16**, 862–873 (2010). [doi:10.1111/j.1472-4642.2010.00682.x](#)
418. J. I. Watling, K. Gerow, M. A. Donnelly, Nested species subsets of amphibians and reptiles on Neotropical forest islands. *Anim. Conserv.* **12**, 467–476 (2009). [doi:10.1111/j.1469-1795.2009.00274.x](#)
419. B. Weller, J. U. Ganzhorn, Carabid beetle community composition, body size, and fluctuating asymmetry along an urban-rural gradient. *Basic Appl. Ecol.* **5**, 193–201 (2004). [doi:10.1078/1439-1791-00220](#)
420. K. Wells, E. K. V. Kalko, M. B. Lakim, M. Pfeiffer, Effects of rain forest logging on species richness and assemblage composition of small mammals in Southeast Asia. *J. Biogeogr.* **34**, 1087–1099 (2007). [doi:10.1111/j.1365-2699.2006.01677.x](#)
421. E. D. Wiafe, R. Amfo-Otu, Forest duiker (*Cephalophus* spp.) abundance and hunting activities in the Kakum conservation area, Ghana. *J. Ecol. Nat. Environ.* **4**, 114–118 (2012).
422. C. D. Williams, J. Sheahan, M. J. Gormally, Hydrology and management of turloughs (temporary lakes) affect marsh fly (Sciomyzidae: Diptera) communities. *Insect Conserv. Divers.* **2**, 270–283 (2009). [doi:10.1111/j.1752-4598.2009.00064.x](#)
423. M. R. Willig, S. J. Presley, C. P. Bloch, C. L. Hice, S. P. Yanoviak, M. M. Díaz, L. A. Chauca, V. Pacheco, S. C. Weaver, Phyllostomid bats of lowland Amazonia: Effects of habitat alteration on abundance. *Biotropica* **39**, 737–746 (2007). [doi:10.1111/j.1744-7429.2007.00322.x](#)

424. R. Winfree, T. Griswold, C. Kremen, Effect of human disturbance on bee communities in a forested ecosystem. *Conserv. Biol.* **21**, 213–223 (2007). [Medline doi:10.1111/j.1523-1739.2006.00574.x](#)
425. J. C. Z. Woinarski, B. Rankmore, B. Hill, A. D. Griffiths, A. Stewart, B. Grace, Fauna assemblages in regrowth vegetation in tropical open forests of the Northern Territory, Australia. *Wildl. Res.* **36**, 675–690 (2009). [doi:10.1071/WR08128](#)
426. B. A. Woodcock, S. G. Potts, E. Pilgrim, A. J. Ramsay, T. Tscheulin, A. Parkinson, R. E. N. Smith, A. L. Gundry, V. K. Brown, J. R. Tallowin, The potential of grass field margin management for enhancing beetle diversity in intensive livestock farms. *J. Appl. Ecol.* **44**, 60–69 (2007). [doi:10.1111/j.1365-2664.2006.01258.x](#)
427. T. Wronski, K. Gilbert, E. Long, B. Micha, R. Quinn, B. Hausdorf, Species richness and meta-community structure of land snails along an altitudinal gradient on Bioko Island, Equatorial Guinea. *J. Molluscan Stud.* **80**, 161–168 (2014). [doi:10.1093/mollus/eyu008](#)
428. J. M. Wunderle Jr., L. M. P. Henriques, M. R. Willig, Short-term responses of birds to forest gaps and understory: An assessment of reduced-impact logging in a lowland Amazon Forest. *Biotropica* **38**, 235–255 (2006). [doi:10.1111/j.1744-7429.2006.00138.x](#)
429. Y. Yamaura, J. A. Royle, N. Shimada, S. Asanuma, T. Sato, H. Taki, S. Makino, Biodiversity of man-made open habitats in an underused country: A class of multispecies abundance models for count data. *Biodivers. Conserv.* **21**, 1365–1380 (2012). [doi:10.1007/s10531-012-0244-z](#)
430. X. L. Yan, W. K. Bao, X. Y. Pang, N. X. Zhang, J. Q. Chen, Regeneration strategies influence ground bryophyte composition and diversity after forest clearcutting. *Ann. For. Sci.* **70**, 845–861 (2013). [doi:10.1007/s13595-013-0323-7](#)
431. Y. Xiaoli, B. Weikai, Evaluation of species composition and development of bryophyte community during early natural recovery progress of high-altitude spruce cutovers. *Biodivers. Sci.* **16**, 110–117 (2008). [doi:10.3724/SP.J.1003.2008.07349](#)
432. X.-L. Yan, W.-K. Bao, Structure and species composition of ground bryophyte community of high-altitude young silvicultural cutovers in Rangtang County, China: Evaluation on effects of clear-cutting and silvicultural management. *Chinese J. Plant Ecol.* **32**, 815–824 (2008).
433. S. Yoshikura, S. Yasui, T. Kamijo, Comparative study of forest-dwelling bats' abundances and species richness between old-growth forests and conifer plantations in Nikko National Park, central Japan. *Mammal Study* **36**, 189–198 (2011). [doi:10.3106/041.036.0402](#)
434. A. S. Zaitsev, M. Chauvat, A. Pflug, V. Wolters, Oribatid mite diversity and community dynamics in a spruce chronosequence. *Soil Biol. Biochem.* **34**, 1919–1927 (2002). [doi:10.1016/S0038-0717\(02\)00208-0](#)
435. A. S. Zaitsev, V. Wolters, R. Waldhardt, J. Dauber, Long-term succession of oribatid mites after conversion of croplands to grasslands. *Appl. Soil Ecol.* **34**, 230–239 (2006). [doi:10.1016/j.apsoil.2006.01.005](#)

436. J. N. Zhang, Q. Li, W. J. Liang, Effect of acetochlor and carbofuran on soil nematode communities in a Chinese soybean field. *African J. Agric. Res.* **5**, 2787–2794 (2010).
437. G. Zimmerman, F. W. Bell, J. Woodcock, A. Palmer, J. Paloniemi, Response of breeding songbirds to vegetation management in conifer plantations established in boreal mixedwoods. *For. Chron.* **87**, 217–224 (2011). [doi:10.5558/tfc2011-009](https://doi.org/10.5558/tfc2011-009)
438. P. A. Murtaugh, Performance of several variable-selection methods applied to real ecological data. *Ecol. Lett.* **12**, 1061–1068 (2009). [Medline](https://pubmed.ncbi.nlm.nih.gov/19244441/) [doi:10.1111/j.1461-0248.2009.01361.x](https://doi.org/10.1111/j.1461-0248.2009.01361.x)
439. D. Bates, M. Maechler, B. Bolker, S. Walker, lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1-7 (2014); available at <http://cran.r-project.org/package=lme4>.
440. R Core Team, R: A Language and Environment for Statistical Computing (2015); available at www.r-project.org.
441. G. C. Hurtt *et al.*, *Clim. Change* **109**, 117–161 (2011).
442. Q. Mu, M. Zhao, S. W. Running, Improvements to a MODIS global terrestrial evapotranspiration algorithm. *Remote Sens. Environ.* **115**, 1781–1800 (2011). [doi:10.1016/j.rse.2011.02.019](https://doi.org/10.1016/j.rse.2011.02.019)
443. R. J. Hijmans, S. E. Cameron, J. L. Parra, P. G. Jones, A. Jarvis, Very high resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**, 1965–1978 (2005). [doi:10.1002/joc.1276](https://doi.org/10.1002/joc.1276)
444. H. I. Reuter, T. Hengl, in *Digital Soil Assessments and Beyond*, B. Minasny, B. P. Malone, A. B. McBratney, Eds. (Taylor & Francis Group, 2012).
445. T. Hengl, J. M. de Jesus, R. A. MacMillan, N. H. Batjes, G. B. Heuvelink, E. Ribeiro, A. Samuel-Rosa, B. Kempen, J. G. Leenaars, M. G. Walsh, M. R. Gonzalez, SoilGrids1km—Global soil information based on automated mapping. *PLOS ONE* **9**, e105992 (2014). [Medline](https://pubmed.ncbi.nlm.nih.gov/25401111/) [doi:10.1371/journal.pone.0105992](https://doi.org/10.1371/journal.pone.0105992)
446. A. Nelson, *Estimated Travel Time to the Nearest City of 50,000 or More People in Year 2000* (Ispra, Italy, 2008); <http://forobs.jrc.ec.europa.eu/products/gam>.
447. M.-N. Tuanmu, W. Jetz, A global 1-km consensus land-cover product for biodiversity and ecosystem modelling. *Glob. Ecol. Biogeogr.* **23**, 1031–1045 (2014). [doi:10.1111/geb.12182](https://doi.org/10.1111/geb.12182)
448. B. Lehner, P. Döll, Development and validation of a global database of lakes, reservoirs and wetlands. *J. Hydrol. (Amst.)* **296**, 1–22 (2004). [doi:10.1016/j.jhydrol.2004.03.028](https://doi.org/10.1016/j.jhydrol.2004.03.028)
449. D. M. Olson *et al.*, Terrestrial ecoregions of the world: A new map of life on Earth. *Bioscience* **51**, 933–938 (2001).
450. S. van Asselen, P. H. Verburg, Land cover change or land-use intensification: Simulating land system change with a global-scale land change model. *Glob. Change Biol.* **19**, 3648–3667 (2013). [doi:10.1111/gcb.12331](https://doi.org/10.1111/gcb.12331)
451. ESRI, ArcGIS Desktop: Release 10. (2011).

452. A. Longacre, L. R. Scott, J. S. Levine, Linear independence of pairwise comparisons of DNA microarray data. *J. Bioinform. Comput. Biol.* **3**, 1243–1262 (2005). [Medline](#) [doi:10.1142/S0219720005001600](https://doi.org/10.1142/S0219720005001600)
453. A. Chao, R. L. Chazdon, R. K. Colwell, T.-J. Shen, A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecol. Lett.* **8**, 148–159 (2005). [doi:10.1111/j.1461-0248.2004.00707.x](https://doi.org/10.1111/j.1461-0248.2004.00707.x)
454. R. A. Mittermeier, C. G. Mittermeier, T. M. Brooks, J. D. Pilgrim, W. R. Konstant, G. A. da Fonseca, C. Kormos, Wilderness and biodiversity conservation. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 10309–10313 (2003). [Medline](#) [doi:10.1073/pnas.1732458100](https://doi.org/10.1073/pnas.1732458100)
455. A. D. Chapman, *Numbers of Living Species in Australia and the World* (Australian Biological Resources Study, Canberra, Australia, 2009; available at www.environment.gov.au/system/files/pages/2ee3f4a1-f130-465b-9c7a-79373680a067/files/nlsaw-2nd-complete.pdf).
456. J. C. Pinheiro, D. M. Bates, *Mixed-Effects Models in S and S-plus* (Springer, 2000).
457. A. F. Zuur, E. N. Ieno, N. J. Walker, A. A. Saveliev, G. M. Smith, *Mixed Effects Models and Extensions in Ecology with R* (Springer, 2009).