

Todo list

■ PC: Need some notions from the sociology literature	4
■ PC: place the measurements of entropy and mutual information in the titles for each panel.	5
■ PC: Move from concave functions to functions whose negatives are convex.	5
■ PC: Locally quadratic, distance-like properties (foreshadowing for later)	5
■ PC: Put in a city here with blockgroup, tract, and maybe ZIP, neighborhood, or county-level aggregation and show how the decomposition works.	7
■ PC: Also bring in an example of generating a boundary via spatial aggregation.	7

Measuring Diversity and Segregation with Convex Functions

Philip S. Chodrow *

July 17, 2026

The study of quantitative measures for residential segregation extends far enough back that a review article in 1955 could express frustration and disappointment with the chaotic state of the methodology. Commenting then on the measures extant at the time, Duncan and Duncan [DD55] wrote:

1. *The proposed indexes of segregation have a number of hitherto unnoticed interrelationships which can be mathematically demonstrated.*
2. *Some of them have mathematical properties of which their proponents were unaware, and which lead to difficulties of interpretation.*
3. *As a consequence, the status of the empirical work already done with segregation indices is questionable, and their validity for further research is undermined.*

Over 70 years later, hints of this chaos remain. This author is aware of two instances in the last ten years of a segregation measure being published as novel despite already existing in the literature under a different name.

This article is not a systematic review of the literature, but rather a guided tour through one particular perspective on the problem of measuring segregation. This perspective begins with a general class of measures of diversity and segregation which

0.1

Broad motivation for the article

- Folks in the math community are turning to quantitative measures of segregation [DMW23, TBK⁺25, HSS25].

*Phil is an assistant professor of computer science at Middlebury College. His email address is pchodrow@middlebury.edu.

- There’s actually a pretty interesting and rich literature in quantitative sociology on this set of topics [Rea, RF02], going back all the way at least to [Sim49].
- MAUP is a thing [Ope84], and we’d like measures that behave in predictable ways as we change analytical scale.
- If you start engaging with these ideas systematically and mathematically, then you come naturally to convexity and Bregman divergences, which are what I am here to sell you today.

Figure 1 is a simple checkerboard pattern. What if we actually oriented the entire piece around the checkerboard¹:

Preliminaries

To make these ideas concrete, we formalize the idea of a *population* and measurements which reflect the diversity or segregation of that population. Our framework for defining populations is intentionally discrete, in recognition of the fact that all Census-type data sets provide demographic information in discrete areal units such as counties, tracts, or blockgroups.

Definition 1. For $\ell \in \mathbb{N}$, the *probability simplex* Δ_ℓ is the set

$$\Delta_\ell = \left\{ \mathbf{y} \in \mathbb{R}^\ell : \sum_{j=1}^{\ell} y_j = 1, y_j \geq 0 \text{ for all } j \in [\ell] \right\}. \quad (1)$$

¹The problem of constructing segregation measures that distinguish between panels (c) and (d) in Figure 1 is often called the “checkerboard problem.”

of probability vectors on ℓ elements.²

Definition 2. A population $\mathcal{P}$ with n locations in d spatial dimensions and ℓ demographic categories is a triple $\mathcal{P} = (\mathcal{X}, \mathbf{Y}, \mu)$, where:

- $\mathcal{X}$ is a finite metric space with n elements labeled x_i , $i = 1, \dots, n$.
- $\mathbf{Y} \in \mathbb{R}^{n \times \ell}$ is a row-stochastic matrix: for each row i , $\mathbf{y}_i \in \Delta_\ell$.
- $\mu \in \Delta_n$.

We denote the set of all populations with n locations, d dimensions, and ℓ demographic categories as $\mathcal{P}_{nd\ell}$.

In this definition, the matrix $\mathcal{X}$ represents space: element x_i of $\mathcal{X}$ can be viewed as giving the location of the i th spatial subunit of the population. The matrix $\mathbf{Y}$ represents the demographic distribution of the population according to some attribute such as gender, race, or education level: the i th row $\mathbf{y}_i$ of $\mathbf{Y}$ gives the proportion of individuals in each demographic category at location i . Finally, the probability vector μ represents the population density across locations; the proportion of individuals living in location i is μ_i . The probability that a single individual, chosen uniformly at random from the population, lives in location i and is in demographic category j is $\mu_i y_{ij}$. The mean population vector $\bar{\mathbf{y}} = \sum_{i=1}^n \mu_i \mathbf{y}_i$ has entries giving the marginal probabilities that an individual chosen uniformly at random belongs to each demographic category.

Definition 3. A population measurement is a function $f : \mathcal{P}_{nd\ell} \rightarrow \mathbb{R}$.³

An example of a population measurement is $f(\mathcal{P}) = \sum_{i=1}^n \mu_i^2$, which measures the degree to which the population is concentrated in space. For fixed n , the measure f is maximized (with value $f(\mathcal{P}) = 1$) when the entire population resides in a single location x_i and minimized (with value $f(\mathcal{P}) = \frac{1}{n}$) when the population is uniformly distributed across all locations.

²Although it is sometimes common to refer to Δ_ℓ as an $(\ell - 1)$ -simplex (this being its manifold dimension), we write Δ_ℓ to emphasize its role as the set of probability distributions over ℓ categories.

³The function η could reasonably be described as a *population measure*, which is an entirely disjoint concept from a population measurement. We will avoid the phrase “population measure” and emphasize “measurement” whenever possible.

Diversity and Convexity

Heuristically, we often say that a population is *diverse* with respect to an attribute when it contains many categories of that attribute in roughly equal proportions. Diversity is usually considered to be a property of a population as a whole, without regard for spatial or demographic subdivisions. In our formal structure from Definitions 2 and 3, a diversity measure $f : \mathcal{P}_{nd\ell} \rightarrow \mathbb{R}$ is a function of the mean population demographic vector: $f(\mathcal{P}) = \psi(\bar{\mathbf{y}})$ for some function $\psi : \Delta_\ell \rightarrow \mathbb{R}$. It remains to specify the properties that ψ should have to capture an intuitive notion of diversity.

Working in mathematical ecology, Pielou [Pie77] gives a proposed set of criteria. For $\ell \in \mathbb{N}$, let $[\ell] = \{1, 2, \dots, \ell\}$. Let $\mathbf{e}_I \in \Delta_\ell$ be the vector with 1 in the indices specified by the multiindex $I \subseteq [\ell]$ and 0 elsewhere.

1. **Uniform maximizer.** “Diversity should be maximized when all groups are present in equal proportions.” In our setup, this means that $\frac{1}{\ell} \mathbf{e}_{[\ell]} = \arg\max_{\mathbf{y} \in \Delta_\ell} \psi(\mathbf{y})$.
2. **Monotone under group addition.** “In two populations in which all groups are equally represented, the population with the greatest number of groups will be more diverse.” Mathematically, if I is a strict subset of $[\ell]$, then $\psi(\mathbf{e}_I / |I|) < \psi(\mathbf{e}_{[\ell]} / \ell)$.
3. **Additive under independent attributes.** “In a multiple classification (two or more qualitative variates) where the classifications are independent, the total diversity should equal the sum of the respective diversities.” This statement can be formalized as follows, interpreting “independent” in the sense of probability distributions: suppose that, for a given population $\mathcal{P} = (\mathcal{X}, \mathbf{Y}, \mu) \in \mathcal{P}_{nd\ell}$, there exist two other populations $\mathcal{P}_1 = (\mathcal{X}, \mathbf{U} \in \mathbb{R}^{n \times \ell_1}, \mu)$ and $\mathcal{P}_2 = (\mathcal{X}, \mathbf{V} \in \mathbb{R}^{n \times \ell_2}, \mu)$ such that $\ell = \ell_1 \ell_2$ and a bijection $\alpha : [\ell_1] \times [\ell_2] \rightarrow [\ell]$ such that $y_{ij} = u_{i\alpha_1(j)} v_{i\alpha_2(j)}$ for all $i \in [n]$ and $j \in [\ell]$. Then $f(\mathcal{P}) = f(\mathcal{P}_1) + f(\mathcal{P}_2)$.

We will soon find that properties (1) and (2) are satisfied by a broad range of functions ψ , while property (3) is more restrictive and essentially characterizes a single diversity measure.

Pielou’s proposed axioms capture some reasonable intuitions, but they don’t offer a recipe for constructing

useful diversity functions ψ . A more constructive intuition is that *diversity should increase on average when populations are mixed*. If $\mathbf{y}$ and $\mathbf{y}'$ are two population demographic vectors and $\lambda \in (0, 1)$ is a mixing parameter, we can form a new demographic vector $\mathbf{y}'' = \lambda\mathbf{y} + (1 - \lambda)\mathbf{y}'$. One mathematization of our proposed intuition is that, if $\mathbf{y} \neq \mathbf{y}'$ then

$$\psi(\mathbf{y}'') > \lambda\psi(\mathbf{y}) + (1 - \lambda)\psi(\mathbf{y}') . \quad (2)$$

Here, the lefthand side gives the diversity of the mixed population $\mathbf{y}''$, while the righthand side gives the λ -weighted average diversity of the original populations $\mathbf{y}$ and $\mathbf{y}'$. This intuition leads us directly to the topic of convexity.

Definition 4. Set $S \in \mathbb{R}^d$ is a *convex set* if for all $x, y \in S$ and $\lambda \in [0, 1]$, we have $\lambda x + (1 - \lambda)y \in S$. If S is a convex set, then $f : S \rightarrow \mathbb{R}$ is a *convex function* if for all $x, y \in S$ and $\lambda \in [0, 1]$, we have $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$. Function f is *strictly convex* if the above inequality is strict whenever $x \neq y$ and $\lambda \in (0, 1)$. Function f is (strictly) *concave* if $-f$ is (strictly) convex.

The simplex Δ_ℓ is a convex set, and so the notion of convexity applies to functions defined on Δ_ℓ . Equation (2) is precisely the condition that ψ is strictly concave. We will abuse notation slightly and refer to a population measurement $f(\mathcal{P}) = \psi(\bar{\mathbf{y}})$ as *concave* if ψ is concave.

Theorem 0.1. *If $\psi : \Delta_\ell \rightarrow \mathbb{R}$ is strictly concave, then the population measurement $f(\mathcal{P}) = \psi(\bar{\mathbf{y}})$ satisfies the monotonicity under group addition property.*

Definition 5. Function f is *symmetric* if for all $\mathbf{y} \in \Delta_\ell$ and all permutations σ of $[\ell]$, we have $f(\mathbf{y}) = f(\sigma(\mathbf{y}))$, where $\sigma(\mathbf{y})$ is the vector obtained by permuting the entries of $\mathbf{y}$ according to σ .

Theorem 0.2. *If $\psi : \Delta_\ell \rightarrow \mathbb{R}$ is a nonnegative, symmetric, and strictly concave function, then the population measurement $f(\mathcal{P}) = \psi(\bar{\mathbf{y}})$ satisfies the uniform maximizer and monotonicity under group addition properties of Pielou's axioms.*

Definition 6. A function $\psi : \Delta_\ell \rightarrow \mathbb{R}$ is *separable* if there exists a function $\phi : [0, 1] \rightarrow \mathbb{R}$ such that $\psi(\mathbf{y}) = \sum_{j=1}^\ell \phi(y_j)$ for all $\mathbf{y} \in \Delta_\ell$.

Theorem 0.3. *If a function $\psi : \Delta_\ell \rightarrow \mathbb{R}$ is nonnegative, continuous⁴, symmetric, strictly concave, separable, and satisfies the additivity under independent attributes property, then⁵*

$$\psi(\mathbf{y}) = -c \sum_{j=1}^\ell y_j \log y_j , \quad (3)$$

for some constant $c > 0$.

A proof is given is given by Chakrabarti and Chakrabarty [CC05], where additivity under independent attributes corresponds to the authors' Axiom 4.

Equation (3) is the Shannon entropy, which is a widely used measure of diversity in ecology, sociology, and information theory. For a systematic treatment of entropy as a measure of diversity, see [Lei21]. The function $\psi(\mathbf{y}) = 1 - \|\mathbf{y}\|_2^2 = 1 - \sum_{j=1}^\ell y_j^2$, the Simpson diversity index [Sim49], is another popular measure of diversity which satisfies the uniform maximizer and monotonicity under group addition properties; it is separable and symmetric but does not satisfy additivity under independent attributes. The Simpson index has an attractive physical interpretation: it is the probability that two individuals chosen uniformly at random from the population belong to different demographic categories.

Segregation

Segregation is

PC: Need some notions from the sociology literature

One natural quantitative intuition is that a population is segregated when individual locations tend to be less diverse than the population as a whole.

Definition 7. Given a concave diversity function $\psi : \Delta_\ell \rightarrow \mathbb{R}$, the *Jensen information* [BGW05] of a population $\mathcal{P} = (\mathcal{X}, \mathbf{Y}, \mu) \in \mathcal{P}_{nd\ell}$ is

$$I_\psi(\mathcal{P}) = \psi(\bar{\mathbf{y}}) - \sum_{i=1}^n \mu_i \psi(\mathbf{y}_i) . \quad (4)$$

⁴Concave functions are continuous on the interior of their domain, so this stipulation requires only that they are also continuous at the boundary.

⁵Throughout this article, we will use the notation $\log$ to denote the natural logarithm, as well as the standard convention that $0 \log 0 = 0$.

Jensen's Inequality, a standard result in convex analysis, implies that $I_\psi(\mathcal{P}) \geq 0$ for all populations $\mathcal{P}$ and concave functions ψ . If ψ is strictly concave, then $I_\psi(\mathcal{P}) = 0$ if and only if $\mathbf{y}_i = \bar{\mathbf{y}}$ for all $i \in [n]$, i.e. every location has the same demographic distribution as the population as a whole.

Figure 1 panels (a)-(c) give some intuition on how the diversity function ψ and the Jensen information I_ψ behave on a simple checkerboard pattern.

PC: place the measurements of entropy and mutual information in the titles for each panel.

Although the Jensen information eq. (4) is nonnegative, individual terms of the form $\psi(\bar{\mathbf{y}}) - \psi(\mathbf{y}_i)$ can be negative. This means that a given location's "contribution to segregation" cannot simply be measured by the difference $\psi(\bar{\mathbf{y}}) - \psi(\mathbf{y}_i)$. It would instead be convenient for analytic purposes to be able to write the Jensen information as a sum of nonnegative contributions from each location:

$$I_\psi(\mathcal{P}) = \sum_{i=1}^n \mu_i g(\mathbf{y}_i, \bar{\mathbf{y}}), \quad (5)$$

PC: Move from concave functions to functions whose negatives are convex.

Theorem 0.4 ([Cho25]). *If $\phi : \Delta_\ell \rightarrow \mathbb{R}$ is a strictly convex function which is differentiable on the relative interior of Δ_ℓ and g is a function such that eq. (5) holds for any population $\mathcal{P} \in \mathcal{P}_{ndt}$, then g is the Bregman divergence induced by ϕ :*

$$g(\mathbf{y}, \bar{\mathbf{y}}) = D_\phi(\mathbf{y} \parallel \bar{\mathbf{y}}) = \phi(\mathbf{y}) - \phi(\bar{\mathbf{y}}) - \nabla \phi(\bar{\mathbf{y}})^\top (\mathbf{y} - \bar{\mathbf{y}}). \quad (6)$$

Literally, the Bregman divergence is a measure of the difference between the value of ϕ at $\mathbf{y}$ and the value of the linearization (first-order Taylor expansion) of ϕ around $\bar{\mathbf{y}}$ evaluated at $\mathbf{y}$. The Bregman divergence can be interpreted as a measure of dissimilarity of $\mathbf{y}$ from $\bar{\mathbf{y}}$. It is always nonnegative and is zero if and only if $\mathbf{y} = \bar{\mathbf{y}}$. The Bregman divergence is not a metric, as it is not symmetric and does not satisfy the triangle inequality, although it does satisfy many other useful properties [BMDG05]. Examples of Bregman divergences include

the squared Euclidean distance $g(\mathbf{y}, \bar{\mathbf{y}}) = \|\mathbf{y} - \bar{\mathbf{y}}\|_2^2$ (corresponding to $\phi(x) = \frac{1}{2}\|x\|_2^2$) and the Kullback-Leibler divergence $g(\mathbf{y}, \bar{\mathbf{y}}) = \sum_{j=1}^\ell y_j \log(y_j/\bar{y}_j)$ (corresponding to $\phi(x) = \sum_{j=1}^\ell x_j \log x_j$).

PC: Locally quadratic, distance-like properties (foreshadowing for later)

0.2 Properties of the Jensen Information for Segregation Measurement

In an influential article, Reardon and Firebaugh [RF02] propose a set of desirable criteria for multigroup segregation measures, and assess a number of existing measures against these criteria with proofs and counterexamples. Of the six measures evaluated by the authors, exactly one of these is a Jensen information: the Shannon mutual information corresponding to $\psi(\mathbf{y}) = -\sum_{j=1}^\ell y_j \log y_j$.⁶ The authors write "We conclude that the [Shannon mutual information] is the most conceptually and mathematically satisfactory index..." due to the fact that the mutual information is the only measure to satisfy all of the criteria deemed clearly desirable by the authors. It turns out this isn't a coincidence: many desirable intuitions about segregation come "for free" from convexity.

Some of the Reardon and Firebaugh criteria are trivially satisfied by any Jensen information; for example, their *size invariance* criterion follows immediately from our construction of I_ψ as a function of *proportions* of populations distributed across groups and in space, rather than absolute counts. Here we'll highlight two nontrivial properties, and consider a third when we discuss the role of space in the next section.

Definition 8. A *migration* of a group a from location i to location j is a map $\text{mig}_{a,i \rightarrow j}^{(\epsilon)} : \mathcal{P}_{ndt} \rightarrow \mathcal{P}_{ndt}$ which takes a population $\mathcal{P} = (\mathcal{X}, \mathbf{Y}, \mu)$ and returns a new population

⁶In the segregation measurement setting the mutual information is sometimes referred to as the *Theil index* or the *information theory index*.

$\mathcal{P}' = (\mathcal{X}, \mathbf{Y}', \mu')$ such that

$$\begin{aligned}\mu' &= \mu + \epsilon(\mathbf{e}_j - \mathbf{e}_i), \\ \mathbf{y}'_i &= \frac{\mu_i}{\mu'_i} (\mathbf{y}_i - \lambda_i \tilde{\mathbf{e}}_a), \\ \mathbf{y}'_j &= \frac{\mu_j}{\mu'_j} (\mathbf{y}_j + \lambda_j \tilde{\mathbf{e}}_a), \\ \mathbf{y}'_k &= \mathbf{y}_k \text{ for all } k \notin \{i, j\}.\end{aligned}$$

Here, $\mathbf{e}_i \in \mathbb{R}^n$ is the i -th standard basis vector in the tangent space to Δ_n , $\tilde{\mathbf{e}}_a \in \mathbb{R}^l$ is the a -th standard basis vector in the tangent space to Δ_ℓ , and $\lambda_i = \frac{\epsilon}{\mu_i}$ is the fraction of the population at location i which is being moved. In this definition, it is required that $\epsilon \leq \mu_i y_{ia}$.

Definition 9. An exchange of groups a and b between locations i and j is a composition of two transfers:

$$\text{exc}_{a,b,i \leftrightarrow j}^{(\epsilon)} = \text{mig}_{a,i \rightarrow j}^{(\epsilon)} \circ \text{mig}_{b,j \rightarrow i}^{(\epsilon)}.$$

For this exchange to be defined, it is required that $\epsilon \leq \min(\mu_i y_{ia}, \mu_j y_{jb})$.

If $\mathcal{P}' = \text{exc}_{a,b,i \leftrightarrow j}^{(\epsilon)}(\mathcal{P})$, then $\mu' = \mu$ and $\mathbf{y}'_k = \mathbf{y}_k$ except in the rows

$$\mathbf{y}'_i = \mathbf{y}_i + \lambda_i(\tilde{\mathbf{e}}_b - \tilde{\mathbf{e}}_a), \quad (7)$$

$$\mathbf{y}'_j = \mathbf{y}_j + \lambda_j(\tilde{\mathbf{e}}_a - \tilde{\mathbf{e}}_b). \quad (8)$$

Definition 10. A segregation measurement S satisfies the *exchanges property of Reardon and Firebaugh* [RF02] if whenever $y_{ia} < y_{ja}$ and $y_{ib} > y_{jb}$, we have $S(\text{exc}_{a,b,i \leftrightarrow j}^{(\epsilon)}(\mathcal{P})) < S(\mathcal{P})$ for all ϵ sufficiently small.

Theorem 0.5. If ψ is separable⁷ and strictly concave, then the Jensen information I_ψ satisfies the exchanges property of Reardon and Firebaugh.⁸

⁷It is possible to weaken the separability requirement slightly; it is sufficient for ψ to have coordinatewise monotone gradients, for which it is in turn sufficient that the Hessian $\nabla^2 \phi$ be strictly diagonally dominant.

⁸Reardon and Firebaugh [RF02] also propose a *transfers property*, a stronger form of the exchanges property, and assert that the Shannon mutual information satisfies this property. This author is unable to follow the proof and has performed computational experiments suggesting the transfers property does not indeed hold for the Shannon mutual information.

Spatializing the Jensen Information: Local Information

The *modifiable areal unit problem* (MAUP) is a longstanding challenge in spatial statistics and quantitative sociology. As Openshaw [Ope84] writes of common areal delimiters like administrative borders,

The definition of these geographical objects is arbitrary and (in theory) modifiable at choice...there are no rules for areal aggregation, no standards, and no international conventions to guide the spatial aggregation process. Quite simply, the areal units (zonal objects) used in many geographical studies are arbitrary, modifiable, and subject to the whims and fancies of whoever is doing, or did, the aggregating.

Figure 1 panel (c) offers a stark illustration of the MAUP. The checkerboard pattern is composed of 32 cells, which are further arranged into 4 blocks of 8 cells each. At the pictured scale, considerable spatial segregation is visible. If, however, we instead aggregated the checkerboard into four mega-blocks of dimensions $(2 \cdot 8) \times (2 \cdot 8)$ cells each, then the resulting population would appear fully integrated, indistinguishable from panel (b). A desirable property of a segregation measure is that it should behave in a predictable fashion as the spatial scale of analysis is changed.

Definition 11 (Spatial Aggregation). Let $n' < n$ and let $\pi : [n] \rightarrow [n']$ be a surjective map. A *spatial aggregation* of a population $\mathcal{P} = (\mathcal{X}, \mathbf{Y}, \mu) \in \mathcal{P}_{ndt}$ by π is a new population $\mathcal{P}' = (\mathcal{X}', \mathbf{Y}', \mu') \in \mathcal{P}_{n'd'\ell}$ such that

$$\begin{aligned}\mu'_j &= \sum_{i \in \pi^{-1}(j)} \mu_i, \\ \mathbf{y}'_j &= \frac{1}{\mu'_j} \sum_{i \in \pi^{-1}(j)} \mu_i \mathbf{y}_i \\ \mathbf{X} &\in \mathbb{R}^{n' \times d}.\end{aligned}$$

The definition of $\mathbf{X}'$ is left intentionally left flexible; one possible choice is to set $x'_j = \frac{1}{\mu'_j} \sum_{i \in \pi^{-1}(j)} \mu_i x_i$, the μ -weighted centroid of the locations in the j th aggregated unit.

Definition 12 (Conditional Jensen Information). Let $\mathcal{P}'$ be a spatial aggregation of $\mathcal{P}$ by π . The *conditional Jensen information* of $\mathcal{P}$ given $\mathcal{P}'$ is

$$I_\psi(\mathcal{P} \mid \mathcal{P}') = \sum_{j=1}^{n'} \mu'_j \underbrace{\left(\phi(\mathbf{y}'_j) - \frac{1}{\mu'_j} \sum_{i \in \pi^{-1}(j)} \mu_i \phi(\mathbf{y}_i) \right)}_{I_\psi(\mathcal{P}_j)}, \quad (9)$$

where $\mathcal{P}_j$ is the subpopulation of $\mathcal{P}$ corresponding to the j th aggregated unit. The conditional Jensen information is the mean Jensen information of the subpopulations defined by the aggregated units, weighted by the population of each aggregated unit.

It follows that $I_\psi(\mathcal{P} \mid \mathcal{P}') \geq 0$ for all populations $\mathcal{P}$ and spatial aggregations $\mathcal{P}'$, with equality if and only if the demographic distribution is uniform within each aggregated unit. A direct calculation shows that

$$I_\psi(\mathcal{P}) = I_\psi(\mathcal{P}') + I_\psi(\mathcal{P} \mid \mathcal{P}'). \quad (10)$$

Equation (10) is a form of decomposition into “between aggregated unit” information (first term) and “within aggregated unit” information (second term). Since both terms are nonnegative, it follows that aggregation cannot increase the Jensen information.

0.3 Multiscale Segregation Measurement

It provides a useful guide for describing the scale of segregation within a population at different levels of spatial aggregation. For example,

PC: Put in a city here with blockgroup, tract, and maybe ZIP, neighborhood, or county-level aggregation and show how the decomposition works.

0.4 Regionalization

PC: Also bring in an example of generating a boundary via spatial aggregation.




Figure 1: A simple checkerboard pattern. Each of the four panels is composed of 32 cells. Panel (c) is further arranged into 4 blocks of 8 cells each.

0.5 Spatial Smoothing Kernels

0.6 Local Information

Extensions

Ordinal measures
[Cho25] [Cho17]
[BMDG05]

References

[BGW05] Arindam Banerjee, Xin Guo, and Hui Wang, *On the Optimality of Conditional Expectation as a Bregman Predictor*,




Figure 2: The smoothing bandwidth is 1 pixels. The mean local information of the checkerboard pattern is 0.228, while the mean local information of the segregated pattern is 0.054.

IEEE Transactions on Information Theory **51** (July 2005), no. 7, 2664–2669.

- [BMDG05] Arindam Banerjee, Srujana Merugu, Inderjit Dhillon, and Joydeep Ghosh, *Clustering with Bregman Divergences*, Journal of Machine Learning Research **6** (April 2005), 1705–1749.
- [CC05] C. G. Chakrabarti and Indranil Chakrabarty, *Shannon entropy: Axiomatic characterization and application*, International Journal of Mathematics and Mathematical Sciences **2005** (January 2005), no. 17, 2847–2854.
- [Cho17] Philip S. Chodrow, *Structure and information in spatial segregation*, Proceedings of the National Academy of Sciences **114** (October 2017), no. 44, 11591–11596.
- [Cho25] ———, *Equivalence of Informations Characterizes Bregman Divergences*, Entropy **27** (2025), no. 7, 766.
- [DD55] Otis Dudley Duncan and Beverly Duncan, *A Methodological Analysis of Segregation Indexes*, American Sociological Review **20** (April 1955), no. 2, 210, available at 2088328.
- [DMW23] Moon Duchin, James M Murphy, and Thomas Weighill, *Measuring segregation via analysis on graphs*, SIAM Journal on Matrix Analysis and Applications **44** (2023), no. 1, 80–105.
- [HSS25] Leah Hoogstra, Katherine Slyman, and Bjorn Sandstede, *Quantitative metrics for trait and identity distributions*, arXiv, 2025.
- [Lei21] Tom Leinster, *Entropy and diversity: The axiomatic approach*, Cambridge University Press, New York, 2021.
- [Ope84] Stan Openshaw, *The modifiable areal unit problem*, Concepts and Techniques in Modern Geography (1984).
- [Pie77] Evelyn C. Pielou, *Mathematical ecology*, 2. ed, A Wiley-Interscience Publication, Wiley, New York, 1977.
- [Rea] Sean F Reardon, *Measures of Income Segregation*.
- [RF02] Sean F. Reardon and Glenn Firebaugh, *Measures of Multi-group Segregation*, Sociological Methodology **32** (August 2002), no. 1, 33–67.
- [Sim49] E. H. Simpson, *Measurement of Diversity*, Nature **163** (April 1949), no. 4148, 688–688.
- [TBK⁺25] Chad M Topaz, Heather Z Brooks, Unchitta Kan, Björn Sandstede, and Christian Michael Smith, *Diversity, identity, and data*, The American Mathematical Monthly **132** (2025), no. 1, 36–47.

A Supplementary Proofs

Proof of Theorem 0.1. Monotonicity under group addition is a consequence of convexity and nonnegativity: if




Figure 3: Test!!

ψ is concave, then for any $I \subseteq [\ell]$ and $\lambda = \frac{|I|}{\ell}$, we have

$$\begin{aligned} \psi\left(\frac{1}{\ell}\mathbf{e}_{[\ell]}\right) &= \psi\left(\lambda\frac{1}{|I|}\mathbf{e}_I + (1-\lambda)\frac{1}{\ell-|I|}\mathbf{e}_{[\ell]\setminus I}\right) \\ &\geq \lambda\psi\left(\frac{1}{|I|}\mathbf{e}_I\right) + (1-\lambda)\psi\left(\frac{1}{\ell-|I|}\mathbf{e}_{[\ell]\setminus I}\right) \\ &\geq \psi\left(\frac{1}{\ell}\mathbf{e}_I\right), \end{aligned}$$

where the first inequality is by convexity and the second is by nonnegativity. $\square$

Proof of Theorem 0.2. To demonstrate maximization under the uniform, for any vector $\mathbf{y} \in \Delta_\ell$,

$$\mathbf{e}_{[\ell]}/\ell = \frac{1}{|\mathcal{S}|} \sum_{\sigma \in \mathcal{S}} \sigma(\mathbf{y}),$$

where $\mathcal{S}$ is the set of all permutations of $[\ell]$. It follows that

$$\begin{aligned} \psi\left(\frac{1}{\ell}\mathbf{e}_{[\ell]}\right) &= \psi\left(\frac{1}{|\mathcal{S}|} \sum_{\sigma \in \mathcal{S}} \sigma(\mathbf{y})\right) \\ &< \frac{1}{|\mathcal{S}|} \sum_{\sigma \in \mathcal{S}} \psi(\sigma(\mathbf{y})) = \psi(\mathbf{y}), \end{aligned}$$

where the inequality is by an inductive application of strict concavity and the final equality is by symmetry. $\square$

Proof of Theorem 0.5. The proof is by direct calculation. Let $\mathcal{P}' = \text{exc}_{a,b,i \leftrightarrow j}^{(\epsilon)}(\mathcal{P})$, and let $\mathbf{y}'_i$ and $\mathbf{y}'_j$ be the demographic vectors at locations i and j in $\mathcal{P}'$. Since the global

demographic vector $\bar{\mathbf{y}}$ is unchanged by the exchange, we have

$$\begin{aligned} I_\psi(\mathcal{P}) - I_\psi(\mathcal{P}') &= \sum_{i=1}^n [\mu_i \psi(\mathbf{y}_i) - \mu'_i \psi(\mathbf{y}'_i)] \\ &= \mu_i [\psi(\mathbf{y}_i) - \psi(\mathbf{y}'_i)] + \mu_j [\psi(\mathbf{y}_j) - \psi(\mathbf{y}'_j)] \\ &= \mu_i \nabla \psi(\mathbf{y}_i)^\top (\mathbf{y}_i - \mathbf{y}'_i) \\ &\quad + \mu_j \nabla \psi(\mathbf{y}_j)^\top (\mathbf{y}_j - \mathbf{y}'_j) \\ &\quad + o(\|\mathbf{y}_i - \mathbf{y}'_i\|) + o(\|\mathbf{y}_j - \mathbf{y}'_j\|). \end{aligned}$$

since $\mathbf{y}_k = \mathbf{y}'_k$ for all $k \notin \{i, j\}$ and $\mu_k = \mu'_k$ for all k . Using now that $\mathbf{y}_i - \mathbf{y}'_i = \lambda_i(\tilde{\mathbf{e}}_a - \tilde{\mathbf{e}}_b)$ and $\mathbf{y}_j - \mathbf{y}'_j = \lambda_j(\tilde{\mathbf{e}}_b - \tilde{\mathbf{e}}_a)$ and ignoring the higher-order terms, we have

$$I_\psi(\mathcal{P}) - I_\psi(\mathcal{P}') \approx \mu_i \lambda_i \nabla \psi(\mathbf{y}_i)^\top (\tilde{\mathbf{e}}_a - \tilde{\mathbf{e}}_b) + \mu_j \lambda_j \nabla \psi(\mathbf{y}_j)^\top (\tilde{\mathbf{e}}_b - \tilde{\mathbf{e}}_a).$$

By the separability assumption, we have $\nabla \psi(\mathbf{y}) = (\phi'(y_1), \dots, \phi'(y_\ell))^\top$, and so

$$\begin{aligned} I_\psi(\mathcal{P}) - I_\psi(\mathcal{P}') &\approx \mu_i \lambda_i (\phi'(y_{ia}) - \phi'(y_{ib})) + \mu_j \lambda_j (\phi'(y_{jb}) - \phi'(y_{ja})) \\ &= \epsilon(\phi'(y_{ia}) - \phi'(y_{ib}) + \phi'(y_{jb}) - \phi'(y_{ja})) \\ &= \epsilon(\phi'(y_{ia}) - \phi'(y_{ja}) + \phi'(y_{jb}) - \phi'(y_{ib})) \\ &> 0. \end{aligned}$$

The last line follows from the assumptions that $y_{ia} < y_{ja}$ and $y_{ib} > y_{jb}$, and the fact that, if ϕ is strictly convex, then ϕ' is strictly increasing, and so $\phi'(y_{ia}) - \phi'(y_{ja}) > 0$ and $\phi'(y_{jb}) - \phi'(y_{ib}) > 0$. Choosing ϵ^* sufficiently small to justify the linear approximation completes the proof. $\square$

B Supplementary Exercises

Exercise B.1. Give a heuristic justification for the convention that $0 \log 0 = 0$ by proving that $\lim_{x \rightarrow 0^+} x \log x = 0$.




Figure 4: (Top): Heatmap showing the proportion of Black residents in Detroit, delimited by nested areal units. Thickest boundaries correspond to county subdivisions, thinner boundaries to Census tracts, and thinnest boundaries to Census block groups. (Bottom): Behavior of the Shannon mutual information between Black and non-Black residents across geographic levels. If $\mathcal{P}$ is the population at the level of blockgroups (far left), the conditional Jensen information $I_{\psi}(\mathcal{P} \mid \mathcal{P}')$ is the difference between the given value of the information and the blockgroup value.