## Urban Diversity Through an Information-Theoretic Lens

Phil Chodrow July 5, 2016

MIT Human Mobility and Networks Laboratory || Operations Research Center

Introduction

Two questions for today:

Two questions for today:

1. What is **urban diversity**, and can we measure it in spatially-aware ways?

#### Two questions for today:

- 1. What is **urban diversity**, and can we measure it in spatially-aware ways?
- 2. What are **neighborhoods**, and how can we distinguish neighborhoods from each other?

#### Two questions for today:

- 1. What is **urban diversity**, and can we measure it in spatially-aware ways?
- 2. What are **neighborhoods**, and how can we distinguish neighborhoods from each other?

#### Central Claim

These two questions are deeply related, and we should approach both using *information theory*.

**Global diversity** is the extent to which different groups are equally represented in the city as a whole.

**Global diversity** is the extent to which different groups are equally represented in the city as a whole.

Spatial evenness "or clustering, refers to the extent to which groups are similarly distributed in residential space." [12]

- **Global diversity** is the extent to which different groups are equally represented in the city as a whole.
- **Spatial evenness** "or clustering, refers to the extent to which groups are similarly distributed in residential space." [12]
- Spatial exposure "refers to the extent that members of one group encounter members of another group (or their own group, in the case of spatial isolation) in their local spatial environments." [12]

- **Global diversity** is the extent to which different groups are equally represented in the city as a whole.
- **Spatial evenness** "or clustering, refers to the extent to which groups are similarly distributed in residential space." [12]
- Spatial exposure "refers to the extent that members of one group encounter members of another group (or their own group, in the case of spatial isolation) in their local spatial environments." [12]


No global diversity  $\implies$  perfect evenness  $\implies$  maximal exposure.

#### Neighborhoods, Exposure, and the Checkerboard

**Claim**: Spatial exposure is about the number, scale, and pattern of neighborhoods.

#### Neighborhoods, Exposure, and the Checkerboard

**Claim**: Spatial exposure is about the number, scale, and pattern of neighborhoods.



#### Neighborhoods, Exposure, and the Checkerboard

**Claim**: Spatial exposure is about the number, scale, and pattern of neighborhoods.



City	Exposure	Neighborhoods		
(c)	Low	Few, large		
(d)	High	Many, small		

**Diversity and Information Theory** 

#### Summary: Information Theory and Diversity

#### **Entropy Measures Diversity**

$$H(Y) \triangleq -\mathbb{E}_{Y}[\log p(Y)]$$

#### Summary: Information Theory and Diversity

#### **Entropy Measures Diversity**

$$H(Y) \triangleq -\mathbb{E}_{Y}[\log p(Y)]$$

#### Mutual Information Measures Evenness

$$I(X, Y) \triangleq \mathbb{E}_{X,Y} \left[ \log \frac{p(X,Y)}{p(X)p(Y)} \right]$$

#### Summary: Information Theory and Diversity

#### **Entropy Measures Diversity**

$$H(Y) \triangleq -\mathbb{E}_{Y}[\log p(Y)]$$

#### Mutual Information Measures Evenness

$$I(X, Y) \triangleq \mathbb{E}_{X,Y} \left[ \log \frac{p(X,Y)}{p(X)p(Y)} \right]$$

#### Mean Local Information Measures Exposure

$$J(X, Y) \triangleq \mathbb{E}_X[\text{trace } J_Y(X)]$$

#### Back to the checkerboard




City	H(Y)	I(X, Y)	J(X, Y)
(a)	0.0	0.0	0.0
(b)	0.7	0.0	0.0
(c)	0.7	0.7	0.6
(d)	0.7	0.7	2.7

#### A Three-Dimensional Characterization of Urban Diversity

City	H(Y)	I(X, Y)	J(X, Y)
Albuquerque	1.08	0.19	0.66
Detroit	1.12	0.57	1.00
Philadelphia	1.31	0.56	1.83

Data accessed from the American Community Survey of the U.S. Census [1].

#### Visualizing Diversity Profiles in Major Cities



 Philadelphia has greater exposure / more neighborhood structure than Detroit.

**Identifying Natural Neighborhoods** 

Suppose we cluster the locations X into groups with labels C.

"Additive Organizational Decomposability" (Chain Rule)

$$I(X,Y) = I(C,Y) + I(X,Y|C).$$

Suppose we cluster the locations X into groups with labels C.

"Additive Organizational Decomposability" (Chain Rule)

$$I(X,Y) = I(C,Y) + I(X,Y|C).$$

We can use this to identify neighborhoods!

Suppose we cluster the locations X into groups with labels C.

#### Recipe for Neighborhood Identification

$$I(X,Y) = \underbrace{I(C,Y)}_{\text{maximize this}} + \underbrace{I(X,Y|C)}_{\text{minimize this}}.$$


Suppose we cluster the locations X into groups with labels C.

#### Recipe for Neighborhood Identification

$$I(X,Y) = \underbrace{I(C,Y)}_{\text{maximize this}} + \underbrace{I(X,Y|C)}_{\text{minimize this}}.$$

This optimization is hard to do exactly, but we can use a greedy approach based on hierarchical clustering.

#### Practical Example: Racial Neighborhoods in Detroit



White	
Black	
Hispanic	

Asian

Other


36	2	7	2	1	1
5	29	4			
2		1			2
2				1	
2	1	1			

Percentage of population by race and neighborhood.

Wrapping Up

#### Generality and future work

- Other demographics: education level, income, occupation type, etc.
- · Other dimensions: time
  - Evolution of urban demographics (timescale: decades)
  - · Dynamics of diversity in daily mobility (timescale: minutes)



Daily commuting flows in Riyad, Saudi Arabia. Image credit: Shan Jiang and MIT HuMNetLab

#### Download our tools!

```
R Package for Information-Theoretic Analysis
https://github.com/PhilChodrow/compx
```

This project: analysis and presentations
https:
//github.com/PhilChodrow/spatial\_complexity

# THANK YOU! Questions? Feedback?

Supplementary Materials

#### References I

2014 American Community Survey 5-Year Estimates, Table B01003, 2016.

Luis M A Bettencourt, Joe Hand, and José Lobo. **Spatial Selection and the Statistics of Neighborhoods.** 2015.

Roger Bivand, Tim Keitt, and Barry Rowlingson. rgdal: bindings for the Geospatial Data Abstraction Library,

2014.

Roger Bivand and Nicholas Lewin-Koh. maptools: Tools for reading and handling spatial objects, 2014.

#### References II

Roger Bivand and Colin Rundel.

rgeos: Interface to Geometry Engine - Open Source (GEOS), 2014.

Thomas M Cover and Joy A Thomas.

Elements of Information Theory.

John Wiley and Sons, New York, 1991.

Imre Csiszar and Paul C Shields.

Information Theory and Statistics: A Tutorial.

Foundations and Trends™ in Communications and Information Theory, 1(4):417–528, 2004.

#### References III

Barret A Lee, Sean F Reardon, Glenn Firebaugh, Chard R Farrell, Stephen A Matthews, and David O'Sullivan.

Beyond the Census Tract: Patterns and Determinants of Racial Segregation at Multiple Geographic Scales.

American Sociological Review, 73(5):766–791, 2008.

Douglas S Massey and Nancy A Denton.

The Dimensions of Residential Segregation.

Social Forces, 67(2):281–315, 1988.

R Core Team.

R: A Language and Environment for Statistical Computing, 2016.

#### References IV

Sean F Reardon and G Firebaugh.

Measures of multigroup segregation.

Sociological Methodology, 32:33–67, 2002.

Sean F. Reardon and David O'Sullivan.

Measures of Spatial Segregation.

Sociological Methodology, 34(1):121–162, 2004.

#### Relation of local and Fisher informations

Suppose that p(y|x) > 0 and that p(y|x) is differentiable as a function of x for all  $x, y \in \mathbb{R}^n \times \mathcal{Y}$ . Define  $B_r \triangleq B_r(x_0) \triangleq \{x \in \mathbb{R}^n \mid |x - x_0| \leq r\}$ . Additionally, define the local mutual information in  $B_r$  as the mutual information between X and Y where X is restricted to  $B_r$ :

$$I_r(X_0) \triangleq \mathbb{E}_X[D[p(\cdot|X)||p(\cdot|X \in B_r)]|X \in B_r]$$
 (1)

$$= \int_{B_r} p(x|X \in B_r) D[p(\cdot|X)||p(\cdot|X \in B_r)] d^n x .$$
 (2)

where  $D[p||q] \triangleq \sum_{y} p(y) \log \frac{p(y)}{q(y)}$  is the Kullback-Leibler divergence of q from p.

#### Relation of local and Fisher informations

#### **Theorem**

Under the stated conditions,

$$\lim_{r \to 0} \frac{I_r(x_0)}{r^2} = \frac{n}{2(n+2)} \text{trace } J_Y(x_0) \ . \tag{3}$$

The Fisher information matrix  $J_Y$  is given by

$$J_{Y}(x) \triangleq \mathbb{E}_{Y} \left[ \nabla_{X} S_{Y}(x) \nabla_{X} S_{Y}(x)^{\mathsf{T}} \right] \tag{4}$$

$$S_y(x) \triangleq \log p(y|x)$$
 (5)

#### Information Measures Check the Boxes

Criterion	H(Y)	I(X, Y)	J(X,Y)
Organizational Equivalence	<b>/</b>	<b>/</b>	/
Size/Density Invariance	<b>/</b>	<b>/</b>	<b>/</b>
Additive Group Decomposability	/	<b>✓</b>	<b>/</b>
Additive Spatial Decomposability	NA	<b>/</b>	*
Scale Interpretability	NA	<b>/</b>	<b>/</b>
Boundary Independence	NA	<b>/</b>	<b>/</b>
Exchanges	NA	<b>/</b>	<b>/</b>

<sup>\*</sup> We contend that [12]'s version is not a workable criterion for explicitly spatial measures.

✓ In the theoretical definition; in practice we are reliant on the data as it is provided.

## Some Comparative Clusters of Equal Information

