


Sharpened CG Iteration Bound for High-contrast Heterogeneous Scalar Elliptic PDEs

Going Beyond Condition Number

P. Soliman¹

EEMCS-DIAM Numerical Analysis MSc.
Thesis Presentation, September, 2025

¹ Delft University of Technology



Opening

Darcy Problem

- Opening question: "What do X, Y, Z have in common?". (Show X, Y and Z with single picture slides and make sure to give short descriptions. Go quickly through the single-picture slides.)

Opening

Darcy Problem

- Opening question: "What do X, Y, Z have in common?". (Show X, Y and Z with single picture slides and make sure to give short descriptions. Go quickly through the single-picture slides.)
- Answer: "They can all be modelled by a..." high-contrast diffusion problem for $u, u_D \in H^1(\Omega)$, $f \in L^2(\Omega)$ and $C \in L^\infty(\Omega)$

$$\begin{aligned} -\nabla \cdot (C \nabla u) &= f \quad \text{in } \Omega \\ u &= u_D \quad \text{on } \partial\Omega \end{aligned}$$

Opening

Darcy Problem

- Opening question: "What do X, Y, Z have in common?". (Show X, Y and Z with single picture slides and make sure to give short descriptions. Go quickly through the single-picture slides.)
- Answer: "They can all be modelled by a..." high-contrast diffusion problem for $u, u_D \in H^1(\Omega)$, $f \in L^2(\Omega)$ and $\mathcal{C} \in L^\infty(\Omega)$

$$\begin{aligned} -\nabla \cdot (\mathcal{C} \nabla u) &= f \quad \text{in } \Omega \\ u &= u_D \quad \text{on } \partial\Omega \end{aligned}$$

- Understanding the problem: We need to find $u \in H^1(\Omega)$ such that the above equation holds.

Opening

Darcy Problem

- Opening question: "What do X, Y, Z have in common?". (Show X, Y and Z with single picture slides and make sure to give short descriptions. Go quickly through the single-picture slides.)
- Answer: "They can all be modelled by a..." high-contrast diffusion problem for $u, u_D \in H^1(\Omega)$, $f \in L^2(\Omega)$ and $C \in L^\infty(\Omega)$

$$\begin{aligned} -\nabla \cdot (C \nabla u) &= f \quad \text{in } \Omega \\ u &= u_D \quad \text{on } \partial\Omega \end{aligned}$$

- Understanding the problem: We need to find $u \in H^1(\Omega)$ such that the above equation holds.
- We can compute approximations $\mathbf{u}_1, \mathbf{u}_2, \dots$ efficiently using the CG method.

Opening

Darcy Problem

- Opening question: "What do X, Y, Z have in common?". (Show X, Y and Z with single picture slides and make sure to give short descriptions. Go quickly through the single-picture slides.)
- Answer: "They can all be modelled by a..." high-contrast diffusion problem for $u, u_D \in H^1(\Omega)$, $f \in L^2(\Omega)$ and $C \in L^\infty(\Omega)$

$$\begin{aligned} -\nabla \cdot (C \nabla u) &= f \quad \text{in } \Omega \\ u &= u_D \quad \text{on } \partial\Omega \end{aligned}$$

- Understanding the problem: We need to find $u \in H^1(\Omega)$ such that the above equation holds.
- We can compute approximations $\mathbf{u}_1, \mathbf{u}_2, \dots$ efficiently using the CG method.
- Main R.Q.: "How can improve existing estimates on the total number of necessary CG iterations?"

Structure

- CG method: under the hood
- Classical iteration bound
- Influence of eigenvalue distribution
- High-contrast coefficients: split eigenspectrum
- Two-cluster bound
- Multi-cluster bound
- Partitioning
- Results on sharpness
- Practical estimation of new bounds