



## **Risk of commercial fisheries to seabird populations within the New Zealand EEZ**

Yvan Richard\*, Edward Abraham\*, and Dominique Filippi†

\*Dragonfly Science

† Sextant Technology Ltd

IAPC - August 2012





## Introduction



### **New Zealand: “seabird capital of the world”**

- 85 species, many of them endemic
- Many species are listed as threatened, and their interaction with fisheries may be one of the major causes
- Level-2 risk assessment, contracted by the Ministry of Fisheries (now MPI)



# Introduction




## Level-2 risk assessment

- Built up on Sharp (2011) and Waugh (2009)
- Explicit parameter uncertainty
- Compared estimated fisheries mortality with estimated population productivity
- Only commercial trawl, bottom and surface longline fishing
- Species and fishery type level (64 species, 16 fishery groups)
- Semi-quantitative (lack of data to build population models)



# Seabird bycatch

- A lot of work on estimating annual captures
- So far on a few species with enough observations
- <http://data.dragonfly.co.nz/psc> publicly available soon
- Demographic impact?
- Poorly observed fisheries?





## Estimating fatalities

### Data

- From bird captures recorded by government observers
- In trawl, surface & bottom longline fisheries
- Data between 2003/04 and 2008/09

### Vulnerability

- Captures assumed to be proportional to the density of birds at each fishing event.
- Species vulnerability to capture, varying among fisheries.

$$C_{fgs} \sim \text{Poisson}(\mu_{fgs})$$
$$\mu_{fgs} = v_{gs} d_{fs} N_s E_f$$


$\mu_{fgs}$  mean captures for fishing event  $f$  in fishery group  $g$  and for species  $s$   
 $v_{gs}$  vulnerability of species  $s$  to captures in fishery group  $g$   
 $d_{fs} N_s$  number of birds of species  $s$  present at fishing event  $f$   
 $E_f$  fishing effort during event  $f$

# Bird density


- Used NABIS, Birdlife Tracking database, data on colonies.
- Two distributions: with and without colonies.


Example: White-capped albatross (*Thalassarche steadi*).

Without colonies



With colonies





## Potential annual fatalities

### **Estimating observable captures**

- Fitted model to predict captures on unobserved fishing events

### **Estimating potential fatalities**

- Included cryptic mortality
- Multipliers based on Watkins et al. (2008) & Brothers et al. (2010)
- Depends on type of seabirds



## Estimating population productivity

### Potential Biological Removal (PBR)

- based only on population size and maximum growth rate
- designed to maintain populations above their Maximum Net Productivity Level (MNPL)
- developed and tested for marine mammals

$$PBR = \frac{1}{2} r_{\max} N_{\min} f$$

|            |  |
|------------|--|
| $N_{\min}$ | conservative estimate of population size |
| $r_{\max}$ | maximum population growth rate           |
| $f$        | recovery factor                          |








## Maximum growth rate - $r_{\max}$

- Life history strategies, allometry. Niel & Lebreton (2005)
- Estimated from survival rate ( $S$ ) and age at first reproduction ( $A$ )

$$\lambda_{\max} =$$

$$\exp \left[ \left( A + \frac{S}{\lambda_{\max} - S} \right)^{-1} \right]$$

$$r_{\max} = \lambda_{\max} - 1$$






## Population size - $N_{\min}$

- Estimated from survival rate ( $S$ ) and age at first reproduction ( $A$ )
- $N_{\min}$  by taking the lower quartile of the distribution of  $N_{BP}$

$$\rho = S^{1-A}$$

$$N_{\text{tot}} = \frac{2N_{BP}}{P_B} \rho$$






## Potential Biological Removal (PBR)

- PBR calculated only from  $S$ ,  $A$ ,  $P_B$ , and  $f$
- Estimates of  $S$ ,  $A$ , and  $P_B$  from literature, groomed to keep best and most recent ones
- 205 final estimates, 65 using proxy species
- $f$  defined according to IUCN red list status, from 0.1 (Critically Endangered) to 0.5 (Least Concern)
- Uncertainties from literature or created to match typical values



## Species at risk



### Sensitivity to uncertainties

- Inshore fisheries poorly observed
- Adult survival rate and number of annual breeding pairs

### Time variation

- Captures in trawl fisheries has decreased, following fishing effort and the use of mitigation devices
- Possible increase in surface longline fisheries





## Limitations

Some intrinsic problems...

- Wrong species identification
- Movement in/out the NZEEZ
- A few fisheries not included
  - e.g. recreational, setnet fisheries
- Other sources of mortality not taken into account underestimate risk
  - e.g. harvest at colonies, pollution, indirect trophic effects
- PBR might often be overestimated
  - $r_{\max}$ , adult ratio



## Conclusions

- The species the most at risk are not the most caught
- Black petrel clearly at risk, urgent action needed
- Some fisheries with obvious lack of observations
  - Inshore fisheries: especially flatfish trawl, small bottom longline
- Some flaws potentially important
- Need international cooperation
- Risk assessments can guide management of research and fisheries



Thank you

## **Acknowledgments**

The observers, the Ministry of Fisheries, the members of the Aquatic Environment Working Group, Ben Sharp, Susan Waugh, Elizabeth Bell, Ralph Powlesland, Phil Bradfield, the contributors to the Birdlife Tracking Database, the Agreement on the Conservation of Albatrosses and Petrels.

Photos: Duncan Wright, Graham Robertson

This research was funded by the Ministry of Fisheries (MPI).

The report can be downloaded from:

<http://fs.fish.govt.nz/Page.aspx?pk=113&dk=22912>

