

Overview talk

Marina Logares

Higgs bundles were introduced in 1987 Nigel Hitchin. They are at the core of many mathematical and physical objects and theories. In physics they appear when working with Yang-Mills equations, mirror symmetry, topological quantum field theory, supergravity. In mathematics they appear in Teichmüller theory, integrable systems, and representations of $\pi_1(X)$.

Definition 0.1. A Higgs bundle (E, φ) where $E \rightarrow X$ is a holomorphic bundle over a compact Riemann Surface. $\varphi \in H^0(X, \text{End}(E \otimes K_X))$ where K_X is the canonical bundle of X .

Let X be an oriented Riemannian manifold, and $P \rightarrow X$ a principal G -bundle. Yang - Mills functional YM is a functional on the space of connections of the associated adjoint bundle, defined by

$$YM(A) = \int_X \text{tr}(F_A \wedge *F_A) \quad (1)$$

where $F_A = dA + [A, A]$ is the curvature of connection A , and $*$ is the Hodge star. $d_A F_A = d_A(*F_A)$ critical points of $D_A(F_A)$ are

where d_A is the covariant derivative and (1) gives the so called Yang Mills equations.

This comes from dimensional reduction on the Yang mills. In 1 dim, this leads to magnetic monopoles studied by Atiyah and Hitchin in 2 dim, get Higgs bundles via the Higgs equations.

$$\bar{\partial}\varphi = 0 \quad (2)$$

$$F_A + [\varphi, \varphi^*] = 0 \quad (3)$$

In the case that $F_A = *F_A$ is an instanton

Let (E, φ) where $E \rightarrow X$ holomorphic, and $E = H^{\mathbb{C}}$ bundle. Let G be a non-maximal component. Then $\varphi \in H^0(X, E(\mathfrak{m}^{\mathbb{C}} \otimes K))$ $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$

Then $G = U(p, q)$ and $E = V \oplus W$, $\varphi = \begin{pmatrix} 0 & \gamma \\ \beta & 0 \end{pmatrix}$

Upcoming: Non-abelian Hodge theory Symplectic structure Hitchin map Singular Higgs bundles (parabolic bundles) Topology of moduli spaces Mirror symmetry for the character varieties.

1 Non abelian Hodge

Let (E, φ) be our Higgs bundle over Riemann surface X .

$\varphi = 0$. $M(r, d)$ is the moduli space of stable vector bundles of rank r and degree d . Narasimha-Seshadri theorem.

$$M(r, 0) \cong \{\pi_1(X) \rightarrow U(n)\}/U(n) \quad (4)$$

Donaldson generalises representation 3 ingredients, v, b , connections $\rho : \pi_1(X) \rightarrow G$

$$H^1(X, G) \cong H_1(X, G) \cong \text{Hom}(\pi/[\pi, \pi], G) \cong \text{Hom}(\pi, G) \quad (5)$$

Non-abelian hodge theory consists of the following technique

$$H^1_{\Delta}(X, G) \cong H^1_{dR}(X, G) \cong H^1_{\partial}(X, G) \quad (6)$$

Relates the singular, d'Rham, and Dolbeaut cohomology. The topological, the differential, and the algebraic

2 Symplectic Geometry

Suppose $H^1(X, \mathbb{R}) = \mathbb{R}$, for X compact Riemann surface of genus g . The cup product provides $H^1(X, \mathbb{R})$ with symplectic structure. Actually, $\mathcal{M}^s(r, d)$ the moduli of stable Higgs bundles is symplectic.

$$T_E M = H^1(X, \text{End}(E)) \quad (7)$$

by Serre duality

$$T_E M^* = H^0(X, E \otimes K_X) \quad (8)$$

M^s is the moduli space of stable vector bundles Say that vector bundle $E \rightarrow X$ is stable if $\forall F \subset E$

$$\deg(F)/\text{rk}(F) < \deg(E)/\text{rk}(E) \quad (9)$$

slope condition. In the case of Higgs bundles require stronger condition

$$\forall F \subset E, \quad \varphi(F) \subset F \otimes K \quad (10)$$

3 Symplectic geometry

$$\dim M = r^2(g-1) + 1 \quad (11)$$

4 Hitchin map

($G = \text{GL}(r, \mathbb{C})$)

$$\mathcal{M}^s \supset T^* M^s \rightarrow M^s \quad (12)$$

The hitchin map

$$\mathcal{M}^s \rightarrow h \bigoplus_{i>0} H^i(X, K^i) = H \quad (13)$$

$$(E, \varphi) \mapsto \text{tr} \left(\bigwedge^i \varphi \right) \quad (14)$$

Proper map, with $\dim H = \frac{1}{2} \dim \mathcal{M}$ (cf spectral curves, mirror symmetry, and integral systems)
SYZ- mirror symmetry

5 Parabolic

[NO IDEA]

6 Topology of moduli of stable Higgs

Define a perfect morse function

$$\mu : \mathcal{M} \rightarrow \mathbb{R} \quad (15)$$

$$(E, \varphi) \mapsto \|\varphi\|^2 \quad (16)$$

$$P_t(\mathcal{M}) = \sum t^{\text{ind}} P_t(\mathcal{N}) \quad (17)$$

where \mathcal{N} are some critical somethings of something.