

Hodge theory

Marille Ong

1 Four worlds of Riemann surfaces

Definition 1.1. A Riemann surface X is a complex manifold of dimension 1.

examples: \mathbb{P}^1, \dots

Denote two sheaves of particular interest. For open subset $U \subset X$,

$$\mathcal{O}_X(U) \text{ holomorphic functions on } U \quad (1)$$

$$\mathcal{M}_X(U) \text{ meromorphic functions on } U \quad (2)$$

$$(3)$$

To be Tabulated

Topological

Classification results | Every closed orientable surface is homeomorphic to a sphere with g handles. Let M be a smooth manifold.

Example: Complex tori relate elliptic curves. $\Gamma(1, \omega)$ and j -invariants

2 Line Bundles

Definition 2.1. A holomorphic line bundle $L \rightarrow X$ is a complex vector bundle. (Axioms stated here)

The set of (holomorphic) line bundles (upto isomorphism) forms a group with tensor power acting as the binary operator. The trivial bundle is then the identity element, and inverse is given by duality. Denote this group $\text{Pic}(X)$.

Definition 2.2. Let L be a line bundle with trivialisation $\{U_i\}$ A holomorphic section s of line bundle ... (Axioms stated) A meromorphic section s of line bundle ... (Axioms stated) \mathcal{O}_L denotes the sheaf of holomorphic sections.

Definition 2.3. A sheaf \mathcal{F} on X is an invertible if for all U_i in the cover $\{U_i\}$ such that \mathcal{F}_i is a free \mathcal{O}_{U_i} module.

Theorem 2.4. There exists a canonical isomorphism

$$\text{Pic}(X) \cong H^1(X, \mathcal{O}_X^*) \quad (4)$$

And isomorphic to the group of invertible sheaf on X under \otimes .

Definition 2.5. The canonical bundle K_X is the determinant of the cotangent bundle.

Theorem 2.6. (Riemann - Roch)

$$h^0(X, \mathcal{O}_L) - h^0(X, \mathcal{O}_L^* \otimes K_X) = \deg(L) + 1 - g \quad (5)$$

Theorem 2.7. (Serre duality)

$$H^k(X, \mathcal{O}_L) \cong H^{n-k}(X, \mathcal{O}_L^* \otimes K)^* \quad (6)$$

3 Divisors

Definition 3.1. A divisor $D : X \rightarrow \mathbb{Z}$ which is 0 for all but finitely many $x \in X$. Represent D as the formal sum $\sum D(x)x$. The set of all divisors is denoted $\text{Div}(X)$. The degree of a divisor D is $\deg(D) = \sum D(x)$. For any function f have the associated divisor $\text{Div}(f) = \sum \text{ord}_x(f)x$

Divisors form a group under formal addition in a natural way. We induce on the set of divisors a partial ordering induced by \mathbb{Z} . There are notable subgroups.

Say two divisors are equivalent if they differ by a principle divisors The class group is defined as

$$\text{Cl}(X) = \frac{\text{Div}(X)}{\text{PDiv}(X)} \quad (7)$$

And the analogue for degree 0 .

For a line bundle we can define the associated divisor, which is defined upto principle divisor. That is ...

Theorem 3.2. There exists

$$\frac{\text{Div}(X)}{\text{PDiv}(X)} \cong \text{Cl}(X) \cong \text{Pic}(X) \quad (8)$$

And the 0 analogue.

Weiestrass problem. Given divisor D

4 Jacobian

Define $H^0(X, \Omega_X^1)$ space of holom 1-forms. Define

$$\lambda_C : H^0(X, \Omega_X^1) \rightarrow \mathbb{C} \quad (9)$$

$$\omega \mapsto \int_C \omega \quad (10)$$

By Stokes λ only depends on the class $[C] \in H_1(X, \mathbb{Z})$ of curve C .

Let Λ be the image of λ .

Definition 4.1. The jacobian of X is given by

$$\text{Jac}(X) = H^0(\Omega_X^1)/\Lambda \quad (11)$$

Thus we identify $H^0(X, \Omega_X^1)^*$ with \mathbb{C}^g

Definition 4.2. Fix a base point $p_0 \in X$ define

$$A : X \rightarrow \text{Jac}(X) \quad (12)$$

$$p \mapsto \int \gamma_p \omega \quad (13)$$

where γ_p is a path p_0 to p . Abel-Jacobi map.

Extend linearly to $A : \text{Div}(X) \rightarrow \text{Jac}(X)$.

Theorem 4.3. (Abel) If $D \in \text{Div}_0$ then $A_0(D) = 0$ iff $D \in \text{PDiv}(X)$

Theorem 4.4. (Jacobi) The map A_0 is surjective and $\text{Pic}_0 \cong \text{Jac}(X)$

5 Hodge theory

Let X now be compact complex manifold. The induced almost complex structure (here defined) We have the decomposition of the exterior algebra into (p, q) forms. Denote the sheaf section of $\Lambda^{p,q} X$ by $A^{p,q}(X)$

We can then decompose d into ∂ and $\bar{\partial}$

Theorem 5.1. (*Dolbeault*)

$$H^{p,q}(X) \cong H^q(X, \Omega_X^p) \quad (14)$$

Really long list defining the operators

Theorem 5.2. (*Hodge theorem*)

Theorem 5.3. $\mathcal{H}_{\bar{\partial}}^{p,q}(X) \cong H^{p,q}(X)$

Theorem 5.4. $H_{\Delta}^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ In the case where $k = 1$ $H_{\Delta}^1(X, \mathbb{C}^*) \cong \text{Jac}(X) \oplus H^0(X, \Omega_X^1)$

Hodge theory gives us the set of 1 dim representations $\pi_1(X)$ is isomorphic to the set of holomorphic line bundles with degree 0