

Quantum Optics, Homework 3

Jinyuan Wu

November 21, 2021

Interference between Gaussian pulses Consider two Gaussian pulses with wave vectors $\mathbf{k}_{1,2} = k(\pm \sin \theta, 0, \cos \theta)$, respectively. They are incident to a plane detector on the surface $z = 0$. The intensity distributions of the two beams are all

$$|\mathcal{E}|^2 \propto e^{-(x^2+y^2)/\sigma^2}, \quad (1)$$

with $\sigma \gg \lambda$. The pulses arrive at the detector simultaneously. The detector absorbs the pulses completely and there is no reflection. Calculate $P^{(1)}(\mathbf{r})$ and $P^{(2)}(\mathbf{r}_1, \mathbf{r}_2)$ for the following states of the optical field:

$$(a) |\psi\rangle = \frac{1}{\sqrt{2^N N!}} (a_1^\dagger + a_2^\dagger)^N |V\rangle.$$

$$(b) |\psi\rangle = \frac{1}{N!} (a_1^\dagger a_2^\dagger)^N |V\rangle.$$

$$(c) |\psi\rangle = \frac{1}{\sqrt{2N!}} \left((a_1^\dagger)^N + (a_2^\dagger)^N \right) |V\rangle.$$

$$(d) |\psi\rangle = D_1(\alpha) D_2(\alpha) |V\rangle, \quad D_j(\alpha) \equiv e^{\alpha a_j^\dagger - \alpha^* a_j}.$$

$$(e) |\psi\rangle = \frac{1}{\sqrt{2}} (D_1(\alpha) + D_2(\alpha)) |V\rangle.$$

Solution The electric field operator is

$$\mathbf{E} = \sum_{i=1,2} \mathcal{E}_i e^{i\mathbf{k}_i \cdot \mathbf{r} - i\omega t} a_i + \text{h.c.} \quad (2)$$

(a) We define

$$b^\dagger = \frac{1}{\sqrt{2}} (a_1^\dagger + a_2^\dagger),$$

and now the wave function is

$$|\psi\rangle = \frac{1}{\sqrt{N!}} (b^\dagger)^N |0\rangle.$$

We have

$$P^{(1)}(\mathbf{r}) = \frac{1}{N!} |\mathcal{E}(\mathbf{r})|^2 \langle 0 | b^N (a_1^\dagger a_1 + a_2^\dagger a_2 + e^{i(\mathbf{k}_2 - \mathbf{k}_1) \cdot \mathbf{r}} a_1^\dagger a_2 + e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} a_2^\dagger a_1) (b^\dagger)^N |0\rangle.$$

Evaluating the terms in the RHS above, we have

$$\begin{aligned} \langle 0 | b^N a_1^\dagger a_1 (b^\dagger)^N |0\rangle &= N \langle 0 | b a_1^\dagger |0\rangle \times N \langle 0 | a_1 b^\dagger |0\rangle \times \text{contraction of } (N-1) \text{ } b\text{'s and } (N-1) \text{ } b^\dagger\text{'s} \\ &= N \langle 0 | b a_1^\dagger |0\rangle \times N \langle 0 | a_1 b^\dagger |0\rangle \times (N-1)! \langle 0 | b b^\dagger |0\rangle \\ &= N \times \frac{1}{\sqrt{2}} \times N \frac{1}{\sqrt{2}} \times (N-1)! \times 1 = \frac{1}{2} N^2 (N-1)!, \end{aligned}$$

and similarly

$$\langle 0 | b^N a_2^\dagger a_2 (b^\dagger)^N |0\rangle = \frac{1}{2} N^2 (N-1)!,$$

and

$$\begin{aligned} \langle 0 | b^N a_1^\dagger a_2 (b^\dagger)^N |0\rangle &= N \langle 0 | b a_1^\dagger |0\rangle \times N \langle 0 | a_2 b^\dagger |0\rangle \times \text{contraction of } (N-1) \text{ } b\text{'s and } (N-1) \text{ } b^\dagger\text{'s} \\ &= N \langle 0 | b a_1^\dagger |0\rangle \times N \langle 0 | a_2 b^\dagger |0\rangle \times (N-1)! \langle 0 | b b^\dagger |0\rangle \\ &= N \times \frac{1}{\sqrt{2}} \times N \frac{1}{\sqrt{2}} \times (N-1)! \times 1 = \frac{1}{2} N^2 (N-1)!, \end{aligned}$$




Figure 1: The two Gaussian beams incident to a detector

and similarly

$$\langle 0 | b^N a_1^\dagger a_2 (b^\dagger)^N | 0 \rangle = \frac{1}{2} N^2 (N-1)!$$

Putting everything together we have

$$\begin{aligned} P^{(1)}(\mathbf{r}) &= \eta \frac{1}{N!} |\mathcal{E}(\mathbf{r})|^2 \times \frac{1}{2} N^2 (N-1)! \times (2 + e^{i(\mathbf{k}_2 - \mathbf{k}_1) \cdot \mathbf{r}} + e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}}) \\ &= \eta N |\mathcal{E}(\mathbf{r})|^2 (1 + \cos(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}) \\ &= \eta N |\mathcal{E}(\mathbf{r})|^2 (1 + \cos(2k \sin \theta x)) \\ &= 2\eta N |\mathcal{E}(\mathbf{r})|^2 \cos^2(kx \sin \theta), \end{aligned}$$

so finally

$$P^{(1)}(\mathbf{r}) = 2N |\mathcal{E}(\mathbf{r})|^2 \cos^2(kx \sin \theta) \propto 2N e^{-(x^2 + y^2)/\sigma^2} \cos^2(kx \sin \theta). \quad (3)$$

The two-photon joint probability is

$$\begin{aligned} P^{(2)}(\mathbf{r}_1, \mathbf{r}_2) &= \eta^2 \frac{1}{N!} |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \langle 0 | b^N (a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_1}) (a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_2}) \\ &\quad \times (a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_1}) (a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_2}) (b^\dagger)^N | 0 \rangle \\ &= \eta^2 \frac{1}{N!} |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \times N \langle 0 | b (a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_1}) | 0 \rangle \\ &\quad \times (N-1) \langle 0 | b (a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_2}) | 0 \rangle \\ &\quad \times N \langle 0 | (a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_1}) b^\dagger | 0 \rangle \\ &\quad \times (N-1) \langle 0 | (a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_2}) b^\dagger | 0 \rangle \\ &\quad \times \text{contraction between } N \text{ } b\text{'s and } b^\dagger\text{'s} \\ &= \eta^2 |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \frac{1}{N!} \times \frac{N}{\sqrt{2}} (e^{-i\mathbf{k}_1 \cdot \mathbf{r}_1} + e^{-i\mathbf{k}_2 \cdot \mathbf{r}_1}) \times \frac{N-1}{\sqrt{2}} (e^{-i\mathbf{k}_1 \cdot \mathbf{r}_2} + e^{-i\mathbf{k}_2 \cdot \mathbf{r}_2}) \\ &\quad \times \frac{N-1}{\sqrt{2}} (e^{i\mathbf{k}_1 \cdot \mathbf{r}_2} + e^{i\mathbf{k}_2 \cdot \mathbf{r}_2}) \times \frac{N}{\sqrt{2}} (e^{i\mathbf{k}_1 \cdot \mathbf{r}_1} + e^{i\mathbf{k}_2 \cdot \mathbf{r}_1}) \times (N-2)! \\ &= \eta^2 |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 N(N-1) (1 + \cos(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}_1) (1 + \cos(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}_2), \end{aligned}$$

so

$$\begin{aligned} P^{(2)}(\mathbf{r}_1, \mathbf{r}_2) &= 4\eta^2 |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 N(N-1) \cos^2(kx_1 \sin \theta) \cos^2(kx_2 \sin \theta) \\ &\propto 4\eta^2 e^{-(x_1^2 + x_2^2 + y_1^2 + y_2^2)/\sigma^2} N(N-1) \cos^2(kx_1 \sin \theta) \cos^2(kx_2 \sin \theta). \end{aligned} \quad (4)$$

(b) We have

$$\begin{aligned} P^{(1)}(\mathbf{r}) &= \frac{\eta}{(N!)^2} |\mathcal{E}(\mathbf{r})|^2 \langle 0 | (a_2 a_1)^N (a_1^\dagger a_1 + a_2^\dagger a_2 + e^{i(\mathbf{k}_2 - \mathbf{k}_1) \cdot \mathbf{r}} a_1^\dagger a_2 + e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} a_2^\dagger a_1) (a_1^\dagger a_2)^\dagger^N | 0 \rangle \\ &= \frac{\eta}{(N!)^2} |\mathcal{E}(\mathbf{r})|^2 \langle 0 | a_2^N a_1^N (a_1^\dagger a_1 + a_2^\dagger a_2 + e^{i(\mathbf{k}_2 - \mathbf{k}_1) \cdot \mathbf{r}} a_1^\dagger a_2 + e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} a_2^\dagger a_1) (a_1^\dagger)^N (a_2^\dagger)^N | 0 \rangle. \end{aligned}$$

Evaluating the terms in the RHS, we have

$$\begin{aligned} \langle 0 | a_1^N a_2^N a_1^\dagger a_1 (a_1^\dagger)^N (a_2^\dagger)^N | 0 \rangle &= N \langle 0 | a_1 a_1^\dagger | 0 \rangle \times N \langle 0 | a_1 a_1^\dagger | 0 \rangle \\ &\quad \times \text{contraction of } (N-1) \text{ } a_1 \text{'s and } (N-1) \text{ } a_1^\dagger \text{'s} \\ &\quad \times \text{contraction of } N \text{ } a_2 \text{'s and } N \text{ } a_2^\dagger \text{'s} \\ &= N \langle 0 | a_1 a_1^\dagger | 0 \rangle \times N \langle 0 | a_1 a_1^\dagger | 0 \rangle \times (N-1)! \langle 0 | a_1 a_1^\dagger | 0 \rangle \times N! \langle 0 | a_2 a_2^\dagger | 0 \rangle \\ &= N^2 N! (N-1)!, \end{aligned}$$

and similarly we have

$$\langle 0 | a_1^N a_2^N a_2^\dagger a_2 (a_1^\dagger)^N (a_2^\dagger)^N | 0 \rangle = N^2 N! (N-1)!.$$

Also we have

$$\begin{aligned} \langle 0 | a_2^N a_1^N a_1^\dagger a_2 (a_1^\dagger)^N (a_2^\dagger)^N | 0 \rangle &= N \langle 0 | a_1 a_1^\dagger | 0 \rangle \times N \langle 0 | a_2 a_2^\dagger | 0 \rangle \\ &\quad \times \text{contraction of } N \text{ } a_2 \text{'s, } (N-1) \text{ } a_1 \text{'s, } N \text{ } a_1^\dagger \text{'s and } (N-1) \text{ } a_2^\dagger \text{'s} \\ &= 0, \end{aligned}$$

so it vanishes, and so does $\langle 0 | a_2^N a_1^N a_2^\dagger a_1 (a_1^\dagger)^N (a_2^\dagger)^N | 0 \rangle$. Putting everything together we have

$$P^{(1)}(\mathbf{r}) = \eta \frac{1}{(N!)^2} |\mathcal{E}(\mathbf{r})|^2 \times 2 \times N^2 N! (N-1)! = 2N |\mathcal{E}(\mathbf{r})|^2,$$

so the single photon probability is

$$P^{(1)}(\mathbf{r}) = 2\eta N |\mathcal{E}(\mathbf{r})|^2 \propto 2\eta N e^{-(x^2+y^2)/\sigma^2}. \quad (5)$$

The two-photon joint probability is

$$\begin{aligned} P^{(2)}(\mathbf{r}_1, \mathbf{r}_2) &= \eta^2 \frac{1}{(N!)^2} |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \langle 0 | a_1^N a_2^N (a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_1}) (a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_2}) \\ &\quad \times (a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_1}) (a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_2}) (a_1^\dagger)^N (a_2^\dagger)^N | 0 \rangle \\ &= \eta^2 \frac{1}{(N!)^2} |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \langle 0 | a_1^N a_2^N (a_1^\dagger a_1^\dagger a_1 a_1 + a_2^\dagger a_2^\dagger a_2 a_2 \\ &\quad + a_1^\dagger a_2^\dagger a_2 a_1 e^{i(\mathbf{k}_1 \cdot \mathbf{r}_2 + \mathbf{k}_2 \cdot \mathbf{r}_1 - \mathbf{k}_1 \cdot \mathbf{r}_1 - \mathbf{k}_2 \cdot \mathbf{r}_2)} + \text{h.c.}) (a_1^\dagger)^N (a_2^\dagger)^N | 0 \rangle \\ &= \eta^2 |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \frac{1}{(N!)^2} (N^2 (N-1)^2 (N-2)! N! \times 2 \\ &\quad + N^4 (N-1)! (N-1)! (e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot (\mathbf{r}_1 - \mathbf{r}_2)} + \text{h.c.})), \end{aligned}$$

where the second equation uses the conservation of particle numbers. So we have

$$\begin{aligned} P^{(2)}(\mathbf{r}_1, \mathbf{r}_2) &= 2\eta^2 |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 (N(N-1) + N^2 \cos(\mathbf{k}_1 - \mathbf{k}_2) \cdot (\mathbf{x}_1 - \mathbf{x}_2)) \\ &= 2\eta^2 |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 (N(N-1) + N^2 \cos(2k(x_1 - x_2) \sin \theta)) \\ &\propto 2\eta^2 e^{-(x_1^2+x_2^2+y_1^2+y_2^2)/\sigma^2} (N(N-1) + N^2 \cos(2k(x_1 - x_2) \sin \theta)). \end{aligned} \quad (6)$$

(c) The single photon probability is now

$$\begin{aligned} P^{(1)} &= \eta \frac{1}{2N!} |\mathcal{E}(\mathbf{r})|^2 \langle 0 | (a_1^N + a_2^N) (a_1^\dagger a_1 + a_2^\dagger a_2 + e^{i(\mathbf{k}_2 - \mathbf{k}_1) \cdot \mathbf{r}} a_1^\dagger a_2 + e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} a_2^\dagger a_1) \\ &\quad \times ((a_1^\dagger)^N + (a_2^\dagger)^N) | 0 \rangle. \end{aligned}$$

Evaluating the terms on the RHS, we have

$$\begin{aligned}\langle 0|(a_1^N + a_2^N)a_1^\dagger a_1((a_1^\dagger)^N + (a_2^\dagger)^N)|0\rangle &= \langle 0|a_1^N a_1^\dagger a_1(a_1^\dagger)^N|0\rangle + \langle 0|a_2^N a_1^\dagger a_1(a_2^\dagger)^N|0\rangle \\ &= N \cdot N \cdot (N-1)! + N!,\end{aligned}$$

as well as

$$\langle 0|(a_1^N + a_2^N)a_2^\dagger a_2((a_1^\dagger)^N + (a_2^\dagger)^N)|0\rangle = N \cdot N \cdot (N-1)! + N!.$$

The third term and fourth term vanish because the photon numbers in the bra and the ket is not the same. So we have

$$\begin{aligned}P^{(1)}(\mathbf{r}) &= \eta \frac{1}{2N!} |\mathcal{E}(\mathbf{r})|^2 \times 2 \times (N^2(N-1)! + N!) \\ &= \eta(N+1) |\mathcal{E}(\mathbf{r})|^2 \propto (N+1) e^{-(x^2+y^2)/\sigma^2}.\end{aligned}\tag{7}$$

The two-photon joint probability is

$$\begin{aligned}P^{(2)}(\mathbf{r}_1, \mathbf{r}_2) &= \eta^2 \frac{1}{2N!} |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \langle 0|(a_1^N + a_2^N)(a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_1}) \\ &\quad \times (a_1^\dagger e^{-i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2^\dagger e^{-i\mathbf{k}_2 \cdot \mathbf{r}_2})(a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_1} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_1}) \\ &\quad \times (a_1 e^{i\mathbf{k}_1 \cdot \mathbf{r}_2} + a_2 e^{i\mathbf{k}_2 \cdot \mathbf{r}_2})((a_1^\dagger)^N + (a_2^\dagger)^N)|0\rangle \\ &= \eta^2 \frac{1}{2N!} |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \langle 0|(a_1^N + a_2^N)(a_1^\dagger a_1^\dagger a_1 a_1 + a_2^\dagger a_2^\dagger a_2 a_2)((a_1^\dagger)^N + (a_2^\dagger)^N)|0\rangle \\ &= \eta^2 \frac{1}{2N!} |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \times N^2(N-1)! \times 2,\end{aligned}$$

where the second equation uses conservation of particle numbers. So we have

$$\begin{aligned}P^{(2)}(\mathbf{r}_1, \mathbf{r}_2) &= \eta^2 N(N-1) |\mathcal{E}(\mathbf{r}_1)|^2 |\mathcal{E}(\mathbf{r}_2)|^2 \\ &\propto \eta^2 N(N-1) e^{-(x_1^2+x_2^2+y_1^2+y_2^2)/\sigma^2}.\end{aligned}\tag{8}$$

(d) We have

$$\begin{aligned}P^{(1)}(\mathbf{r}) &= \eta |\mathcal{E}(\mathbf{r})|^2 \langle \alpha, \alpha | (a_1^\dagger a_1 + a_2^\dagger a_2 + e^{i(\mathbf{k}_2 - \mathbf{k}_1) \cdot \mathbf{r}} a_1^\dagger a_2 + e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} a_2^\dagger a_1) | \alpha, \alpha \rangle \\ &= \eta |\mathcal{E}(\mathbf{r})|^2 (2 + 2 \cos(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}) \\ &= \eta |\mathcal{E}(\mathbf{r})|^2 \cos^2(kx \sin \theta) \propto \eta e^{-(x^2+y^2)/\sigma^2} \cos^2(kx \sin \theta).\end{aligned}\tag{9}$$

Similarly by the definition of coherent states we have

$$P^{(2)}(\mathbf{r}_1, \mathbf{r}_2) = P^{(1)}(\mathbf{r}_1) P^{(1)}(\mathbf{r}_2).\tag{10}$$

(e) We have

$$\begin{aligned}P^{(1)}(\mathbf{r}) &= \frac{1}{2} \eta |\mathcal{E}(\mathbf{r})|^2 (\langle \alpha, 0 | + \langle 0, \alpha |) (a_1^\dagger a_1 + a_2^\dagger a_2 \\ &\quad + e^{i(\mathbf{k}_2 - \mathbf{k}_1) \cdot \mathbf{r}} a_1^\dagger a_2 + e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} a_2^\dagger a_1) (| \alpha, 0 \rangle + | 0, \alpha \rangle) \\ &= \frac{1}{2} \eta (|\alpha|^2 + |\alpha|^2),\end{aligned}$$

so

$$P^{(1)}(\mathbf{r}) = \eta |\mathcal{E}(\mathbf{r})|^2 |\alpha|^2 \propto \eta e^{-(x^2+y^2)/\sigma^2} |\alpha|^2.\tag{11}$$

Similarly, all terms involving both a_1 and a_2 vanish because either of them gives 0 when acting on $|\alpha, 0\rangle$ or $|0, \alpha\rangle$, and we just have (10) as well.




Figure 2: Surface profile measuring using lasers

Laser surface measurement

(a) Consider the Michaelson interferometer in Figure 2. Suppose that there is a step on the test surface with height $h \ll \lambda$, and that the step has no scattering effects and there is no interference between the left and the right reflected light beam. Describe the output, and estimate the necessary power P to achieve $\delta h/h = 0.1$ within time duration T .

(b) Replace the laser by a series of single photon pulses.

(c) Replace the laser by a thermal light source where

$$\bar{n} = \frac{1}{e^{\beta\hbar\omega} - 1} \gg 1. \quad (12)$$

Discuss the relation between this case and the case of coherent light.

Solution Consider Figure 3. The transformation matrix of the light propagating in the space is

$$\begin{pmatrix} e^{i\varphi/2} & \\ & e^{-i\varphi/2} \end{pmatrix},$$

where

$$\varphi = k(x_1 - x_2) =: kx \ll 1. \quad (13)$$

The transformation matrix is

$$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} e^{i\varphi/2} & \\ & e^{-i\varphi/2} \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} \cos \varphi/2 & -i \sin \varphi/2 \\ -i \sin \varphi/2 & \cos \varphi/2 \end{pmatrix}.$$

Therefore we have

$$\begin{pmatrix} b_1^\dagger \\ b_2^\dagger \end{pmatrix} = \begin{pmatrix} \cos \varphi/2 & -i \sin \varphi/2 \\ -i \sin \varphi/2 & \cos \varphi/2 \end{pmatrix} \begin{pmatrix} a_1^\dagger \\ a_2^\dagger \end{pmatrix}. \quad (14)$$

By detecting $\langle b_2^\dagger b_2 \rangle$ i.e.

$$\begin{aligned} \langle n_2 \rangle &= \langle b_2^\dagger b_2 \rangle = \langle (-i \sin \varphi/2 a_1^\dagger + \cos \varphi/2 a_2^\dagger)(i \sin \varphi/2 a_1 + \cos \varphi/2 a_2) \rangle \\ &= \sin^2 \varphi/2 \langle a_1^\dagger a_1 \rangle \end{aligned} \quad (15)$$

we can measure x . Also we have

$$\begin{aligned} \langle n_2^2 \rangle &= \langle ((-i \sin \varphi/2 a_1^\dagger + \cos \varphi/2 a_2^\dagger)(i \sin \varphi/2 a_1 + \cos \varphi/2 a_2))^2 \rangle \\ &= \langle (\sin^2 \varphi/2 a_1^\dagger a_1 - i \sin \varphi/2 \cos \varphi/2 a_1^\dagger a_2)(\sin^2 \varphi/2 a_1^\dagger a_1 + i \sin \varphi/2 \cos \varphi/2 a_2^\dagger a_1) \rangle \\ &= \sin^4 \varphi/2 \langle a_1^\dagger a_1 a_1^\dagger a_1 \rangle + \sin^2 \varphi/2 \cos^2 \varphi/2 \langle a_1^\dagger a_2 a_2^\dagger a_1 \rangle \\ &= \sin^4 \varphi/2 \langle (a_1^\dagger)^2 a_1^2 \rangle + \sin^4 \varphi/2 \langle a_1^\dagger a_1 \rangle + \sin^2 \varphi \cos^2 \varphi/2 \langle a_1^\dagger a_1 \rangle \\ &= \sin^4 \varphi/2 \langle (a_1^\dagger)^2 a_1^2 \rangle + \sin^2 \varphi/2 \langle a_1^\dagger a_1 \rangle, \end{aligned}$$

and the error is

$$\delta n_2 = \sqrt{\sin^4 \varphi/2 \langle (a_1^\dagger)^2 a_1^2 \rangle + \sin^2 \varphi/2 \langle a_1^\dagger a_1 \rangle - \sin^4 \varphi/2 \langle a_1^\dagger a_1 \rangle^2}. \quad (16)$$

(a) For a coherent input on a_1^\dagger mode, the measurement result is

$$\langle n_2 \rangle = |\alpha|^2 \sin^2 \varphi/2. \quad (17)$$

By (16), the fluctuation of $\langle n_2 \rangle$ is

$$\delta n_2 = |\alpha| \sin \varphi/2,$$

and since φ is small, we have $\langle n_2 \rangle \propto \varphi^2$, and therefore

$$\frac{\delta \varphi}{\varphi} = \frac{1}{2} \frac{\delta n_2}{\langle n_2 \rangle} = \frac{1}{2|\alpha| \sin \varphi/2}. \quad (18)$$

Note that

$$|\alpha|^2 = \langle \text{numbers output photons} \rangle = \frac{PT}{\hbar \omega},$$

and again by using the fact that φ is small, we obtain

$$\delta \varphi \approx \sqrt{\frac{\hbar \omega}{PT}} = \sqrt{\frac{2\pi \hbar c}{PT\lambda}}. \quad (19)$$

Now we want to measure h , which is

$$h = \frac{\varphi_L - \varphi_R}{k}, \quad (20)$$

so we have

$$\begin{aligned} \frac{\delta h}{h} &= \frac{\sqrt{\delta \varphi_L^2 + \delta \varphi_R^2}}{kh} = \frac{\sqrt{2} \delta \varphi}{kh} \\ &= \frac{\lambda}{2\pi h} \sqrt{\frac{4\pi \hbar c}{PT\lambda}}, \end{aligned} \quad (21)$$

where $\varphi = \varphi_L \approx \varphi_R$. The condition $\delta h/h < 0.1$ is equivalent to

$$P > \frac{100\hbar\lambda c}{\pi T h^2}. \quad (22)$$

(b) This time the input state is

$$|\psi\rangle = a_1^\dagger |0\rangle, \quad (23)$$

so

$$\langle n_2 \rangle = \sin^2 \varphi/2, \quad (24)$$

and

$$\delta n_2 = \sin \varphi/2. \quad (25)$$

Therefore, after N pulses being measured we have

$$\frac{\delta \varphi}{\varphi} = \frac{1}{\sqrt{N}} \frac{1}{2 \sin \varphi/2}. \quad (26)$$

It can be seen that the form of the equation is the same as (18). Since the derivation in (a) after (18) has nothing to do with the exact meaning of $|\alpha|$, everything should be same for (18) and (26), and we have

$$P > \frac{100\hbar\lambda c}{\pi T h^2} \quad (27)$$

again.




Figure 3: A standard Michaelson interferometer

(c) For a thermal optical field, we have

$$\langle n_2 \rangle = \bar{n} \sin^2 \varphi / 2, \quad (28)$$

and

$$\begin{aligned} \langle (a_1^\dagger)^2 a_1^2 \rangle &= \sum_{n=0}^{\infty} \frac{\bar{n}^n}{(1 + \bar{n})^{1+n}} \langle n | (a_1^\dagger)^2 a_1^2 | n \rangle \\ &= \sum_{n=0}^{\infty} \frac{\bar{n}^n}{(1 + \bar{n})^{1+n}} n(n-1) \\ &= \frac{\alpha^2}{1 + \bar{n}} \sum_{n \geq 2} \alpha^{n-2} n(n-1) \quad (\alpha := \frac{\bar{n}}{1 + \bar{n}}) \\ &= \frac{\alpha^2}{1 + \bar{n}} \frac{d^2}{d\alpha^2} \sum_{n \geq 0} \alpha^n \\ &= \frac{\alpha^2}{1 + \bar{n}} \frac{d^2}{d\alpha^2} \frac{1}{1 - \alpha} = \frac{\alpha^2}{1 + \bar{n}} \frac{2}{(1 - \alpha)^3} \\ &= 2\bar{n}^2, \end{aligned}$$

so

$$\begin{aligned} \delta n_2 &= \sqrt{\sin^4 \varphi / 2 \cdot 2\bar{n}^2 + \sin^2 \varphi / 2 \bar{n} - \sin^4 \varphi / 2 \bar{n}^2} \\ &= \sin \varphi / 2 \sqrt{\bar{n}^2 \sin^2 \varphi / 2 + \bar{n}}. \end{aligned} \quad (29)$$

From (28) and the fact that φ is small we also have

$$\frac{\delta \varphi}{\varphi} = \frac{1}{2} \frac{\delta n_2}{\langle n_2 \rangle} = \frac{1}{2 \sin \varphi / 2} \sqrt{\sin^2 \varphi / 2 + \frac{1}{\bar{n}}}.$$

Since \bar{n} is large, we have

$$\frac{\delta \varphi}{\varphi} = \frac{1}{2}. \quad (30)$$

It can be seen that the precision of a single measurement cannot be improved unboundedly even when considering solely the shot noise. Using a thermal light source is not an effective idea.

Michaelson atomic clock The Michaelson interferometer (see again Figure 3) can also be used to measure photon frequency when $\Delta x = x_1 - x_2$ is already known. Derive its precision and compare the result with the Ramsey atomic clock.

Solution We can reuse results in the last problem. We have

$$\omega = \frac{c}{\Delta x} \varphi, \quad (31)$$

and φ is measured from $\langle n_2 \rangle$. If the input light is laser, we have (18), and from (31) we have

$$\frac{\delta\omega}{\omega} = \frac{\delta\varphi}{\varphi} = \frac{1}{2|\alpha| \sin\left(\frac{\omega\Delta x}{2c}\right)}, \quad (32)$$

or

$$\delta\omega \approx \frac{\omega}{2|\alpha| \frac{\omega\Delta x}{2c}} = \frac{1}{|\alpha| \Delta x/c}. \quad (33)$$

This is similar to the case in the Ramsey atomic clock, which is

$$\delta\omega = \frac{1}{\sqrt{NT}}, \quad T = \Delta x/v, \quad (34)$$

but here v is the speed of atoms instead of light. Therefore using Michaelson interferometer as a atomic clock is not a good idea since the precision is poor compared to a Ramsey atomic clock.