

# QFT I, Homework 4

Jinyuan Wu

December 7, 2021

**Scalar QED** Consider the theory of a complex scalar field  $\phi$  interacting with the electromagnetic field  $A^\mu$ . The Lagrangian is

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (D_\mu\phi)^* D^\mu\phi - m^2\phi^*\phi. \quad (1)$$

where  $D_\mu = \partial_\mu + ieA_\mu$  is the usual gauge covariant derivative.

(a) Show the Lagrangian is invariant under the gauge transformations

$$\phi(x) \rightarrow e^{-i\alpha(x)}\phi(x), \quad A_\mu(x) \rightarrow A_\mu(x) + \frac{1}{e}\partial_\mu\alpha(x). \quad (2)$$

(b) Derive the Feynman rules for the interaction between photons and scalar particles.

(c) Draw all the leading-order Feynman diagrams and compute the amplitude for the process  $\gamma\gamma \rightarrow \phi\phi^*$ .

(d) Compute the differential cross section  $d\sigma/d\cos\theta$ . You can take an average over all initial state polarizations. For simplicity, you can restrict your calculation in the limit  $m = 0$ .

(e) Draw all leading order Feynman diagrams, that contribute to the Compton scattering process  $\gamma\phi \rightarrow \gamma\phi$  and compute the differential cross section  $d\sigma/d\cos\theta$  with  $m = 0$ .

## Solution

(a) Under the gauge transformation (2), we have

$$F_{\mu\nu} \rightarrow F'_{\mu\nu} = \partial_\mu A'_\nu - \partial_\nu A'_\mu = \partial_\mu \left( A_\nu + \frac{1}{e}\partial_\nu\alpha \right) - \partial_\nu \left( A_\mu + \frac{1}{e}\partial_\mu\alpha \right) = \partial_\mu A_\nu - \partial_\nu A_\mu = F_{\mu\nu},$$

so the first term in (1) remains the same. It is obvious that under (2)

$$\phi^*\phi \rightarrow \phi'^*\phi' = e^{i\alpha}\phi^*e^{-i\alpha}\phi = \phi^*\phi,$$

so the third term in (1) is also invariant. Also we have

$$\begin{aligned} D^\mu\phi &\rightarrow (\partial^\mu + ieA'^\mu)\phi' = (\partial^\mu + ieA^\mu + i\partial^\mu\alpha)e^{-i\alpha}\phi \\ &= e^{-i\alpha}(\partial^\mu - i\partial^\mu\alpha + ieA^\mu + i\partial^\mu\alpha)\phi \\ &= e^{-i\alpha}D^\mu\phi, \end{aligned}$$

and also

$$(D^\mu\phi)^* = e^{i\alpha}D^\mu\phi^*,$$

so  $D^\mu\phi(D^\mu\phi)^*$  is also invariant. Therefore (1) is invariant under (2).

(b) We make the following expansion of Fourier transformation. For the complex scalar field we have

$$\phi(x) = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (a_{\mathbf{p}}e^{-ip\cdot x} + b_{\mathbf{p}}^\dagger e^{ip\cdot x}). \quad (3)$$

which was proved in (10) in Homework 2. The vector field is expanded as

$$A_\mu(x) = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} \sum_{r=1}^2 \epsilon_\mu^r(\mathbf{p}) (a_{\mathbf{p},r}^\dagger e^{ip\cdot x} + a_{\mathbf{p},r} e^{-ip\cdot x}). \quad (4)$$


Expanding (2) we have

$$\mathcal{L} = \mathcal{L}_{\text{scalar}} + \mathcal{L}_{\text{vector}} + \mathcal{L}_{\text{scalarQED}}, \quad (5)$$

where  $\mathcal{L}_{\text{scalar}}$  and  $\mathcal{L}_{\text{vector}}$  are Lagrangians of free scalar field and free massless vector field, and


$$\begin{aligned}\mathcal{L}_{\text{scalarQED}} &= (D_\mu \phi)^* D^\mu \phi - (\partial_\mu \phi)^* \partial^\mu \phi \\ &= e^2 \eta_{\mu\nu} A^\mu A^\nu \phi^* \phi - ie A_\mu \phi^* \partial^\mu \phi + ie \partial_\mu \phi^* A^\mu \phi.\end{aligned}\quad (6)$$

The first term has no derivatives. Therefore it gives the following (momentum space) vertex:



$$= 2ie^2 \eta_{\mu\nu}, \quad (7)$$

where the factor  $i$  comes from the time evolution operator and the factor 2 comes from the fact that there are two identical photon lines. The two  $\phi$  lines can be any of the following four:




The second term gives

$$-ie A_\mu \phi^* \partial^\mu \phi \sim -ie A_\mu (a_{\mathbf{p}}^\dagger e^{ip \cdot x} + b_{\mathbf{p}} e^{-ip \cdot x}) (-i(p' \cdot x) a_{\mathbf{p}'} e^{-ip' \cdot x} + i(p' \cdot x) b_{\mathbf{p}'}^\dagger e^{ip' \cdot x}),$$

and the third term is its complex conjugate. Therefore, the  $a^\dagger a$  term in the Lagrangian is


$$\sim -e(p_1 + p_2)_\mu A^\mu a_{\mathbf{p}_1}^\dagger a_{\mathbf{p}_2},$$

so after adding the  $i$  factor from the time evolution operator we have



$$= -ie(p_\mu + q_\mu), \quad (8)$$

and we can change the direction of a momentum line and a  $\phi$ -particle line arbitrarily; if a momentum line goes in contrast to the corresponding particle line, then we need to add a minus sign to the corresponding momentum. For example we have




$$= ie(p_\mu + q_\mu). \quad (9)$$

There are four vertices in this type in total.

#### Note


Here we follow the notation of Peskin, i.e. using the *momentum* arrow to denote whether this line represents creation or annihilation and using the arrow *on* a particle line to show whether this line represents a particle (if the direction of the particle line is parallel to the direction of the momentum line) or a antiparticle (otherwise). The real direction of a 4-momentum is *not* represented in any arrow.

(c) We enumerate over all possible diagrams. The vertex (7) itself is a diagram:




$$= (\epsilon^\sigma)^\mu (\epsilon^{\sigma'})^\nu \times 2ie^2 \eta_{\mu\nu} =: i\mathcal{M}_4. \quad (10)$$

Combining two (8)-type vertices we have a  $t$ -channel



$$\begin{aligned} &= \epsilon_\mu^\sigma \epsilon_\nu^{\sigma'} \times ie(p - (k - p))^\mu \times ie(-(k - p) - q)^\nu \times \frac{i}{(k - p)^2 - m^2 + i0^+} \\ &= -ie^2 \epsilon_\mu^\sigma (2p - k)^\mu \epsilon_\nu^{\sigma'} (-k + p - q)^\nu \frac{1}{(k - p)^2 - m^2 + i0^+} =: i\mathcal{M}_t, \end{aligned} \quad (11)$$

and a  $u$ -channel



$$\begin{aligned} &= \epsilon_\mu^\sigma \epsilon_\nu^{\sigma'} \times ie(p - (q - k))^\nu \times ie(-q - (q - k))^\mu \times \frac{i}{(q - k)^2 - m^2 + i0^+} \\ &= ie^2 \epsilon_\mu^\sigma (2q - k)^\mu \epsilon_\nu^{\sigma'} (p - q + k)^\nu \frac{1}{(q - k)^2 - m^2 + i0^+} =: i\mathcal{M}_u. \end{aligned} \quad (12)$$

#### Note

We *do not* need to distinguish the direction of the  $q'$  momentum line. This line can be either a particle line or an antiparticle line, but since the ordinary propagator  $i/(p^2 - m^2 + i0^+)$  is obtained by summing up the two cases, when we write down this propagator, we have automatically considered both processes.

Summing everything up, we have

$$\begin{aligned} i\mathcal{M}(\gamma\gamma \rightarrow \phi\phi^*) &= i(\mathcal{M}_4 + \mathcal{M}_t + \mathcal{M}_u) \\ &= ie^2 (\epsilon^\sigma)^\mu (\epsilon^{\sigma'})^\nu \left( 2\eta_{\mu\nu} + \frac{(k - 2p)_\mu (k' - 2q)_\nu}{t - m^2} + \frac{(k - 2q)_\mu (k' - 2p)_\nu}{u - m^2} \right), \end{aligned} \quad (13)$$

where

$$t = (k - p)^2, \quad u = (q - k)^2. \quad (14)$$

(d) The massless limit can be calculated with Eq. (4.85) in Peskin, which is

$$\left( \frac{d\sigma}{d\Omega} \right)_{\text{CM}} = \frac{|\mathcal{M}|^2}{64\pi^2 E_{\text{CM}}^2}, \quad (15)$$

What we need is  $|\mathcal{M}|^2$ . We have