

# **Phylogenetic Biology**

## **Week 13**

Biology 1425

Professor: Casey Dunn, [dunnlab.org](http://dunnlab.org)

Brown University

2013

# Front matter...

All original content in this document is distributed under the following license:



Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License  
([http://creativecommons.org/licenses/by-nc-sa/3.0/deed.en\\_US](http://creativecommons.org/licenses/by-nc-sa/3.0/deed.en_US))


See sources for copyright of non-original content

# Sources


Some non-original content is drawn from:



Baum, D and S. Smith (2012) Tree Thinking: and Introduction to Phylogenetic Biology. Roberts and Company Publishers. ISBN 9781936221165




Swofford, D. L., Olsen, G. J., Waddell, P. J., & Hillis, D. M. (1996). Phylogenetic inference. In: Molecular Systematics, Second Edition. eds: D. M. Hillis, C Moritz, & B. K. Mable. Sinauer Associates. ISBN 9780878932825



Felsenstein, J. (2003) Inferring Phylogenies. Sinauer Associates. ISBN 978-0878931774


Other non-original content is referenced by url.

# Phylogenies in community biogeography



Baum and Smith 2012, Figure 11.1

# Phylogenies in community ecology




When searching for a tree under a molecular clock, can just apply the additional constraint that the tree is ultrametric.

Clock models use node depth as a parameter, there are  $n-1$  internal nodes.

Non-clock models use branch length as a parameter. There are  $2n-3$  branches

So clock models are actually simpler.





Clock models are often found to be violated - there really are changes in rates.

Relaxed clock models can accommodate these rate changes.

To determine actual node ages,  
need fossils...