Outline

- HH Model Revisited
- Nonlinear Dynamics in HH Model

HH Model Revisited

Introduction



(figure from Internet)

In HH model, we use 3 additional ODEs to replace the 2 imposed eqs of IF model. And its first ODE is also gotten from its circuit diagram.

$$
\begin{aligned}
I &= C_m \frac{\mathrm{d}V_m}{\mathrm{d}t} + \bar{g}_{\mathrm{K}} n^4 (V_m - V_K) + \bar{g}_{\mathrm{Na}} m^3 h (V_m - V_{Na}) + \bar{g}_l (V_m - V_l) \\
\frac{dn}{dt} &= \alpha_n(V_m)(1-n) - \beta_n(V_m) n \\
\frac{dm}{dt} &= \alpha_m(V_m)(1-m) - \beta_m(V_m) m \\
\frac{dh}{dt} &= \alpha_h(V_m)(1-h) - \beta_h(V_m) h
\end{aligned}
$$

(figure from Internet)

Physical Perspective

- What is the meaning of α and β ?
  - Transition probability in the 2 state Markov chain.
- Why the exponent is $4, 3, 1$ for n, m, h in the first ODE?
  - They come from fitting the experiment data.

Biological Perspective

There are a few things you need to know:

- How to calculate the V_K and V_{Na}? (Sometimes they are called Nernst Equilibrium Potential)
  - Method 1: using diffusion equation
    - 

  - Method 2: using Boltzmann distribution
- What is the quantity of Nernst Equilibrium Potential of typical ions?



Equilibrium Potentials

$$
\begin{aligned}
Na^+ \quad & 62 \log \tfrac{145}{5} = 90 \text{ mV} \\
& 62 \log \tfrac{145}{15} = 61 \text{ mV} \\
K^+ \quad & 62 \log \tfrac{5}{140} = -90 \text{ mV} \\
Cl^- \quad & -62 \log \tfrac{110}{4} = -89 \text{ mV} \\
Ca^{2+} \quad & 31 \log \tfrac{2.5}{10^{-4}} = 136 \text{ mV} \\
& 31 \log \tfrac{5}{10^{-4}} = 146 \text{ mV}
\end{aligned}
$$

Ion	Concentration outside (in mM)	Concentration inside (in mM)	Ratio Out : In	E_{ion} (at 37∞C)
K^+	5	100	1 : 20	−80 mV
Na^+	150	15	10 : 1	62 mV
Ca^{2+}	2	0.0002	10,000 : 1	123 mV
Cl^-	150	13	11.5 : 1	−65 mV

- What are the concentration of Na^+, K^+, Cl^-, Ca^{2+} inside the neuron and outside the neuron?
  - see above.
- How much will they change during the action potential?
  - K^+
    - **For a cell with a 50 μm diameter, containing 100 mM K^+, it can be calculated that the concentration change required to take the membrane from 0 to -80 mV is about 0.00001 mM.**

      That is, when the channels were inserted and the K^+ flowed out until equilibrium was reached, the internal K^+ concentration went from 100 to 99.99999.



  - Ca^{2+}: ???
- What does the $Na^+ - K^+$ *pump* do in the action potential? Is it the same thing as the ion channel?
  - hyper-polarization.
  - no.
- 3 ways for Na^+, K^+ to pass the membrane
  - leaky
  - ion channel
  - ion exchanger/ion pump
- Why HH only used Na^+ and K^+?



(Figure 12.7 of *Biological Physics*)



left panel: an axon filled with potassium sulfate solution.

right panel: an normal axon.

(Figure 12.13 of *Biological Physics*)

Nonlinear Dynamics in HH Model

Introduction

Eg1: harmonic oscillator



Eg2: simple pendulum

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \pm\sqrt{2a^2\cos x - C_1}.$$



图 5-5

Fixed Point

- nullcline
- vector field

Limit Cycle

- Poincare-Bendixson TH