

Contents

1	Numeric series	3
1.1	Series convergence	4
1.2	Non-negative terms series	4
1.3	Alternating series	6
1.4	Absolute convergence and rearrangement of series	6
2	Sequences and series of functions	9
2.1	Sequences of functions	10
2.2	Series of functions	10
2.3	Power series	12
2.4	Stone-Weierstraßapproximation theorem	13
3	Improper integrals	15
3.1	Locally integrable functions	16
3.2	Improper integrals of non-negative functions	16
3.3	Absolute convergence of improper integrals	17
3.4	Differentiation under integral sign	17
3.5	Gamma function	19
4	Fourier series	21
4.1	Preliminaries	22
4.2	Orthogonal and orthonormal systems of functions	22
4.3	Fourier coefficients. Fourier series	23
4.3.1	Exponential form	23
4.3.2	Fourier series in terms of sine and cosine	24
4.3.3	Fourier series in terms of sine or cosine	24
4.3.4	Change of period	24
4.4	Punctual convergence of Fourier series	25
4.4.1	Dirichlet nucleus	25

Chapter 1

Numeric series

1.1 Series convergence

Definition 1.1.1. Let (a_n) be a sequence of real numbers. A *numeric series* is an expression of the form

$$\sum_{n=1}^{\infty} a_n.$$

We call a_n *general term of the series* and $S_N = \sum_{n=1}^N a_n$, for all $N \in \mathbb{N}$, *N-th partial sum of the series*¹.

Definition 1.1.2. We say the series $\sum a_n$ is *convergent* if the sequence of partial sums is convergent, that is, if $S = \lim_{N \rightarrow \infty} S_N$ exist and it's finite. In that case, S is called the *sum of the series*. If the previous limit doesn't exist or it is infinite we say the series is *divergent*².

Proposition 1.1.1. Let (a_n) be a sequence such that $\sum a_n < \infty$. Then $\forall \varepsilon > 0 \exists n_0 \in \mathbb{N}$ such that

$$\left| \sum_{n=1}^N a_n - \sum_{n=1}^{\infty} a_n \right| < \varepsilon$$

if $N \geq n_0$.

Theorem 1.1.2 (Cauchy's test). Let (a_n) be a sequence. $\sum a_n < \infty$ if and only if $\forall \varepsilon > 0 \exists n_0 \in \mathbb{N}$ such that

$$\left| \sum_{n=N}^M a_n \right| < \varepsilon$$

if $M \geq N \geq n_0$.

Corollary 1.1.3. Changing a finite number of terms in a series has no effect on the convergence or divergence of the series.

Corollary 1.1.4. If $\sum a_n < \infty$, then $\lim_{n \rightarrow \infty} a_n = 0$.

Theorem 1.1.5 (Linearity). Let $\sum a_n, \sum b_n$ be two convergent series with sums A and B respectively and let λ be a real number. The series

$$\sum_{n=1}^{\infty} (a_n + \lambda b_n)$$

is convergent and has sum $A + \lambda B$.

Theorem 1.1.6 (Associative property). Let $\sum a_n$ be a convergent series with sum A . Suppose (n_k) is a strictly increasing sequence of natural numbers. The series $\sum b_n$, with $b_k = a_{n_{k-1}+1} + \dots + a_{n_k}$ for all $i \in \mathbb{N}$, is convergent and its sum is A .

1.2 Non-negative terms series

Theorem 1.2.1. Let $\sum a_n$ be a series of non-negative terms $a_n \geq 0$ ³. The series converges if and only if the sequence (S_N) of partial sums is bounded.

Theorem 1.2.2 (Comparison test). Let $(a_n), (b_n) \geq 0$ be two sequences of real numbers. Suppose that exists a constant $C > 0$ and a number $n_0 \in \mathbb{N}$ such that $a_n \leq C b_n$ for all $n \geq n_0$.

¹From now on we will write $\sum a_n$ to refer $\sum_{n=1}^{\infty} a_n$.

²We will use the notation $\sum a_n < \infty$ or $\sum a_n = +\infty$ to express that the series converges or diverges, respectively.

³Obviously the following results are also valid if the series is of non-positive terms or has a finite number of negative or positive terms.

1. If $\sum b_n < \infty \implies \sum a_n < \infty$.
2. If $\sum a_n = +\infty \implies \sum b_n = +\infty$.

Theorem 1.2.3 (Limit comparison test). *Let $(a_n), (b_n) \geq 0$ be two sequences of real numbers. Suppose that the limit $\ell = \lim_{n \rightarrow \infty} \frac{a_n}{b_n}$ exists.*

1. If $0 < \ell < \infty \implies \sum a_n < \infty \iff \sum b_n < \infty$.
2. If $\ell = 0$ and $\sum b_n < \infty \implies \sum a_n < \infty$.
3. If $\ell = \infty$ and $\sum a_n < \infty \implies \sum b_n < \infty$.

Theorem 1.2.4 (Root test). *Let $(a_n) \geq 0$. Suppose that the limit $\ell = \lim_{n \rightarrow \infty} \sqrt[n]{a_n}$ exists.*

1. If $\ell < 1 \implies \sum a_n < \infty$.
2. If $\ell > 1 \implies \sum a_n = +\infty$.

Theorem 1.2.5 (Ratio test). *Let $(a_n) \geq 0$. Suppose that the limit $\ell = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$ exists.*

1. If $\ell < 1 \implies \sum a_n < \infty$.
2. If $\ell > 1 \implies \sum a_n = +\infty$.

Theorem 1.2.6 (Raabe's test). *Let $(a_n) \geq 0$. Suppose that the limit $\ell = \lim_{n \rightarrow \infty} n \left(1 - \frac{a_{n+1}}{a_n} \right)$ exists.*

1. If $\ell > 1 \implies \sum a_n < \infty$.
2. If $\ell < 1 \implies \sum a_n = +\infty$.

Theorem 1.2.7 (Condensation test). *Let $(a_n) \geq 0$ be a decreasing sequence. Then:*

$$\sum a_n < \infty \iff \sum 2^n a_{2^n} < \infty.$$

Theorem 1.2.8 (Logarithmic test). *Let $(a_n) \geq 0$. Suppose that the limit $\ell = \lim_{n \rightarrow \infty} \frac{\log \frac{1}{a_n}}{\log n}$ exists.*

1. If $\ell > 1 \implies \sum a_n < \infty$.
2. If $\ell < 1 \implies \sum a_n = +\infty$.

Theorem 1.2.9 (Integral test). *Let $f : [1, \infty) \rightarrow (0, \infty)$ be a decreasing function. Then:*

$$\sum f(n) < \infty \iff \iff \exists C > 0 \text{ such that } \int_1^n f(x) dx \leq C \forall n.$$

⁴Later we will see that this is equivalent to say that if $f : [1, \infty) \rightarrow (0, \infty)$ is a locally integrable decreasing function, then:

$$\sum f(n) < \infty \iff \int_1^\infty f(x) dx < \infty.$$

1.3 Alternating series

Definition 1.3.1. An *alternating series* is a series of the form $\sum (-1)^n a_n$, with $(a_n) \geq 0$.

Theorem 1.3.1 (Leibnitz's test). *Let $(a_n) \geq 0$ be a decreasing sequence such that $\lim_{n \rightarrow \infty} a_n = 0$. Then, $\sum (-1)^n a_n$ is convergent.*

Theorem 1.3.2 (Abel's summation formula). *Let $(a_n), (b_n)$ be two sequences of real numbers. Then,*

$$\sum_{n=N}^M a_n (b_{n+1} - b_n) = a_{M+1} b_{M+1} - a_N b_N - \sum_{n=N}^M b_{n+1} (a_{n+1} - a_n).$$

Theorem 1.3.3 (Dirichlet's test). *Let $(a_n), (b_n)$ be two sequences of real numbers such that:*

1. $\exists C > 0$ such that $\left| \sum_{n=1}^N a_n \right| \leq C$ for all $N \in \mathbb{N}$.
2. (b_n) is monotone and $\lim_{n \rightarrow \infty} b_n = 0$.

Then, $\sum a_n b_n$ is convergent.

Theorem 1.3.4 (Abel's test). *Let $(a_n), (b_n)$ be two sequences of real numbers such that:*

1. *The series $\sum a_n$ is convergent.*
2. *(b_n) is monotone and bounded.*

Then, $\sum a_n b_n$ is convergent.

1.4 Absolute convergence and rearrangement of series

Definition 1.4.1. We say a series $\sum a_n$ is *absolutely convergent* if $\sum |a_n|$ is convergent.

Theorem 1.4.1. *If a series converges absolutely, it converges.*

Definition 1.4.2. We say a sequence (b_n) is a *rearrangement of the sequence (a_n)* if exists a bijective map $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $b_n = a_{\sigma(n)}$. A *rearrangement of the series $\sum a_n$* is the series $\sum a_{\sigma(n)}$ for some bijection $\sigma : \mathbb{N} \rightarrow \mathbb{N}$.

Definition 1.4.3. Let $x \in \mathbb{R}$. We define the *positive part of x* as

$$x^+ = \begin{cases} x & \text{si } x \geq 0 \\ 0 & \text{si } x < 0 \end{cases}$$

Analogously, we define the *negative part of x* as

$$x^- = \begin{cases} 0 & \text{si } x \geq 0 \\ -x & \text{si } x < 0 \end{cases}$$

Note that we can write $x = x^+ - x^-$ and $|x| = x^+ + x^-$.

Theorem 1.4.2. *A series $\sum a_n$ is absolutely convergent if and only if positive and negative terms series, $\sum a_n^+$ and $\sum a_n^-$, converge. In this case,*

$$\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} a_n^+ - \sum_{n=1}^{\infty} a_n^-.$$

Theorem 1.4.3. *Let $\sum a_n$ be an absolutely convergent series. Then, for all bijection $\sigma : \mathbb{N} \rightarrow \mathbb{N}$, the rearranged series $\sum a_{\sigma(n)}$ is absolutely convergent and $\sum a_n = \sum a_{\sigma(n)}$.*

Theorem 1.4.4 (Riemann's theorem). *Let $\sum a_n$ be a convergent series but not absolutely convergent. Then, $\forall \alpha \in \mathbb{R} \cup \{\infty\}$, there exists a bijective map $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $\sum a_{\sigma(n)}$ converges and $\sum a_{\sigma(n)} = \alpha$.*

Theorem 1.4.5. *A series $\sum a_n$ is absolutely convergent if and only if any rearranged series converges to the same value of $\sum a_n$.*

Chapter 2

Sequences and series of functions

2.1 Sequences of functions

Definition 2.1.1. Let $D \subseteq \mathbb{R}$. A set

$$(f_n(x)) = \{f_1(x), f_2(x), \dots, f_n(x), \dots\}$$

is a *sequence of real functions* if $f_i : D \rightarrow \mathbb{R}$ is a real-valued function. In this case we say the sequence $(f_n(x))$, or simply (f_n) , is well-defined on D .

Definition 2.1.2. Let (f_n) be a sequence of functions defined on $D \subseteq \mathbb{R}$ and $f : D \rightarrow \mathbb{R}$. We say (f_n) *converges pointwise to f on D* if $\forall x \in D, \lim_{n \rightarrow \infty} f_n(x) = f(x)$

Definition 2.1.3. Let (f_n) be a sequence of functions defined on $D \subseteq \mathbb{R}$ and $f : D \rightarrow \mathbb{R}$. We say (f_n) *converges uniformly to f on D* if $\forall \varepsilon > 0, \exists n_0 : |f_n(x) - f(x)| < \varepsilon \forall n \geq n_0$ and $\forall x \in D$.

Lemma 2.1.1. Let (f_n) be an uniform convergent sequence of functions defined on $D \subseteq \mathbb{R}$ and let f be a function such that (f_n) converges pointwise to f . Then, (f_n) converges uniformly f on D .

Lemma 2.1.2. Let (f_n) be a sequence of functions defined on $D \subseteq \mathbb{R}$. (f_n) converges uniformly a f en D if and only if $\lim_{n \rightarrow \infty} \sup \{|f_n(x) - f(x)| : x \in D\} = 0$.

Corollary 2.1.3. A sequence of functions (f_n) converges uniformly to f on $D \subseteq \mathbb{R}$ if and only if there is a sequence (a_n) , with $a_n \geq 0$ and $\lim_{n \rightarrow \infty} a_n = 0$, and a number $n_0 \in \mathbb{N}$ such that $\sup \{|f_n(x) - f(x)| : x \in D\} \leq a_n, \forall n \geq n_0$.

Theorem 2.1.4 (Cauchy's test). A sequence of functions (f_n) converges uniformly to f on $D \subseteq \mathbb{R}$ if and only if $\forall \varepsilon > 0 \exists n_0 : \sup \{|f_n(x) - f_m(x)| : x \in D\} < \varepsilon$ if $n, m \geq n_0$.

Theorem 2.1.5. Let (f_n) be a sequence of continuous functions defined on $D \subseteq \mathbb{R}$. If (f_n) converges uniformly to f on D , then f is continuous on D , that is, for any $x_0 \in D$, it satisfies:

$$\lim_{n \rightarrow \infty} \left(\lim_{x \rightarrow x_0} f_n(x) \right) = \lim_{x \rightarrow x_0} f(x).$$

Theorem 2.1.6. Let (f_n) be a sequence of functions defined on $I = [a, b] \subseteq \mathbb{R}$. If (f_n) are Riemann-integrable on I and (f_n) converges uniformly to f on I , then f is Riemann-integrable on I and

$$\int_a^b \lim_{n \rightarrow \infty} f_n(x) dx = \lim_{n \rightarrow \infty} \int_a^b f_n(x) dx.$$

Theorem 2.1.7. Let (f_n) be a sequence of functions defined on $I = (a, b) \subset \mathbb{R}$. If (f_n) are derivable on I , $(f'_n(x))$ converges uniformly on I and $\exists x_0 \in I : \lim_{n \rightarrow \infty} f_n(x_0) \in \mathbb{R}$, then there is a function f such that (f_n) converges uniformly to f on I , f is derivable on I and $(f'_n(x))$ converges uniformly to f' on I .

2.2 Series of functions

Definition 2.2.1. Let (f_n) be a sequence of functions defined on $D \subseteq \mathbb{R}$. The expression

$$\sum_{n=1}^{\infty} f_n(x)$$

is the *series of functions associated with (f_n)* .

Definition 2.2.2. A series of functions $\sum f_n(x)$ defined on $D \subseteq \mathbb{R}$ *converges pointwise on D* if the sequence of partials sums

$$F_N(x) = \sum_{n=1}^N f_n(x)$$

converges pointwise. If the pointwise limit of (F_N) is $F(x)$, we say F is the *sum of the series in a pointwise sense*.

Definition 2.2.3. A series of functions $\sum f_n(x)$ defined on $D \subseteq \mathbb{R}$ converges uniformly on D if the sequence of partials sums

$$F_N(x) = \sum_{n=1}^N f_n(x)$$

converges uniformly. If the uniform limit of (F_N) is $F(x)$, we say F is the sum of the series in an uniform sense.

Theorem 2.2.1 (Cauchy's test). A series of functions $\sum f_n(x)$ defined on $D \subseteq \mathbb{R}$ converges uniformly if and only if $\forall \varepsilon > 0 \exists n_0$ such that

$$\sup \left\{ \left| \sum_{n=N}^M f_n(x) \right| : x \in D \right\} < \varepsilon$$

if $M \geq N \geq n_0$.

Corollary 2.2.2. If $\sum f_n(x)$ is a series of continuous functions on $D \subseteq \mathbb{R}$, then (f_n) converges uniformly to zero on D .

Theorem 2.2.3. If $\sum f_n(x)$ is uniformly convergent series of functions on $D \subseteq \mathbb{R}$, then its sum function is also continuous on D .

Theorem 2.2.4. Let (f_n) be a sequence of functions defined on $I = [a, b] \subseteq \mathbb{R}$. If (f_n) are Riemann-integrable on I and $\sum f_n(x)$ converges uniformly on I , then $\sum f_n(x)$ is Riemann-integrable on I and

$$\int_a^b \sum_{n=1}^{\infty} f_n(x) dx = \sum_{n=1}^{\infty} \int_a^b f_n(x) dx.$$

Theorem 2.2.5. Let (f_n) be a sequence of functions defined on $I = (a, b) \subset \mathbb{R}$. If (f_n) are derivable on I , $\sum f'_n(x)$ converges uniformly on I and $\exists c \in I : \sum f_n(c) < \infty$, then $\sum f_n(x)$ converges uniformly on I , $\sum f_n(x)$ is derivable on I and

$$\left(\sum_{n=1}^{\infty} f_n(x) \right)' = \sum_{n=1}^{\infty} f'_n(x).$$

Theorem 2.2.6 (Weierstraß M-test). Let (f_n) be a sequence of functions defined on $D \subseteq \mathbb{R}$ such that $|f_n(x)| \leq M_n \forall x \in D$ and suppose that $\sum M_n$ is a convergent numeric series. Then, $\sum f_n(x)$ is converges uniformly on D .

Theorem 2.2.7 (Dirichlet's test). Let $(f_n), (g_n)$ be two sequences of functions defined on $D \subseteq \mathbb{R}$. Suppose:

1. $\exists C > 0 : \sup \left\{ \left| \sum_{n=1}^N f_n(x) \right| : x \in D \right\} \leq C, \forall N$.
2. $(g_n(x))$ is a monotone sequence for all $x \in D$ and $\lim_{n \rightarrow \infty} \sup \{|g_n(x)| : x \in D\} = 0$.

Then, $\sum f_n(x)g_n(x)$ converges uniformly on D .

Theorem 2.2.8 (Abel's test). Let $(f_n), (g_n)$ be two sequences of functions defined on $D \subseteq \mathbb{R}$. Suppose:

1. The series $\sum f_n(x)$ converges uniformly on D .
2. $(g_n(x))$ is a monotone and bounded sequence for all $x \in D$.

Then, $\sum f_n(x)g_n(x)$ converges uniformly on D .

2.3 Power series

Definition 2.3.1. Let (a_n) be a sequence of real numbers and $x_0 \in \mathbb{R}$. A *power series centred on x_0* is a series of functions of the form

$$\sum_{n=0}^{\infty} a_n (x - x_0)^n.$$

Proposition 2.3.1. Let $\sum a_n (x - x_0)^n$ be a power series. Suppose there exists an $x_1 \in \mathbb{R}$ such that $\sum a_n (x_1 - x_0)^n < \infty$. Then, $\sum a_n (x - x_0)^n$ converges uniformly on any closed interval $I \subset A = \{x \in \mathbb{R} : |x - x_0| < |x_1 - x_0|\}$.

Theorem 2.3.2. Let $\sum a_n (x - x_0)^n$ be a power series and consider

$$R = \left(\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|} \right)^{-1} \in [0, \infty].$$

Then:

1. If $|x - x_0| < R \implies \sum a_n (x - x_0)^n$ converges absolutely.
2. If $0 \leq r < R \implies \sum a_n (x - x_0)^n$ converges uniformly on $[x_0 - r, x_0 + r]$.
3. If $|x - x_0| > R \implies \sum a_n (x - x_0)^n$ diverges.

The number R is called *radius of convergence of the power series*.

Theorem 2.3.3 (Abel's theorem). Let $\sum a_n x^n$ be a power series¹ with radius of convergence R satisfying $\sum a_n R^n < \infty$. Then the series $\sum a_n x^n$ converges uniformly on $[0, R]$. In particular, if $f(x) = \sum a_n x^n$,

$$\lim_{x \rightarrow R^-} f(x) = \sum_{n=0}^{\infty} a_n R^n.$$

Corollary 2.3.4. Let f be the sum function of a power series $\sum a_n x^n$. Then f is continuous on the domain of convergence of the series.

Corollary 2.3.5. If the series $\sum a_n x^n$ has radius of convergence $R \neq 0$ and f is its sum function, then f is Riemann-integrable on any closed subinterval on the domain of convergence of the series. In particular, for $|x| < R$,

$$\int_0^x f(t) dt = \sum_{n=0}^{\infty} a_n \frac{x^{n+1}}{n+1}.$$

Corollary 2.3.6. Let f be the sum function of the power series $\sum a_n x^n$. Then f is derivable within the domain of convergence of the series and

$$f'(x) = \sum_{n=0}^{\infty} n a_n x^{n-1}.$$

Corollary 2.3.7. Any function f defined as a sum of a power series $\sum a_n x^n$ is indefinitely derivable within the domain of convergence of the series and

$$f^{(k)}(x) = \sum_{n=k}^{\infty} n(n-1) \cdots (n-k+1) a_n x^{n-k},$$

for all $k \in \mathbb{N} \cup \{0\}$. In particular $f^{(k)}(0) = k! a_k$.

Definition 2.3.2. A function is *analytic* if it can be expressed locally as a power series.

¹From now on we will suppose, for simplicity, $x_0 = 0$.

²The formula is also valid for $|x| = R$ if the series $\sum a_n R^n$ (or $\sum a_n (-R)^n$) is convergent.

2.4 Stone-Weierstraß approximation theorem

Definition 2.4.1. Let f be a real-valued function. We say f has *compact support*³ if exists an $M > 0$ such that $f(x) = 0$ for all $x \in \mathbb{R} \setminus [-M, M]$.

Definition 2.4.2. Let f, g be real-valued functions with compact support. We define the convolution of f with g as

$$(f * g)(x) = \int_{\mathbb{R}} f(t)g(x-t)dt^4.$$

Definition 2.4.3. We say a sequence of functions (ϕ_ε) with compact support is an *approximation of unity* if

1. $\phi_\varepsilon \geq 0$.
2. $\int_{\mathbb{R}} \phi_\varepsilon = 1$.
3. For all $\delta > 0$, $\phi_\varepsilon(t)$ converges uniformly to zero when $\varepsilon \rightarrow 0$ if $|t| > \delta$.

Lemma 2.4.1. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function with compact support. Let (ϕ_ε) be an approximation of unity. Then $(f * \phi_\varepsilon)$ converges uniformly to f on \mathbb{R} when $\varepsilon \rightarrow 0$.

Theorem 2.4.2 (Stone-Weierstraß theorem). Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then, there exists polynomials $p_n \in \mathbb{R}[x]$ such that the sequence (p_n) converge uniformly to f on $[a, b]$.

³In general, the support of a function is the adherence of the set of points which are not mapped to zero.

⁴Alternatively if f, g are Riemann-integrable functions on $[a, b]$ we can define the convolution of f and g as

$$(f * g)(x) = \int_a^b f(t)g(x-t)dt.$$

Chapter 3

Improper integrals

3.1 Locally integrable functions

Definition 3.1.1. Let $f : [a, b) \rightarrow \mathbb{R}$, with $b \in \mathbb{R} \cup \{\infty\}$. We say f is *locally integrable* on $[a, b)$ if f is Riemann-integrable on $[a, x]$ for all $a \leq x < b$.

Definition 3.1.2. Let $f : [a, b) \rightarrow \mathbb{R}$ be a locally integrable function. If there exists

$$\lim_{x \rightarrow b^-} \int_a^x f$$

and it's finite, we say that the *improper integral* of f on $[a, b)$, $\int_a^b f$, is *convergent*.

Theorem 3.1.1 (Cauchy's test). *Let $f : [a, b) \rightarrow \mathbb{R}$ be a locally integrable function. The improper integral $\int_a^b f$ is convergent if and only if $\forall \varepsilon > 0 \exists b_0, a < b_0 < b$, such that*

$$\left| \int_{x_1}^{x_2} f \right| < \varepsilon$$

if $b_0 < x_1 < x_2 < b$.

3.2 Improper integrals of non-negative functions

Theorem 3.2.1. *Let $f : [a, b) \rightarrow \mathbb{R}$ be a locally integrable non-negative function. A necessary and sufficient condition for $\int_a^b f$ to be convergent is that the function*

$$F(x) = \int_a^x f(t) dt$$

must be bounded for all $x < b$.

Theorem 3.2.2 (Comparison test). *Let $f, g : [a, b) \rightarrow [0, +\infty)$ be two locally integrable non-negative functions. Then:*

1. *If $\exists C > 0$ such that $f(x) \leq Cg(x) \forall x$ on a neighborhood of b and $\int_a^b g < \infty \implies \int_a^b f < \infty$.*
2. *Suppose the limit $\ell = \lim_{x \rightarrow b} \frac{f(x)}{g(x)}$ exists. Then,*

$$(a) \text{ If } \ell \in (0, \infty) \implies \int_a^b f < \infty \iff \int_a^b g < \infty.$$

$$(b) \text{ If } \ell = 0 \text{ and } \int_a^b g < \infty \implies \int_a^b f < \infty.$$

$$(c) \text{ If } \ell = \infty \text{ and } \int_a^b f < \infty \implies \int_a^b g < \infty.$$

3.3 Absolute convergence of improper integrals

Definition 3.3.1. Let $f : [a, b) \rightarrow (0, \infty)$ be a locally integrable function. We say $\int_a^b f$ converges absolutely if $\int_a^b |f|$ is convergent.

Theorem 3.3.1 (Dirichlet's test). Let $f, g : [a, b) \rightarrow \mathbb{R}$ be two locally integrable functions. Suppose:

1. $\exists C > 0$ such that $|\int_a^x f(t)dt| \leq C$ for all $x \in [a, b)$.

2. g is monotone and $\lim_{x \rightarrow b} g(x) = 0$.

Then, $\int_a^b fg$ is convergent.

Theorem 3.3.2 (Abel's test). Let $f, g : [a, b) \rightarrow \mathbb{R}$ be two locally integrable functions. Suppose:

1. $\int_a^b f$ is convergent.

2. g is monotone and bounded.

Then, $\int_a^b fg$ is convergent.

3.4 Differentiation under integral sign

Theorem 3.4.1. Let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a continuous function on $[a, b] \times [c, d]$. Consider the function $F(y) = \int_a^b f(x, y)dx$ defined on $[c, d]$. Then, F is continuous, that is, if $c < y_0 < d$,

$$\begin{aligned} \lim_{y \rightarrow y_0} F(y) &= \lim_{y \rightarrow y_0} \int_a^b f(x, y)dx = \int_a^b \lim_{y \rightarrow y_0} f(x, y)dx = \\ &= \int_a^b f(x, y_0)dx = F(y_0). \end{aligned}$$

Theorem 3.4.2. Let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a Riemann-integrable function and let $F(y) = \int_a^b f(x, y)dx$. If f is differentiable with respect to y and $\partial f / \partial y$ is continuous on $[a, b] \times [c, d]$, then $F(y)$ is derivable on (c, d) and its derivative is

$$F'(y) = \int_a^b \frac{\partial f}{\partial y}(x, y)dx,$$

for all $y \in (c, d)$.

Theorem 3.4.3. Let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a continuous function on $[a, b] \times [c, d]$. Let $a, b : [c, d] \rightarrow \mathbb{R}$ be two differentiable functions satisfying $a \leq a(y) \leq b(y) \leq b$ for every $y \in [c, d]$. Suppose that

$\partial f / \partial y$ is continuous on $\{(x, y) \in \mathbb{R}^2 : a(y) \leq x \leq b(y), c \leq y \leq d\}$. Then $F(y) = \int_{a(y)}^{b(y)} f(x, y) dx$ is

derivable on (c, d) and its derivative is

$$F'(y) = b'(y)f(b(y), y) - a'(y)f(a(y), y) + \int_{a(y)}^{b(y)} \frac{\partial f}{\partial y}(x, y) dx,$$

for all $y \in (c, d)$.

Theorem 3.4.4. Let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a continuous function on $[a, b] \times [c, d]$. We consider

$F(y) = \int_a^b f(x, y) dx$. Suppose that:

1. $\frac{\partial f}{\partial y}$ is continuous on $[a, b] \times [c, d]$.
2. Given $y_0 \in [c, d]$, $\exists \delta > 0$ such that the integral

$$\int_a^b \sup \left\{ \left| \frac{\partial f}{\partial y}(x, y) \right| : y \in (y_0 - \delta, y_0 + \delta) \right\} dx$$

exists and it's finite on $[a, b]$.

Then, $F(y)$ is derivable at y_0 and

$$F'(y_0) = \int_a^b \frac{\partial f}{\partial y}(x, y_0) dx.$$

Theorem 3.4.5. Let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a continuous function on $[a, b] \times [c, d]$. Let $a, b : [c, d] \rightarrow \mathbb{R}$ be two differentiable functions satisfying $a \leq a(y) \leq b(y) \leq b$ for every $y \in [c, d]$. We

consider $F(y) = \int_{a(y)}^{b(y)} f(x, y) dx$. Suppose that:

1. $\frac{\partial f}{\partial y}$ is continuous on $\{(x, y) \in \mathbb{R}^2 : a(y) \leq x \leq b(y), c \leq y \leq d\}$.
2. Given $y_0 \in [c, d]$, $\exists \delta > 0$ such that the integral

$$\int_{a(y)}^{b(y)} \sup \left\{ \left| \frac{\partial f}{\partial y}(x, y) \right| : y \in (y_0 - \delta, y_0 + \delta) \right\} dx$$

exists and it's finite on $[a, b]$.

The, $F(y)$ is derivable at y_0 and

$$F'(y_0) = b'(y_0)f(b(y_0), y_0) - a'(y_0)f(a(y_0), y_0) + \int_{a(y_0)}^{b(y_0)} \frac{\partial f}{\partial y}(x, y_0) dx.$$

3.5 Gamma function

Definition 3.5.1. For $x > 0$, *Gamma function* is defined as

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt.$$

Theorem 3.5.1. *Gamma function is a generalization of the factorial. In fact, for $x > 0$ we have*

$$\Gamma(x+1) = x\Gamma(x).$$

In particular, $\Gamma(n+1) = n!$ for all $n \in \mathbb{N}$.

Theorem 3.5.2. *Gamma function satisfies:*

$$\lim_{x \rightarrow \infty} \frac{\Gamma(x+1)}{(x/e)^x \sqrt{2\pi x}} = 1.$$

Corollary 3.5.3 (Stirling's formula).

$$\lim_{n \rightarrow \infty} \frac{n!}{n^n e^{-n} \sqrt{2\pi n}} = 1.$$

Chapter 4

Fourier series

4.1 Preliminaries

Definition 4.1.1. Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be a function. We say f has a period T or f is T -periodic with $T > 0$ if and only if $f(x + T) = f(x)$ for all $x \in \mathbb{R}$.

Lemma 4.1.1. Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be a T -periodic function. Then, $f(x + T') = f(x)$ for all $x \in \mathbb{R}$ if and only if $T' = kT$ for some $k \in \mathbb{Z}$.

Proposition 4.1.2. Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be a T -periodic function and $a \in \mathbb{R}$ a number. Then,

$$\int_a^{a+T} f \, dx = \int_0^T f \, dx. \quad (4.1)$$

Lemma 4.1.3. Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be a T -periodic function continuous in \mathbb{R} . Then, $|f|$ is bounded.

Proposition 4.1.4. Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be a T -periodic function. Then, there is no a power series that converges uniformly to f in \mathbb{R} .

4.2 Orthogonal and orthonormal systems of functions

Definition 4.2.1. Let $f, g : [a, b] \rightarrow \mathbb{C}$ be two integrable functions. We define the *inner product* of f and g as

$$\langle f, g \rangle_2 := \int_a^b f(x) \overline{g(x)} \, dx, \quad (4.2)$$

where \bar{g} is the conjugate of g .

Definition 4.2.2. Let $f, g : [a, b] \rightarrow \mathbb{C}$ two integrable functions. We define the *2-norm* of f as

$$\|f\|_2 := \sqrt{\langle f, f \rangle_2} = \left(\int_a^b |f(x)|^2 \, dx \right)^{1/2}. \quad (4.3)$$

Definition 4.2.3. Let $f, g : [a, b] \rightarrow \mathbb{C}$ two integrable functions. We define the *distance between f and g* as

$$d(f, g) := \|f - g\|. \quad (4.4)$$

Proposition 4.2.1. Let $f, g : [a, b] \rightarrow \mathbb{C}$ two integrable functions. Then,

1. $\langle f, f \rangle_2 \geq 0$.
2. $\langle f + g, h \rangle_2 = \langle f, h \rangle_2 + \langle g, h \rangle_2$ and $\langle f, g + h \rangle_2 = \langle f, g \rangle_2 + \langle f, h \rangle_2$.
3. $\langle f, g \rangle_2 = \overline{\langle g, f \rangle_2}$.
4. For $\alpha \in \mathbb{C}$, $\langle \alpha f, g \rangle_2 = \alpha \langle f, g \rangle_2$ and $\langle f, \alpha g \rangle_2 = \bar{\alpha} \langle f, g \rangle_2$.

Note that for Riemann integrable functions it is not true that $\langle f, f \rangle_2 = 0 \Leftrightarrow f = 0$, since a function that is not zero at some point will satisfy $\langle f, f \rangle_2 = 0$. However, if we deal with the space of continuous functions in $[a, b]$ then it is true.

Theorem 4.2.2. Let $f, g : [a, b] \rightarrow \mathbb{C}$ two Riemann integrable functions. Then,

$$|\langle f, g \rangle_2| \leq \|f\|_2 \|g\|_2. \quad (4.5)$$

Theorem 4.2.3. Let $f, g : [a, b] \rightarrow \mathbb{C}$ two Riemann integrable functions. Then,

$$\|f + g\|_2 \leq \|f\|_2 + \|g\|_2. \quad (4.6)$$

Definition 4.2.4. Let $f, g : [a, b] \rightarrow \mathbb{C}$ two Riemann integrable functions with $f \neq g$. Then,

1. we say f and g are orthogonal if and only if $\langle f, g \rangle_2 = 0$.
2. We say f and g are orthonormal if and only if $\langle f, g \rangle_2 = 0$ and $\|f\|_2 = \|g\|_2 = 1$.
3. Let $S = \{\phi_0, \phi_1, \dots\}$ be a collection of Riemann integrable functions in $[a, b]$. We say S is an orthonormal system if and only if $\|\phi_i\|_2 = 1 \forall i$ and $\langle \phi_i, \phi_j \rangle_2 = 0 \forall i \neq j$.

Definition 4.2.5. Let $\{\phi_0, \dots, \phi_n\}$ be a collection of Riemann integrable functions in $[a, b]$. We say the collection is linearly dependent in $[a, b]$ if and only if there exist c_0, \dots, c_n with not all being zero such that

$$c_0\phi_0(x) + \dots + c_n\phi_n(x) = 0, \forall x \in [a, b]. \quad (4.7)$$

Theorem 4.2.4. Let $S = \{\phi_0, \phi_1, \dots\}$ be an orthonormal system in $[a, b]$ such that $\sum_{n=0}^{\infty} c_n\phi_n(x)$ converges uniformly in $[a, b]$. Let f be the function that defines the series in $[a, b]$. Then f is Riemann integrable in $[a, b]$ and

$$c_n = \langle f, \phi_n \rangle_2 = \int_a^b f(x) \overline{\phi_n(x)} dx, n \geq 0. \quad (4.8)$$

4.3 Fourier coefficients. Fourier series

4.3.1 Exponential form

Definition 4.3.1. Let f be a Riemann integrable function in $[0, 1]$. We define the n -th Fourier coefficient with $n \in \mathbb{Z}$ as

$$\hat{f}(n) = \langle f, e_n \rangle_2 = \int_0^1 f(x) e^{-2\pi i n x} dx. \quad (4.9)$$

The series

$$Sf(x) = \sum_{n=0}^{\infty} \hat{f}(n) e^{2\pi i n x} \quad (4.10)$$

constructed by these coefficients is called the *Fourier series of f* .

Proposition 4.3.1. Let f be a Riemann integrable function in $[0, 1]$ and $\lambda, \mu \in \mathbb{C}$ two numbers. In relation to Fourier coefficients, the following statements are true.

1. $\widehat{\lambda f + \mu g}(n) = \lambda \hat{f}(n) + \mu \hat{g}(n)$.
2. If $\tau \in (0, 1)$ and $f_\tau(x) := f(x - \tau)$, then $\hat{f}_\tau(n) = e^{-2\pi i n \tau} \hat{f}(n)$.
3. If f is even, then $\hat{f}(n) = \hat{f}(-n) \forall n$ and if f is odd, $\hat{f}(n) = -\hat{f}(-n) \forall n$.
4. If f' exists and it is continuous, then $\hat{f}'(n) = 2\pi i n \hat{f}(n)$.

Definition 4.3.2. Let f, g be two Riemann integrable functions in $[0, 1]$. We define the *convolution of f and g* as

$$(f * g)(x) = \int_0^1 f(t) g(x - t) dt, \quad (4.11)$$

defined for $x \in \mathbb{R}$.

Proposition 4.3.2. Let f, g be two Riemann integrable functions in $[0, 1]$. Then, $\widehat{f * g}(n) = \hat{f}(n) \hat{g}(n)$.

4.3.2 Fourier series in terms of sine and cosine

We can write the Fourier series as

$$Sf(x) = A_0 + 2 \sum_{n=0}^{\infty} A_n \cos(2\pi nx) + B_n \sin(2\pi nx), \quad (4.12)$$

$$A_0 = \int_0^1 f(x) dx, \quad A_n = \int_0^1 f(x) \cos(2\pi nx) dx, \quad B_n = \int_0^1 f(x) \sin(2\pi nx) dx. \quad (4.13)$$

Proposition 4.3.3. *Let f be a Riemann integrable function in $[0, 1]$. If f is even, then its Fourier series is written as*

$$Sf(x) = A_0 + 2 \sum_{n=0}^{\infty} A_n \cos(2\pi nx). \quad (4.14)$$

If f is odd, then its Fourier series is written as

$$Sf(x) = 2 \sum_{n=0}^{\infty} B_n \sin(2\pi nx), \quad (4.15)$$

4.3.3 Fourier series in terms of sine or cosine

Definition 4.3.3. Let f be a Riemann integrable function in $[0, 1/2]$. We define the *even extension* and *odd extension*, respectively, as

$$\tilde{f}(x) = \begin{cases} f(x), & \text{if } x \in [0, 1/2] \\ f(-x), & \text{if } x \in [-1/2, 0] \end{cases}, \quad \tilde{f}(x) = \begin{cases} f(x), & \text{if } x \in [0, 1/2] \\ -f(-x), & \text{if } x \in [-1/2, 0] \end{cases}. \quad (4.16)$$

Proposition 4.3.4. *Let $f : [0, 1/2] \rightarrow \mathbb{C}$ be a Riemann integrable function. If we make the even extension of f , then*

$$Sf(x) = 2A_0 + 4 \sum_{n=0}^{\infty} A_n \cos(2\pi nx), \quad A_0 = \int_0^{1/2} f(x) dx, \quad A_n = \int_0^{1/2} f(x) \cos(2\pi nx) dx. \quad (4.17)$$

If we make the odd extension of f , then

$$Sf(x) = 4 \sum_{n=0}^{\infty} B_n \sin(2\pi nx), \quad B_n = \int_0^{1/2} f(x) \sin(2\pi nx) dx. \quad (4.18)$$

4.3.4 Change of period

For a function $f : \mathbb{R} \rightarrow \mathbb{C}$ T -periodic,

$$Sf(x) = \sum_{n=-\infty}^{\infty} \hat{f}(n) e^{2\pi n x i / T}, \quad \hat{f}(n) = \int_{-T/2}^{T/2} f(x) e^{-2\pi i n x / T} dx. \quad (4.19)$$

or

$$Sf(x) = A_0 + 2 \sum_{n=1}^{\infty} A_n \cos \left[\frac{2\pi n x}{T} \right] + B_n \sin \left[\frac{2\pi n x}{T} \right], \quad (4.20)$$

$$A_0 = \frac{1}{T} \int_{-T/2}^{T/2} f(x) dx, \quad A_n = \frac{1}{T} \int_{-T/2}^{T/2} f(x) \cos \left[\frac{2\pi n x}{T} \right] dx, \quad B_n = \frac{1}{T} \int_{-T/2}^{T/2} f(x) \sin \left[\frac{2\pi n x}{T} \right] dx. \quad (4.21)$$

4.4 Punctual convergence of Fourier series

We denote the N -th partial sum as

$$S_N f := \sum_{n=-N}^N \hat{f}(n) e^{2\pi i n x}. \quad (4.22)$$

4.4.1 Dirichlet nucleus

$$S_N f(x) = \int_0^1 f(t) \sum_{n=-N}^N e^{e\pi i n(x-t)} dt \quad (4.23)$$

Definition 4.4.1. We define the Dirichlet nucleus of N -th order as

$$D_N(t) = \sum_{n=-N}^N e^{2\pi i n t}. \quad (4.24)$$

This way, we have

$$S_N f(x) = (f * D_N)(x). \quad (4.25)$$

$$D_N(t) = \frac{\sin[(2N+1)\pi t]}{\sin \pi t}. \quad (4.26)$$

Proposition 4.4.1. *The Dirichlet nucleus has the following elemental properties.*

1. D_N is a periodic function of period 1.
2. D_N is an even function.
3. $\int_0^1 D_N(t) dt = 1$.

Definition 4.4.2 (Dirichlet kernel). We define the *Dirichlet kernel of order N* as

$$D_N(t) = \frac{1}{T} \sum_{n=-N}^N e^{\frac{2\pi i n t}{T}} = \frac{1}{T} \frac{\sin\left(\frac{(2N+1)\pi t}{T}\right)}{\sin\left(\frac{\pi t}{T}\right)}.$$

Proposition 4.4.2. *The Dirichlet kernel has the following properties:*

1. D_N is a T -periodic and even function.
2. $\int_0^T D_N(t) dt = 1$ for all N .

Proposition 4.4.3. *Suppose $f \in L^1([-T/2, T/2])$. Then*

$$\begin{aligned} S_N f(x) &= (f * D_N)(x) = \int_{-T/2}^{T/2} f(x-t) D_N(t) dt = \\ &= \int_0^{T/2} [f(x+t) + f(x-t)] D_N(t) dt. \end{aligned}$$

Lemma 4.4.4 (Riemann-Lebesgue lemma). *Let $f \in L^1([-T/2, T/2])$ and $\lambda \in \mathbb{R}$. Then:*

$$\lim_{\lambda \rightarrow \infty} \int_{-T/2}^{T/2} f(t) \sin(\lambda t) dt = \lim_{\lambda \rightarrow \infty} \int_{-T/2}^{T/2} f(t) \cos(\lambda t) dt = 0.$$

In particular, $\lim_{|n| \rightarrow \infty} \widehat{f}(n) = 0$.

Theorem 4.4.5. *Let $f \in L^1([-T/2, T/2])$ be a function having left and right-hand side derivatives at x_0 , that is, there exists the following limits*

$$f'(x_0^+) = \lim_{t \rightarrow 0^+} \frac{f(x_0 + t) - f(x_0^+)}{t},$$

$$f'(x_0^-) = \lim_{t \rightarrow 0^-} \frac{f(x_0 + t) - f(x_0^-)}{t},$$

(supposing left and right-hand side exists). Then,

$$\lim_{N \rightarrow \infty} S_N f(x_0) = \frac{f(x_0^+) + f(x_0^-)}{2}.$$

Theorem 4.4.6 (Dini's theorem). *Let $f \in L^1([-T/2, T/2])$, $x_0 \in (-T/2, T/2)$ and $\ell \in \mathbb{R}$ such that*

$$\int_0^\delta \frac{|f(x_0 + t) + f(x_0 - t) - 2\ell|}{t} dt < \infty$$

for some $\delta > 0$. Then $\lim_{N \rightarrow \infty} S_N f(x_0) = \ell$.

Theorem 4.4.7 (Lipschitz's theorem). *Let $f \in L^1([-T/2, T/2])$ such that at a point $x_0 \in (-T/2, T/2)$ it satisfies $|f(x_0 + t) - f(x_0)| \leq L|t|$ for some constant L and for $|t| < \delta$. Then*

$$\lim_{N \rightarrow \infty} S_N f(x_0) = f(x_0).$$

4.5 Uniform convergence

Definition 4.5.1. Let $\sum a_n$ be a series with partial sums S_k . The series $\sum a_n$ is called *Cesàro summable* with sum S if

$$\lim_{N \rightarrow \infty} \frac{S_1 + \cdots + S_N}{N} = S.$$

Definition 4.5.2 (Fejer kernel). We define the *Fejer kernel of order N* as

$$K_N(t) = \frac{1}{N+1} \sum_{k=0}^N D_k(t) = \frac{1}{T(N+1)} \frac{\sin^2\left(\frac{(N+1)\pi t}{T}\right)}{\sin^2\left(\frac{\pi t}{T}\right)},$$

being $D_k(t)$ the Dirichlet kernel of order k , $0 \leq k \leq N$.

Proposition 4.5.1. *The Fejer kernel has the following properties:*

1. K_N is a T -periodic, even and non-negative function.

2. $\int_{-T/2}^{T/2} K_N(t) dt = 1$ for all N .

3. For all $\delta > 0$, $\lim_{N \rightarrow \infty} \sup\{|K_N(t)| : \delta \leq |t| \leq T/2\} = 0$.

Definition 4.5.3. Let $f \in L^1([-T/2, T/2])$. We define the *Fejér means* $\sigma_N f$, for all $N \in \mathbb{N}$, as

$$\sigma_N f(x) = \frac{S_0 f(x) + \cdots + S_N f(x)}{N+1}.$$

Proposition 4.5.2. Let $f \in L^1([-T/2, T/2])$. Then

$$\begin{aligned} \sigma_N f(x) &= (f * K_N)(x) = \int_{-T/2}^{T/2} f(x-t) K_N(t) dt = \\ &= \int_0^{T/2} [f(x+t) + f(x-t)] K_N(t) dt. \end{aligned}$$

Theorem 4.5.3 (Fejér's theorem). Let $f \in L^1([-T/2, T/2])$ be a function having left and right-hand side limits at point x_0 . Then,

$$\lim_{N \rightarrow \infty} \sigma_N f(x_0) = \frac{f(x_0^+) + f(x_0^-)}{2}.$$

In particular, if f is continuous at x_0 , $\lim_{N \rightarrow \infty} \sigma_N f(x_0) = f(x_0)$.

Theorem 4.5.4 (Fejér's theorem). Let f be continuous on $[-T/2, T/2]$. Then $\sigma_N f$ converges uniformly to f on $[-T/2, T/2]$.

Corollary 4.5.5. Let f be continuous on $[-T/2, T/2]$. Then there exists a sequence of trigonometric polynomials that converge uniformly to f on $[-T/2, T/2]$. In fact,

$$\sigma_N f(x) = \sum_{k=-N}^N \left(1 - \frac{|k|}{N+1}\right) \hat{f}(k) e^{2\pi i k x}.$$

Corollary 4.5.6. Let f and g be continuous functions on $[-T/2, T/2]$ such that $Sf(x) = Sg(x)$. Then $f = g$.

4.6 Convergence in norm

Definition 4.6.1. We say a sequence (f_N) converge to f in norm $L^2([-T/2, T/2])$ if $\lim_{N \rightarrow \infty} \|f_N - f\| = 0$.

Theorem 4.6.1. Let $f \in L^2([-T/2, T/2])$. Then, $\lim_{N \rightarrow \infty} \|\sigma_N f - f\| = 0$.

Corollary 4.6.2. Let $f \in L^1([-T/2, T/2])$. Then $\lim_{N \rightarrow \infty} \|\sigma_N f - f\|_1 = 0$.

Corollary 4.6.3. Let $f, g \in L^1([-T/2, T/2])$ be functions such that $Sf(x) = Sg(x)$. Then $\lim_{N \rightarrow \infty} \|g - f\|_1 = 0$.

Theorem 4.6.4 (Bessel's inequality). Let $f \in L^2(I)$, where I is any interval on the real line. Then:

$$\begin{aligned} T \sum_{n=-N}^N |\hat{f}(n)|^2 &\leq \|f\|^2, \\ \frac{T}{2} \left(\frac{|a_0|^2}{2} + \sum_{n=1}^N |a_n|^2 + |b_n|^2 \right) &\leq \|f\|^2, \end{aligned}$$

for all $N \in \mathbb{N}$.

Theorem 4.6.5. $S_N f$ is the trigonometric polynomial of degree N that best approximates f in norm L^2 .

Corollary 4.6.6. Let $f \in L^2([-T/2, T/2])$. Then, $\lim_{N \rightarrow \infty} \|S_N f - f\| = 0$.

Theorem 4.6.7 (Parseval's identity). Let $f, g \in L^2([-T/2, T/2])$ be bounded functions. Then

$$\langle f, g \rangle = T \sum_{n \in \mathbb{Z}} \widehat{f}(n) \overline{\widehat{g}(n)}.$$

In particular, if $f = g$, then:

$$\begin{aligned} \|f\|^2 &= T \sum_{n \in \mathbb{Z}} |\widehat{f}(n)|^2, \\ \|f\|^2 &= \frac{T}{2} \left(\frac{|a_0|^2}{2} + \sum_{n=1}^{\infty} |a_n|^2 + |b_n|^2 \right). \end{aligned}$$

4.7 Applications of Fourier series

Theorem 4.7.1 (Wirtinger's inequality). Let f be a function such that $f(0) = f(T)$, $f' \in L^2([0, T])$ and $\int_a^b f(t) dt = 0$. Then,

$$\int_0^T |f(x)|^2 dx \leq \frac{T^2}{4\pi^2} \int_0^T |f'(x)|^2 dx,$$

with equality if and only if $f(x) = A \cos\left(\frac{2\pi x}{T}\right) + B \sin\left(\frac{2\pi x}{T}\right)$.

Theorem 4.7.2 (Wirtinger's inequality). Let $f \in C^1([a, b])$ with $f(a) = f(b) = 0$. Then,

$$\int_a^b |f(x)|^2 dx \leq \frac{(b-a)^2}{\pi^2} \int_a^b |f'(x)|^2 dx.$$

Theorem 4.7.3 (Isoperimetric inequality). Let c be a simple and closed curve of class \mathcal{C}^1 whose length is L . If A_c is the area enclosed by c , then

$$A_c \leq \frac{L^2}{4\pi},$$

with equality if and only if c is a circle.