Module `fermi_eos`

Functions

```
def beta_equilibrium_condition(k_p, k_n)
```

Condition of beta equilibrium in order to compute k_p as a function of k_n. If it returns zero, then the Fermi momentum of the proton is the expected one, since it means that chemical potentials of neutrons, protons and electrons match the equilibrum condition for the (beta) weak interactions.

Parameters

`k_p` : `float`
Fermi momentum of protons in g cm / s.

`k_n` : `float`
Fermi momentum of neutrons in g cm / s.

Returns

`float`
Chemical potential in erg.

```
def compute_energy_density_emp(p)
```

Compute energy density from pressure.

Parameters

`p` : `float`
Pressure in dyne/cm^2.

Raises

`ValueError`
If energy density is negative.

Returns

`float`
Energy density in erg/cm^3.

```
def compute_energy_density_from_u_emp(u)
```

Compute energy density from u = n / n_0.

Parameters

u : float
Dimensionless u = n / n_0.

Returns

float
Energy density in erg/cm^3.

```
def compute_energy_density_from_u_sky(u)
```

Compute energy density from u = n / n_0.

Parameters

u : float
Dimensionless u = n / n_0.

Returns

float
Energy density in erg/cm^3.

```
def compute_energy_density_from_x_gen(x)
```

Compute energy density from x = k / (m c).

Parameters

x : float
Dimensionless x = k / (m c).

Raises

ValueError
If energy density is negative.

Returns

float
Energy density in erg/cm^3.

```
def compute_energy_density_from_x_npe(x, e_0)
```

Compute energy density from x = k / (mc) and prefactor e_0.

Parameters

x : float
Dimensionless x.

e_0 : float
Prefactor in erg/cm^3.

Returns

float
Energy density in erg/cm^3.

```
def compute_energy_density_gen(p)
```

Compute energy density from pressure.

Parameters: - p : float Pressure in dyne/cm^2.

Returns: - float Energy density in erg/cm^3.

```
def compute_energy_density_npe(k_n)
```

In presence of protons, electrons and neutrons, compute energy density from Fermi momentum of neutrons.

Parameters

k_n : float
Fermi momentum of neutrons in g cm / s.

Raises

ValueError
If energy density is negative.

Returns:

float
Energy density in erg/cm^3.

```
def compute_energy_density_pe(k_p)
```

In presence of protons and electrons (no neutrons), compute energy density from Fermi momentum of protons.

Parameters

k_n : float
Fermi momentum of protons in g cm / s.

Raises

`ValueError`
: If energy density is negative.

Returns

`float`
: Energy density in erg/cm^3.

```
def compute_energy_density_sky(p)
```

Compute energy density from pressure.

Parameters

`p` : `float`
: Pressure in dyne/cm^2.

Raises:

ValueError If energy density is negative.

Returns

`float`
: Number density in erg/cm^3.

```
def compute_fermi_momentum_nonrel(p)
```

Compute Fermi momentum from pressure in the non-relativistic limit.

Parameters

`p` : `float`
: Pressure in dyne/cm^2.

Returns

`float`
: Fermi momentum in g cm/s.

```
def compute_k_n(n)
```

Compute Fermi momentum of neutrons from number density.

Parameters

`n` : `float`
: Number density in 1/cm^3

Returns

`float`
Fermi momentum of neutrons in g cm / s.

```
def compute_k_p(k_n)
```

Compute k_p from k_n. he fsolve algorithm is used to find the right value of k_p that matches the beta equilibrium condition for chemical potential.

Parameters

`k_n` : `float`
Fermi momentum of neutrons in g cm / s.

Raises

`ValueError`
If Fermi momentum of protons is negative.

Returns

`float`
Fermi momentum of protons in g cm / s.

```
def compute_number_density_emp(p)
```

Compute number density from pressure.

Parameters

`p` : `float`
Pressure in dyne/cm^2.

Raises

`ValueError`
If number density is negative.

Returns

`float`
Number density in erg/cm^3.

```
def compute_number_density_gen(p)
```

Compute number density from pressure.

Compute number density from pressure.

Parameters: - p : float Pressure in dyne/cm^2.

Returns: - float Number density in erg/cm^3.

```
def compute_number_density_nonrel(p)
```

Compute number density from pressure in the non-relativistic limit.

Parameters

p : `float`
Pressure in dyne/cm^2.

Returns

`float`
Number density in 1/cm^3.

```
def compute_number_density_sky(p)
```

Compute number density from pressure.

Parameters

p : `float`
Pressure in dyne/cm^2.

Raises

`ValueError`
If number density is negative.

Returns:

float Energy density in erg/cm^3.

```
def compute_pressure_from_u_emp(u, p)
```

Compute calculated pressure minus expected pressure from the latter and u = n / n_0. If it returns zero, then the pressure is exactly the desidered one.

Parameters

u : `float`
Dimensionless u = n / n_0.

p : `float`
Pressure in dyne/cm^2.

Returns

`float`

Calculated pressure minus expected.

```
def compute_pressure_from_u_sky(u, p)
```

Compute calculated pressure minus expected pressure from expected pressure and u = n / n_0. If it returns zero, then the pressure is exactly the desidered one.

Parameters

u : `float`
Dimensionless u = n / n_0.

p : `float`
Pressure in dyne/cm^2.

Returns

`float`
Calculated pressure minus expected.

```
def compute_pressure_from_x_npe(x, e_0)
```

Compute pressure from x = k / (mc) and prefactor e_0.

Parameters

x : `float`
Dimensionless x.

e_0 : `float`
Prefactor in erg/cm^3.

Returns

`float`
Pressure in dyne/cm^2.

```
def compute_pressure_gen(x, p)
```

Compute calculated pressure minus expected pressure given the latter and x = k / (m c). If it returns zero, then the pressure is exactly the desidered one.

Parameters

x : `float`
Dimensionless x = k / (m c).

p : `float`
Pressure in dyne/cm^2.

Returns

`float`
Calculated pressure minus expected.

```
def compute_pressure_npe(k_n)
```

In presence of protons, electrons and neutrons, compute pressure from Fermi momentum of neutrons.

Parameters

`k_n` : `float`
Fermi momentum of neutrons in g cm / s.

Raises

`ValueError`
If pressure is negative.

Returns

`float`
Pressure in dyne/cm^2.

```
def compute_pressure_pe(k_p)
```

In presence of protons and electrons (no neutrons), compute pressure from Fermi momentum of protons.

Parameters

`k_p` : `float`
Fermi momentum of protons in g cm / s.

Raises

`ValueError`
If pressure is negative.

Returns

`float`
Pressure in dyne/cm^2.

```
def compute_u_emp(p)
```

Compute u = n / n_0 from pressure. The toms748 algorithm is used to find the right value of u that matches the expected pressure with the calculated one.

Parameters

p : `float`
Pressure in dyne/cm^2.

Raises

`ValueError`
If u is negative.

Returns

`float`
Dimensionless u = n / n_0.

```
def compute_u_sky(p)
```

Compute u = n / n_0 from pressure. The toms748 algorithm is used to find the right value of u that matches the expected pressure with the calculated one.

Parameters

u : `float`
Dimensionless u = n / n_0.

Raises

`ValueError`
If u is negative.

Returns

`float`
Energy density in erg/cm^3.

```
def compute_x_gen(p)
```

Compute x = k / (m c) from pressure. The toms748 algorithm is used to find the right value of x that matches the expected pressure with the calculated one.

Parameters

p : `float`
Pressure in dyne/cm^3.

Raises

`ValueError`
If x is negative.

Returns

`float`
Dimensionless x = k / (m c).

```
def equation_of_state_emp(pressures)
```

Compute and save to file the equation of state for empirical interactions. The equation of state is in the form energy density as function of pressure. Also number densities are computed and saved to file.

Parameters

`pressures` : `NumPy array`
Numpy array of pressures in dyne/cm^2.

Raises: - ValueError If pressure is negative.

```
def equation_of_state_gen(pressures)
```

Compute and save to file the equation of state for a generic degenerate ideal Fermi gas of neutrons. The equation of state is in the form energy density as function of pressure. Also number densities are computed and saved to file.

Parameters

`pressures` : `NumPy array`
Numpy array of pressures in dyne/cm^2.

Raises: - ValueError If pressure is negative.

```
def equation_of_state_nonrel(pressures)
```

Compute and save to file the equation of state for a non-relativistic degenerate ideal Fermi gas of neutrons. The equation of state is in the form energy density as function of pressure. Also number densities are computed and saved to file.

Parameters

`pressures` : `NumPy array`
Numpy array of pressures in dyne/cm^2.

Raises: - ValueError If pressure is negative.

```
def equation_of_state_npe(number_densities)
```

Compute and save to file the equation of state for a generic degenerate ideal Fermi gas of neutrons, protons and electrons in beta equilibrium. The equation of state is in the form energy density as function of pressure.

Parameters

`number_densities` : `NumPy array`
Numpy array of number densities in 1/cm^3.

Raises: - ValueError If number density is negative.

```
def equation_of_state_sky(pressures)
```

Compute and save to file the equation of state for Skyrme Hartee-Fock interactions. The equation of state is in the form energy density as function of pressure. Also number densities are computed and saved to file.

Parameters

pressures : NumPy array
Numpy array of pressures in dyne/cm^2.

Raises: - ValueError If pressure is negative.

```
def polytropic_equation_of_state(p, k,
gamma)
```

Compute energy density from pressure using polytropic equation of state. K is the prefactor and gamma is the exponent of energy density.

Parameters

p : float
Pressure in dyne/cm^2.

k : float
K in (dyne/cm^2)^(1 - gamma).

gamma : float
Dimensionless Gamma.

Returns

float
Energy density in erg/cm^3.

Functions

- compute_k_p
- compute_number_density_emp
- compute_number_density_gen
- compute_number_density_nonrel
- compute_number_density_sky
- compute_pressure_from_u_emp
- compute_pressure_from_u_sky
- compute_pressure_from_x_npe
- compute_pressure_gen
- compute_pressure_npe
- compute_pressure_pe
- compute_u_emp
- compute_u_sky
- compute_x_gen
- equation_of_state_emp
- equation_of_state_gen
- equation_of_state_nonrel
- equation_of_state_npe
- equation_of_state_sky
- polytropic_equation_of_state