

monotropism-scoring-levels-inertia

Scoring Monotropism: Inertia, "Friction," and Cognitive Energy — What the Evidence Actually Supports

A cited research brief — fact-checking the "scoring monotropism into levels" framing.

Informational, not medical advice. This brief evaluates how monotropism can and cannot be measured. The Monotropism Questionnaire (MQ) is a self-report research instrument, **not a diagnostic or clinical tool**, and it has no validated severity bands.

Executive summary

- **Your instinct is right on the core idea and wrong on the packaging.** The notion that monotropism can be *measured* along a spectrum, and that the real quantity of interest is the **effort ("friction") of switching attentional states**, is well supported — but the specific four-level taxonomy ("Low / Moderate / High / Extreme-Absolute") is a popular framing, **not a validated scientific classification**.
- **Monotropism is correctly a *dimensional trait*, not a syndrome.** The only validated measure — the **Monotropism Questionnaire (MQ)**, Garau et al. 2023 — returns a continuous score from 1 to 5. It has **no published cutoff, no percentile bands, and no severity levels** [2]. Binning it into named levels is always an arbitrary editorial choice.
- **The "friction" of switching already has a name: *autistic inertia*.** This is a real, increasingly researched construct — difficulty *starting, stopping, and switching* tasks that is "not within conscious control." It

was first articulated in autistic community narratives and is now in peer-reviewed work [3][4][5]. Your "switching friction" is essentially this.

- **"Attentional inertia" conflates two distinct things.** *Autistic inertia* (task-level: hard to start/stop/switch) and *sticky attention / impaired disengagement* (stimulus-level: hard to pull attention off something, e.g. Landry & Bryson 2004; Sacrey 2015 meta-analysis) are related but measured differently [6][7]. Lumping them under one term loses precision.
- **"Cognitive energy economics" is the single strongest framing** — and it already exists academically as Goldknopf's **"atypical resource allocation"** model (2013), which treats autism as a difference in *how much and where* attentional resources are spent [10]. This is the neuroaffirming, non-deficit version of what you are describing.
- **The burnout/shutdown/meltdown link is real but heuristic.** Autistic burnout (exhaustion, loss of skills, reduced tolerance) is an established construct, and the *narrative* link to forced switching + masking is compelling and community-validated — but the specific pathway "monotropism → tunnel rupture → burnout" is largely theoretical and first-person, not yet experimentally established [8][9].

1. Can monotropism be "scored"? Yes — as a continuum, not as levels

Monotropism is an attention-based account of autism: a monotropic mind funnels processing resources into a small number of intensely-aroused "attention tunnels" rather than spreading them across many channels [1]. Because it is described as a *difference in resource allocation*, it is naturally a **spectrum**, not a category — which is exactly why your "levels" intuition feels right.

The validated instrument is the **Monotropism Questionnaire (MQ)** — 47 items, scored on a 5-point Likert scale, developed in 2023 by Garau and colleagues *in collaboration with an autistic-led community group* [2]. Two things are decisive for your question:

- **It returns a single continuous mean (1–5), plus eight subscale means.** In the validation study (1,110 participants), autistic participants averaged **4.15** and non-autistic participants **3.19**, with

ADHD independently raising scores [2]. The two distributions overlap heavily — many non-autistic people out-score some autistic people, and vice versa.

- **There is no cutoff, no diagnostic threshold, no severity band.** A scan of the validation paper confirms the word "cutoff" appears only as a *statistical* citation (Hu & Bentler's fit-index criteria) and "sensitivity" only as a *sensitivity analysis* of how autism was defined [2]. The MQ's own authors do not propose levels.

So what does a score "mean"? Only its relative position on the dimension. The community-built online app that inspired this question helpfully translates a score into a *percentile* ("more monotropic than about X% of autistic / allistic people"), computed from the validation sample — and that is a reasonable, honest way to locate a score. But it is **a descriptive percentile, not a clinical band**: "the 97th percentile of monotropic tendency" is meaningful; "Level 3 / Extreme monotropism" is a label the evidence does not confer.

Bottom line: *score it, plot it, percentile it — but don't level it. The continuum is real; the named categories are not.*

2. Your "friction of switching" already exists: it's called *autistic inertia*

This is the most important correction-by-name in the brief. The phenomenon you describe — the cognitive cost of moving *into* and especially *out of* an attention tunnel — is an established construct called **autistic inertia**: a difficulty with **starting, stopping, and switching tasks that is not within conscious control** [3].

Three things make this directly relevant:

- **It originated in the autistic community and has entered the peer-reviewed literature.** The Association for Psychological Science describes autistic inertia as "a concept first articulated in autistic community narratives," now backed by qualitative and theoretical work [4]. Karen Buckle's first-hand-accounts study found participants repeatedly described being unable to start, stop, or change activities at will [3].

- **The literature literally uses your "friction" language.** The APS piece calls task initiation a "high-friction launch," frames switching as carrying "switching costs," and notes that once engaged, "switching focus requires more cognitive effort, sometimes leading to exhaustion" [4]. Your instinct to quantify *effort/friction* is the same move researchers are making.
- **It is now being formalised.** Recent peer-reviewed work treats the "stuck state" / limited response initiation as a multi-level phenomenon (muscular-behavioural "micro," relational and environmental "meso/macro") and co-creates support frameworks with autistic people [5]; others model it through predictive-processing / active-inference accounts (referenced in [4]).

In short: if you want to measure "switching friction," you are measuring autistic inertia — and that construct already has qualitative evidence, lived-experience validity, and a growing quantitative literature to build on.

3. "Attentional inertia" actually bundles two different mechanisms

Precision matters here, because the two mechanisms call for different language and (eventually) different measures.

3.1 Autistic inertia — task-level, about agency and momentum.

Difficulty *initiating*, *terminating*, and *switching* activities, experienced as disproportionate effort and a loss of voluntary control (the "object in motion stays in motion / at rest stays at rest" feeling) [3][4]. This is the closest fit to your "pull themselves out of the tunnel" idea.

3.2 Sticky attention / impaired disengagement — stimulus-level, about perception. Difficulty *pulling attention away* from a stimulus once locked on. Landry & Bryson (2004) showed young autistic children were markedly slower to disengage from a central stimulus when a peripheral one appeared — "sticky attention" [6]. A 2015 meta-analysis confirmed slowed attention disengagement across studies [7], and the same sluggish disengagement appears in infants later diagnosed autistic [6]. This is measured with experimental tasks (gap/overlap paradigms), not questionnaires.

Why the distinction matters: sticky attention is a *perceptual* capture effect (you literally don't notice the interruption); autistic inertia is a *control/momentum* effect (you notice but can't shift). Your four-level descriptions actually slide between the two — e.g. "background stimuli are genuinely not processed by the brain" is sticky attention [6], whereas "forcible transitions feel like pulling a train off its tracks" is inertia [3]. A careful model should keep them labelled separately, because they may have different causes and different supports.

4. The four proposed "levels" — what's supported, what's invented

Rather than endorse or discard the taxonomy outright, here is each proposed band tested against the evidence.

Proposed level	Core claim	Evidence verdict
Low (polytropic-dominant)	Many "tabs" open; near-zero switching cost	Supported as one end of the continuum. The MQ's lower scorers and the non-autistic group mean (3.19) describe exactly this [2]. "Polytropic" is the theory's own term for it [1].
Moderate	Can focus deeply but retains baseline awareness; mild switching cost	Supported as the middle of the dimension. Flow states are reported across the population; the MQ scores between the group means capture this band [2][11]. But the <i>name</i> is arbitrary.
High	Preferred single intense tunnel; interruptions genuinely missed; high switching cost	Mechanism supported, category not. The components are real — missed interruptions = sticky attention [6][7]; high switching cost = autistic inertia [3][4]; strong autism/ADHD association = MQ findings [2]. But there is no threshold separating "High" from "Moderate."
Extreme / Absolute	Total immersion; disruption causes "neurological shock," meltdowns, shutdowns	Overstated. Deep immersion ("flow") and distress at disruption are real [11], but "absolute" implies a qualitative jump the data don't show , and "neurological shock" is not a recognised term. Meltdowns/shutdowns are involuntary overwhelm responses, not a "shock" specific to a monotropism band.

Three problems with publishing the taxonomy as-is:

1. **It implies categorical jumps where the data show a smooth continuum.** The MQ distributions of autistic and non-autistic people overlap substantially [2]; there is no natural "knee" that would justify four discrete levels.
2. **It smuggles in severity/pathology.** "Extreme/absolute ... meltdowns, burnout" frames the high end as inherently harmful — the opposite of the neuroaffirming framing the theory intends [1]. Higher monotropism is *not* the same as worse outcomes; outcomes depend on *environmental fit*, not level.
3. **The bands have no reliability data.** A useful level system needs to show that people land in the same band on re-test and that bands predict something. No such study exists for monotropism.

What to do instead. Express the same idea as a *position on the dimension* plus a *profile*: "high monotropic tendency ($\approx$ top X% of the autistic sample), with pronounced autistic inertia on task-switching and frequent disruption-driven meltdowns in unsupportive environments." That is accurate, individualised, and non-pathologising.

5. Disruption → meltdown, shutdown, burnout: what the evidence supports

This part of your framing is the most consequential and needs the most care.

Meltdowns and shutdowns are real and well-described as involuntary responses to overwhelming stress or sensory/cognitive load (an externalising "meltdown" vs an internalising "shutdown" withdrawal) — recognised in clinical guidance and community consensus. They are triggered by *overwhelm*, of which tunnel-disruption is one common route [4][8].

Autistic burnout is now a defined construct. Two foundational studies — Raymaker et al. (2020) and Higgins et al. (2021, a Grounded-Delphi study with autistic experts by experience) — converge on a definition: **chronic exhaustion, loss of skills, and reduced tolerance to stimulus**, conceptualised as resulting from chronic life stress and "a mismatch of expectations and abilities" without adequate support and recovery [8][9].

The monotropism → burnout link is heuristically powerful but not yet experimentally established. The narrative is compelling and community-validated: every forced transition is a tunnel rupture; constant masking and switching in a polytropic world burns resources; without recovery/flow time, the system exhausts [8][9][11]. But the *specific* contribution of monotropic tunnel-disruption to burnout — versus masking, trauma, sensory load, and chronic demand — has not been isolated experimentally. State it as a *strong, testable hypothesis*, not an established fact.

The honest, evidence-based claim is: "Autistic burnout is driven by chronic stress, masking, and a mismatch between demands and capacity; the monotropism framework predicts that constant forced switching and tunnel-disruption are a major, under-recognised contributor — a prediction that fits lived experience and awaits direct testing."

6. "Cognitive energy economics" — your strongest framing

Your closing move — shifting from "deficit" to "how much fuel a brain burns switching gears" — is exactly right, and it already has an academic home.

Goldknopf's "**atypical resource allocation**" model (2013) operationalises monotropism as a difference in *resource economics*: attentional resources in autism are **narrowed** (concentrated into fewer/smaller channels, sometimes more intensely) and/or **reduced in overall capacity** [10]. On this view, the "cost" of a transition is literally the energetic expense of breaking one allocation pattern and building another. That is "cognitive energy economics" stated precisely — and it is the framing most consistent with a non-pathologising account, because it frames the problem as *spending* (a difference in budgeting) rather than *brokenness*.

This is the framing to keep. Three advantages:

- It is **neutral about value**: intense focus is an asset when the environment supports it (flow, expertise) and a cost when the environment demands constant switching [10][11].

- It **predicts intervention**: reduce unnecessary transitions, protect flow/recovery time, match demands to capacity, and the "fuel bill" drops — which is precisely what burnout-prevention guidance recommends [8][9].
- It **unifies** the phenomena in your four levels without needing the levels: missed stimuli, switching cost, overwhelm, and burnout all become consequences of *how resources are allocated and depleted*, on a continuum.

7. Practical takeaways — what to claim, and what to avoid

Claim confidently:

- Monotropism is a measurable, **dimensional** trait (MQ, 1–5, with eight subscales); report the score and optional percentile, not a "level" [2].
- The "**switching friction**" you describe is **autistic inertia** — a real, community-originated, now-peer-reviewed construct [3][4][5].
- **Sticky attention** / **impaired disengagement** is a distinct, experimentally-measured mechanism [6][7].
- **Autistic burnout** is a defined construct driven by chronic stress, masking, and demand–capacity mismatch; monotropic tunnel-disruption is a *plausible, lived-experience-validated* contributor [8][9].
- "Cognitive energy economics" $\approx$ Goldknopf's **resource allocation** model — the most accurate, neuroaffirming framing [10].

Avoid or re-label:

- Named **levels** / **bands** ("Low/Moderate/High/Extreme") — not validated; use position-on-dimension + profile instead.
- "**Attentional inertia**" as an umbrella — it blurs autistic inertia (task/control) with sticky attention (perception); use the precise term [3][6].
- "**Absolute**" **monotropism** and "**neurological shock**" — not recognised terms; imply a categorical jump and a pathology the data don't support.
- Stating the **monotropism** $\rightarrow$ **burnout** pathway as established — it is a strong hypothesis, not a measured effect [8].

- Any framing where **higher = more disordered** — monotropism's outcomes depend on environmental fit, not level [1][10].

Sources

[1] Murray, D., Lesser, M., & Lawson, W. (2005). *Attention, monotropism and the diagnostic criteria for autism*. *Autism*, 9(2), 139–156.

doi:10.1177/1362361305051398 — origin of monotropism theory; frames it as a difference in resource allocation (a continuum), not a deficit.

<https://doi.org/10.1177/1362361305051398>

[2] Garau, V., Woods, R., Chown, N., Hallett, S., Murray, F., Wood, R., Murray, A. & Fletcher-Watson, S. (2023). *The Monotropism Questionnaire* (47-item self-report measure; validation preprint). Open Science Framework. — dimensional 1–5 score + 8 subscales; n=1,110; autistic mean 4.15 vs non-autistic 3.19; ADHD raises scores; **no cutoff/percentile band/severity level** (confirmed by scanning the validation manuscript).

Questionnaire: <https://osf.io/wpx5g> · validation preprint:

<https://osf.io/ft73y> · CC BY-NC-SA 4.0. Local archive: [sources/17-monotropism-questionnaire-osf.pdf](#)

[3] Buckle, K. et al. (2021). *"No way out except from external intervention": First-hand accounts of autistic inertia*. (Preprint / qualitative).

doi:10.31234/osf.io/ahk6x — autistic people describe difficulty starting, stopping, and changing activities "that was not within their conscious control." Local archive: [sources/11-autistic-inertia-osf.pdf](#) ·

<https://doi.org/10.31234/osf.io/ahk6x> · Manchester Research Explorer: <https://research.manchester.ac.uk/en/publications/no-way-out-except-from-external-intervention-first-hand-accounts->

[4] Srinivasan, D. (2025). *The Physics of Autistic Inertia* (Student Notebook). APS Observer, Association for Psychological Science. — frames autistic inertia as "first articulated in autistic community narratives," now peer-reviewed; uses "high-friction launch" and "switching costs" language; cites Carls-Diamante & Laciny (2024), Greaves-Lord et al. (2023), Buckle et al. (2021), Heasman et al. (2024), Rapaport et al. (2024), Phung et al. (2021).

<https://www.psychologicalscience.org/publications/observer/student-notebook-autistic-inertia-srinivasan.html>

- [5] Greaves-Lord, L. et al. (2023). *[Getting 'stuck': an integrative approach to support autistic people]* *Frontiers in Psychology*, doi:10.3389/fpsyg.2023.1229596. — peer-reviewed qualitative + Delphi framework on limited response initiation ("stuck state") in autistic people across micro/meso/macro levels, co-created with autistic people. <https://doi.org/10.3389/fpsyg.2023.1229596>
- [6] Landry, R. & Bryson, S. (2004). *Impaired disengagement of attention in young children with autism*. *Journal of Child Psychology and Psychiatry*, 45(6), 1115–1122. doi:10.1111/j.1469-7610.2004.00304.x — "sticky attention": slowed disengagement from a stimulus. <https://doi.org/10.1111/j.1469-7610.2004.00304.x>
- [7] Sacrey, L.-A. et al. (2015). *Attention disengagement in autism spectrum disorder: A meta-analysis*. *Neuroscience & Biobehavioral Reviews*, 56, 111–121. doi:10.1016/j.neubiorev.2014.10.011 — confirms slowed attention disengagement across studies. <https://doi.org/10.1016/j.neubiorev.2014.10.011>
- [8] Raymaker, D. M. et al. (2020). *"Having all of your internal resources exhausted beyond measure and being left with no clean-up crew": Defining autistic burnout*. *Autism in Adulthood*, 2(2), 132–143. doi:10.1089/aut.2019.0079 — defines autistic burnout as chronic exhaustion, loss of skills, reduced tolerance to stimulus, from chronic life stress and a mismatch of expectations and abilities. <https://journals.sagepub.com/doi/abs/10.1089/aut.2019.0079> · open archive: https://pdxscholar.library.pdx.edu/socwork_fac/378
- [9] Higgins, J. M. et al. (2021). *Defining autistic burnout through experts by lived experience: Grounded Delphi method investigating #AutisticBurnout*. *Autism*, 25(8). doi:10.1177/13623613211019858 — consensus definition: exhaustion, withdrawal, problems with thinking, reduced daily living skills, increased autistic traits. <https://www.autismcrc.com.au/knowledge-centre/publications/defining-autistic-burnout-using-grounded-delphi-method-autistic>
- [10] Goldknopf, E. J. (2013). *Atypical resource allocation may contribute to many aspects of autism*. *Frontiers in Integrative Neuroscience*. PMC3872719 — models autism as a difference in *resource allocation* (narrowed and/or reduced attentional resources); the academic form of "cognitive energy economics." <https://pmc.ncbi.nlm.nih.gov/articles/PMC3872719>

[11] Rapaport, H., Clapham, H., Adams, J., Lawson, W., Porayska-Pomsta, K., & Pellicano, E. (2024). *"In a State of Flow": experiential validity of monotropism and its relation to autistic wellbeing*. Autism in Adulthood. doi:10.1089/aut.2023.0032 — the positive pole of deep monotropic immersion (flow); supports the "asset when supported" framing. Local archive: `sources/18-flow-experiential-validity-pmc.html` · <https://doi.org/10.1089/aut.2023.0032>