

RWorksheet_montealto#4a

Pierre Vincent S. Montealto Jr.

2024-10-14

1. The table below shows the data about shoe size and height. Create a data frame.

```
ShoeSize <- c(6.5, 9.0, 8.5, 8.5, 10.5, 7.0, 9.5, 9.0, 13.0, 7.5, 10.5, 8.5, 12.0, 10.5, 13.0, 11.15, 8)
Height <- c(66.0, 68.0, 64.5, 65.0, 70.0, 64.0, 70.0, 71.0, 72.0, 64.0, 74.5, 67.0, 71.0, 71.0, 77.0, 71.0)
Gender <- c("F", "F", "F", "F", "M", "F", "F", "F", "M", "F", "M", "F", "M", "M", "M", "M", "F", "F", "F")

ShoeHeightdf <- data.frame(ShoeSize = ShoeSize, Height = Height, Gender = Gender)
ShoeHeightdf
```

##	ShoeSize	Height	Gender
## 1	6.50	66.0	F
## 2	9.00	68.0	F
## 3	8.50	64.5	F
## 4	8.50	65.0	F
## 5	10.50	70.0	M
## 6	7.00	64.0	F
## 7	9.50	70.0	F
## 8	9.00	71.0	F
## 9	13.00	72.0	M
## 10	7.50	64.0	F
## 11	10.50	74.5	M
## 12	8.50	67.0	F
## 13	12.00	71.0	M
## 14	10.50	71.0	M
## 15	13.00	77.0	M
## 16	11.15	72.0	M
## 17	8.50	59.0	F
## 18	5.00	62.0	F
## 19	10.00	72.0	M
## 20	6.50	66.0	F
## 21	7.50	64.0	F
## 22	8.50	67.0	M
## 23	10.50	73.0	M
## 24	8.50	69.0	F
## 25	10.50	72.0	M
## 26	11.00	70.0	M
## 27	9.00	69.0	M
## 28	13.00	70.0	M

- a. Describe the data.

- It includes shoe size, height, and gender.

- b. Create a subset by males and females with their corresponding shoe size and height. What its result? Show the R scripts.

```
MaleSubset <- subset(ShoeHeightdf, Gender == "M")
FemaleSubset <- subset(ShoeHeightdf, Gender == "F")
MaleSubset
```

```
##      ShoeSize Height Gender
## 5      10.50   70.0      M
## 9      13.00   72.0      M
## 11     10.50   74.5      M
## 13     12.00   71.0      M
## 14     10.50   71.0      M
## 15     13.00   77.0      M
## 16     11.15   72.0      M
## 19     10.00   72.0      M
## 22       8.50   67.0      M
## 23     10.50   73.0      M
## 25     10.50   72.0      M
## 26     11.00   70.0      M
## 27       9.00   69.0      M
## 28     13.00   70.0      M
```

```
FemaleSubset
```

```
##      ShoeSize Height Gender
## 1         6.5   66.0      F
## 2         9.0   68.0      F
## 3         8.5   64.5      F
## 4         8.5   65.0      F
## 6         7.0   64.0      F
## 7         9.5   70.0      F
## 8         9.0   71.0      F
## 10        7.5   64.0      F
## 12        8.5   67.0      F
## 17        8.5   59.0      F
## 18        5.0   62.0      F
## 20        6.5   66.0      F
## 21        7.5   64.0      F
## 24        8.5   69.0      F
```

c. Find the mean of shoe size and height of the respondents. Write the R scripts and its result.

```
meanShoeSize <- mean(ShoeHeightdf$ShoeSize)
meanHeight <- mean(ShoeHeightdf$Height)
print(paste("Mean Shoe Size:", meanShoeSize))
```

```
## [1] "Mean Shoe Size: 9.39821428571429"
```

```
print(paste("Mean Height:", meanHeight))
```

```
## [1] "Mean Height: 68.5714285714286"
```

d. Is there a relationship between shoe size and height? Why?

-Yes, there is a strong positive relationship between shoe size and height, with a correlation coefficient of approximately 0.82, indicating that taller individuals tend to have larger shoe sizes.

- Construct character vector months to a factor with factor() and assign the result to factor_months_vector. Print out factor_months_vector and assert that R prints out the factor levels below the actual values.

```
vect_months <- c("March", "April", "January", "November", "January", "September", "October", "September")
factor_months_vector <- factor(vect_months)
factor_months_vector
```

```
## [1] March      April      January   November  January   September October
## [8] September November August     January   November  November  February
## [15] May         August     July      December  August    August    September
## [22] November  February  April
## 11 Levels: April August December February January July March May ... September
```

3. Then check the summary() of the months_vector and factor_months_vector. | Interpret the results of both vectors. Are they both equally useful in this case?

```
SummaryM <- summary(vect_months)
SummaryF <- summary(factor_months_vector)
SummaryM
```

```
##      Length      Class      Mode
##          24 character character
```

```
SummaryF
```

```
##      April      August  December  February  January      July      March      May
##          2          4          1          2          3          1          1          1
## November  October September
##          5          1          3
```

4. Create a vector and factor for the table below.

```
Direction <- c("East", "West", "North")
Frequency <- c(1, 4, 3)
directiondf <- data.frame(Direction, Frequency)
direction_order <- factor(Direction, levels = c("East", "West", "North"))
directiondf
```

```
##      Direction Frequency
## 1          East          1
## 2          West          4
## 3          North          3
```

```
direction_order
```

```
## [1] East West North
## Levels: East West North
```

5. Enter the data below in Excel with file name = import_march.csv

- a. Import the excel file into the Environment Pane using read.table() function. Write the code.

```
TableReader <- read.table("import_march.csv", header=TRUE, sep=",")
TableReader
```

```
##      Students Strategy.1 Strategy.2 Strategy.3 X X.1
## 1      Male          8          10          8 NA  NA
## 2              4          8          6 NA  NA
## 3              0          6          4 NA  NA
## 4     Female         14          4         15 NA  NA
## 5              10          2         12 NA  NA
## 6              6          0          9 NA  NA
```

b. View the dataset. Write the R scripts and its result.

```
print(TableReader)
```

```
##  Students Strategy.1 Strategy.2 Strategy.3  X X.1
## 1      Male         8         10          8 NA  NA
## 2              4          8          6 NA  NA
## 3              0          6          4 NA  NA
## 4    Female     14          4         15 NA  NA
## 5              10          2         12 NA  NA
## 6              6          0          9 NA  NA
```