

# Existence of a least increasing surjection from an artinian poset onto a set of ordinals<sup>1</sup>

Pierre-Yves Gaillard

Let  $X$  be a set. If  $g : X \rightarrow S$  and  $h : X \rightarrow T$  are surjections and  $S$  and  $T$  are sets of ordinals, say that  $g \leq h$  if  $g(x) \leq h(x)$  for all  $x$ . If  $X$  is a poset, say that  $g$  is **increasing** if  $x < y$  implies  $g(x) < g(y)$ .

**Theorem 1.** *If  $X$  is an artinian poset, then there is a least increasing surjection  $f$  from  $X$  onto a set of ordinals. For  $x$  in  $X$ , set  $X_x := \{y \in X \mid y < x\}$ . Then  $f$  is the unique surjection from  $X$  onto a set  $S$  of ordinals such that, setting  $\alpha_x := \sup f(X_x)$ , we have*

$$f(x) = \begin{cases} \alpha_x + 1 & \text{if } \alpha_x \in f(X_x) \\ \alpha_x & \text{if } \alpha_x \notin f(X_x) \end{cases} \quad (1)$$

for all  $x$  in  $X$ .

**Lemma 2.** *There is at most one such surjection  $f$ .*

*Proof.* Let  $f$  and  $g$  satisfy (1) and let  $x$  be a minimal element of  $X$  such that  $f(x) \neq g(x)$ . Then (1) yields a contradiction.  $\square$

**Lemma 3.** *Let  $A \subset X$  be the set of all  $x$  in  $X$  such that there is a surjection  $f_x : X_x \rightarrow S(x)$  s.t.  $S(x)$  is a set of ordinals, and, setting  $\alpha_{xy} := \sup f_x(X_y)$  for all  $y$  in  $X_x$ , we have*

$$f_x(y) = \begin{cases} \alpha_{xy} + 1 & \text{if } \alpha_{xy} \in f_x(X_y) \\ \alpha_{xy} & \text{if } \alpha_{xy} \notin f_x(X_y) \end{cases} \quad (2)$$

for all  $y$  in  $X_x$ . Then  $A = X$ .

*Proof.* We assume by contradiction  $A \neq X$ . Let  $x$  be minimal in  $X \setminus A$ . So  $f_x$  doesn't exist, but  $f_y$  exists for all  $y < x$ . We will define  $f_x$ . Let  $y$  be less than  $x$ . We must define  $f_x(y)$ . If  $y < z < x$  for some  $z$ , we set  $f_x(y) := f_z(y)$ . By Lemma 2, this is well-defined. Assume now that no such  $z$  exists. Then we set  $\beta_y = \sup f_y(X_y)$  and

$$f_x(y) = \begin{cases} \beta_y + 1 & \text{if } \beta_y \in f_y(X_y) \\ \beta_y & \text{if } \beta_y \notin f_y(X_y). \end{cases} \quad (3)$$

Now  $f_x$  is defined, and we must show that it satisfies (2). Let  $y$  be in  $X_x$ .

If  $y < z < x$  for some  $z$ , then  $f_x(y)$  is defined by  $f_x(y) := f_z(y)$ , and we have  $\alpha_{xy} = \alpha_{zy}$ . Since  $f_z(y)$  satisfies (2) with  $z$  instead of  $x$  by assumption, we are done.

If no such  $z$  exists, then  $f_x(y)$  is defined by (3), and it suffices to show  $\beta_y = \alpha_{xy}$ . Let  $t$  be in  $X_y$ . It suffices to show  $f_x(t) = f_y(t)$ , but this follows immediately from the definition of  $f_x(t)$ .

This shows that  $f_x$  satisfies (2), contradiction.  $\square$

*Proof of Theorem 1.* It suffices to show that there is a surjection  $f$  from  $X$  onto a set  $S$  of ordinals satisfying (1). Set  $Y := X \cup \{\infty\}$  with  $\infty \notin X$  and  $\infty > x$  for all  $x$  in  $X$  (and  $x \leq x'$  in  $Y$  iff  $x \leq x'$  in  $X$  for all  $x, x'$  in  $X$ ). In particular,  $Y$  is artinian, and we have  $Y_\infty = X$ . Applying Lemma 3 to  $Y$ , we get a surjection  $f_\infty : X \rightarrow S$  satisfying (1).  $\square$

---

<sup>1</sup>This text is available at <https://github.com/Pierre-Yves-Gaillard/Artin-poset-to-ordinals>. Here is a related text: <https://github.com/Pierre-Yves-Gaillard/The-Transfinite-Recursion-Theorem>