

Technical Data Sheet

Purified Rat Anti-Mouse PNAd Carbohydrate Epitope

Product Information

Material Number:	553863
Alternate Name:	CD62L Ligand
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	MECA-79
Immunogen:	Collagenase-dispersed BALB/c lymph node stroma
Isotype:	Rat (WF) IgM, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The MECA-79 antibody reacts with sulfate-dependent carbohydrate epitopes of peripheral lymph node addressin (PNAd). The MECA-79-reactive antigen is closely associated with the carbohydrate ligands for L-selectin (eg, CD34, GlyCAM-1, MAdCAM-1), which are expressed on high endothelial venules (HEV) in lymphoid tissues and at sites of chronic inflammation. Cross-reactivity with human, sheep, cow, primate, and pig tissues has been observed. MECA-79 antibody inhibits L-selectin-dependent lymphocyte and platelet homing to lymph nodes *in vivo*, and *in vitro* adhesion to lymphoid tissue HEV and immobilized PNAd.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.
Store undiluted at 4°C.

Application Notes

Application

Immunohistochemistry-paraffin	Routinely Tested
Immunohistochemistry-frozen	Reported
Immunoprecipitation	Reported
Western blot	Reported
Blocking	Reported
Immunohistochemistry-zinc-fixed	Reported

Recommended Assay Procedure:

This antibody has been tested by immunohistochemical staining (IHC) of citrate-pretreated formalin-fixed paraffin-embedded sections (5 - 20 μ g/ml) to assure specificity and reactivity. Other reported applications include IHC of acetone-fixed frozen sections, immunoprecipitation, western blot analysis, and *in vitro* and *in vivo* adhesion blocking.

Suggested Companion Products

Catalog Number	Name	Size	Clone
553940	Purified Rat IgM, κ Isotype Control	0.5 mg	R4-22
554016	FITC Goat Anti-Rat Ig	0.5 mg	Polyclonal

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Sodium azide is a reversible inhibitor of oxidative metabolism; therefore, antibody preparations containing this preservative agent must not be used in cell cultures nor injected into animals. Sodium azide may be removed by washing stained cells or plate-bound antibody or dialyzing soluble antibody in sodium azide-free buffer. Since endotoxin may also affect the results of functional studies, we recommend the NA/LE (No Azide/Low Endotoxin) antibody format, if available, for *in vitro* and *in vivo* use.

References

Berg EL, Robinson MK, Warnock RA, Butcher EC. The human peripheral lymph node vascular addressin is a ligand for LECAM-1, the peripheral lymph node homing receptor. *J Cell Biol.* 1991; 114(2):343-349. (Clone-specific: Blocking, Immunoprecipitation)
 Binns RM, Whyte A, Licence ST. The role of E-selectin in lymphocyte and polymorphonuclear cell recruitment into cutaneous delayed hypersensitivity reactions in sensitized pigs. *J Immunol.* 1996; 157(9):4094-4099. (Biology)
 Diacovo TG, Puri KD, Warnock RA, Springer TA, von Andrian UH. Platelet-mediated lymphocyte delivery to high endothelial venules. *Science.* 1996; 273(5272):252-255. (Clone-specific: Blocking)

BD Biosciences

bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean
 877.232.8995 866.979.9408 32.2.400.98.95 0120.8555.90 65.6861.0633 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

© 2017 BD. BD, the BD Logo and all other trademarks are property of Becton, Dickinson and Company.

553863 Rev. 7



Faveeuw C, Gagnerault MC, Lepault F. Expression of homing and adhesion molecules in infiltrated islets of Langerhans and salivary glands of nonobese diabetic mice. *J Immunol.* 1994; 152(12):5969-5978. (Biology)

Hemmerich S, Butcher EC, Rosen SD. Sulfation-dependent recognition of high endothelial venules (HEV)-ligands by L-selectin and MECA 79, and adhesion-blocking monoclonal antibody. *J Exp Med.* 1994; 180(6):2219-2226. (Clone-specific: Immunoprecipitation)

Malý P, Thall A, Petryniak B, et al. The alpha(1,3)fucosyltransferase Fuc-TVII controls leukocyte trafficking through an essential role in L-, E-, and P-selectin ligand biosynthesis. *Cell.* 1996; 86(4):643-653. (Biology)

Michie SA, Streeter PR, Bolt PA, Butcher EC, Picker LJ. The human peripheral lymph node vascular addressin. An inducible endothelial antigen involved in lymphocyte homing. *Am J Pathol.* 1993; 143(6):1688-1698. (Clone-specific: Blocking)

Puri KD, Finger EB, Gaudernack G, Springer TA. Sialomucin CD34 is the major L-selectin ligand in human tonsil high endothelial venules. *J Cell Biol.* 1995; 131(1):261-270. (Clone-specific: Blocking, Western blot)

Streeter PR, Rouse BT, Butcher EC. Immunohistologic and functional characterization of a vascular addressin involved in lymphocyte homing into peripheral lymph nodes. *J Cell Biol.* 1988; 107(5):1853-1862. (Immunogen: Blocking, Immunohistochemistry)