

# Parametric estimation via MMD optimization: robustness to outliers and to dependence

Pierre Alquier



Center for  
Advanced Intelligence Project

CDT in Modern Statistics and Statistical Machine Learning  
Imperial College London and University of Oxford  
August 18, 2020

# RIKEN AIP : ABI team



Approximate Bayesian  
Inference team (ABI), lead  
by Emtiyaz Khan



Please visit the team website

<https://emtiyaz.github.io/>

# Co-authors



Badr-Eddine Chérif-Abdellatif

ENSAE Paris → Oxford



Mathieu Gerber

Univ. of Bristol



Alexis Derumigny

Univ. of Twente



Jean-David Fermanian

ENSAE Paris

# The Maximum Likelihood Estimator (MLE)

Let  $X_1, \dots, X_n$  be i.i.d in  $\mathcal{X}$  from a probability distribution  $P_0$ .

# The Maximum Likelihood Estimator (MLE)

Let  $X_1, \dots, X_n$  be i.i.d in  $\mathcal{X}$  from a probability distribution  $P_0$ .

Statistical inference :

- propose a model  $(P_\theta, \theta \in \Theta)$ , assume  $P_0 = P_{\theta_0}$ .
- compute  $\hat{\theta}_n = \hat{\theta}_n(X_1, \dots, X_n)$ .

# The Maximum Likelihood Estimator (MLE)

Let  $X_1, \dots, X_n$  be i.i.d in  $\mathcal{X}$  from a probability distribution  $P_0$ .

Statistical inference :

- propose a model  $(P_\theta, \theta \in \Theta)$ , assume  $P_0 = P_{\theta_0}$ .
- compute  $\hat{\theta}_n = \hat{\theta}_n(X_1, \dots, X_n)$ .

Letting  $p_\theta$  denote the density of  $P_\theta$ , then

$$\hat{\theta}_n^{MLE} = \arg \max_{\theta \in \Theta} L(\theta), \text{ where } L(\theta) = \prod_{i=1}^n p_\theta(X_i).$$

# The Maximum Likelihood Estimator (MLE)

Let  $X_1, \dots, X_n$  be i.i.d in  $\mathcal{X}$  from a probability distribution  $P_0$ .

Statistical inference :

- propose a **model** ( $P_\theta, \theta \in \Theta$ ), assume  $P_0 = P_{\theta_0}$ .
- compute  $\hat{\theta}_n = \hat{\theta}_n(X_1, \dots, X_n)$ .

Letting  $p_\theta$  denote the density of  $P_\theta$ , then

$$\hat{\theta}_n^{MLE} = \arg \max_{\theta \in \Theta} L(\theta), \text{ where } L(\theta) = \prod_{i=1}^n p_\theta(X_i).$$


**Example :**  $P_{(m,\sigma)} = \mathcal{N}(m, \sigma^2)$  then

$$\hat{m} = \frac{1}{n} \sum_{i=1}^n X_i \text{ and } \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \hat{m})^2.$$

# MLE not unique / not consistent

Example :

$$p_{\theta}(x) = \frac{\exp(-|x - \theta|)}{2\sqrt{\pi}|x - \theta|},$$







# MLE not unique / not consistent

Example :

$$p_{\theta}(x) = \frac{\exp(-|x - \theta|)}{2\sqrt{\pi}|x - \theta|},$$

$$L(\theta) = \frac{\exp(-\sum_{i=1}^n |X_i - \theta|)}{(2\sqrt{\pi})^n \prod_{i=1}^n \sqrt{|X_i - \theta|}}.$$



# MLE fails in the presence of outliers

What is an outlier ?

# MLE fails in the presence of outliers

## What is an outlier?

Huber proposed the **contamination** model : with probability  $\varepsilon$ ,  $X_i$  is not drawn from  $P_{\theta_0}$  but from  $Q$  that can be **anything** :

$$P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q.$$

# MLE fails in the presence of outliers

## What is an outlier?

Huber proposed the **contamination** model : with probability  $\varepsilon$ ,  $X_i$  is not drawn from  $P_{\theta_0}$  but from  $Q$  that can be **anything** :

$$P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q.$$

**Example** :  $P_\theta = \text{Unif}[0, \theta]$ , then

# MLE fails in the presence of outliers

## What is an outlier ?

Huber proposed the **contamination** model : with probability  $\varepsilon$ ,  $X_i$  is not drawn from  $P_{\theta_0}$  but from  $Q$  that can be **anything** :

$$P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q.$$

**Example** :  $P_\theta = \text{Unif}[0, \theta]$ , then

$$L(\theta) = \frac{1}{\theta^n} \prod_{i=1}^n 1_{\{0 \leq X_i \leq \theta\}} \Rightarrow \hat{\theta} = \max_{1 \leq i \leq n} X_i.$$

# MLE fails in the presence of outliers

## What is an outlier?

Huber proposed the **contamination** model : with probability  $\varepsilon$ ,  $X_i$  is not drawn from  $P_{\theta_0}$  but from  $Q$  that can be **anything** :

$$P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q.$$

**Example** :  $P_\theta = \mathcal{U}nif[0, \theta]$ , then

$$L(\theta) = \frac{1}{\theta^n} \prod_{i=1}^n 1_{\{0 \leq X_i \leq \theta\}} \Rightarrow \hat{\theta} = \max_{1 \leq i \leq n} X_i.$$

In the case of the following contamination, the MLE is extremely far from the truth :

$$P_0 = (1 - \varepsilon).\mathcal{U}nif[0, 1] + \varepsilon.\mathcal{N}(10^{10}, 1)...$$

# Requirements for a “good” estimator

A **universal** estimator  $\hat{\theta}_n$  must be such that, for **some distance  $d$  on probability distributions**,

# Requirements for a “good” estimator

A **universal** estimator  $\hat{\theta}_n$  must be such that, for **some distance  $d$  on probability distributions**,

- 1 when the model is well specified, that is,  $P_0 = P_{\theta_0}$ ,

$$\mathbb{E} [d(P_{\hat{\theta}_n}, P_0)] \leq r_n(\Theta) \xrightarrow{n \rightarrow \infty} 0.$$



# Requirements for a “good” estimator

A **universal** estimator  $\hat{\theta}_n$  must be such that, for **some distance  $d$  on probability distributions**,

- 1 when the model is well specified, that is,  $P_0 = P_{\theta_0}$ ,

$$\mathbb{E} [d(P_{\hat{\theta}_n}, P_0)] \leq r_n(\Theta) \xrightarrow{n \rightarrow \infty} 0.$$

- 2 in the misspecified case  $P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q$ , **for any  $Q$** ,

$$\mathbb{E} [d(P_{\hat{\theta}_n}, P_0)] \leq \underbrace{d(P_0, P_{\theta_0})}_{\xrightarrow{\varepsilon \rightarrow \infty} 0} + \underbrace{r_n(\Theta)}_{\xrightarrow{n \rightarrow \infty} 0}.$$

# Requirements for a “good” estimator

A **universal** estimator  $\hat{\theta}_n$  must be such that, for **some distance  $d$  on probability distributions**,

- 1 when the model is well specified, that is,  $P_0 = P_{\theta_0}$ ,

$$\mathbb{E} [d(P_{\hat{\theta}_n}, P_0)] \leq r_n(\Theta) \xrightarrow{n \rightarrow \infty} 0.$$

- 2 in the misspecified case  $P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q$ , **for any  $Q$** ,

$$\mathbb{E} [d(P_{\hat{\theta}_n}, P_0)] \leq \underbrace{d(P_0, P_{\theta_0})}_{\xrightarrow{\varepsilon \rightarrow 0} 0} + \underbrace{r_n(\Theta)}_{\xrightarrow{n \rightarrow \infty} 0}.$$

The MLE does not satisfy these requirements.

# Some examples

Yatracos' skeleton estimate  $\hat{\theta}_n^Y$  :

$$\mathbb{E} \left[ d_{TV}(P_{\hat{\theta}_n^Y}, P_0) \right] \leq 3d_{TV}(P_0, P_{\theta_0}) + C \cdot \sqrt{\frac{\dim(\Theta)}{n}}$$

where

$$d_{TV}(P, Q) = \sup_E |P(E) - Q(E)|.$$



Yatracos, Y. G. (1985). Rates of convergence of minimum distance estimators and Kolmogorov's entropy. *Annals of Statistics*.

# Some examples

Yatracos' skeleton estimate  $\hat{\theta}_n^Y$  :

$$\mathbb{E} \left[ d_{TV}(P_{\hat{\theta}_n^Y}, P_0) \right] \leq 3d_{TV}(P_0, P_{\theta_0}) + C \cdot \sqrt{\frac{\dim(\Theta)}{n}}$$

where

$$d_{TV}(P, Q) = \sup_E |P(E) - Q(E)|.$$



Yatracos, Y. G. (1985). Rates of convergence of minimum distance estimators and Kolmogorov's entropy. *Annals of Statistics*.

More recent work with the Hellinger distance :



Baraud, Y., Birgé, L., & Sart, M. (2017). A new method for estimation and model selection :  $\rho$ -estimation. *Inventiones mathematicae*.

But...

# But...

Problem with the aforementioned estimators : they cannot be computed in practice.

# But...

Problem with the aforementioned estimators : they cannot be computed in practice.

Additional requirement : an estimator must be computable!!!

# Overview of the talk

## 1 Estimation via MMD optimization

- Definition of the estimator
- Basic properties
- References and further works

## 2 Robustness to outliers

- Application to Huber contamination model
- Example : estimation of the mean of a Gaussian
- Numerical experiments

## 3 Robustness to dependence

- Extension to non-independent observations
- A (new?) dependence coefficient
- Example : auto-regressive observations



## 1 Estimation via MMD optimization

- Definition of the estimator
- Basic properties
- References and further works

## 2 Robustness to outliers

- Application to Huber contamination model
- Example : estimation of the mean of a Gaussian
- Numerical experiments

## 3 Robustness to dependence

- Extension to non-independent observations
- A (new ?) dependence coefficient
- Example : auto-regressive observations

## Reminder : kernels

Let  $\mathcal{H}$  be a Hilbert space and any continuous function  $\Phi : \mathcal{X} \rightarrow \mathcal{H}$ . The function

$$K(x, y) = \langle \Phi(x), \Phi(y) \rangle_{\mathcal{H}}$$

is called a **kernel**.

## Reminder : kernels

Let  $\mathcal{H}$  be a Hilbert space and any continuous function  $\Phi : \mathcal{X} \rightarrow \mathcal{H}$ . The function

$$K(x, y) = \langle \Phi(x), \Phi(y) \rangle_{\mathcal{H}}$$

is called a **kernel**. Conversely :

### Mercer's theorem

Let  $K(x, y)$  be a continuous function such that for any  $(x_1, \dots, x_n) \in \mathcal{X}^n$  and  $(c_1, \dots, c_n) \neq (0, \dots, 0) \in \mathbb{R}^n$ ,

$$\sum_{i=1}^n \sum_{j=1}^n c_i c_j K(x_i, x_j) > 0,$$

then there is  $\mathcal{H}$  and  $\Phi$  such that  $K(x, y) = \langle \Phi(x), \Phi(y) \rangle_{\mathcal{H}}$ .

## Reminder : MMD

Assume that the kernel is bounded :  $0 \leq K(x, y) \leq 1$ .

## Reminder : MMD

Assume that the kernel is bounded :  $0 \leq K(x, y) \leq 1$ .

Consider, for any probability distribution  $P$  on  $\mathcal{X}$ ,

$$\mu_K(P) = \mathbb{E}_{X \sim P} [\Phi(x)] .$$

## Reminder : MMD

Assume that the kernel is bounded :  $0 \leq K(x, y) \leq 1$ .

Consider, for any probability distribution  $P$  on  $\mathcal{X}$ ,

$$\mu_K(P) = \mathbb{E}_{X \sim P} [\Phi(X)].$$

The kernel  $K$  is said to be **characteristic** if

$$\mu_K(P) = \mu_K(Q) \Rightarrow P = Q.$$

## Reminder : MMD

Assume that the kernel is bounded :  $0 \leq K(x, y) \leq 1$ .

Consider, for any probability distribution  $P$  on  $\mathcal{X}$ ,

$$\mu_K(P) = \mathbb{E}_{X \sim P} [\Phi(x)].$$

The kernel  $K$  is said to be **characteristic** if

$$\mu_K(P) = \mu_K(Q) \Rightarrow P = Q.$$

### Theorem

$K(x, y) = \exp(-\frac{\|x-y\|^2}{\gamma^2})$  and  $\exp(-\frac{\|x-y\|}{\gamma})$  are char. kernels.

## Reminder : MMD

Assume that the kernel is bounded :  $0 \leq K(x, y) \leq 1$ .

Consider, for any probability distribution  $P$  on  $\mathcal{X}$ ,

$$\mu_K(P) = \mathbb{E}_{X \sim P} [\Phi(x)] .$$

The kernel  $K$  is said to be **characteristic** if

$$\mu_K(P) = \mu_K(Q) \Rightarrow P = Q .$$

### Theorem

$K(x, y) = \exp(-\frac{\|x-y\|^2}{\gamma^2})$  and  $\exp(-\frac{\|x-y\|}{\gamma})$  are char. kernels.

### Definition : the MMD distance

$$\mathbb{D}_K(P, Q) = \|\mu_K(P) - \mu_K(Q)\|_{\mathcal{H}} .$$



# MMD-based estimator

Reminder of the context :

- 1  $X_1, \dots, X_n$  be i.i.d in  $\mathcal{X}$  from a probability distribution  $P_0$ ,
- 2 model  $(P_\theta, \theta \in \Theta)$ .

# MMD-based estimator

Reminder of the context :

- 1  $X_1, \dots, X_n$  be i.i.d in  $\mathcal{X}$  from a probability distribution  $P_0$ ,
- 2 model  $(P_\theta, \theta \in \Theta)$ .

## Definition - MMD based estimator

$$\hat{\theta}_n^{MMD} = \arg \min_{\theta \in \Theta} \mathbb{D}_K \left( P_\theta, \hat{P}_n \right) \text{ where } \hat{P}_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}.$$

# A preliminary lemma

## Lemma

For any  $P_0$ , when  $X_1, \dots, X_n$  are i.i.d from  $P_0$ ,

$$\mathbb{E} \left[ \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \right] \leq \frac{1}{\sqrt{n}}.$$

# A preliminary lemma

## Lemma

For any  $P_0$ , when  $X_1, \dots, X_n$  are i.i.d from  $P_0$ ,

$$\mathbb{E} \left[ \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \right] \leq \frac{1}{\sqrt{n}}.$$

$$\begin{aligned} \left\{ \mathbb{E} \left[ \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \right] \right\}^2 &\leq \mathbb{E} \left[ \mathbb{D}_K^2 \left( \hat{P}_n, P^0 \right) \right] \\ &= \mathbb{E} \left[ \left\| (1/n) \sum (\mu(\delta_{X_i}) - \mu(P_0)) \right\|_{\mathcal{H}}^2 \right] \\ &= (1/n) \mathbb{E} \left[ \left\| \mu(\delta_{X_1}) - \mu(P_0) \right\|_{\mathcal{H}}^2 \right] \\ &\leq 1/n. \end{aligned}$$

# A bound in expectation

$$\begin{aligned}\forall \theta, \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P^0 \right) &\leq \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, \hat{P}_n \right) + \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \\ &\leq \mathbb{D}_K \left( P_\theta, \hat{P}_n \right) + \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \\ &\leq \mathbb{D}_K \left( P_\theta, P^0 \right) + 2\mathbb{D}_K \left( \hat{P}_n, P^0 \right)\end{aligned}$$

# A bound in expectation

$$\begin{aligned}\forall \theta, \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P^0 \right) &\leq \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, \hat{P}_n \right) + \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \\ &\leq \mathbb{D}_K \left( P_\theta, \hat{P}_n \right) + \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \\ &\leq \mathbb{D}_K \left( P_\theta, P^0 \right) + 2\mathbb{D}_K \left( \hat{P}_n, P^0 \right)\end{aligned}$$

## Theorem

For any  $P_0$ , when  $X_1, \dots, X_n$  are i.i.d from  $P_0$ ,

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2}{\sqrt{n}}.$$

# A bound in probability

We can replace the control on the expectation of  $\mathbb{D}_K \left( \hat{P}_n, P^0 \right)$  by a bound that holds with large probability, thanks to McDiarmid's inequality.

# A bound in probability

We can replace the control on the expectation of  $\mathbb{D}_K(\hat{P}_n, P^0)$  by a bound that holds with large probability, thanks to McDiarmid's inequality.

## Theorem

For any  $P_0$ , when  $X_1, \dots, X_n$  are i.i.d from  $P_0$ , with probability at least  $1 - \delta$ ,

$$\mathbb{D}_K(P_{\hat{\theta}_n}, P^0) \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_{\theta}, P^0) + \frac{2 + 2\sqrt{2 \log\left(\frac{1}{\delta}\right)}}{\sqrt{n}}.$$



# How to compute $\hat{\theta}_n^{MMD}$ ?

We actually have

$$\begin{aligned} \mathbb{D}_K^2(P_\theta, \hat{P}_n) &= \mathbb{E}_{X, X' \sim P_\theta} [K(X, X')] - \frac{2}{n} \sum_{i=1}^n \mathbb{E}_{X \sim P_\theta} [K(X_i, X)] \\ &\quad + \frac{1}{n^2} \sum_{1 \leq i, j \leq n} K(X_i, X_j) \end{aligned}$$

# How to compute $\hat{\theta}_n^{MMD}$ ?

We actually have

$$\mathbb{D}_K^2(P_\theta, \hat{P}_n) = \mathbb{E}_{X, X' \sim P_\theta} [K(X, X')] - \frac{2}{n} \sum_{i=1}^n \mathbb{E}_{X \sim P_\theta} [K(X_i, X)] + \frac{1}{n^2} \sum_{1 \leq i, j \leq n} K(X_i, X_j)$$

and so

$$\begin{aligned} & \nabla_\theta \mathbb{D}_K^2(P_\theta, \hat{P}_n) \\ &= 2 \mathbb{E}_{X, X' \sim P_\theta} \left\{ \left[ K(X, X') - \frac{1}{n} \sum_{i=1}^n K(X_i, X) \right] \nabla_\theta [\log p_\theta(X)] \right\} \end{aligned}$$

that can be approximated by sampling from  $P_\theta$ .

# Short bibliography



Dziugaite, G. K., Roy, D. M., & Ghahramani, Z. (2015). Training generative neural networks via maximum mean discrepancy optimization. *UAI 2015*.

define the estimator and used it to train GANs.

# Short bibliography



Dziugaite, G. K., Roy, D. M., & Ghahramani, Z. (2015). Training generative neural networks via maximum mean discrepancy optimization. *UAI 2015*.

define the estimator and used it to train GANs.



# Short bibliography



Dziugaite, G. K., Roy, D. M., & Ghahramani, Z. (2015). Training generative neural networks via maximum mean discrepancy optimization. *UAI 2015*.

define the estimator and used it to train GANs.



Briol, F. X., Barp, A., Duncan, A. B., & Girolami, M. (2019). Statistical Inference for Generative Models with Maximum Mean Discrepancy. *Preprint arXiv :1906.05944*.

# Short bibliography



Dziugaite, G. K., Roy, D. M., & Ghahramani, Z. (2015). Training generative neural networks via maximum mean discrepancy optimization. *UAI 2015*.

define the estimator and used it to train GANs.



Briol, F. X., Barp, A., Duncan, A. B., & Girolami, M. (2019). Statistical Inference for Generative Models with Maximum Mean Discrepancy. *Preprint arXiv :1906.05944*.

provided the first theoretical study : asymptotic distribution.



Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.



Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.

the results I will present in the following Bayes.





Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.

the results I will present in the following Bayes.



Chérie-Abdellatif, B.-E. and Alquier, P. (2020). MMD-Bayes : Robust Bayesian Estimation via Maximum Mean Discrepancy. *Proceedings of AABI*.



Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.

the results I will present in the following Bayes.



Chérie-Abdellatif, B.-E. and Alquier, P. (2020). MMD-Bayes : Robust Bayesian Estimation via Maximum Mean Discrepancy. *Proceedings of AABI*.

studies a Bayesian version of this estimator.



Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.

the results I will present in the following Bayes.



Chérie-Abdellatif, B.-E. and Alquier, P. (2020). MMD-Bayes : Robust Bayesian Estimation via Maximum Mean Discrepancy. *Proceedings of AABI*.

studies a Bayesian version of this estimator.



Alquier, P. and Gerber, M. (2020). Universal Robust Regression via Maximum Mean Discrepancy. *Preprint arxiv :2006.00840*.



Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.

the results I will present in the following Bayes.



Chérie-Abdellatif, B.-E. and Alquier, P. (2020). MMD-Bayes : Robust Bayesian Estimation via Maximum Mean Discrepancy. *Proceedings of AABI*.

studies a Bayesian version of this estimator.



Alquier, P. and Gerber, M. (2020). Universal Robust Regression via Maximum Mean Discrepancy. *Preprint arxiv :2006.00840*.

application to (linear, logistic, Poisson) regression.



Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.

the results I will present in the following Bayes.



Chérie-Abdellatif, B.-E. and Alquier, P. (2020). MMD-Bayes : Robust Bayesian Estimation via Maximum Mean Discrepancy. *Proceedings of AABI*.

studies a Bayesian version of this estimator.



Alquier, P. and Gerber, M. (2020). Universal Robust Regression via Maximum Mean Discrepancy. *Preprint arxiv :2006.00840*.

application to (linear, logistic, Poisson) regression.



Alquier, P., Chérif-Abdellatif, B.-E., Derumigny, A. and Fermanian, J.-D. (2020). Estimation of copulas via Maximum Mean Discrepancy. *Coming soon !*



Chérie-Abdellatif, B.-E. and Alquier, P. (2019). Finite Sample Properties of Parametric MMD Estimation : Robustness to Misspecification and Dependence. *Preprint arxiv :1912.05737*.

the results I will present in the following Bayes.



Chérie-Abdellatif, B.-E. and Alquier, P. (2020). MMD-Bayes : Robust Bayesian Estimation via Maximum Mean Discrepancy. *Proceedings of AABI*.

studies a Bayesian version of this estimator.



Alquier, P. and Gerber, M. (2020). Universal Robust Regression via Maximum Mean Discrepancy. *Preprint arxiv :2006.00840*.

application to (linear, logistic, Poisson) regression.



Alquier, P., Chérif-Abdellatif, B.-E., Derumigny, A. and Fermanian, J.-D. (2020). Estimation of copulas via Maximum Mean Discrepancy. *Coming soon !*

application to the estimation of copulas.

- 1 Estimation via MMD optimization
  - Definition of the estimator
  - Basic properties
  - References and further works
- 2 Robustness to outliers
  - Application to Huber contamination model
  - Example : estimation of the mean of a Gaussian
  - Numerical experiments
- 3 Robustness to dependence
  - Extension to non-independent observations
  - A (new ?) dependence coefficient
  - Example : auto-regressive observations

# Huber contamination model

## Reminder

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2}{\sqrt{n}}.$$



# Huber contamination model

## Reminder

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2}{\sqrt{n}}.$$

Huber contamination model :  $P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q$ .

# Huber contamination model

## Reminder

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2}{\sqrt{n}}.$$

Huber contamination model :  $P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q$ .

$$\begin{aligned} \mathbb{D}_K(P_{\theta_0}, P_0) &= \|P_{\theta_0} - [(1 - \varepsilon)P_{\theta_0} + \varepsilon Q]\|_{\mathcal{H}} \\ &\leq \varepsilon \|P_{\theta_0}\|_{\mathcal{H}} + \varepsilon \|Q\|_{\mathcal{H}} \\ &= 2\varepsilon. \end{aligned}$$

# Huber contamination model

## Reminder

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2}{\sqrt{n}}.$$

Huber contamination model :  $P_0 = (1 - \varepsilon)P_{\theta_0} + \varepsilon Q$ .

$$\mathbb{D}_K(P_{\theta_0}, P_0) \leq 2\varepsilon.$$

## Corollary

When  $X_1, \dots, X_n$  are i.i.d from  $(1 - \varepsilon)P_{\theta_0} + \varepsilon Q$ ,

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_{\theta_0} \right) \right] \leq 4\varepsilon + \frac{2}{\sqrt{n}}.$$

## Example : Gaussian mean estimation

Example : the model is given by  $p_\theta = \mathcal{N}(\theta, \sigma^2 I)$  for  $\theta \in \mathbb{R}^d$ .

## Example : Gaussian mean estimation

**Example :** the model is given by  $p_\theta = \mathcal{N}(\theta, \sigma^2 I)$  for  $\theta \in \mathbb{R}^d$ .

Using a Gaussian kernel  $K(x, y) = \exp(-\|x - y\|^2/\gamma^2)$ , from the previous theorem and from the equality

$$\mathbb{D}_K^2(P_\theta, P_{\theta'}) = 2 \left( \frac{\gamma^2}{4\sigma^2 + \gamma^2} \right)^{\frac{d}{2}} \left[ 1 - \exp \left( -\frac{\|\theta - \theta'\|^2}{4\sigma^2 + \gamma^2} \right) \right]$$

we obtain

$$\begin{aligned} & \mathbb{E} \left[ \|\hat{\theta}_n^{MMD} - \theta_0\|^2 \right] \\ & \leq -(4\sigma^2 + \gamma^2) \log \left[ 1 - 4 \left( \frac{1}{n} + \varepsilon^2 \right) \left( \frac{4\sigma^2 + \gamma^2}{\gamma^2} \right)^{\frac{d}{2}} \right]. \end{aligned}$$

## Example : Gaussian mean estimation

**Example :** the model is given by  $p_\theta = \mathcal{N}(\theta, \sigma^2 I)$  for  $\theta \in \mathbb{R}^d$ .

Using a Gaussian kernel  $K(x, y) = \exp(-\|x - y\|^2/\gamma^2)$ , from the previous theorem and from the equality

$$\mathbb{D}_K^2(P_\theta, P_{\theta'}) = 2 \left( \frac{\gamma^2}{4\sigma^2 + \gamma^2} \right)^{\frac{d}{2}} \left[ 1 - \exp \left( -\frac{\|\theta - \theta'\|^2}{4\sigma^2 + \gamma^2} \right) \right]$$

we obtain

$$\begin{aligned} \mathbb{E} \left[ \|\hat{\theta}_n^{MMD} - \theta_0\|^2 \right] & \text{ take } \gamma = 2d\sigma^2 \\ & \leq -(4\sigma^2 + \gamma^2) \log \left[ 1 - 4 \left( \frac{1}{n} + \varepsilon^2 \right) \left( \frac{4\sigma^2 + \gamma^2}{\gamma^2} \right)^{\frac{d}{2}} \right]. \end{aligned}$$

## Example : Gaussian mean estimation

**Example :** the model is given by  $p_\theta = \mathcal{N}(\theta, \sigma^2 I)$  for  $\theta \in \mathbb{R}^d$ .

Using a Gaussian kernel  $K(x, y) = \exp(-\|x - y\|^2/\gamma^2)$ , from the previous theorem and from the equality

$$\mathbb{D}_K^2(P_\theta, P_{\theta'}) = 2 \left( \frac{\gamma^2}{4\sigma^2 + \gamma^2} \right)^{\frac{d}{2}} \left[ 1 - \exp \left( -\frac{\|\theta - \theta'\|^2}{4\sigma^2 + \gamma^2} \right) \right]$$

we obtain

$$\mathbb{E} \left[ \|\hat{\theta}_n^{MMD} - \theta_0\|^2 \right] \lesssim d\sigma^2 \left( \frac{1}{n} + \varepsilon^2 \right).$$

## Example : Gaussian mean estimation, simulations

Model :  $\mathcal{N}(\theta, 1)$ , and  $X_1, \dots, X_n$  i.i.d  $\mathcal{N}(\theta_0, 1)$ ,  $n = 100$  and we repeat the experiment 200 times.

	$\hat{\theta}_n^{MLE}$	$\hat{\theta}_n^{MMD}$
mean absolute error	0.0722	0.0838



## Example : Gaussian mean estimation, simulations

Model :  $\mathcal{N}(\theta, 1)$ , and  $X_1, \dots, X_n$  i.i.d  $\mathcal{N}(\theta_0, 1)$ ,  $n = 100$  and we repeat the experiment 200 times.

	$\hat{\theta}_n^{MLE}$	$\hat{\theta}_n^{MMD}$
mean absolute error	0.0722	0.0838

Now,  $\varepsilon = 2\%$  of the observations drawn from a Cauchy.

mean absolute error	0.2349	0.0953
---------------------	--------	--------

# Example : Gaussian mean estimation, simulations

Model :  $\mathcal{N}(\theta, 1)$ , and  $X_1, \dots, X_n$  i.i.d  $\mathcal{N}(\theta_0, 1)$ ,  $n = 100$  and we repeat the experiment 200 times.

	$\hat{\theta}_n^{MLE}$	$\hat{\theta}_n^{MMD}$
mean absolute error	0.0722	0.0838

Now,  $\varepsilon = 2\%$  of the observations drawn from a Cauchy.

mean absolute error	0.2349	0.0953
---------------------	--------	--------

Now,  $\varepsilon = 1\%$  are replaced by 1,000.

mean absolute error	10.018	0.0903
---------------------	--------	--------

## 1 Estimation via MMD optimization

- Definition of the estimator
- Basic properties
- References and further works

## 2 Robustness to outliers

- Application to Huber contamination model
- Example : estimation of the mean of a Gaussian
- Numerical experiments

## 3 Robustness to dependence

- Extension to non-independent observations
- A (new ?) dependence coefficient
- Example : auto-regressive observations

# And now, non-independent observations

## Lemma

When  $X_1, \dots, X_n$  are identically distributed from  $P_0$ ,

$$\mathbb{E} \left[ \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \right] \leq ?$$

# And now, non-independent observations

## Lemma

When  $X_1, \dots, X_n$  are identically distributed from  $P_0$ ,

$$\mathbb{E} \left[ \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \right] \leq ?$$

$$\begin{aligned} & \mathbb{E} \left[ \mathbb{D}_K^2 \left( \hat{P}_n, P^0 \right) \right] \\ &= \mathbb{E} \left[ \left\| (1/n) \sum (\mu(\delta_{X_i}) - \mu(P_0)) \right\|_{\mathcal{H}}^2 \right] \\ &= \frac{1}{n} + \frac{2}{n^2} \sum_{1 \leq i < j \leq n} \mathbb{E} \langle \mu(\delta_{X_i}) - \mu(P_0), \mu(\delta_{X_j}) - \mu(P_0) \rangle_{\mathcal{H}} \end{aligned}$$

# Measure of dependence via covariance in $\mathcal{H}$

## Definition

When  $(X_1, \dots, X_n, \dots)$  is a stationary process with marginal distribution  $P_0$ , we put :

$$\varrho_h = \left| \mathbb{E} \left\langle \mu(\delta_{X_{t+h}}) - \mu(P_0), \mu(\delta_{X_t}) - \mu(P_0) \right\rangle_{\mathcal{H}} \right|.$$

# Measure of dependence via covariance in $\mathcal{H}$

## Definition

When  $(X_1, \dots, X_n, \dots)$  is a stationary process with marginal distribution  $P_0$ , we put :

$$\varrho_h = \left| \mathbb{E} \left\langle \mu(\delta_{X_{t+h}}) - \mu(P_0), \mu(\delta_{X_t}) - \mu(P_0) \right\rangle_{\mathcal{H}} \right|.$$

## Lemma - dependent case

When  $X_1, \dots, X_n$  are identically distributed from  $P_0$ ,

$$\mathbb{E} \left[ \mathbb{D}_K \left( \hat{P}_n, P^0 \right) \right] \leq \frac{1}{n} \left[ 1 + \sum_{h=1}^n \varrho_h \right]$$

# Mesure of dependence via covariance in $\mathcal{H}$

## Theorem - dependent case

When  $(X_1, \dots, X_n, \dots)$  is a stationary process with marginal distribution  $P_0$

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2 + 2 \sum_{h=1}^n \varrho_h}{\sqrt{n}}.$$



# Mesure of dependence via covariance in $\mathcal{H}$

## Theorem - dependent case

When  $(X_1, \dots, X_n, \dots)$  is a stationary process with marginal distribution  $P_0$

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2 + 2 \sum_{h=1}^n \varrho_h}{\sqrt{n}}.$$

① assume that  $\sum_{h=1}^{\infty} \varrho_h = \Sigma < +\infty$  then

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2 + 2\Sigma}{\sqrt{n}}.$$

# Measure of dependence via covariance in $\mathcal{H}$

## Theorem - dependent case

When  $(X_1, \dots, X_n, \dots)$  is a stationary process with marginal distribution  $P_0$

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2 + 2 \sum_{h=1}^n \varrho_h}{\sqrt{n}}.$$

- 1 assume that  $\sum_{h=1}^{\infty} \varrho_h = \Sigma < +\infty$  then

$$\mathbb{E} \left[ \mathbb{D}_K \left( P_{\hat{\theta}_n^{MMD}}, P_0 \right) \right] \leq \inf_{\theta \in \Theta} \mathbb{D}_K(P_\theta, P_0) + \frac{2 + 2\Sigma}{\sqrt{n}}.$$

- 2 we also have a bound in probability, based on Rio's version of Hoeffding's inequality ; it requires more assumptions.

# An example : auto-regressive processes

## Proposition

Assume that  $X_t$  takes values in  $\mathbb{R}^d$  and that  $K(x, y) = F(\|x - y\|)$  where  $F$  is an  $L$ -Lipschitz function. Assume that

$$X_{t+1} = AX_t + \varepsilon_{t+1}$$

where the  $(\varepsilon_t)$  are i.i.d with  $\mathbb{E}\|\varepsilon_0\| < \infty$ , and  $A$  is a matrix with  $\|A\| = \sup_{\|x\|=1} \|Ax\| < 1$ .

# An example : auto-regressive processes

## Proposition

Assume that  $X_t$  takes values in  $\mathbb{R}^d$  and that  $K(x, y) = F(\|x - y\|)$  where  $F$  is an  $L$ -Lipschitz function. Assume that

$$X_{t+1} = AX_t + \varepsilon_{t+1}$$

where the  $(\varepsilon_t)$  are i.i.d with  $\mathbb{E}\|\varepsilon_0\| < \infty$ , and  $A$  is a matrix with  $\|A\| = \sup_{\|x\|=1} \|Ax\| < 1$ .

Then

$$\varrho_t \leq \|A\|^t \frac{2L\mathbb{E}\|\varepsilon_0\|}{1 - \|A\|} \text{ and } \Sigma = \sum_{t=1}^{\infty} \varrho_t = \frac{2\|A\|L\mathbb{E}\|\varepsilon_0\|}{(1 - \|A\|)^2}.$$

# A non-mixing process with $\Sigma < +\infty$

Example : consider  $X_0 \sim \mathcal{U}([0, 1])$ ,  $\eta_t$  i.i.d  $\mathcal{Be}(1/2)$  and

$$X_{t+1} = \frac{X_t + \eta_{t+1}}{2}.$$

# A non-mixing process with $\Sigma < +\infty$

**Example :** consider  $X_0 \sim \mathcal{U}([0, 1])$ ,  $\eta_t$  i.i.d  $\mathcal{Be}(1/2)$  and

$$X_{t+1} = \frac{X_t + \eta_{t+1}}{2}.$$

It satisfies the assumptions of the previous proposition, we have  $\varrho_t \leq L/2^t$  and  $\Sigma = 2L$ .

## A non-mixing process with $\Sigma < +\infty$

**Example :** consider  $X_0 \sim \mathcal{U}([0, 1])$ ,  $\eta_t$  i.i.d  $\mathcal{Be}(1/2)$  and

$$X_{t+1} = \frac{X_t + \eta_{t+1}}{2}.$$

It satisfies the assumptions of the previous proposition, we have  $\varrho_t \leq L/2^t$  and  $\Sigma = 2L$ .

Note however that this process is known to be non-mixing.

## A non-mixing process with $\Sigma < +\infty$

**Example :** consider  $X_0 \sim \mathcal{U}([0, 1])$ ,  $\eta_t$  i.i.d  $\mathcal{Be}(1/2)$  and

$$X_{t+1} = \frac{X_t + \eta_{t+1}}{2}.$$

It satisfies the assumptions of the previous proposition, we have  $\varrho_t \leq L/2^t$  and  $\Sigma = 2L$ .

Note however that this process is known to be non-mixing.

More generally, we prove the following result :

### Proposition

Under some (non-restrictive) assumption on the kernel  $K$ ,

$$\varrho_t \leq c_K \cdot \beta_t \text{ (the } \beta\text{-mixing coef.)}$$



Thank you !