

1 Limited predictability of amino acid substitutions in seasonal influenza viruses

2 Pierre Barrat-Charlaix,^{1,2} John Huddleston,³ Trevor Bedford,⁴ and Richard A. Neher^{1,2,*}

3 ¹*Biozentrum, Universität Basel,*

4 *Switzerland*

5 ²*Swiss Institute of Bioinformatics, Basel,*

6 *Switzerland*

7 ³*Molecular Cell Biology,*

8 *University of Washington, Seattle, WA,*

9 *USA*

10 ⁴*Vaccine and Infectious Disease Division,*

11 *Fred Hutchinson Cancer Research Center, Seattle, WA,*

12 *USA*

13 (Dated: November 19, 2020)

14 Seasonal influenza viruses repeatedly infect humans in part because they rapidly change
15 their antigenic properties and evade host immune responses, necessitating frequent
16 updates of the vaccine composition. Accurate predictions of strains circulating in the
17 future could therefore improve the vaccine match. Here, we studied the predictability of
18 frequency dynamics and fixation of amino acid substitutions. Current frequency was the
19 strongest predictor of eventual fixation, as expected in neutral evolution. Other properties,
20 such as occurrence in previously characterized epitopes or high *Local Branching Index*
21 (LBI) had little predictive power. Parallel evolution was found to be moderately predictive
22 of fixation. While the LBI had little power to predict frequency dynamics, it was still
23 successful at picking strains representative of future populations. The latter is due to a
24 tendency of the LBI to be high for consensus-like sequences that are closer to the future
25 than the average sequence. Simulations of models of adapting populations, in contrast,
26 show clear signals of predictability. This indicates that the evolution of influenza HA and
27 NA, while driven by strong selection pressure to change, is poorly described by common
28 models of directional selection such as travelling fitness waves.

29 INTRODUCTION

30 Seasonal influenza A viruses (IAV) infect about 10% of 53
31 the global population every year, resulting in hundreds 54
32 of thousands of deaths (Organization, 2018; Petrova and 55
33 Russell, 2017). Vaccination is the primary measure to 56
34 reduce influenza morbidity. However, the surface proteins 57
35 hemagglutinin (HA) and neuraminidase (NA) continu- 58
36 ously accumulate mutations at a high rate, leading to fre- 59
37 quent antigenic changes (Petrova and Russell, 2017; Shih 60
38 et al., 2007; Bhatt et al., 2011; Koel et al., 2013). While 61
39 a vaccine targeting a particular strain may be efficient for 62
40 some time, antigenic drift will sooner or later render it ob- 63
41 solete. The World Health Organization (WHO) regularly 64
42 updates influenza vaccine recommendations to best match 65
43 the circulating strains. Since developing, manufacturing, 66
44 and distributing the vaccine takes many months, forecast- 67
45 ing the evolution of influenza is of essential interest to 68
46 public health (Morris et al., 2018; Klingen et al., 2018). 69

47 The number of available high quality HA and 70
48 NA sequences has increased rapidly over the last 20 71
49 years (Bogner et al., 2006; Shu and McCauley, 2017) 72
50 and virus evolution and dynamics can be now be tracked 73

51 at high temporal and spatial resolution (Rambaut et al.,
52 2008). This wealth of data has given rise to an active
53 field of predicting influenza virus evolution (Morris et al.,
54 2018; Klingen et al., 2018). These models predict the
55 future population of influenza viruses by estimating strain
56 fitness or proxies of fitness. Luksza and Lässig (2014),
57 for example, train a fitness model to capture antigenic
58 drift and protein stability on patterns of epitope and non-
59 epitope mutations. Other approaches by Steinbrück et al.
60 (2014); Neher et al. (2016) predict fitness by using hemag-
61 glutination inhibition (HI) data to determine possible
62 antigenic drift of clades in the genealogy of the HA pro-
63 tein. Finally, Neher et al. (2014) use branching patterns
64 of HA phylogenies as a proxy for fitness. These branching
65 patterns are summarized by the Local Branching Index
66 (LBI), which was shown to be a proxy of relative fitness
67 in mathematical models of rapidly adapting populations
68 (Neher et al., 2014).

69 The underlying assumption of all these methods is that
70 (i) differences in growth rate between strains can be es-
71 timated from sequence or antigenic data and (ii) that
72 these growth rate differences persist for long enough to be
73 predictive of future success. Specific positions in surface
74 proteins are of particular interest in this context. The
75 surface proteins are under a strong positive selection and
76 change their amino acid sequence much more rapidly than
77 other IAV proteins or than expected under neutral evo-
78 lution (Bhatt et al., 2011; Strelkowa and Lässig, 2012).

* Correspondence to: Richard Neher, Biozentrum, Klingelbergstrasse 70, 4056, Basel, Switzerland.
richard.neher@unibas.ch

Epitope positions, i.e., positions targeted by human antibodies, are expected to change particularly often since viruses with altered epitopes can evade existing immune responses (Shih et al., 2007; Koel et al., 2013; Wolf et al., 2006). It therefore seems plausible that mutations at these positions have a tendency to increase fitness and a higher probability of fixation (Strelkowa and Lässig, 2012). But one has to be careful to account for the fact that these positions are often ascertained post-hoc (Shih et al., 2007) and human immune responses are diverse with substantial inter-individual variation (Lee et al., 2019).

In this work, we use HA and NA sequences of A/H3N2 and A/H1N1pdm influenza from year 2000 to 2019 to perform a retrospective analysis of frequency trajectories of amino acid mutations. We quantify how rapidly mutations at different frequencies are lost or fixed and how rapidly they spread through the population. We further investigate whether any properties or statistics are predictive of whether a particular mutation fixes or not. To our surprise, we find that the predictability of these trajectories is very limited: The probability that a mutation fixes differs little from its current frequency, as would be expected if fixation happened purely by chance. This observation holds for many different categories of mutations, including mutations at epitope positions. This weak predictability is not attributable solely to clonal interference and genetic linkage, as simulation of models including even strong interference retain clear signatures of predictability. Consistent with these observations, we show that a simple predictor uninformed by fitness, the consensus sequence, performs as well as the Local Branching Index (LBI), the growth measure based on the genealogy used in (Neher et al., 2014). This suggests that although LBI has predictive power, the reason for its success may not be related to it approximating fitness of strains.

RESULTS

The main underlying question asked in this work is the following: given a mutation X in the genome of influenza that we observe at a frequency f in the population at a given date, what can we say about the future of X ? The trajectory of a mutation will depend on its own effect on fitness, the contribution of the genetic background on the same segment, and the effect of the remaining seven segments. Here, we investigate properties of broad categories of mutations effectively averaging over different genetic backgrounds to isolate the effects intrinsic to the mutation.

First, we ask whether we can quantitatively predict the frequency of X at future times $f(t)$. In other words, having observed a mutation at frequencies (f_1, f_2, \dots, f_n) at dates (t_1, t_2, \dots, t_n) , what can we say about its frequency at future dates $(t_{n+1}, t_{n+2}, \dots)$? A simpler, more

qualitative question, is to ask whether X will fix in the population, will disappear, or whether the site will stay polymorphic.

We use amino-acid sequences of the HA and NA genes of A/H3N2 (since the year 2000) and A/H1N1pdm (since the year 2009) influenza available in GISAID (Shu and McCauley, 2017) (see supplementary materials for an acknowledgment of all data contributors). This amounts to 44 976 HA and 36 300 NA sequences for A/H3N2 and 45 350 HA and 40 412 NA sequences for A/H1N1, with a minimum of 100 per year. These sequences are binned in non-overlapping intervals of one month. Each single-month time bin and the sequences that it contains represent a (noisy) snapshot of the influenza population at a given date. The number of sequences per time bin varies strongly both with year and according to the season, with earlier time bins containing around 10 sequences while more recent bins contain several hundreds (see figures S5 and S6 in SM for details).

The central quantities that we derived from this data are *frequency trajectories* of amino acids at each position in the sequences. If an amino acid X_i is found at position i at a frequency between 5% and 95% in the population of a given time bin t , then the population is considered polymorphic at position i and at time t . This polymorphism is characterized by the frequency of X_i , $f_{X_i}(t)$, and also by frequencies of other amino acids at i . The series of values $f_{X_i}(t)$ for contiguous time bins constitutes the frequency trajectory of X_i . A trajectory is terminated if the corresponding frequency is measured above 95% (resp. below 5%) for two time bins in a row, in which case amino acid X_i is considered as *fixed* (resp. *absent*) in the population. Otherwise, the trajectory is considered *active*. Examples of trajectories can be seen in figure S7 of the Supplement.

In the rest of this work, we will focus on frequency trajectories that are starting at a zero (low) frequency, i.e. $f(t=0) = 0$. These represent new amino acid variants which were absent in the population at the time bin when the trajectory started and are currently rising in the population (see Methods). Such distinction in novel and ancestral variants is necessary to meaningfully interrogate predictability. Each rising trajectory of a new mutation implies the existence of another decreasing one at the same position, since frequencies of all amino acids at a given position must sum to one. If novel variants arise by selection, we expect to see a stronger signal of selection after conditioning on these novel variants. In classic models of population genetics, strongly advantageous variants undergo rapid selective *sweeps*, i.e., the rapid rise and fixation. The sweep of a mutation can be due to its own fitness effect, to the genetic background or to the effect of the seven other segments. By considering the ensemble of novel variants that are rising in frequency, we effectively average over backgrounds, obtaining a set of mutations that we expect to be beneficial on average. If such sweeps

are common in the evolution of HA and NA, the restriction to trajectories that start at low frequency should thus enrich for mutations that are positively selected and on their way to fixation.

Predicting future frequencies

Having observed the frequency trajectory $f(t)$ of a mutation until a given date t_0 , how much can we say about the future values of f after t_0 ? We consider the idealized case sketched in panel **A** of figure 1: given the trajectory of a *new* mutation, *i.e.* that started at a frequency of 0, and that we observe at frequency f_0 at time t_0 , what is the probability $P_{\Delta t}(f)$ of observing it at a value f at time $t_0 + \Delta t$?

To answer this question retrospectively, we use all frequency trajectories extracted from HA and NA sequences that satisfy these conditions for a given f_0 . The number of trajectories is limited and the frequency estimates themselves are based on a finite sample and are hence imprecise. Therefore, we consider trajectories in an interval $[f_0 - \delta f, f_0 + \delta f]$ with $\delta f = 0.05$.

For $f_0 = 0.3$, we found 120 such trajectories in the case of A/H3N2 influenza, represented on the panel **B** of figure 1, where time is shifted such that $t_0 = 0$. The same analysis was performed for A/H1N1pdm, with the 89 found trajectories displayed in figure S9. Some trajectories fall in the frequency bin around f_0 while decreasing, even though they crossed that bin at an earlier time. This is due to the fact that some trajectories “skipped” the interval f_0 in question on their initial rise due to sparse sampling. These trajectories are nevertheless rising in the sense that they start at frequency 0 for $t \rightarrow -\infty$. Removing them does not change results significantly.

Since rapid sequence evolution of influenza HA and NA mediates immune evasion, one could expect that a significant fraction of new amino acid mutations on rising trajectories in figure 1 are *adaptive*. We could thus expect that most of these trajectories continue to rise after reaching frequency f_0 , at least for some time. A fraction of those would then sweep through the population and fix.

To quantify the extent to which this preception of sweeping adaptive mutations is true, we estimated the probability distribution $P_{\Delta t}(f|f_0)$ of finding a trajectory at frequency f after a time Δt given that it was observed at f_0 at time 0. The results for different Δt are shown in figure 1C. Initially, *i.e.* at time $t_0 = 0$, this distribution is by construction peaked around f_0 . If a large fraction of the trajectories keep increasing after this time, we should see the “mass” of $P_{\Delta t}(f|f_0)$ move to the right towards higher frequencies as time progresses.

However, future distributions for $\Delta t > 0$ do not seem to follow a pattern compatible with selective sweeps. The thick black line in Figure 1B shows the average frequency

of all trajectories. This average makes a sharp turn at $t = 0$ and is essentially flat for $t > 0$ in the case of A/H3N2, and slightly increasing for A/H1N1pdm (see supplement). Hence, the fact that this average rose for $t < 0$ gives little information for $t > 0$, and is due to the conditions by which these trajectories were selected. This shows that sweep-like trajectories rising steadily from frequency 0 to 1 are not common enough to dominate the average trajectory.

Consistent with the average, the frequency distribution of the selected trajectories broadens in time without a significant shift of the mean as time passes. After 60 days, the distribution is rather symmetrical around the initial $f_0 = 0.3$ value, suggesting that the knowledge that the trajectories were rising is lost after two months. On a timescale of 60 to 120 days, the only possible prediction is that trajectories are likely to be found in a broad interval around the initial frequency f_0 . After one year the distribution becomes almost flat (excluding mutations that have disappeared or fixed), and the initial peak at f_0 is not visible anymore. The only information remaining from the initial frequency is the fraction that fixed or was lost (see below). This behavior is expected in neutral models of evolution (Kimura, 1964) but incompatible with a dynamic dominated by sweeps taking over the population.

While this observation does not rule out that signatures exist that predict future frequency dynamics, past dynamics alone is weakly informative.

Prediction of fixation or loss

Instead of predicting future frequency, let's consider the long-term goal of predicting the probability that a mutation fixes in the population. We first estimate the fraction of frequency trajectories that either fix in the population or are lost, as well as the time it takes for one or the other to happen. Panels **A** and **B** of figure 2 shows the fraction of frequency trajectories in HA and NA that either have fixed, were lost or remained active as a function of the time elapsed since they were first seen above 25% frequency. Most mutations are either lost or become fixed after 2-3 years, with very few trajectories remaining active after 5 years. This time scale of 2-3 years is consistent with the typical coalescence time observed in phylogenetic trees of A/H3N2 influenza (Rambaut et al., 2008; Yan et al., 2019). We also note that the fraction of lost trajectories increases sharply at small times with 40% of mutations observed above 25% frequency being lost within one year for A/H3N2, while it takes longer to fix a mutation in the whole population.

We then examined the probability of mutations to fix in the population as a function of the frequency at which they are seen. For different values of frequency f , we consider all trajectories that started at a null frequency




FIG. 1 **A:** Sketch of the idea behind the short term prediction of frequency trajectories. Given a mutation that we have seen increasing in frequency and that we “catch” at frequency f_0 at time t_0 , what can we say about the distribution of future frequencies $P_{\Delta t}(f|f_0)$? **B:** All frequency trajectories of amino acid mutations in the A/H3N2 HA and NA genes that were absent in the past, are seen around $f_0 = 30\%$ frequency at time $t_0 = 0$, and are based on more than 10 sequences at each time point. Red curves represent mutations that will ultimately fix, blue the ones that will be lost, and black the ones for which we do not know the final status. Dashed horizontal lines (blue and red) represent loss and fixation thresholds. The thick black line is the average of all trajectories, counting those that fix (resp. disappear) as being at frequency 1 (resp. 0). Figure S8 shows equivalent figures for other values of f_0 . **C:** Distribution of future frequencies $P_{\Delta t}(f|f_0)$ for the trajectories shown in panel **B** and for specific values of Δt .




FIG. 2 **A:** Activity of all rising frequency trajectories seen above 25% frequency for A/H3N2 HA and NA. **B:** Same as **A** for A/H1N1. **C:** Probability of fixation of a mutation (amino acid or synonymous) $P_{fix}(f)$ as a function of the frequency f at which it is measured, for A/H3N2 HA and NA. Only new mutations are considered, *i.e.* mutations that were absent in the past. The diagonal dashed line is the expectation from a neutrally evolving population. Colored dashed lines represent synonymous mutations. Colored solid lines represent amino acid mutations. Error bars represent a 95% confidence interval. **D:** Same as **C** for A/H1N1.

and are seen in the interval $[f - 7.5\%, f + 7.5\%]$ at any given time. The probability of a mutation fixing given that it is seen at frequency f , $P_{fix}(f)$, is then estimated by the fraction of those trajectories which terminate at a frequency larger than 95%, *i.e.* our fixation threshold. Panels **C** and **D** of figure 2 show $P_{fix}(f)$ as a function of f for NA and HA. For both proteins, the probability of fixation of a new mutation at frequency f is close to f itself, that is $P_{fix}(f) \simeq f$. This result is exactly what is expected in a population evolving in the *absence* of selection. A mutation or trait appearing at frequency f is shared by $f \cdot N$ individuals, and the probability for one of them to become the ancestor of all the future population is $f \cdot N/N = f$. Thus, the probability of this mutation or trait to fix in the population is equal to its current frequency, a case which we will refer to as the neutral expectation. Panel **C** of figure 2 indicates that

mutations in the surface proteins of A/H3N2 influenza are in good agreement with the neutral expectation, while those in A/H1N1pdm show only small deviations from it. In both cases, the probability of fixation seems to be mainly dictated by the current frequency f at which the mutation is observed.

This dynamics is in apparent contradiction with evidence that influenza surface proteins are under strong selective pressure to evade human immune responses (Bhatt et al., 2011). If strong selection was present, we would expect rising amino acid mutations to fix at a distinctively higher frequency than the one at which they are measured. In an extreme case where most trajectories would be clean sweeps, $P_{fix}(f)$ should be close to 1 for all but very small values of f .

Next, we searched for features of mutations that allow prediction of fixation beyond frequency by dividing

328 frequencies into categories that deviate from the diagonal in panels **C** and **D** of figure 2. We first turn to the Local Branching Index (LBI), a quantity calculated for each node in a phylogenetic tree that indicates how dense the branching of the tree is around that node. LBI has previously been successfully used as a predictor of future population of influenza (Neher et al., 2014), and was shown to be a proxy for fitness of leaves or ancestral nodes in mathematical models of evolution. Here, we define the LBI of a mutation at date t as the average LBI of strains that carry this mutation and that were sampled in the time bin corresponding to t . Panel A of figure 3 shows fixation probability for HA mutations with LBI in the top or bottom half of the distribution. Both groups have identical probability of fixation, suggesting that LBI carries very little information on the probability of fixation of a mutation.

345 Next, we focused on previously reported antigenic sites in the A/H3N2 HA protein, referred to as *epitope* positions. Mutations at these position might mediate immune escape and are therefore likely under strong selection and show sweep-like behavior. We used four lists of relevant epitope positions from different sources comprising from 7 to 129 positions in the sequence of the HA1 protein (Shih et al., 2007; Koel et al., 2013; Luksza and Lässig, 2014; Wolf et al., 2006). Panel Fig. 3B shows fixation probability as a function of frequency for the four lists of epitopes. Only mutations at the 7 epitope sites reported in (Koel et al., 2013) have higher chances of fixation than expected by chance. No clear difference is found for the lists by Luksza and Lässig (2014); Wolf et al. (2006), while positions from Shih et al. (2007) show lower chances of fixation. One should also note that many of these positions were determined post-hoc and might be enriched for positions that experienced rapid substitutions before the publication of the respective studies.

364 Two ways of categorising mutations, however, suggest some power to predict fixation. In panel Fig. 3C, we split trajectories into those occurring at binary positions where only two amino acid variants co-circulate and non-binary positions with more than two variants. Novel variants at non-binary positions, *i.e.* ones for which competition between three amino acids or more has occurred at least once, have a higher chance of fixation. In panel D, we separated mutations that appear more than once or only once in the reconstructed tree (see methods), and found that the former fix more often. Panels C and D show that it is possible to gain some information on the chance of fixation of a particular mutation, as was done in panel B. However, the predictive power remains small, with the “top” curves in panels C&D being very close to the diagonal.

380 We conduct the same analysis on A/H1N1pdm influenza, with results shown in figure S11. Results are qualitatively similar to those obtained for A/H3N2, with LBI giving little information and mutations at non-binary

positions having a higher chance of fixation. Panel D differs between figures 3 and S11, with convergent evolution giving less information on fixation in the latter case. However, this could be due to the shorter time period over which A/H1N1pdm evolved, resulting in a shorter tree and less possibilities of convergent evolution. Indeed, error bars for mutations appearing multiple times in D of figure S11 are relatively large, indicating a lower amount of trajectories.

Since influenza is seasonal in temperate regions, geographic spread and persistence might be predictive of the success of mutations. We quantify geographic spread of a mutation by the entropy of its frequency distribution across regions (see methods) and its persistence by the age of the trajectory by the time it reaches frequency f . Figures S12 and S13 show the fixation probabilities as a function of observed frequency for mutations classified according to these scores. The two scores also allow a quantitatively moderate distinction between mutations: for a given frequency f , mutations found in many regions or those that are older (in the sense that they have taken more time to reach frequency f) tend to fix more often than geographically localized mutations or more recent ones, but the effect is small. These two scores are in fact correlated, with older trajectories representing mutations that are more geographically spread, as can be seen in figure S14 of SM. However, it is important to note that sampling biases and heterogeneity across time and space (see supplementary figures S5 and S6) make answering such specific hypothesis challenging. Frequency of mutations might thus be amplified through different sampling biases, making the connection between geographic spread, seasonality and mutation frequency non-trivial to measure.

Simulations of models of adaptation

The results shown in figures 2 and 3 are difficult to reconcile with the idea that seasonal influenza virus evolution is driven by rapid directed positive selection. One possible explanation for the weakly predictable behaviour of mutations (beyond their current frequency) might be tight genetic linkage inside each segment and strong competition between different adaptive mutations (Strelkowa and Lässig, 2012; Neher and Shraiman, 2011). We design a simple model of population evolution based on the `ffpopsim` simulation software to test this hypothesis (Zanini and Neher, 2012). The model represents a population of binary genomes of length $L = 200$ evolving in a fitness landscape that changes through time.

First, we use an additive fitness function, with sequence $(x_1 \dots x_L)$ having a fitness $\sum_i h_i x_i$. This implies that for a given genome position i , the trait $x_i = 1$ is favored if $h_i > 0$ whereas $x_i = -1$ is favored if $h_i < 0$. All h_i 's




FIG. 3 Fixation probability $P_{fix}(f)$ as a function of frequency, for A/H3N2 influenza. Figure S11 shows the same analysis for A/H1N1. **A:** HA mutations with higher or lower LBI values, based on their position with respect to the median LBI value. **B:** Different lists of epitope positions in the HA protein. The authors and the number of positions is indicated in the legend. **C:** HA and NA mutations for binary positions, *i.e.* positions for which we never see more than two amino acids in the same time bin. **D:** HA and NA mutations that appear once or more than once in the tree for a given time bin.

437 have the same magnitude, and only their signs matter 457
 438 Every Δt generations, we randomly choose a position i 458
 439 and flip the sign of h_i , effectively changing the fitness 459
 440 landscape. Individuals in the population now have the 460
 441 opportunity to make an adaptive mutation at site i giving 461
 442 them a fitness advantage $2|h|$. A “flip” at position i of the 462
 443 fitness landscape will decrease fitness of all individuals 463
 444 that carried the adapted variant at position i and increases 464
 445 the fitness of those that happened to carry a deleterious 465
 446 variant.

447 To increase competition between genomes, we designed 467
 448 a second model that includes epistasis. Once again, the 468
 449 baseline fitness of a genome is an additive function, this 469
 450 time with values of h_i that do not change through time 470
 451 In addition, we added a component that mimics immune 471
 452 selection. Every Δt generation, we now introduce “an- 472
 453 tibodies” that target a specific sub-sequence of length 473
 454 $l = 5$, noted $(x_{i_1}^{ab}, \dots, x_{i_l}^{ab})$. The positions $(i_1 \dots i_l)$ are 474
 455 chosen at random, while the targeted sub-sequence is the 475
 456 dominant state at each position. Genomes that include 476

the *exact* sub-sequence targeted by the antibody suffer a strong fitness penalty. However, a single mutation away from that sub-sequence removes this penalty completely, resulting in a fitness landscape with very strong epistasis. This has the effect of triggering a strong competition between adaptive mutations: for a given antibody, $l = 5$ possible mutations are now adaptive, but combinations of these mutations do not bring any fitness advantage.

Having simulated populations in these two fitness landscapes, we perform the same analysis of frequency trajectories as for the real influenza data. Figure S16 of the SM shows the $P_{fix}(f)$ as a function of f for the two models and for different values of the inverse rate of change Δt of the fitness landscape. For all models, this curve deviates significantly from the diagonal. This is most evident for the case of a simple additive fitness landscape that changes rarely $\Delta t = 1000$: rising mutations almost always fix in the population, with $P_{fix}(f) \simeq 1$ for any f larger than a few percent. This is corroborated by visual inspection of the trajectories, which shows that evolution in this

regime is driven by regular selective sweeps that take a₅₃₀
typical time of ~ 400 generations. In other regimes, with₅₃₁
smaller Δt or with strong epistatic competition, $P_{fix}(f)$ ₅₃₂
is reduced and closer to the diagonal. However, it takes₅₃₃
an extremely fast changing fitness landscape to push P_{fix} ₅₃₄
close to the diagonal: with $\Delta t = 10$, that is about 40₅₃₅
changes to the fitness landscape in the time it would take₅₃₆
a selective sweep to go from 0% to fixation, $P_{fix}(f)$ differs₅₃₇
from f in a way that is comparable to what is observed₅₃₈
in A/H1N1pdm influenza.₅₃₉

These models are not meant to be accurate models of in₅₄₀
fluenza viruses evolution. But figure S16 does show is that₅₄₁
the patterns observed in influenza virus evolution are only₅₄₂
reproduced by models of adapting populations when push₅₄₃
ing clonal competition to extreme values. We conclude₅₄₄
that the pattern in figure 2 may not be a straightforward₅₄₅
manifestation of genetic linkage and clonal interference₅₄₆
but that some more intricate interplay of epidemiology₅₄₇
seasonality, human immunity and chance gives rise to₅₄₈
the weakly predictable yet strongly selected evolutionary₅₄₉
dynamics of IAVs.₅₅₀

498 Why do predictions work?

The statistics of frequency trajectories seem to be in₅₅₅
conflict with the notion that influenza evolution is pre₅₅₆
dictable. Likewise, the LBI, a quantity that correlates₅₅₇
with fitness in mathematical models and is used to predict₅₅₈
future influenza populations (Neher et al., 2014), does not₅₅₉
seem to contain any information on whether a specific₅₆₀
mutation is going to fix or not, see figure 3. To resolve₅₆₁
this conundrum, we first note that the criterion by which₅₆₂
predictive power for influenza was measured in (Neher₅₆₃
et al., 2014) was the distance between the strain with the₅₆₄
highest LBI and the future population, not the ability of₅₆₅
the LBI to predict dynamics. The distance was compared₅₆₆
to the average distance between the present and future₅₆₇
population, as well as the post-hoc optimal representative₅₆₈
and the future.₅₆₉

To quantify the ability of the LBI and other measures₅₇₀
to pick good representatives of the future, we construct₅₇₁
a large tree of HA sequences with 100 sequences in non₅₇₂
overlapping time bins of 4 months from year 2003 to 2019
(a total of 4402 as some 4 month intervals contain less
than 100 sequences). Each time bin is considered as a₅₇₃
snapshot of the A/H3N2 influenza population and we will
refer to sequences in time bin t as the population of the₅₇₄
present. From this present population, we predict future₅₇₅
populations in time bin $t + \Delta t$, using only sequences in₅₇₆
time bin t and before.₅₇₇

To assess the ability of the LBI to pick a close represen₅₇₈
tative of the future, we compute the LBI of each node of₅₇₉
one time bin in the tree using only the leaves that belong₅₈₀
to that time bin. The top panel in figure 4 shows the₅₈₁
hamming distance of the strain with the highest LBI to₅₈₂

future populations at different Δt along with the same
distance for a randomly chosen strain. The figure shows
the distance averaged over all possible values of t for Δt
between 0 and 32 months, giving us an average efficiency
of a predictor over 16 years of influenza evolution.

The strain with the highest LBI is consistently closer
to the future than the average strain by about 1-2 amino
acids, while the overall distance increases linearly due to
the continuous evolution of the population. We hence
reproduce previous results showing that the LBI picks
closer than average representatives (Neher et al., 2014).
To investigate whether this apparent success is due to the
ability of the LBI to predict fitness or not, we explored a
different predictor: the amino acid consensus sequence of
the present population (see Methods for a definition of
the consensus sequence). The choice is motivated by the
fact that it can be shown to be the best possible long term
predictor for a neutrally evolving population in terms of
Hamming distance (see SM section 1). Figure 4 shows
that the consensus sequence is in fact a equally good
or even slightly better representative of the future than
the sequence with highest LBI (note that the consensus
sequence does *not* necessarily exist in the population).

This near equivalence of the consensus and the strain
with highest LBI can be explained as follows: The LBI
tends to be high for nodes in a tree that are close to
the root of a dense and large clade. A typical sample
of influenza HA sequences fall into a small number of
recognizable clades, and the strains with maximal LBI
will often be close to the root of the largest of those clades.
This root of the largest clade will often be close to the
consensus of the whole population, explaining the similar
distance patterns. To test that hypothesis, we measure the
hamming distance from the sequence of the top LBI strain
to the consensus sequence for populations of all time bins.
Panel **B** of figure 4 shows these distances, scaled with
respect to an average strain (details in caption). It clearly
shows that the top-LBI strain and the consensus sequence
are indeed quite similar: out of 48 time bins, only once
is the sequence of the top-LBI strain farther away from
the consensus than the average sequence is. Moreover,
the sequence of the top-LBI strain *exactly* matches the
consensus in 19 cases.

DISCUSSION

Predicting the trajectory of a mutation requires (i)
significant fitness difference between genomes carrying
different variants at the site and (ii) a selection pressure
that changes slowly over time. Under such conditions, it is
expected that frequency trajectories will show a persistent
behavior which would make them predictable for some
time. However, we could find only limited evidence for
such persistent behavior in the past 19 years of IAV
evolution. This lead us to conclude that (i) influenza




FIG. 4 **A:** Average Hamming distance of the sequences of different predictors to HA sequences of future influenza populations, themselves averaged over all “present” populations from year 2003 to 2019. Predictors are: a randomly picked sequence in the present population; the sequence of the strain with the highest LBI in the present population; the consensus sequence of the present population. **B:** Scaled Hamming distance between the sequence of the top LBI strain and the consensus sequence for populations at different dates. The scaling is such that for each date, the Hamming distance between a strain from the population and the consensus is on average 1. The strain with the highest LBI is almost always closer to the consensus sequence than the average strain.

virus evolution is qualitatively different from models of rapidly adapting population (despite clear evidence for frequent positive selection), and (ii) previous methods to predict influenza evolution work primarily because they pick strains that represent the future well, not because they predict future dynamics.

The primary focus in this work was the investigation of frequency trajectories of new amino acid mutations. In the short term, we found that on average the direction of trajectory does not persist for longer than a few months. Indeed, the average trajectory in figure 1 takes a sharp turn when going from $t < 0$ to $t > 0$, instead of showing “inertia”. This suggests that selective sweeps are not representative of typical trajectories. **Pierre: We could cite the Illingworth paper here. They do show that as well, in a different way. The fact many trajectories are not sweeps is expected when there is strong linkage. What cannot be easily explained by linkage only is the complete**

lack of inertia on average (at least I think?), and the $P_{fix}(f) = f$.

On a longer timescale, we investigated the probability that a novel mutation observed at frequency f fixes. In neutral models of evolution this probability equals f , while it should be higher or lower than f for mutations with a beneficial or deleterious effect on fitness, respectively. However, in the case of influenza, this probability differs little from f , making current frequency the best predictor for fixation. In figure 3, we split trajectories into groups for which we expected P_{fix} to deviate from f . Many of these splits, such as high/low LBI or epitope/non-epitope positions, did not result in an increased predictability, while others gave limited information on fixation. Despite the lack of predictability of mutation frequency trajectories, influenza surface proteins show strong signatures of selection (Bhatt et al., 2011; Strelkowa and Lässig, 2012).

Methods for predicting the future evolution of influenza

either construct explicit fitness models (Luksza and Lässig, 2014; ?), use historical patterns of evolution (Luksza and Lässig, 2014; Bush et al., 1999), phenotypic assays (Neher et al., 2016; Steinbrück and McHardy, 2012), or dynamic or phylogenetic patterns (Neher et al., 2014; Klingen et al., 2018). The goal of these methods is to pick strains that are good representatives of future populations and could serve as vaccine candidates (Morris et al., 2018).

The low power to predict frequency dynamics or fixation naturally triggers the question why the above methods have been found to work. Picking representatives of the future and predicting frequency dynamics are distinct objectives and success at the former (as compared to random picks) is not necessarily inconsistent with a lack of predictable dynamics. In fact, (?) reports that the rate at which the frequency of a strain changes is often a poor predictor – consistent with our observations here. But despite the fact that future frequencies are not predicted by the LBI, the strain with the highest LBI in the population is a better predictor of the future population than a randomly picked one. While the LBI was shown to be a correlate of relative fitness and be predictive of fixation in mathematical models of evolution (Neher et al., 2014), it does not seem to be [predictive of influenza evolution](#) because it measures fitness from genealogical structure. Instead, we believe it picks closer than average strains simply because it has the tendency to be maximal at the base of large and dense clades. These basal genotypes are closer to the future populations than the current tips of the tree and hence a better predictor on average. The consensus sequence of all present strains performs slightly but consistently better than picking the strain with the highest LBI. The consensus sequence is the best possible predictor for a neutrally evolving population, and does not attempt to model fitness in any way.

At the same time, influenza virus phylogenies show clear deviations from those expected from the neutral Kingman coalescent, similar to those expected under Bolthausen-Sznitman coalescent (BSC) processes that are generated by traveling wave models of rapid evolution (Neher and Hallatschek, 2013; Desai et al., 2013). The correspondence between the BSC and traveling wave models comes from transient exponential amplification of fit strains before these fitness differences are wiped out by further mutation. This exponential amplification generates long-tailed effective offspring distributions which in turn can lead to genealogies described by the BSC (Neher and Hallatschek, 2013; Schweinsberg, 2003). Many processes other than selection, including seasonality and spatio-temporal heterogeneity, can generate effective long tailed offspring distributions even in absence of bona-fide fitness differences, which might explain ladder-like non-Kingman phylogenetic trees.

A recent preprint proposed that influenza virus evolution is primarily limited by an asynchrony between population level selection and generation of new variants

within infected hosts (Morris et al., 2020). Along these lines, it is possible that the A/H3N2 population readily responds once population level selection is high enough by giving rise to essentially equivalent variants. Furthermore, selection might cause the rapid rise of a novel variant to macroscopic frequencies (observable in a global sample) but its benefit rapidly “expires” because competing variants catch up and/or it mediates immune escape only to a small fraction of the population. These considerations might explain the disconnect between models of rapid adaptation and the frequency dynamics observed in influenza virus populations.

METHODS

Data and code availability

The sequences used are obtained from the GISAID database (Shu and McCauley, 2017). Strain names and accession numbers are given as tables in two supplementary files. Outliers strains listed at <https://github.com/PierreBarrat/FluPredictability/src/config> were removed.

The code used to generate the figures presented here is available at <https://github.com/PierreBarrat/FluPredictability>.

Frequency trajectories

For a set of sequences in a given time bin, we compute frequencies of amino acids at each position by simple counting. We make the choice of not applying any smoothing method in an attempt to be as close to the data and “model-less” as possible. This is especially important for the short term prediction of frequency trajectories, as estimations of the “persistence time” of a trajectory might be biased by a smoothing method.

We compute frequency trajectories based on the frequencies of amino acids. A trajectory begins at time t if an amino acid is seen under the lower frequency threshold of 5% (resp. above the higher threshold of 95%) for the two time bins preceding t , and above this lower threshold (resp. below the higher threshold) for time bin t . It ends in the reciprocal situation, that is when the frequency is measured below the lower threshold (resp. above the higher threshold) for two time bins in a row.

In order to avoid estimates of frequencies that are too noisy, we only keep trajectories that are based on a population of at least 10 sequences for *each* time bin. As said in the Results section, we also restrict the analysis to trajectories that begin at a 0 frequency, in part to avoid double counting. We find a total of 460 such trajectories. However, only 106 reach a frequency of 20%, on which figure 2 is based for instance.

Note that the fact that we use samples of relatively small

725 sizes – at least for some time bins – leads to biases in⁷⁶²
 726 the estimation of frequencies. We show in Supplementary⁷⁶³
 727 Material that these biases are generally small and do not⁷⁶⁴
 728 induce any qualitative changes to results presented here.⁷⁶⁵

X is concentrated in only one region.
 Region used are the ones defined in the *Nextstrain*
 tool (Hadfield et al., 2018). Those are North America,
 South America, Europe, China, Oceania, Southeast Asia,
 Japan & Korea, South Asia, West Asia, and Africa.

729 Local Branching Index

730 LBI was introduced in (Neher et al., 2014) as an approx-
 731 imation of fitness in populations evolving under persistent⁷⁶⁸
 732 selective pressure that is fully based on a phylogenetic tree.
 733 It relies on the intuition that the tree below high-fitness
 734 individuals will show dense branching events, whereas
 735 absence of branching is a sign of low-fitness individuals.
 736 Quantitatively, the LBI $\lambda_i(\tau)$ of a node i is the integral
 737 of all of the tree’s branch length around i , with an expo-
 738 nentially decreasing weight $e^{-t/\tau}$ with t being the branch
 739 length. When considering a time binned population, the
 740 LBI is computed once for each time bin by considering
 741 only the leaves of the tree that belong to the time bin.
 742 This means that only branches that ultimately lead to
 743 a leaf that belongs to the time bin are considered in the
 744 integration.

745 τ is the time scale for which the tree is informative of
 746 the fitness of a particular node. Here, we use a value of
 747 τ equal to a tenth of $T_C \simeq 6$ years, the coalescence time
 748 for influenza A/H3N2 strains, converted to units of tree
 749 branch length through the average nucleotide substitution⁷⁶⁹
 750 rate ($\simeq 4 \cdot 10^{-3}$ substitutions per site per year for HA)⁷⁷⁰
 751 We have observed that given our method to predict the⁷⁷¹
 752 future from present populations corresponding to time
 753 bins of 4 months, changing the value of τ has little effect
 754 on the pick of the top LBI strain. By retrospectively⁷⁷²
 755 optimizing its value, it is possible to reduce the average
 756 distance to the population 2 years ahead by ~ 0.25 amino
 757 acids on average, making the LBI method almost as good
 758 as the consensus on figure 4.

759 Measuring the geographical spread of a mutation

For a mutation X we define its regional distribution using the numbers $n_r(X)$ that represent the number of sequences sampled in region r that carry X . Regional weights are then defined as

$$w_r(X) = \frac{n_r(X)}{\sum_r n_r(X)}.$$

We can then measure the geographical spread $G(X)$ of X by using the Shannon entropy of the probability distribution $w_r(X)$:

$$G(X) = \sum_r w_r(X) \log(w_r(X)).$$

760 $G(X)$ is a positive quantity that is larger when X is
 761 equally present in many regions, and equal to zero when

767 Assigning a fitness to trajectories

Consensus sequence

Given a set of N sequences $(\sigma^1, \dots, \sigma^N)$ based on an alphabet \mathcal{A} (e.g. \mathcal{A} has 20 elements for amino acids, 4 for nucleotides), we can define a *profile* distribution $p_i(a)$ by the following expression:

$$p_i(a) = \sum_{n=1}^N \delta_{\sigma_i^n, a}$$

where i is a position in the sequence, σ_i^n the character appearing at position i in sequence σ^n , a a character of the alphabet and δ the Kronecker delta. The profile $p_i(a)$ simply represents the fraction of sequences which have character a at position i .

We then simply define the consensus sequence σ^{cons} such that

$$\sigma_i^{cons} = \operatorname{argmax}_a p_i(a).$$

In other words, the consensus sequence is the one that has the dominant character of the initial set of sequences at each position.

Earth Mover’s Distance

In order to measure the distance of several predictor sequences to the future population, we rely on the *Earth Mover’s Distance* (EMD), a metric commonly applied in machine learning to compare collections of pixels or words (Rubner et al., 1998; Kusner et al., 2015). Here, we apply it to compute the distance between the sequences of two populations, noted as $\mathcal{X} = \{(x^n, p^n)\}$ and $\mathcal{Y} = \{(y^m, q^m)\}$ with $n \in \{1 \dots N\}$ and $m \in \{1 \dots M\}$. In this notation, x^n and y^m are sequences, and p^n and q^m are the frequencies at which these sequences are found in their respective populations. For convenience, we also define $d_{mn} = H(x^n, y^m)$ as the Hamming distance between pairs of sequences in the two populations.

We now introduce the following functional

$$F(\mathbf{w}) = \sum_{n,m} d_{nm} w_{nm},$$

with $\mathbf{w} = \{w_{nm}\}$ being a matrix of positive weights. The EMD between the two populations \mathcal{X} and \mathcal{Y} is now defined as the minimum value of function F under the conditions

$$\sum_{n=1}^N w_{nm} = q^m, \quad \sum_{m=1}^M w_{nm} = p^n, \text{ and } w_{nm} \geq 0$$

773 Intuitively, the weight w_{nm} tells us how much of sequence⁸²⁸
 774 x^n is “moved” to sequence y^m . The functional F sums⁸²⁹
 775 all of these moves and attributes them a cost equal to⁸³⁰
 776 the Hamming distance d_{nm} . The conditions on weights⁸³¹
 777 in \mathbf{w} ensure that all the weight p^n of x^n is “moved” to⁸³²
 778 elements in \mathcal{Y} and vice versa.⁸³³

779 The minimization is easily performed by standard linear⁸³⁴
 780 optimization libraries. Here, we use the Julia library⁸³⁵
 781 JuMP (Dunning et al., 2017).⁸³⁶

782 I. FUNDING

783 This work was funded in part by NIAID (R01AI127893-⁸⁴³
 784 01) and the SNF (310030_188547).⁸⁴⁴

785 REFERENCES

- 786 World Health Organization, 2018. URL [https://www.who.int/fr/news-room/fact-sheets/detail/influenza-\(seasonal\)](https://www.who.int/fr/news-room/fact-sheets/detail/influenza-(seasonal)).⁸⁵⁰
- 787 Velislava N. Petrova and Colin A. Russell. The evolution of⁸⁵³
 788 seasonal influenza viruses. *Nature Reviews Microbiology*, 16:⁸⁵⁴
 789 (1):47–60, October 2017. ISSN 1740-1526, 1740-1534. doi:⁸⁵⁵
 790 10.1038/nrmicro.2017.118. URL <http://www.nature.com/doifinder/10.1038/nrmicro.2017.118>.⁸⁵⁶
- 791 Arthur Chun-Chieh Shih, Tzu-Chang Hsiao, Mei-Shang Ho,⁸⁵⁸
 792 and Wen-Hsiung Li. Simultaneous amino acid substitutions⁸⁵⁹
 793 at antigenic sites drive influenza a hemagglutinin⁸⁶⁰
 794 evolution. *Proceedings of the National Academy of Sciences*, 104(15):6283–6288, 2007. ISSN 0027-8424. doi:⁸⁶¹
 795 10.1073/pnas.0701396104. URL <https://www.pnas.org/content/104/15/6283>.⁸⁶³
- 796 Samir Bhatt, Edward C. Holmes, and Oliver G. Pybus⁸⁶⁵
 797 The Genomic Rate of Molecular Adaptation of the Human⁸⁶⁶
 798 Influenza A Virus. *Molecular Biology and Evolution*, 28(9):2443–2451, September 2011. ISSN 0737-4038.⁸⁶⁷
 799 doi:10.1093/molbev/msr044. URL <https://academic.oup.com/mbe/article/28/9/2443/1007907>.⁸⁶⁹
- 800 Björn F. Koel, David F. Burke, Theo M. Bestebroer, Stefan⁸⁷¹
 801 van der Vliet, Gerben C. M. Zondag, Gaby Vervaeet, Eugene⁸⁷²
 802 Slepnev, Nicola S. Lewis, Monique I. J. Spronken,⁸⁷³
 803 Colin A. Russell, Mikhail Y. Eropkin, Aeron C. Hurt,⁸⁷⁴
 804 Ian G. Barr, Jan C. de Jong, Guus F. Rimmelzwaan, Albert⁸⁷⁵
 805 D. M. E. Osterhaus, Ron A. M. Fouchier, and Derek⁸⁷⁶
 806 Smith. Substitutions near the receptor binding site determine⁸⁷⁷
 807 major antigenic change during influenza virus evolution.⁸⁷⁸
 808 *Science*, 342(6161):976–979, 2013. ISSN 0036-8075. doi:10.1126/science.1244730. URL <https://science.sciencemag.org/content/342/6161/976>.⁸⁸¹
- 809 Dylan H Morris, Katelyn M Gostic, Simone Pompei, Trevor⁸⁸²
 810 Bedford, Marta Luksza, Richard A Neher, Bryan T Grenfell,⁸⁸³
 811 Michael Lässig, and John W McCauley. Predictive modeling⁸⁸⁴
 812 of influenza shows the promise of applied evolutionary⁸⁸⁵
 813 biology. *Trends in microbiology*, 26(2):102–118, 2018.⁸⁸⁶
- 814 Thorsten R. Klingen, Susanne Reimering, Carlos A⁸⁸⁷
 815 Guzmán, and Alice C. McHardy. In Silico Vaccine⁸⁸⁸
 816 Strain Prediction for Human Influenza Viruses⁸⁸⁹
 817 *Trends in Microbiology*, 26(2):119–131, February 2018a.⁸⁹⁰
 818 ISSN 0966-842X. doi:10.1016/j.tim.2017.09.001. URL⁸⁹¹
- 819 <http://www.sciencedirect.com/science/article/pii/S0966842X17302068>.
- 820 Peter Bogner, Ilaria Capua, David J Lipman, and Nancy J Cox. A global initiative on sharing avian flu data. *Nature*, 442(7106):981–981, 2006.
- 821 Yuelong Shu and John McCauley. Gisaid: Global initiative⁸⁹²
 822 on sharing all influenza data—from vision to reality. *Euro-surveillance*, 22(13), 2017.
- 823 Andrew Rambaut, Oliver G. Pybus, Martha I. Nelson, Cecile Viboud, Jeffery K. Taubenberger, and Edward C. Holmes. The genomic and epidemiological dynamics of human influenza A virus. *Nature*, 453(7195):615–619, May 2008. ISSN 1476-4687. doi:10.1038/nature06945. URL <https://www.nature.com/articles/nature06945>.
- 824 Marta Luksza and Michael Lässig. A predictive fitness model for influenza. *Nature*, 507(7490):57–61, March 2014. ISSN 1476-4687. doi:10.1038/nature13087. URL <https://www.nature.com/articles/nature13087>. Number: 7490 Publisher: Nature Publishing Group.
- 825 L. Steinbrück, T. R. Klingen, and A. C. McHardy. Computational prediction of vaccine strains for human influenza a (h3n2) viruses. *Journal of Virology*, 88(20):12123–12132, 2014. ISSN 0022-538X. doi:10.1128/JVI.01861-14. URL <https://jvi.asm.org/content/88/20/12123>.
- 826 Richard A. Neher, Trevor Bedford, Rodney S. Daniels, Colin A. Russell, and Boris I. Shraiman. Prediction, dynamics, and visualization of antigenic phenotypes of seasonal influenza viruses. *Proceedings of the National Academy of Sciences of the United States of America*, 113(12):E1701–1709, March 2016. ISSN 1091-6490 0027-8424. doi:10.1073/pnas.1525578113.
- 827 Richard A. Neher, Colin A. Russell, and Boris I. Shraiman. Predicting evolution from the shape of genealogical trees. *eLife*, 3, November 2014. ISSN 2050-084X. doi:10.7554/eLife.03568. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4227306/>.
- 828 Natalja Strelkowa and Michael Lässig. Clonal Interference in the Evolution of Influenza. *Genetics*, 192(2):671–682, October 2012. ISSN 0016-6731, 1943-2631. doi:10.1534/genetics.112.143396. URL <http://www.genetics.org/content/192/2/671>.
- 829 Yuri I Wolf, Cecile Viboud, Edward C Holmes, Eugene V Koonin, and David J Lipman. Long intervals of stasis punctuated by bursts of positive selection in the seasonal evolution of influenza A virus. *Biology Direct*, 1:34, October 2006. ISSN 1745-6150. doi:10.1186/1745-6150-1-34. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1647279/>.
- 830 Juhye M Lee, Rachel Eguia, Seth J Zost, Saket Choudhary, Patrick C Wilson, Trevor Bedford, Terry Stevens-Ayers, Michael Boeckh, Aeron C Hurt, Seema S Lakdawala, Scott E Hensley, and Jesse D Bloom. Mapping person-to-person variation in viral mutations that escape polyclonal serum targeting influenza hemagglutinin. *eLife*, 8:e49324, August 2019. ISSN 2050-084X. doi:10.7554/eLife.49324. URL <https://doi.org/10.7554/eLife.49324>. Publisher: eLife Sciences Publications, Ltd.
- 831 Motoo Kimura. Diffusion Models in Population Genetics. *Journal of Applied Probability*, 1(2):177–232, 1964. ISSN 0021-9002. doi:10.2307/3211856. URL <http://www.jstor.org/stable/3211856>.
- 832 Le Yan, Richard A Neher, and Boris I Shraiman. Phylogenetic theory of persistence, extinction and speciation of rapidly adapting pathogens. *eLife*, 8:e44205, September 2019. ISSN 2050-084X. doi:10.7554/eLife.44205. URL

- 892 <https://doi.org/10.7554/eLife.44205>. Publisher: eLife⁹⁵⁶
 893 Sciences Publications, Ltd.⁹⁵⁷
- 894 R. A. Neher and B. I. Shraiman. Genetic draft and quasi-⁹⁵⁸
 895 neutrality in large facultatively sexual populations. *Genetics*⁹⁵⁹
 896 188(4):975–996, August 2011. ISSN 1943-2631 0016-6731⁹⁶⁰
 897 doi:10.1534/genetics.111.128876.⁹⁶¹
- 898 Fabio Zanini and Richard A. Neher. FFPopSim: an effi-⁹⁶²
 899 cient forward simulation package for the evolution of large⁹⁶³
 900 populations. *Bioinformatics*, 28(24):3332–3333, 10 2012⁹⁶⁴
 901 ISSN 1367-4803. doi:10.1093/bioinformatics/bts633. URL⁹⁶⁵
 902 <https://doi.org/10.1093/bioinformatics/bts633>.⁹⁶⁶
- 903 R. M. Bush, C. A. Bender, K. Subbarao, N. J. Cox, and⁹⁶⁷
 904 W. M. Fitch. Predicting the evolution of human influenza A⁹⁶⁸
 905 *Science (New York, N.Y.)*, 286(5446):1921–1925, December⁹⁶⁹
 906 1999. ISSN 0036-8075.⁹⁷⁰
- 907 Katherine S. Xue, Louise H. Moncla, Trevor Bedford,⁹⁷¹
 908 and Jesse D. Bloom. Within-Host Evolution of Hu-⁹⁷²
 909 man Influenza Virus. *Trends in Microbiology*. ISSN⁹⁷³
 910 0966-842X. doi:10.1016/j.tim.2018.02.007. URL⁹⁷⁴
 911 <https://www.sciencedirect.com/science/article/pii/S0966842X1830043X>.⁹⁷⁵
- 912 Dylan H. Morris, Velislava Petrova, Fernando W. Rossine,⁹⁷⁷
 913 Edyth Parker, Bryan Grenfell, Richard Neher, Simon Levin,⁹⁷⁸
 914 and Colin Russell. Asynchrony between virus diversity and
 915 antibody selection limits influenza virus evolution. preprint,
 916 Open Science Framework, April 2020. URL <https://osf.io/847p2>.⁹⁷⁹
- 917 Lars Steinbrück and Alice Carolyn McHardy. Inference
 918 of Genotype–Phenotype Relationships in the Antigenic
 919 Evolution of Human Influenza A (H3N2) Viruses. *PLOS
 920 Computational Biology*, 8(4):e1002492, April 2012. ISSN
 921 1553-7358. doi:10.1371/journal.pcbi.1002492. URL
 922 <http://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1002492>.
- 923 Thorsten R. Klingen, Susanne Reimering, Jens Loers, Kyra
 924 Mooren, Frank Klawonn, Thomas Krey, Gülsah Gabriel,
 925 and Alice C. McHardy. Sweep Dynamics (SD) plots: Com-
 926 putational identification of selective sweeps to monitor the
 927 adaptation of influenza A viruses. *Scientific Reports*, 8(1):
 928 373, January 2018b. ISSN 2045-2322. doi:10.1038/s41598-
 929 017-18791-z. URL <https://www.nature.com/articles/s41598-017-18791-z>.
- 930 Richard A. Neher and Oskar Hallatschek. Genealogies of
 931 rapidly adapting populations. *Proceedings of the National
 932 Academy of Sciences of the United States of America*, 110
 933 (2):437–442, January 2013. ISSN 1091-6490 0027-8424. doi:
 934 10.1073/pnas.1213113110.
- 935 Michael M. Desai, Aleksandra M. Walczak, and Daniel S.
 936 Fisher. Genetic Diversity and the Structure of Genealo-
 937 gies in Rapidly Adapting Populations. *Genetics*, 193(2):
 938 565–585, February 2013. ISSN 0016-6731, 1943-2631. doi:
 939 10.1534/genetics.112.147157. URL <http://www.genetics.org/content/193/2/565>.
- 940 Jason Schweinsberg. Coalescent processes obtained from
 941 supercritical Galton–Watson processes. *Stochastic Pro-
 942 cesses and their Applications*, 106(1):107–139, July 2003.
 943 ISSN 0304-4149. doi:10.1016/S0304-4149(03)00028-0.
 944 URL <http://www.sciencedirect.com/science/article/pii/S0304414903000280>.
- 945 James Hadfield, Colin Megill, Sidney M Bell, John Hud-
 946 dleston, Barney Potter, Charlton Callender, Pavel Sag-
 947 ulenko, Trevor Bedford, and Richard A Neher. Nextstrain:
 948 real-time tracking of pathogen evolution. *Bioinformat-
 949 ics*, 34(23):4121–4123, 05 2018. ISSN 1367-4803. doi:
 950 10.1093/bioinformatics/bty407. URL <https://doi.org/10.1093/bioinformatics/bty407>.
- 951 Y. Rubner, C. Tomasi, and L. J. Guibas. A metric
 952 for distributions with applications to image databases.
 953 In *Sixth International Conference on Computer Vision
 954 (IEEE Cat. No.98CH36271)*, pages 59–66, Jan 1998. doi:
 955 10.1109/ICCV.1998.710701.
- 956 Matt J. Kusner, Yu Sun, Nicholas I. Kolkin, and Kilian Q.
 957 Weinberger. From word embeddings to document distances.
 958 In *Proceedings of the 32nd International Conference on
 959 International Conference on Machine Learning - Volume
 960 37, ICML'15*, page 957–966. JMLR.org, 2015.
- 961 Iain Dunning, Joey Huchette, and Miles Lubin. Jump: A
 962 modeling language for mathematical optimization. *SIAM
 963 Review*, 59(2):295–320, 2017. doi:10.1137/15M1020575.
- 964 Julia Sigwart. Gene Genealogies, Variation and Evolution:
 965 A Primer in Coalescent Theory.—Jotun Hein, Mikkel H.
 966 Schierup, and Carsten Wiuf. 2004. Oxford University Press,
 967 Oxford. xiii + 276 pp. ISBN 0-19-852996-1, £29.95 (paper-
 968 back); ISBN 0-19-852995-3, £65.00 (hardback). *Systematic
 969 Biology*, 54(6):986–987, 12 2005. ISSN 1063-5157. doi:
 970 10.1080/10635150500354860. URL <https://doi.org/10.1080/10635150500354860>.

979 **SUPPLEMENTARY MATERIAL**980 **1. Consensus sequence as a predictor for neutrally evolving populations**

981 We consider the case of a neutrally evolving and structure-less population, such as the one in the Wright-Fisher
 982 model of evolution (Sigwart, 2005). At an initial time $t = 0$, the population consists of N individuals with genomes
 983 $(\sigma^1 \dots \sigma^N)$ of length L (not necessarily distinct).

984 We make two hypotheses about this population. We first suppose that *no* mutations occur during the evolution of this
 985 population. This may seem surprising and is of course not true in the case of influenza. This assumption is however in
 986 line with the fact that the object of this work is to predict the outcome of *already existing* mutations in the influenza
 987 population. The prediction of mutations that we have not yet seen is not in its scope. Thus, assuming that no new
 988 mutations take place can be seen as a simple way to model the fact that we have no information about such events.
 989 The second assumption is that the population evolves in a completely neutral way, meaning that the average number
 990 of descendants of each genome σ^n is the same. Let us now consider the population after it has evolved for a long time
 991 $t \gg T$ where T is the typical coalescence time (for the Wright-Fisher model, $T = 2N$). At this point, all individuals in
 992 the future population will descend from a unique individual n_0 in the $t = 0$ population. Our two hypotheses now allow
 993 us to make two statements. First, since no new mutations are allowed, the population at $t \gg T$ will be clonal, with all
 994 individuals having genome σ^{n_0} . Second, since the evolution is neutral and does not favour any genome in particular,
 995 the probability that σ^{n_0} is equal to a given genome σ is $1/N$. In other words, the probability that a genome at $t = 0$
 996 ultimately becomes the ancestor of all the future population is equal to its frequency in the $t = 0$ population.

997

998 We now try to find the genome σ that best predicts the future population on the long run, that is for $t \gg T$. Here,
 999 we take best to mean that the predictor minimizes $H(\sigma, \sigma^{n_0})$ where H is the Hamming distance defined by

$$H(\sigma^a, \sigma^b) = \sum_{i=1}^L (1 - \delta_{\sigma_i^a, \sigma_i^b}), \quad (1)$$

1000 with σ_i being the character appearing at position i of genome σ and δ the Kronecker delta. Since we do not know n_0 ,
 1001 we have to average over all its possible values. σ must thus minimize the following quantity:

$$\begin{aligned} \langle H(\sigma, \sigma^{n_0}) \rangle_{n_0} &= \sum_{n=1}^N H(\sigma, \sigma^n) \\ &= \sum_{i=1}^L \sum_{n=1}^N (1 - \delta_{\sigma_i, \sigma_i^n}) \end{aligned} \quad (2)$$

by using the definition of the Hamming distance. We now assume that characters at each positions of the genomes can
 be indexed by an integer a running from 1 to q . For instance, if these were amino acid sequences, we could index the
 20 amino acids by a running from 1 to $q = 20$. We rewrite the Kronecker delta in the previous expression using this
 indexation:

$$\delta_{\sigma_i, \sigma_i^n} = \sum_{a=1}^q \delta_{\sigma_i, a} \delta_{\sigma_i^n, a}.$$

1002 We also introduce the *profile* frequencies $p_i(a)$ of the population at time $t = 0$:

$$p_i(a) = \sum_{n=1}^N \delta_{\sigma_i^n, a}. \quad (3)$$

1003 $p_i(a)$ represents the frequency at which character a appears at position i in genomes of the initial population.

1004 Equation 2 now becomes

$$\begin{aligned}
 \langle H(\sigma, \sigma^{n_0}) \rangle_{n_0} &= \sum_{i=1}^L \sum_{n=1}^N \left(1 - \sum_{a=1}^q \delta_{\sigma_i, a} \delta_{\sigma_i^n, a} \right) \\
 &= \sum_{i=1}^N \left(1 - \sum_{a=1}^q \delta_{\sigma_i, a} p_i(a) \right) \\
 &= \sum_{i=1}^L (1 - p_i(\sigma_i))
 \end{aligned} \tag{4}$$

1005 This means that the genome $\sigma = (\sigma_1 \dots \sigma_L)$ which best predicts the future population according to our definition is
 1006 the one that minimizes the quantity $(1 - p_i(\sigma_i))$ for all positions i . This obviously implies that each σ_i must be chosen
 1007 as to maximize $p_i(a)$, that is σ_i must be the character that appears the most frequently at position i . Thus, σ must be
 1008 the *consensus* sequence of the initial population.

1009 **2. Predictor based on the local LBI maxima**

In figure 15, we use several sequences as a predictor of the future population. Distance between two sets of sequences, *i.e.* the predictor sequences and the ones of the future population, is defined as the Earth Mover's Distance (EMD). Here, we show that for a population evolving under the same hypotheses as in section 1, the best *multiple* sequence long term predictor is again the consensus sequence with weight 1.

Let the predictor be a set of weighted sequences $\{(s^\alpha, q_\alpha)\}$. We again use the fact that in the long term, a unique sequence σ^{n_0} from the present will be the ancestor of the entire population. We want to compute the EMD from the predictor to σ^{n_0} , that is the EMD between the sets $\mathcal{X} = \{(s^\alpha, q_\alpha)\}$ and $\mathcal{Y} = \{\sigma^{n_0}, 1\}$. Applying the definition of the Methods section, it follows that the weights \mathbf{w} are in this case equal to the q_α s. By averaging over all values of n_0 , we now obtain

$$\langle \text{EMD}(\{(s^\alpha, q_\alpha)\}) \rangle_{n_0} = \sum_{n=1}^N \sum_{\alpha} H(s^\alpha, \sigma^n) \cdot q_\alpha.$$

By the same calculation procedure as in the previous section, this expression simplifies to

$$\langle \text{EMD}(\{(s^\alpha, q_\alpha)\}) \rangle_{n_0} = \sum_{i=1}^L \left(1 - \sum_{a=1}^q p_i(a) q_i(a) \right),$$

where the profile of the present population $p_i(a)$ has already been defined, and $q_i(a)$ stands for the profile of the predictor, that is

$$q_i(a) = \sum_{\alpha} \delta_{s_i^\alpha, a} q_\alpha.$$

1010 To minimize this distance, we find a profile $q_i(a)$ that maximizes the quantity $\sum_{\alpha} \delta_{s_i^\alpha, a} q_\alpha$ for each position i . It is
 1011 clear that this is done by assigning a value $q_i(a) = 1$ if a maximizes $p_i(a)$, and $q_i(a) = 0$ otherwise. Thus, the profile of
 1012 the predictor must be that of the consensus sequence, which is only possible if the predictor becomes $\{\sigma^{cons}, 1\}$.

1013 **3. Correcting for nested mutations**

1014 The analysis of the main text computes probabilities of fixation assuming that all trajectories are independent.
 1015 However, it is well-known that mutations in influenza viruses are nested: they appear on backgrounds that already
 1016 carry other mutations. Since mutations appearing on the same genomes will jointly fix or disappear, many frequency
 1017 trajectories are not independent but correlated. In order to compensate for potential biases due to this effect, we
 1018 attempted to cluster trajectories based on similarity in their strain composition. Our aim is that two trajectories
 1019 corresponding to mutations appearing mostly on the same genomes will be grouped in the same cluster. We then
 1020 conduct the same analysis as in the main text on a set of *effectively independent* trajectories constructed by taking
 1021 one trajectory from each cluster.

In order to perform clustering, we define a distance between trajectories. A frequency trajectory X is characterized by a series of frequency values and a time interval T . $f(t)$ for $t \in T$ corresponds to the frequency at which a given mutation x appears in the population at date t , and we define $S(t)$ as the strains that carry mutation x at date t . With this notation, $f(t)$ is the ratio of the number of elements in $S(t)$ to the total number of strains at date t . Let us now consider two frequency trajectories X_1 and X_2 .

We define the distance $d(X_1, X_2)$ between these two trajectories based on the average similarity of the strains S_1 and S_2 that compose them:

$$d(X_1, X_2) = \frac{1}{|T_1 \cap T_2|} \sum_{t \in T_1 \cap T_2} \frac{|S_1(t) \cap S_2(t)|}{|S_1(t) \cup S_2(t)|},$$

where $T_1 \cap T_2$ is the time interval where both trajectories are active, and $|\cdot|$ denotes the number of elements of a set. The quantity summed corresponds to the Jaccard index between strains composing X_1 and X_2 at a given date. It is 1 if the two trajectories share exactly the same strains for this date, and 0 if they share no strain at all. This leads to the two following properties of d :

- if $d(X_1, X_2) = 0$, then X_1 and X_2 represent the same frequency trajectory. The mutations x_1 and x_2 that they correspond to always appear on the same strains and are totally linked.
- if $d(X_1, X_2) = 1$, then X_1 and X_2 can be considered completely independent. This can be the case if the two trajectories do not occur at the same dates, *i.e.* $|T_1 \cap T_2| = 0$, or if their respective mutations are never present on the same genomes.

we attempt to reduce the potential statistical bias due to the nesting of trajectories by grouping them based on the above defined distance. Given a set of trajectories $\{X\}$, we perform a decomposition of $\{X\}$ into disjoint clusters $C_1(d^*) \cup \dots \cup C_n(d^*) = \{X\}$ where d^* is an arbitrary threshold distance. Clusters are built in such a way that given two trajectories X_i and X_j

$$d(X_i, X_j) \leq d^* \Rightarrow \exists k : X_i, X_j \in C_k$$

and

$$X_i \in C_k \Rightarrow \exists j \in C_k : d(X_i, X_j) \leq d^*.$$

These condition imply that the clusters formed are the minimal ones that guarantee that any two trajectories closer than the threshold distance d^* belong to the same cluster. The number of clusters n depends on the chosen value for d^* .

We compute clusters for different values of d^* for the case of the HA gene in A/H3N2. The top panel of figure S?? shows the number cluster n as a function of d^* . In the $d^* = 0$ case, only trajectories that are exactly identical in terms of strain composition are clustered together. In this case, our clustering amounts to counting mutations that appear on exactly the same strains as one, reducing the number of effective trajectories from 800 to slightly less than 700. For higher values of d^* , the number of clusters steadily goes down until it reaches 1 for $d^* = 1$, which is the maximum value of the distance $d(X_1, X_2)$. The sharp drop in n for $d^* = 0.5$ is explained by the high number of very short (typically one time point) and low frequency trajectories that share one out of two strains.

Since the choice of d^* is arbitrary and since no particular value can be chosen based on the number of clusters $n(d^*)$, we decide to test our clustering strategy for five values, namely $d^* \in \{0, 0.05, 0.1, 0.2, 0.49\}$. The bottom panel of figure S?? shows examples of a cluster for the four non-zero values of d^* . The cluster displayed in each case is the one containing the mutation HA1:33R. As d^* increases, more and more unlike trajectories are grouped together. In the case $d^* = 0.49$, the cluster consists of 13 trajectories, 5 of which end up dying while the rest fix. Since such a high value of d^* results in grouping trajectories that do not have the same fate (fixation or death), we decide to exclude it from the rest of the analysis, resulting in four remaining values $d^* \in \{0, 0.05, 0.1, 0.2\}$.

Once clustering is performed, we re-conduct the analysis of the main text on a set of effective trajectories. This set is constructed by taking one trajectory at random from each cluster. Effective trajectories are then considered independent from each other. The left panel of figure S?? shows the fixation probability of trajectories as a function of their frequency for different values of d^* , for the HA gene of A/H3N2. The result obtained in panel A of figure




Figure S 1 **Top:** Left-axis: number n of clusters as a function of the threshold distance d^* . The total number of trajectories is shown as a flat orange line. Right-axis: average size of clusters as a function of d^* . **Bottom:** Examples of clusters for four values of d^* . The four clusters displayed are the ones to which mutation HA1:33R belongs to.

2 of the main text is also showed as a reference. For the three lower values of d^* , results do not differ from the one obtained in the main text, even though the number of trajectories in each frequency bin has dropped as can be seen in the right panel of figure S???. This indicates that grouping together trajectories that share most of their strains, and are thus very correlated, does not modify the computed fixation probability in any way. For the higher value $d^* = 0.2$, fixation probability drops slightly across all frequency bins, suggesting that fixating trajectories tend to be grouped together more frequently. However, this drop remains of limited amplitude.

Overall, this analysis leads us to think that even though mutations in influenza may be nested, considering trajectories as independent does not result in strong statistical biases. Indeed, clustering similar trajectories together does significantly modify results presented in the main text.




Figure S 2 **Left:** Fixation probability of trajectories as a function of probability for four values of d^* . The reference curve is the same as in panel A of figure 2 for the HA gene. For readability, error bars are only displayed for the reference curve. **Bottom:** Number of trajectories in each frequency bin corresponding to the left panel.

1062 4. Ability of fitness models to predict fixation

1063 In a companion article (?), authors developed several models to predict the future HA population for A/H3N2
 1064 influenza. The premise of these models is to assign a fitness score to the HA gene of each A/H3N2 strain in a given
 1065 year, and to then apply deterministic evolutionary equations to obtain the future strain composition. The fitness
 1066 scores are derived from several quantities that have been shown to be connected with viral fitness.

1067 Here, we assess the ability of four fitness scores computed in (?) to predict fixation: mutational load and
 1068 epitope antigenic novelty introduced in (Luksza and Lässig, 2014), hemagglutination inhibition (HI) antigenic novelty
 1069 (Neher et al., 2016) and the previously mentioned Local Branching Index (LBI) (Neher et al., 2014). Figure S??
 1070 shows fixation probability for HA mutations with fitness scores in the top or bottom half of the fitness distribution
 1071 for the four fitness measures. *Pierre:* Maybe use delta-frequency instead of epitope mutations? They behave weirdly
 1072 (high fitness is worse), and delta-frequency uses more than the last time point in the trajectory, which may partly
 1073 answer one point of Rev1.

1074 Next, we use two fitness scores based on linear combinations of the following quantities: mutational load + LBI
 1075 and mutational load + HI antigenic novelty. These two combinations were found in (?) to be the best predictors of
 1076 future population. The coefficients of the linear combinations were trained each year using the five previous years of
 1077 data, and here we use their average value over a 25 years training period.

1078 5. Biases in frequency estimations

1079 The frequency of mutations in a given time-bin is simply performed by computing their frequency in sequences
 1080 sampled in that time bin. This leads to potential biases in estimating frequencies, that arise for two reasons:

- 1081 (i) A mutation present at frequency p in the population might be observed at another frequency $f \neq p$ if f is
 1082 estimated using a sub-sample of the population.
- 1083 (ii) For a neutrally evolving population, the distribution of frequencies of alleles is of the form $P(p) \propto 1/p$. This




Figure S 3 Ability of fitness scores based on **A** LBI, **B** HI titer, **C** mutational load and **D** epitope antigenic novelty to predict fixation of mutation frequency trajectories in the HA gene of A/H3N2. Similar to figure 3 of the main text.




Figure S 4 Ability of fitness scores based on two combinations of the four fitness scores mentioned above: **A** mutational load + LBI; **B** mutational load + HI titer.




Figure S 5 **Left:** For a mutation present at frequency p in the population, probability of being observed in the frequency bin $[0.1, 0.2]$ as a function of p and for different sample sizes n . The dashed black line sketches the (non-normalized) background distribution $P_b(p)$. **Right:** Expected “real” average frequency of mutations found in frequency bin $[f_1, f_2]$ as a function of the centre of the bin $(f_1 + f_2)/2$, for different sample sizes.

means that the amount of alleles at frequency p is lower when p is higher.

To illustrate (i), let us compute the probability that a mutation present at “real” frequency p in the population is found to be in a given frequency bin $[f_1, f_2]$ when p is estimated from a sample of size n . The sample consists of n observations $\{x_i\}$ with $1 \leq i \leq n$, with $x_i = 1$ if sequence sequence i of the sample bears the mutation, and $x_i = 0$ if not. If n is small with regard to the total population size, we can consider the x_i as random variables with a binomial distribution, meaning that $P(x_i = 1) = p$ and $P(x_i = 0) = 1 - p$. The empirical frequency f is then estimated by taking the average of the x_i variables, that is $f = (x_1 + \dots + x_n)/n$. If those are independently sampled and n is large enough, the probability of measuring value f is given by the Central Limit Theorem:

$$P_{n,p}(f) \propto e^{(f-p)^2/2\sigma^2}, \text{ where } \sigma^2 = \frac{p(1-p)}{n}. \quad (5)$$

To compute the probability that this mutation is found in a given frequency bin $[f_1, f_2]$, we integrate this distribution:

$$P_{f_1, f_2}(p, n) = \int_{f_1}^{f_2} dx P_{n,p}(x). \quad (6)$$

Function $P_{f_1, f_2}(p, n)$ is shown as a function of p for a fixed interval and for different values of n in the first panel of figure S1. Note the asymmetry of it: the variance of a binomial distribution of parameter p is small when p is close to 0 or 1, and goes through a maximum at $p = 0.5$. For this reason, mutations present at frequency p close to 0.5 have a higher probability of being observed in other frequency bins. On the contrary, this is unlikely for very rare or very frequent mutations.

We now try to estimate biases in frequency estimation due this phenomenon. Given a set of mutations that have been measured in frequency bin $[f_1, f_2]$, what is the average *real* frequency of these mutations? To compute this, we need to sum $P_{f_1, f_2}(p, n)$ over all possible real frequencies p , giving us the amount of mutations that are observed in interval $[f_1, f_2]$, and weigh this sum by the frequency value p as well as by the background distribution of frequencies $P_b(p) \propto 1/p$. This last quantity represents the expected amount of mutations that are present at frequency p in the population. Note that there is no divergence problem as the smallest non zero frequency is $1/N$, where N is the

1106 population size. This leads us to the following expression for the average of “real” frequencies:

$$\begin{aligned}\langle p \rangle(f_1, f_2, n) &= \int_{1/N}^{1-1/N} dp P_{f_1, f_2}(p, n) P_b(p) p \\ &= \int_{1/N}^{1-1/N} dp P_{f_1, f_2}(p, n).\end{aligned}\tag{7}$$

1107 We have not made normalization explicit in these equations. It is simply achieved by dividing the above expression by
1108 $\int dp P_{f_1, f_2}(p, n) P_b(p)$.

1109 In the second panel of figure S1, $\langle p \rangle(f_1, f_2, n)$ is plotted as a function of the centre of the interval $[f_1, f_2]$ and for
1110 different values of n . For sample sizes $n > 100$, the biases due to this effect are almost non existent. For smaller
1111 samples, for instance $n = 10$, they are small but non negligible. However, we argue that this is not a significant problem
1112 with respect to the main results presented in this article. First, figure S6 shows that sample sizes of the order of $n = 10$
1113 are only the case for a few months in the period going from year 2000 to 2018. From 2010 and onwards, more than a
1114 hundred sequences are available per month for most months. Secondly, even if most samples were in the $n = 10$ case,
1115 deviations shown in figure S1 are small enough that results shown in figures 2 and 3 would be *qualitatively* unchanged.
1116 Note that using the centre of the interval as a reference in figure S1, *i.e.* $(f_1 + f_2)/2$, would be correct in the case of
1117 a very large n and a flat background distribution $P_b(p)$. For figures 2 and 3 of the main text however, the average
1118 frequency of mutations found in an interval $[f_1, f_2]$ is computed by taking the average of the observed frequencies, and
1119 not the centre of the interval. This partially takes into account biases considered here, as the background distribution
1120 $P_b(p)$ is then accounted for, even though it is equivalent to assuming infinite sample sizes.

1121 6. Cutting off the HA1 159S branch



Figure S 6 Tree used for this study, based on a random selection of 100 strains per month from year 2002 to 2018. Nodes and branches are colored according to the amino acid found at position HA1:159. The HA1 159S mutation is visible as a thin but long light-greened color branch, coalescing with the “trunk” around 2013.

1122 The analysis of the main text is in a large part based on the probability of fixation of mutations. The motivation
 1123 underlying this choice is the relatively short coalescence time of the A/H3N2 influenza population, typically around
 1124 three years. This can be seen in figure 2 of the main text, which shows the typical lifetime of frequency trajectories,
 1125 ending in fixation or loss after at most 3 years in most cases. The tree in figure S2 is another illustration of this: for
 1126 the most part of it, a “trunk” is clearly identifiable, and lineages that depart from it have a relatively short lifetime.
 1127 This is no longer the case since the year ~ 2013: two clades have been competing since then, with no definite way to
 1128 identify a trunk in the tree. The clade defined by the HA1 159S mutation, colored in light green on figure S2, is one of
 1129 these two competing lineages. Because of this particular situation, the number of mutations fixating in the population
 1130 is strongly reduced, as a mutation must appear in both clades to reach a frequency of 1. This is a potential flaw in our
 1131 analysis, which concentrates on mutations fixating.

1132 For this reason, we decided to re-run our analysis after having cut off the HA1 159S clade. In other words, we remove
 1133 from the set of sequences those that carry the HA1 159S mutation. Results are shown in figures , equivalent to figures 2
 1134 and 3 of the main text. It is clear that qualitative results are left unchanged when this competing clade is removed.
 1135 This can be surprising, as almost no complete fixation of an amino acid mutation has occurred since 2013. Cutting off
 1136 the HA1 159S branch should thus result in many new fixations, changing the analysis. The reason for the similarity
 1137 of results can be explained: fixation (resp. loss) of a mutation are defined here as the frequency of this mutation
 1138 being measured above 95% (resp. 5%) frequency for two months in a row. As the HA1 159S clade is rather sparsely
 1139 populated, it reaches frequencies lower than 5% two times (in 2015 and 2017), allowing mutations in the competing
 1140 clade to “fix” as defined here. Thus, removing strains carrying HA1 159S does not introduce a significant amount of
 1141 “new” fixation events.




Figure S 7 Equivalent to figure 2 of the main text, but with strains carrying the HA1 159S mutation removed.




Figure S 8 Equivalent to figure 3 of the main text, but with strains carrying the HA1 159S mutation removed.

1142 **7. Probability of fixation in single locus model of evolution**

1143 In (Kimura, 1964), Kimura investigates a simple model of evolution with a single locus and a population of size
 1144 N . In this framework, a mutation at this locus with fitness effect s and observed at frequency f has the following
 1145 probability of fixation:

$$P_{fix}(f|s, N) = \frac{1 - e^{-sNf}}{1 - e^{-sN}}. \quad (8)$$

1146 Expanding this formula for $sN \ll 1$, that is in the weak selection regime, yields at the first order

$$P_{fix}(f|s, N) = f + f(1 - f) \frac{sN}{2}. \quad (9)$$

1147 Equation 9 tells us two things. First, when the mutation is neutral, that is $s = 0$, we have $P_{fix}(f) = f$. This naturally
 1148 confirms the result obtained for a neutral model of evolution. Seconds, when $sN \neq 0$, we can expect deviations from
 1149 the diagonal in a P_{fix} against f plot. The sign of these deviations is determined by the sign of s , with beneficial
 1150 mutations being found above diagonal while deleterious one are found below. The amplitude of these deviations
 1151 depends on the strength of selection sN , as well as on the frequency through the $f(1 - f)$ term, making them larger
 1152 for $f \sim 0.5$.

1153 8. Mutation tables

Gene	Position	AA	Start date	End date	Shih	Luksza	Koel	Tree counts
HA1	144	D	2001-06-09	2002-02-04	true	true	false	0
HA1	189	N	2003-07-29	2004-05-24	false	true	true	2
HA1	159	F	2003-08-28	2004-05-24	false	true	true	2
HA1	226	I	2003-09-27	2004-09-21	true	true	false	3
HA1	145	N	2003-12-26	2004-11-20	false	true	true	2
HA1	227	P	2003-05-30	2005-04-19	false	true	false	2
HA2	32	I	2004-06-23	2005-07-18	false	false	false	1
HA1	193	F	2004-12-20	2006-03-15	false	true	true	1
HA2	46	D	2006-06-13	2007-05-09	false	false	false	2
HA2	121	K	2006-06-13	2007-06-08	false	false	false	1
HA1	50	E	2006-09-11	2007-06-08	false	true	false	2
HA1	140	I	2006-11-10	2007-11-05	true	false	false	1
HA1	173	Q	2007-07-08	2009-01-28	true	true	false	2
HA2	32	R	2007-07-08	2009-01-28	false	false	false	1
HA1	158	N	2009-01-28	2009-07-27	true	true	true	2
HA1	189	K	2009-01-28	2009-07-27	false	true	true	2
HA1	212	A	2009-03-29	2011-01-18	false	false	false	2
HA1	45	N	2010-03-24	2013-02-06	false	false	false	3
HA1	223	I	2010-12-19	2013-02-06	false	false	false	2
HA1	48	I	2011-03-19	2013-02-06	false	false	false	1
HA1	198	S	2011-03-19	2013-02-06	false	false	false	1
HA1	312	S	2009-08-26	2013-03-08	false	false	false	3
HA1	278	K	2011-06-17	2013-03-08	false	true	false	1
HA1	145	S	2011-04-18	2013-04-07	false	true	true	4
HA1	33	R	2011-06-17	2013-06-06	false	false	false	2
HA2	160	N	2012-07-11	2015-09-24	false	false	false	3
HA1	225	D	2013-08-05	2015-09-24	false	false	false	3
HA1	3	I	2013-08-05	2016-11-17	false	false	false	2
HA1	159	Y	2014-02-01	2016-11-17	false	true	true	2
HA1	160	T	2014-01-02	2017-07-15	false	true	false	2

Table S I The 30 trajectories that took place between year 2000 and year 2018 and resulted in fixation. Columns **Shih**, **Luksza** and **Koel** respectively indicate whether the position is found in the epitopes lists in (respectively) (Shih et al., 2007), (Luksza and Lässig, 2014) and (Koel et al., 2013). The **Tree counts** column indicates the number of times the mutation corresponding to the trajectory can be found in the phylogenetic tree. Note that a trajectory is only shown in the table if the sequenced population counts more than 10 strains at its time of fixation. This explains that only 30 trajectories are displayed, whereas more mutations did fix in this period of time.

Gene	Position	AA	Start date	End date	Fixation	Max. freq.
HA1	106	A	2001-02-09	2002-02-04	lost	1.0
HA1	144	D	2001-06-09	2002-02-04	fixed	1.0
HA1	105	H	2003-04-30	2003-10-27	lost	1.0
HA1	126	D	2003-04-30	2004-05-24	lost	1.0
HA1	140	Q	2004-01-25	2004-06-23	lost	0.31
HA1	226	I	2003-09-27	2004-09-21	fixed	1.0
HA1	173	E	2004-12-20	2006-03-15	lost	0.63
HA1	142	G	2006-06-13	2007-05-09	lost	0.71
HA1	144	D	2006-07-13	2007-05-09	lost	0.67
HA1	128	A	2006-09-11	2007-05-09	lost	0.25
HA1	157	S	2006-09-11	2007-05-09	lost	0.59
HA1	140	I	2006-11-10	2007-11-05	fixed	1.0
HA1	173	N	2007-12-05	2008-07-02	lost	0.3
HA1	157	S	2007-12-05	2008-09-30	lost	0.31
HA1	173	E	2006-06-13	2008-12-29	lost	0.67
HA1	173	Q	2007-07-08	2009-01-28	fixed	0.96
HA1	158	N	2009-01-28	2009-07-27	fixed	0.96
HA1	62	K	2009-01-28	2011-05-18	lost	0.73
HA1	144	K	2009-01-28	2011-05-18	lost	0.75
HA1	62	V	2011-04-18	2011-09-15	lost	0.34
HA1	157	S	2013-05-07	2015-09-24	lost	0.35
HA1	128	A	2012-08-10	2016-11-17	lost	0.81
HA1	197	K	2015-11-23	2016-11-17	lost	0.27
HA1	142	R	2018-05-11	2018-10-08	lost	0.38
HA1	142	G	2012-03-13		poly	0.86
HA1	144	S	2013-12-03		poly	0.96
HA1	121	K	2015-12-23		poly	0.82
HA1	142	K	2016-05-21		poly	0.77
HA1	62	G	2017-03-17		poly	0.75
HA1	128	A	2018-01-11		poly	0.56

Table S II Trajectories of mutations at epitope positions in (Shih et al., 2007) (*Shih et. al.*) that have been observed at least once above frequency 0.25. The **Fixation** column indicates whether the mutation has fixed, disappeared, or is still polymorphic as of October 2018. The **Max.freq.** column indicates the maximum frequency reached by the trajectory. A maximum frequency of 1 for mutations that finally disappear is explained by trajectories reaching frequency 1 for one time bin and going back to lower values for following ones (a frequency above 0.95 for two time bins in a row defines fixation).

Gene	Position	AA	Start date	End date	Fixation	Max. freq.
HA1	50	G	2001-02-09	2002-02-04	lost	1.0
HA1	144	D	2001-06-09	2002-02-04	fixed	1.0
HA1	126	D	2003-04-30	2004-05-24	lost	1.0
HA1	189	N	2003-07-29	2004-05-24	fixed	1.0
HA1	159	F	2003-08-28	2004-05-24	fixed	1.0
HA1	226	I	2003-09-27	2004-09-21	fixed	1.0
HA1	145	N	2003-12-26	2004-11-20	fixed	1.0
HA1	188	N	2004-07-23	2005-02-18	lost	0.36
HA1	227	P	2003-05-30	2005-04-19	fixed	1.0
HA1	173	E	2004-12-20	2006-03-15	lost	0.63
HA1	193	F	2004-12-20	2006-03-15	fixed	0.97
HA1	142	G	2006-06-13	2007-05-09	lost	0.71
HA1	144	D	2006-07-13	2007-05-09	lost	0.67
HA1	157	S	2006-09-11	2007-05-09	lost	0.59
HA1	50	E	2006-09-11	2007-06-08	fixed	0.95
HA1	173	N	2007-12-05	2008-07-02	lost	0.3
HA1	157	S	2007-12-05	2008-09-30	lost	0.31
HA1	173	E	2006-06-13	2008-12-29	lost	0.67
HA1	173	Q	2007-07-08	2009-01-28	fixed	0.96
HA1	158	N	2009-01-28	2009-07-27	fixed	0.96
HA1	189	K	2009-01-28	2009-07-27	fixed	0.96
HA1	213	A	2009-01-28	2010-02-22	lost	0.68
HA1	144	K	2009-01-28	2011-05-18	lost	0.75
HA1	53	N	2009-11-24	2013-02-06	lost	0.72
HA1	278	K	2011-06-17	2013-03-08	fixed	0.98
HA1	145	S	2011-04-18	2013-04-07	fixed	0.99
HA1	159	S	2013-11-03	2015-08-25	lost	0.46
HA1	157	S	2013-05-07	2015-09-24	lost	0.35
HA1	159	Y	2014-02-01	2016-11-17	fixed	0.97
HA1	159	S	2015-10-24	2016-11-17	lost	0.4
HA1	197	K	2015-11-23	2016-11-17	lost	0.27
HA1	160	T	2014-01-02	2017-07-15	fixed	0.96
HA1	142	R	2018-05-11	2018-10-08	lost	0.38
HA1	135	N	2018-06-10	2018-10-08	lost	0.38
HA1	142	G	2012-03-13		poly	0.86
HA1	144	S	2013-12-03		poly	0.96
HA1	121	K	2015-12-23		poly	0.82
HA1	142	K	2016-05-21		poly	0.77
HA1	131	K	2016-09-18		poly	0.77
HA1	135	K	2016-11-17		poly	0.47

Table S III Same as table SII, for (Luksza and Lässig, 2014) (*Luksza et. al.*).

Gene	Position	AA	Start date	End date	Fixation	Max. freq.
HA1	189	N	2003-07-29	2004-05-24	fixed	1.0
HA1	159	F	2003-08-28	2004-05-24	fixed	1.0
HA1	145	N	2003-12-26	2004-11-20	fixed	1.0
HA1	193	F	2004-12-20	2006-03-15	fixed	0.97
HA1	158	N	2009-01-28	2009-07-27	fixed	0.96
HA1	189	K	2009-01-28	2009-07-27	fixed	0.96
HA1	145	S	2011-04-18	2013-04-07	fixed	0.99
HA1	159	S	2013-11-03	2015-08-25	lost	0.46
HA1	159	Y	2014-02-01	2016-11-17	fixed	0.97
HA1	159	S	2015-10-24	2016-11-17	lost	0.4

Table S IV Same as table SII, for (Koel et al., 2013) (*Koel et. al.*).

1154 9. Supplementary figures




Figure S 9 Number of A/H3N2 HA sequences per year from year 1990.




Figure S 10 Number of H3N2 HA and NA sequences per month from year 2000.




Figure S 11 Frequency trajectories for the 9 most entropic positions in the A/H3N2 HA protein.




Figure S 12 Equivalent to panel B of figure 1 of the main text for A/H3N2, with f_0 equal to 0.5 in **A** (76 trajectories), and 0.7 in **B** (63 trajectories).




Figure S 13 Equivalent to panels **B** and **C** of figure 1 of the main text for A/H1N1pdm influenza. 89 trajectories are shown and participate to the mean (thick black line).




Figure S 14 Equivalent to panel **B** of figure 1 of the main text for A/H1N1pdm, with f_0 equal 0.5 in **A** (50 trajectories), and 0.7 in **B** (41 trajectories).




Figure S 15 Equivalent of figure 3 of the main text for the HA gene of A/H1N1pdm influenza. Fixation probability $P_{fix}(f)$ as a function of frequency. **A:** Mutation with higher or lower LBI values, based on their position with respect to the median LBI value. **B:** Different lists of epitope positions in the HA protein. The authors and the number of positions is indicated in the legend. **C:** Mutations for binary positions, *i.e.* positions for which we never see more than two amino acids in the same time bin. **D:** Mutations that appear once or more than once in the tree for a given time bin.





Figure S 17 Based on A/H1N1pdm HA and NA. **A:** Mutations with a higher or lower geographical spread, based on the median value of the score used (see Methods). Note: the words *local* and *global* only reflect the position of the geographic spread of the mutation relative to the median value computed for all mutations found at this frequency. As this median value may change with the considered frequency bin, so does the definition of local and global mutations. **B:** Mutations whose trajectories are older or more recent, based on the median age of trajectories when reaching the considered frequency f .




Figure S 18 Geographic spread of mutations as a function of the time for which they have been present in the population above a frequency of 5%. Points represent individual mutations and for a population in a given time bin. The line is the average of dots for a given value on the x -axis. Based on data for A/H3N2 HA.




Figure S 19 Earth mover's distance to the future population for different predictors. A present population consists of all A/H3N2 HA sequences sampled in a 4 months time window. Quantities are averaged over all possible "present" populations from the year 2002. Predictors are: **Global consensus**: Consensus sequence of the present population. Best long-term predictor for a structure-less neutrally evolving population. **All present population**: All sequences in the present population. Perfect predictor if the population does not change at all through time. **Cluster-wise consensus**: Consensus sequence for each cluster in the present population. Clusters are based on local maxima of the LBI. Sequences are assigned to a given cluster based on their tree branch-length distance to the corresponding local maximum.




Figure S 20 Fixation probability as a function of frequency for the simulations discussed in the main text. **Top:** Simulation without antibodies. The three colored curves reflect different rate of change for the fitness landscape. Visual inspection of the frequency trajectories indicates a typical sweep time of ~ 400 generations. **Bottom:** Simulation with antibodies. The different colored curves indicate the rate at which antibodies are introduced.




Figure S 21 Fixation probability as a function of frequency for the simulations discussed in the main text, with trajectories stratified according to real fitness values. “High” and “low” fitness classes are defined with respect to the median value. **Top:** Simulation without antibodies and with changes to the fitness landscape every $dt = 10$ generations. **Bottom:** Simulation with antibodies, with a new antibody every $dt = 10$ generations.