

Limited predictability of amino acid substitutions in seasonal influenza viruses

Pierre Barrat-Charlaix,^{1,2} John Huddleston,³ Trevor Bedford,⁴ and Richard A. Neher^{1,2,*}

¹*Biozentrum, Universität Basel, Switzerland*

²*Swiss Institute of Bioinformatics, Basel, Switzerland*

³*Molecular Cell Biology, University of Washington, Seattle, WA, USA*

⁴*Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Research Center, Seattle, WA, USA*

(Dated:)

Seasonal influenza viruses repeatedly infect humans in part because they rapidly change their antigenic properties and evade host immune responses, necessitating frequent updates of the vaccine composition. Accurate predictions of strains circulating in the future could therefore improve the vaccine match. Here, we studied the predictability of frequency dynamics and fixation of amino acid substitutions. Current frequency was the strongest predictor of eventual fixation, as expected in neutral evolution. Other properties, such as occurrence in previously characterized epitopes or high *Local Branching Index* (LBI) had little predictive power. Parallel evolution was found to be moderately predictive of fixation. While the LBI had little power to predict frequency dynamics, it was still successful at picking strains representative of future populations. The latter is due to a tendency of the LBI to be high for consensus-like sequences that are closer to the future than the average sequence. These results are in contrast to simulations of models of adapting populations, where we find the clear signals of predictability. This indicates that the evolution of influenza HA and NA, while driven by strong selection pressure to change, is poorly described by common models of directional selection such as travelling fitness waves.

INTRODUCTION

Seasonal influenza A viruses (IAV) infect about 10% of the global population every year, resulting in hundreds of thousands of deaths [1, 2]. Vaccination is the primary measure to reduce influenza morbidity. However, the surface proteins hemagglutinin (HA) and neuraminidase (NA) continuously accumulate mutations at a high rate, leading to frequent antigenic changes [2–5]. While a vaccine targeting a particular strain may be efficient for some time, antigenic drift will sooner or later render it obsolete. The World Health Organization (WHO) regularly updates influenza vaccine recommendations to best match the circulating strains. Since developing, manufacturing, and distributing the vaccine takes many months, forecasting the evolution of influenza is of essential interest to public health [6, 7].

The number of available high quality HA and NA sequences has increased rapidly over the last 20 years [8, 9] and virus evolution and dynamics can be now be tracked at high temporal and spatial resolution [10]. This wealth of data has given rise to an active field of predicting influenza virus evolution [6, 7]. These models predict the future population of influenza viruses by estimating strain fitness or proxies of fitness. Luksza and Lässig [11], for example, train a fitness model to capture antigenic drift and protein stability on patterns of epitope and non-epitope mutations. Other approaches by Steinbrück et al. [12], Neher et al. [13] predict fitness by using hemagglutination inhibition (HI) data to determine possible

antigenic drift of clades in the genealogy of the HA protein. Finally, Neher et al. [14] use branching patterns of HA phylogenies as a proxy for fitness. These branching patterns are summarized by the Local Branching Index (LBI), which was shown to be a proxy of relative fitness in mathematical models of rapidly adapting populations [14].

The underlying assumption of all these methods is that (i) differences in growth rate between strains can be estimated from sequence or antigenic data and (ii) that these growth rate differences persist for long enough to be predictive of future success. Specific positions in surface proteins are of particular interest in this context. The surface proteins are under a strong positive selection and change their amino acid sequence much more rapidly than other IAV proteins or than expected under neutral evolution [4, 15]. Epitope positions, i.e., positions targeted by human antibodies, are expected to change particularly often since viruses with altered epitopes can evade existing immune responses [3, 5, 16]. It therefore seems plausible that mutations at these positions have a tendency to increase fitness and a higher probability of fixation [15]. But one has to be careful to account for the fact that these positions are often ascertained post-hoc [3] and human immune responses are diverse with substantial inter-individual variation [17].

In this work, we use HA and NA sequences of A/H3N2 and A/H1N1pdm influenza from year 2000 to 2019 to perform a retrospective analysis of frequency trajectories of amino acid mutations. We quantify how rapidly mutations at different frequencies are lost or fixed and how rapidly they spread through the population. We further investigate whether any properties or statistics are predictive of whether a particular mutation fixes or not. To our surprise, we find that the predictability of

* Correspondence to: Richard Neher, Biozentrum, Klingelbergstrasse 70, 4056, Basel, Switzerland.
richard.neher@unibas.ch

these trajectories is very limited: The probability that a mutation fixes differs little from its current frequency, as would be expected if fixation happened purely by chance. This observation holds for many different categories of mutations, including mutations at epitope positions. This weak predictability is not attributable solely to clonal interference and genetic linkage, as simulation of models including even strong interference retain clear signatures of predictability. Consistent with these observations, we show that a simple predictor uninformed by fitness, the consensus sequence, performs as well as the Local Branching Index (LBI), the growth measure based on the genealogy used in [14]. This suggests that although LBI has predictive power, the reason for its success may not be related to it approximating fitness of strains.

RESULTS

The main underlying question asked in this work is the following: given a mutation X in the genome of influenza that we observe at a frequency f in the population at a given date, what can we say about the future of X ? There are different ways to make this question more specific. First, one can try to quantitatively predict the frequency of X at future times $f(t)$. In other words, having observed a mutation at frequencies (f_1, f_2, \dots, f_n) at dates (t_1, t_2, \dots, t_n) , what can we say about its frequency at future dates $(t_{n+1}, t_{n+2}, \dots)$? A simpler, more qualitative question, is to ask whether X will fix in the population, will disappear, or whether the site will stay polymorphic.

In this work, we use amino-acid sequences of the HA and NA genes of A/H3N2 (since the year 2000) and A/H1N1pdm (since the year 2009) influenza available in GISAID [9] (see supplementary materials for an acknowledgment of all data contributors). This amounts to 44 976 HA and 36 300 NA sequences for A/H3N2 and 45 350 HA and 40 412 NA sequences for A/H1N1, with a minimum of 100 per year. These sequences are binned in non-overlapping intervals of one month. Each single-month time bin and the sequences that it contains represent a (noisy) snapshot of the influenza population at a given date. The number of sequences per time bin varies strongly both with year and according to the season, with earlier time bins containing around 10 sequences while more recent bins contain several hundreds (see figures S5 and S6 in SM for details).

The central quantities that we derived from this data are *frequency trajectories* of amino acids at each position in the sequences. If an amino acid X_i is found at position i at a frequency between 5% and 95% in the population of a given time bin t , then the population is considered polymorphic at position i and at time t . This polymorphism is characterized by the frequency of X_i , $f_{X_i}(t)$, and also by frequencies of other amino acids at i . The series of values $f_{X_i}(t)$ for contiguous time bins constitutes the frequency trajectory of X_i . A trajectory is terminated if the corresponding frequency is measured above 95%

(resp. below 5%) for two time bins in a row, in which case amino acid X_i is considered as *fixed* (resp. *absent*) in the population. Otherwise, the trajectory is considered *active*. Examples of trajectories can be seen in figure S7 of the Supplement.

In the rest of this work, we will focus on frequency trajectories that are starting at a zero (low) frequency, i.e. $f(t=0) = 0$. These represent new amino acid variants which were absent in the population at the time bin when the trajectory started and are currently rising in the population (see Methods). Such distinction in novel and ancestral variants is necessary to meaningfully interrogate predictability. Each rising trajectory of a new mutation implies the existence of another decreasing one at the same position, since frequencies of all amino acids at a given position must sum to one. If novel variants arise by selection, we expect to see a stronger signal of selection after conditioning on these novel variants. In classic models of population genetics, strongly advantageous variants undergo rapid selective *sweeps*, i.e., the rapid rise and fixation. If such sweeps are common in the evolution of HA and NA, the restriction to trajectories that start at low frequency, should enrich for mutations that are positively selected and on their way to fixation.

Predicting future frequencies

Having observed the frequency trajectory $f(t)$ of a mutation until a given date t_0 , how much can we say about the future values of f after t_0 ? We consider the idealized case sketched in panel **A** of figure 1: given the trajectory of a *new* mutation, i.e. that started at a frequency of 0, and that we observe at frequency f_0 at time t_0 , what is the probability $P_{\Delta t}(f)$ of observing it at a value f at time $t_0 + \Delta t$?

To answer this question retrospectively, we use all frequency trajectories extracted from HA and NA sequences that satisfy these conditions for a given f_0 . The number of trajectories is limited and the frequency estimates themselves are based on a finite sample and are hence imprecise. Therefore, we consider trajectories in an interval $[f_0 - \delta f, f_0 + \delta f]$ with $\delta f = 0.05$.

For $f_0 = 0.3$, we found 120 such trajectories in the case of A/H3N2 influenza, represented on the panel **B** of figure 1, where time is shifted such that $t_0 = 0$. The same analysis was performed for A/H1N1pdm, with the 89 found trajectories displayed in figure S9. Some trajectories fall in the frequency bin around f_0 while decreasing, even though they crossed that bin at an earlier time. This is due to the fact that some trajectories “skipped” the interval f_0 in question on their initial rise due to sparse sampling. These trajectories are nevertheless rising in the sense that they start at frequency 0 for $t \rightarrow -\infty$. Removing them does not change results significantly.

Since rapid sequence evolution of influenza HA and NA mediates immune evasion, one could expect that a significant fraction of new amino acid mutations on

rising trajectories in figure 1 are *adaptive*. We could thus expect that most of these trajectories continue to rise after reaching frequency f_0 , at least for some time. A fraction of those would then sweep through the population and fix.

To quantify the extent to which this preconception of sweeping adaptive mutations is true, we estimated the probability distribution $P_{\Delta t}(f|f_0)$ of finding a trajectory at frequency f after a time Δt given that it was observed at f_0 at time 0. The results for different Δt are shown in figure 1C. Initially, *i.e.* at time $t_0 = 0$, this distribution is by construction peaked around f_0 . If a large fraction of the trajectories keep increasing after this time, we should see the “mass” of $P_{\Delta t}(f|f_0)$ move to the right towards higher frequencies as time progresses.

However, future distributions for $\Delta t > 0$ do not seem to follow a pattern compatible with selective sweeps. The thick black line in Figure 1B shows the average frequency of all trajectories. This average makes a sharp turn at $t = 0$ and is essentially flat for $t > 0$ in the case of A/H3N2, and slightly increasing for A/H1N1pdm (see supplement). Hence, the fact that this average rose for $t < 0$ gives little information for $t > 0$, and is due to the conditions by which these trajectories were selected. This shows that sweep-like trajectories rising steadily from frequency 0 to 1 are not common enough to dominate the average trajectory.

Consistent with the average, the frequency distribution of the selected trajectories broadens in time without significant shift of the mean as time passes. After 60 days the distribution is rather symmetrical around the initial $f_0 = 0.3$ value, suggesting that the knowledge that the trajectories were rising is lost after two months. On a timescale of 60 to 120 days, the only possible prediction is that trajectories are likely to be found in a broad interval around the initial frequency f_0 . After one year the distribution becomes almost flat (excluding mutations that have disappeared or fixed), and the initial peak at f_0 is not visible anymore. The only information remaining from the initial frequency is the fraction that fixed or was lost (see below). This behavior is expected in neutral models of evolution [18] but incompatible with a dynamic dominated by sweeps taking over the population.

While this observation does not rule out that signatures exist that predict future frequency dynamics, past dynamics alone is weakly informative.

a function of the time elapsed since they were first seen above 25% frequency. Most mutations are either lost or become fixed after 2-3 years, with very few trajectories remaining active after 5 years. This time scale of 2-3 years is consistent with the typical coalescence time observed in phylogenetic trees of A/H3N2 influenza [10, 19]. We also note that the fraction of lost trajectories increases sharply at small times with 40% of mutations observed above 25% frequency being lost within one year for A/H3N2, while it takes longer to fix a mutation in the whole population.

We then examined the probability of mutations to fix in the population as a function of the frequency at which they are seen. For different values of frequency f , we consider all trajectories that started at a null frequency and are seen in the interval $[f - 7.5\%, f + 7.5\%]$ at any given time. The probability of a mutation fixing given that it is seen at frequency f , $P_{fix}(f)$, is then estimated by the fraction of those trajectories which terminate at a frequency larger than 95%, *i.e.* our fixation threshold. Panels C and D of figure 2 show $P_{fix}(f)$ as a function of f for NA and HA. For both proteins, the probability of fixation of a new mutation at frequency f is close to f itself, that is $P_{fix}(f) \simeq f$. This result is exactly what is expected in a population evolving in the *absence* of selection. A mutation or trait appearing at frequency f is shared by $f \cdot N$ individuals, and the probability for one of them to become the ancestor of all the future population is $f \cdot N/N = f$. Thus, the probability of this mutation or trait to fix in the population is equal to its current frequency, a case which we will refer to as the neutral expectation. Panel C of figure 2 indicates that mutations in the surface proteins of A/H3N2 influenza are in good agreement with the neutral expectation, while those in A/H1N1pdm show only small deviations from it. In both cases, the probability of fixation seems to be mainly dictated by the current frequency f at which the mutation is observed.

This dynamics is in apparent contradiction with evidence that influenza surface proteins are under strong selective pressure to evade human immune responses [4]. If strong selection was present, we would expect rising amino acid mutations to fix at a distinctively higher frequency than the one at which they are measured. In an extreme case where most trajectories would be clean sweeps, $P_{fix}(f)$ should be close to 1 for all but very small values of f .

Next, we searched for features of mutations that allow prediction of fixation beyond frequency by dividing frequencies into categories that deviate from the diagonal in panels C and D of figure 2. We first turn to the *Local Branching Index* (LBI), a quantity calculated for each node in a phylogenetic tree that indicates how dense the branching of the tree is around that node. LBI has previously been successfully used as a predictor of the future population of influenza [14], and was shown to be a proxy for fitness of leaves or ancestral nodes in mathematical models of evolution. Here, we define the LBI of a mutation at date t as the average LBI of strains

Prediction of fixation or loss

Instead of predicting future frequency, let's consider the long-term goal of predicting the probability that a mutation fixes in the population. We first estimate the fraction of frequency trajectories that either fix in the population or are lost, as well as the time it takes for one or the other to happen. Panels A and B of figure 2 show the fraction of frequency trajectories in HA and NA that either have fixed, were lost or remained active as




FIG. 1. **A:** Sketch of the idea behind the short term prediction of frequency trajectories. Given a mutation that we have seen increasing in frequency and that we “catch” at frequency f_0 at time t_0 , what can we say about the distribution of future frequencies $P_{\Delta t}(f|f_0)$? **B:** All frequency trajectories of amino acid mutations in the A/H3N2 HA and NA genes that were absent in the past, are seen around $f_0 = 30\%$ frequency at time $t_0 = 0$, and are based on more than 10 sequences at each time point. Red curves represent mutations that will ultimately fix, blue the ones that will be lost, and black the ones for which we do not know the final status. Dashed horizontal lines (blue and red) represent loss and fixation thresholds. The thick black line is the average of all trajectories, counting those that fix (resp. disappear) as being at frequency 1 (resp. 0). Figure S8 shows equivalent figures for other values of f_0 . **C:** Distribution of future frequencies $P_{\Delta t}(f|f_0)$ for the trajectories shown in panel **B** and for specific values of Δt .

that carry this mutation and that were sampled in the³¹⁶
time bin corresponding to t . Panel A of figure 3 shows³¹⁷
fixation probability for HA mutations with LBI in the³¹⁸
top or bottom half of the distribution. Both groups have³¹⁹
identical probability of fixation, suggesting that LBI car-³²⁰
ries very little information on the probability of fixation³²¹
of a mutation.³²²

Next, we focused on previously reported antigenic sites in the A/H3N2 HA protein, referred to as *epitope* positions. Mutations at these position might mediate immune escape and are therefore likely under strong selection and show sweep-like behavior. We used four lists of relevant epitope positions from different sources comprising from 7 to 129 positions in the sequence of the HA1 pro-




FIG. 2. **A:** Activity of all rising frequency trajectories seen above 25% frequency for A/H3N2 HA and NA. **B:** Same as **A** for A/H1N1. **C:** Probability of fixation of a mutation (amino acid or synonymous) $P_{fix}(f)$ as a function of the frequency f at which it is measured, for A/H3N2 HA and NA. Only new mutations are considered, *i.e.* mutations that were absent in the past. The diagonal dashed line is the expectation from a neutrally evolving population. Colored dashed lines represent synonymous mutations. Colored solid lines represent amino acid mutations. Error bars represent a 95% confidence interval. **D:** Same as **C** for A/H1N1.

tein [3, 5, 11, 16]. Panel Fig. 3B shows fixation probability³⁴⁰ as a function of frequency for the four lists of epitopes³⁴¹. Only mutations at the 7 epitope sites reported in [5] have³⁴² higher chances of fixation than expected by chance. No³⁴³ clear difference is found for the lists by Luksza and Lässig³⁴⁴ [11], Wolf et al. [16], while positions from Shih et al. [3]³⁴⁵ show lower chances of fixation. One should also note that³⁴⁶ many of these positions were determined post-hoc and³⁴⁷ might be enriched for positions that experienced rapid³⁴⁸ substitutions before the publication of the respective stud-³⁴⁹ies.³⁵⁰

Two ways of categorising mutations, however, suggest³⁵¹ some power to predict fixation. In panel Fig. 3C, we split³⁵² trajectories into those occurring at binary positions where³⁵³ only two amino acid variants co-circulate and non-binary³⁵⁴ positions with more than two variants. Novel variants³⁵⁵ at non-binary positions, *i.e.* ones for which competition³⁵⁶

between three amino acids or more has occurred at least once, have a higher chance of fixation. In panel D, we separate mutations that appear more than once or only once in the reconstructed tree (see methods), and found that the former fix more often. Panels C and D show that it is possible to gain some information on the chance of fixation of a particular mutation, as was done in panel B. However, the predictive power remains small, with the “top” curves in panels C&D being very close to the diagonal.

We conduct the same analysis on A/H1N1pdm influenza, with results shown in figure S11. Results are qualitatively similar to those obtained for A/H3N2, with LBI giving little information and mutations at non-binary positions having a higher chance of fixation. Panel D differs between figures 3 and S11, with convergent evolution giving less information on fixation in the latter

case. However, this could be due to the shorter time⁴¹² period over which A/H1N1pdm evolved, resulting in a⁴¹³ shorter tree and less possibilities of convergent evolution⁴¹⁴. Indeed, error bars for mutations appearing multiple times⁴¹⁵ in **D** of figure S11 are relatively large, indicating a lower amount of trajectories.⁴¹⁶

Since influenza is seasonal in temperate regions, geographic spread and persistence might be predictive of the success of mutations. We quantify geographic spread of a mutation by the entropy of its frequency distribution across regions (see methods) and its persistence by the age of the trajectory by the time it reaches frequency f . Figures S12 and S13 show the fixation probabilities as a function of observed frequency for mutations classified according to these scores. The two scores also allow a quantitatively moderate distinction between mutations for a given frequency f , mutations found in many regions or those that are older (in the sense that they have taken more time to reach frequency f) tend to fix more often than geographically localized mutations or more recent ones, but the effect is small. These two scores are in fact correlated, with older trajectories representing mutations that are more geographically spread, as can be seen in figure S14 of SM. However, it is important to note that sampling biases and heterogeneity across time and space (see supplementary figures S5 and S6) make answering such specific hypothesis challenging. Frequency of mutations might thus be amplified through different sampling biases, making the connection between geographic spread, seasonality and mutation frequency non-trivial to measure.⁴⁴¹

Simulations of models of adaptation

The results shown in figures 2 and 3 are difficult to reconcile with the idea that seasonal influenza virus evolution is driven by rapid directed positive selection. One possible explanation for the weakly predictable behaviour of mutations (beyond their current frequency) might be tight genetic linkage inside each segment and strong competition between different adaptive mutations [15, 20]. We design a simple model of population evolution based on the `ffpopsim` simulation software to test this hypothesis [21]. The model represents a population of binary genomes of length $L = 200$ evolving in a fitness landscape that changes through time.⁴⁴⁶

First, we use an additive fitness function, with sequence $(x_1 \dots x_L)$ having a fitness $\sum_i h_i x_i$. This implies that for a given genome position i , the trait $x_i = 1$ is favored if $h_i > 0$ whereas $x_i = -1$ is favored if $h_i < 0$. All h_i 's have the same magnitude, and only their signs matter. Every Δt generations, we randomly choose a position i and flip the sign of h_i , effectively changing the fitness landscape. Individuals in the population now have the opportunity to make an adaptive mutation at site i giving them a fitness advantage $2|h|$. A “flip” at position i of the

fitness landscape will decrease fitness of all individuals that carried the adapted variant at position i and increases the fitness of those that happened to carry a deleterious variant.

To increase competition between genomes, we designed a second model that includes epistasis. Once again, the baseline fitness of a genome is an additive function, this time with values of h_i that do not change through time. In addition, we added a component that mimics immune selection. Every Δt generation, we now introduce “antibodies” that target a specific sub-sequence of length $l = 5$, noted $(x_{i_1}^{ab}, \dots, x_{i_l}^{ab})$. The positions $(i_1 \dots i_l)$ are chosen at random, while the targeted sub-sequence is the dominant state at each position. Genomes that include the *exact* sub-sequence targeted by the antibody suffer a strong fitness penalty. However, a single mutation away from that sub-sequence removes this penalty completely, resulting in a fitness landscape with very strong epistasis. This has the effect of triggering a strong competition between adaptive mutations: for a given antibody, $l = 5$ possible mutations are now adaptive, but combinations of these mutations do not bring any fitness advantage.

Having simulated populations in these two fitness landscapes, we perform the same analysis of frequency trajectories as for the real influenza data. Figure S16 of the SM shows the $P_{fix}(f)$ as a function of f for the two models and for different values of the inverse rate of change Δt of the fitness landscape. For all models, this curve deviates significantly from the diagonal. This is most evident for the case of a simple additive fitness landscape that changes rarely $\Delta t = 1000$: rising mutations almost always fix in the population, with $P_{fix}(f) \simeq 1$ for any f larger than a few percent. This is corroborated by visual inspection of the trajectories, which shows that evolution in this regime is driven by regular selective sweeps that take a typical time of ~ 400 generations. In other regimes, with smaller Δt or with strong epistatic competition, $P_{fix}(f)$ is reduced and closer to the diagonal. However, it takes an extremely fast changing fitness landscape to push P_{fix} close to the diagonal: with $\Delta t = 10$, that is about 40 changes to the fitness landscape in the time it would take a selective sweep to go from 0% to fixation, $P_{fix}(f)$ differs from f in a way that is comparable to what is observed in A/H1N1pdm influenza.

These models are not meant to be accurate models of influenza viruses evolution. But figure S16 does show is that the patterns observed in influenza virus evolution are only reproduced by models of adapting populations when pushing clonal competition to extreme values. We conclude that the pattern in figure 2 may not be a straightforward manifestation of genetic linkage and clonal interference, but that some more intricate interplay of epidemiology, seasonality, human immunity and chance gives rise to the weakly predictable yet strongly selected evolutionary dynamics of IAVs.




FIG. 3. Fixation probability $P_{fix}(f)$ as a function of frequency, for A/H3N2 influenza. Figure S11 shows the same analysis for A/H1N1. **A:** HA mutations with higher or lower LBI values, based on their position with respect to the median LBI value. **B:** Different lists of epitope positions in the HA protein. The authors and the number of positions is indicated in the legend. **C:** HA and NA mutations for binary positions, *i.e.* positions for which we never see more than two amino acids in the same time bin. **D:** HA and NA mutations that appear once or more than once in the tree for a given time bin.

Why do predictions work?

The statistics of frequency trajectories seem to be in conflict with the notion that influenza evolution is predictable. Likewise, the LBI, a quantity that correlates with fitness in mathematical models and is used to predict future influenza populations [14], does not seem to contain any information on whether a specific mutation is going to fix or not, see figure 3. To resolve this conundrum, we first note that the criterion by which predictive power for influenza was measured in [14] was the distance between the strain with the highest LBI and the future population, not the ability of the LBI to predict dynamics. The distance was compared to the average distance between the present and future population, as well as the post-hoc optimal representative and the future.

To quantify the ability of the LBI and other measures to pick good representatives of the future, we construct a large tree of HA sequences with 100 sequences in non-overlapping time bins of 4 months from year 2003 to 2019.

(a total of 4402 as some 4 month intervals contain less than 100 sequences). Each time bin is considered as a snapshot of the A/H3N2 influenza population and we will refer to sequences in time bin t as the population of the *present*. From this present population, we predict *future* populations in time bin $t + \Delta t$, using only sequences in time bin t and before.

To assess the ability of the LBI to pick a close representative of the future, we compute the LBI of each node of one time bin in the tree using only the leaves that belong to that time bin. The top panel in figure 4 shows the hamming distance of the strain with the highest LBI to future populations at different Δt along with the same distance for a randomly chosen strain. The figure shows the distance averaged over all possible values of t for Δt between 0 and 32 months, giving us an average efficiency of a predictor over 16 years of influenza evolution.

The strain with the highest LBI is consistently closer to the future than the average strain by about 1-2 amino acids, while the overall distance increases linearly due to the continuous evolution of the population. We hence

reproduce previous results showing that the LBI picks closer than average representatives [14]. To investigate whether this apparent success is due to the ability of the LBI to predict fitness or not, we explored a different predictor: the amino acid consensus sequence of the present population (see Methods for a definition of the consensus sequence). The choice is motivated by the fact that it can be shown to be the best possible long term predictor for a neutrally evolving population in terms of Hamming distance (see SM section 1). Figure 4 shows that the consensus sequence is in fact a equally good or even slightly better representative of the future than the sequence with highest LBI (note that the consensus sequence does not necessarily exist in the population).

This near equivalence of the consensus and the strain with highest LBI can be explained as follows: The LBI tends to be high for nodes in a tree that are close to the root of a dense and large clade. A typical sample of influenza HA sequences fall into a small number of recognizable clades, and the strains with maximal LBI will often be close to the root of the largest of those clades. This root of the largest clade will often be close to the consensus of the whole population, explaining the similar distance patterns. To test that hypothesis, we measure the hamming distance from the sequence of the top LBI strain to the consensus sequence for populations of all time bins. Panel B of figure 4 shows these distances, scaled with respect to an average strain (details in caption). It clearly shows that the top-LBI strain and the consensus sequence are indeed quite similar: out of 48 time bins, only once is the sequence of the top-LBI strain farther away from the consensus than the average sequence is. Moreover, the sequence of the top-LBI strain exactly matches the consensus in 19 cases.

DISCUSSION

Predicting the trajectory of a mutation requires significant fitness difference between genomes carrying different variants at the site and a selection pressure that changes slowly over time. Under such conditions, it is expected that frequency trajectories will show a persistent behavior which would make them predictable for some time. However, we could find only limited evidence for such persistent behavior in the past 19 years of IAV evolution. This lead us to conclude that (i) influenza virus evolution is qualitatively different from models of rapidly adapting population (despite clear evidence for frequent positive selection), and (ii) previous methods to predict influenza evolution work primarily because they pick strains that represent the future well, not because they predict future dynamics.

The primary focus in this work was the investigation of frequency trajectories of new amino acid mutations. In the short term, we found that on average the direction of trajectory does not persist for longer than a few months. Indeed, the average trajectory in figure 1 takes a sharp

turn when going from $t < 0$ to $t > 0$, instead of showing “inertia”. This suggests that selective sweeps are not representative of typical trajectories.

On a longer timescale, we investigated the probability that a novel mutation observed at frequency f fixes. In neutral models of evolution this probability equals f , while it should be higher or lower than f for mutations with a beneficial or deleterious effect on fitness, respectively. However, in the case of influenza, this probability differs little from f , making current frequency the best predictor for fixation. In figure 3, we split trajectories into groups for which we expected P_{fix} to deviate from f . Many of these splits, such as high/low LBI or epitope/non-epitope positions, did not result in an increased predictability, while others gave limited information on fixation. Despite the lack of predictability of mutation frequency trajectories, influenza surface proteins show strong signatures of selection [4, 15].

Methods for predicting the future evolution of influenza either construct explicit fitness models [11?], use historical patterns of evolution [11, 22], phenotypic assays [13, 23], or dynamic or phylogenetic patterns [14, 24]. The goal of these methods is to pick strains that are good representatives of future populations and could serve as vaccine candidates [6].

The low power to predict frequency dynamics or fixation naturally triggers the question why the above methods have been found to work. Picking representatives of the future and predicting frequency dynamics are distinct objectives and success at the former (as compared to random picks) is not necessarily inconsistent with a lack of predictable dynamics. In fact, [?] reports that the rate at which the frequency of a strain changes is often a poor predictor – consistent with our observations here. But despite the fact that future frequencies are not predicted by the LBI, the strain with the highest LBI in the population is a better predictor of the future population than a randomly picked one. While the LBI was shown to be a correlate of relative fitness and be predictive of fixation in mathematical models of evolution [14], it does not seem to be predict influenza evolution because it measures fitness from genealogical structure. Instead, we believe it picks closer than average strains simply because it has the tendency to be maximal at the base of large and dense clades. These basal genotypes are closer to the future populations than the current tips of the tree and hence a better predictor on average. The consensus sequence of all present strains performs slightly but consistently better than picking the strain with the highest LBI. The consensus sequence is the best possible predictor for a neutrally evolving population, and does not attempt to model fitness in any way.

At the same time, influenza virus phylogenies show clear deviations from those expected from the neutral Kingman coalescent, similar to those expected under Bolthausen-Sznitman coalescent (BSC) processes that are generated by traveling wave models of rapid evolution [25, 26]. The correspondence between the BSC and traveling wave mod-




FIG. 4. **A:** Average Hamming distance of the sequences of different predictors to HA sequences of future influenza populations, themselves averaged over all “present” populations from year 2003 to 2019. Predictors are: a randomly picked sequence in the present population; the sequence of the strain with the highest LBI in the present population; the consensus sequence of the present population. **B:** Scaled Hamming distance between the sequence of the top LBI strain and the consensus sequence for populations at different dates. The scaling is such that for each date, the Hamming distance between a strain from the population and the consensus is on average 1. The strain with the highest LBI is almost always closer to the consensus sequence than the average strain.

els comes from transient exponential amplification of fit₆₃₉ strains before these fitness differences are wiped out by₆₄₀ further mutation. This exponential amplification gen-₆₄₁ erates long-tailed effective offspring distributions which₆₄₂ in turn can leads to genealogies described by the BSC₆₄₃ [25, 27]. Many processes other than selection, includ-₆₄₄ ing seasonality and spatio-temporal heterogeneity, can generate effective long tailed offspring distributions even in absence of bona-fide fitness differences, which might₆₄₅ explain ladder-like non-Kingman phylogenetic trees.

A recent preprint proposed that influenza virus evo-₆₄₆ lution is primarily limited by an asynchrony between population level selection and generation of new variants₆₄₇ within infected hosts [28]. Along these lines, it is possi-₆₄₈ ble that the A/H3N2 population readily responds once₆₄₉ population level selection is high enough by giving rise₆₅₀ to essentially equivalent variants. Furthermore, selection₆₅₁ might cause the rapid rise of a novel variant to macro-₆₅₂ scopic frequencies (observable in a global sample) but₆₅₃

its benefit rapidly “expires” because competing variants catch up and/or it mediates immune escape only to a small fraction of the population. These considerations might explain the disconnect between models of rapid adaptation and the frequency dynamics observed in influenza virus populations.

METHODS

Data and code availability

The sequences used are obtained from the GISAID database [9]. Strain names and accession numbers are given as tables in two supplementary files. Outliers strains listed at <https://github.com/nextstrain/seasonal-flu/tree/081be2f487182252627418491be36cd01d645954/config> were removed.

654 The code used to generate the figures presented here⁷⁰⁶
 655 is available at <https://github.com/PierreBarrat/FluPredictability>.
 656

657 Frequency trajectories

658 For a set of sequences in a given time bin, we compute frequencies of amino acids at each position by simple counting.⁷¹⁴
 659 We make the choice of not applying any smoothing method in an attempt to be as close to the data and “model-less” as possible.⁷¹⁵
 660 This is especially important for the short term prediction of frequency trajectories, as estimations of the “persistence time” of a trajectory might
 661 be biased by a smoothing method.⁷¹⁶

662 We compute frequency trajectories based on the frequencies of amino acids. A trajectory begins at time t if an amino acid is seen under the lower frequency threshold of 5% (resp. above the higher threshold of 95%) for the two time bins preceding t , and above this lower threshold (resp. below the higher threshold) for time bin t . It ends in the reciprocal situation, that is when the frequency is measured below the lower threshold (resp. above the higher threshold) for two time bins in a row.

663 In order to avoid estimates of frequencies that are too noisy, we only keep trajectories that are based on a population of at least 10 sequences for *each* time bin. As said in the Results section, we also restrict the analysis to trajectories that begin at a 0 frequency, in part to avoid double counting. We find a total of 460 such trajectories. However, only 106 reach a frequency of 20%, on which figure 2 is based for instance.

664 Note that the fact that we use samples of relatively small sizes – at least for some time bins – leads to biases in the estimation of frequencies. We show in Supplementary Material that these biases are generally small and do not induce any qualitative changes to results presented here.⁷²⁴

708 τ equal to a tenth of $T_C \simeq 6$ years, the coalescence time for influenza A/H3N2 strains, converted to units of tree branch length through the average nucleotide substitution rate ($\simeq 4 \cdot 10^{-3}$ substitutions per site per year for HA). We have observed that given our method to predict the future from present populations corresponding to time bins of 4 months, changing the value of τ has little effect on the pick of the top LBI strain. By retrospectively optimizing its value, it is possible to reduce the average distance to the population 2 years ahead by ~ 0.25 amino acids on average, making the LBI method almost as good as the consensus on figure 4.

718 Measuring the geographical spread of a mutation

For a mutation X we define its regional distribution using the numbers $n_r(X)$ that represent the number of sequences sampled in region r that carry X . Regional weights are then defined as

$$w_r(X) = \frac{n_r(X)}{\sum_r n_r(X)}.$$

We can then measure the geographical spread $G(X)$ of X by using the Shannon entropy of the probability distribution $w_r(X)$:

$$G(X) = \sum_r w_r(X) \log(w_r(X)).$$

$G(X)$ is a positive quantity that is larger when X is equally present in many regions, and equal to zero when X is concentrated in only one region.

Region used are the ones defined in the Nextstrain tool [29]. Those are North America, South America, Europe, China, Oceania, Southeast Asia, Japan & Korea, South Asia, West Asia, and Africa.

688 Local Branching Index

726 LBI was introduced in [14] as an approximation of fitness in populations evolving under persistent selective pressure that is fully based on a phylogenetic tree. It relies on the intuition that the tree below high-fitness individuals will show dense branching events, whereas absence of branching is a sign of low-fitness individuals. Quantitatively, the LBI $\lambda_i(\tau)$ of a node i is the integral of all of the tree’s branch length around i , with an exponentially decreasing weight $e^{-t/\tau}$ with t being the branch length. When considering a time binned population, the LBI is computed once for each time bin by considering only the leaves of the tree that belong to the time bin. This means that only branches that ultimately lead to a leaf that belongs to the time bin are considered in the integration.

704 τ is the time scale for which the tree is informative of the fitness of a particular node. Here, we use a value of

Assigning a fitness to trajectories

Consensus sequence

Given a set of N sequences $(\sigma^1, \dots, \sigma^N)$ based on an alphabet \mathcal{A} (e.g. \mathcal{A} has 20 elements for amino acids, 4 for nucleotides), we can define a *profile* distribution $p_i(a)$ by the following expression:

$$p_i(a) = \sum_{n=1}^N \delta_{\sigma_i^n, a}$$

where i is a position in the sequence, σ_i^n the character appearing at position i in sequence σ^n , a a character of the alphabet and δ the Kronecker delta. The profile $p_i(a)$ simply represents the fraction of sequences which have character a at position i .

We then simply define the consensus sequence σ^{cons} such that

$$\sigma_i^{cons} = \operatorname{argmax}_a p_i(a).$$

In other words, the consensus sequence is the one that has the dominant character of the initial set of sequences at each position.

731 Earth Mover's Distance

In order to measure the distance of several predictor sequences to the future population, we rely on the *Earth Mover's Distance* (EMD), a metric commonly applied in machine learning to compare collections of pixels or words [30, 31]. Here, we apply it to compute the distance between the sequences of two populations, noted as $\mathcal{X} = \{(x^n, p^n)\}$ and $\mathcal{Y} = \{(y^m, q^m)\}$ with $n \in \{1 \dots N\}$ and $m \in \{1 \dots M\}$. In this notation, x^n and y^m are sequences, and p^n and q^m are the frequencies at which these sequences are found in their respective populations. For convenience, we also define $d_{mn} = H(x^n, y^m)$ as the

Hamming distance between pairs of sequences in the two populations.

We now introduce the following functional

$$F(\mathbf{w}) = \sum_{n,m} d_{nm} w_{nm},$$

with $\mathbf{w} = \{w_{nm}\}$ being a matrix of positive weights. The EMD between the two populations \mathcal{X} and \mathcal{Y} is now defined as the minimum value of function F under the conditions

$$\sum_{n=1}^N w_{nm} = q^m, \quad \sum_{m=1}^M w_{nm} = p^n, \text{ and } w_{nm} \geq 0$$

Intuitively, the weight w_{nm} tells us how much of sequence x^n is “moved” to sequence y^m . The functional F sums all of these moves and attributes them a cost equal to the Hamming distance d_{nm} . The conditions on weights in \mathbf{w} ensure that all the weight p^n of x^n is “moved” to elements in \mathcal{Y} and vice versa.

The minimization is easily performed by standard linear optimization libraries. Here, we use the Julia library JuMP [32].

- [1] World Health Organization, 2018. URL [https://www.who.int/fr/news-room/fact-sheets/detail/influenza-\(seasonal\)](https://www.who.int/fr/news-room/fact-sheets/detail/influenza-(seasonal)).
- [2] Velislava N. Petrova and Colin A. Russell. The evolution of seasonal influenza viruses. *Nature Reviews Microbiology* 16(1):47–60, October 2017. ISSN 1740-1526, 1740-1534. doi:10.1038/nrmicro.2017.118. URL <http://www.nature.com/doifinder/10.1038/nrmicro.2017.118>.
- [3] Arthur Chun-Chieh Shih, Tzu-Chang Hsiao, Mei-Shang Ho, and Wen-Hsiung Li. Simultaneous amino acid substitutions at antigenic sites drive influenza a hemagglutinin evolution. *Proceedings of the National Academy of Sciences*, 104(15):6283–6288, 2007. ISSN 0027-8424. doi:10.1073/pnas.0701396104. URL <https://www.pnas.org/content/104/15/6283>.
- [4] Samir Bhatt, Edward C. Holmes, and Oliver G. Pybus. The Genomic Rate of Molecular Adaptation of the Human Influenza A Virus. *Molecular Biology and Evolution*, 28(9):2443–2451, September 2011. ISSN 0737-4038. doi:10.1093/molbev/msr044. URL <https://academic.oup.com/mbe/article/28/9/2443/1007907>.
- [5] Björn F. Koel, David F. Burke, Theo M. Bestebroer, Stefan van der Vliet, Gerben C. M. Zondag, Gaby Vervaet, Eugene Skepner, Nicola S. Lewis, Monique I. J. Spronken, Colin A. Russell, Mikhail Y. Eropkin, Aeron C. Hurt, Ian G. Barr, Jan C. de Jong, Guus F. Rimmelzwaan, Albert D. M. E. Osterhaus, Ron A. M. Fouchier, and Derek J. Smith. Substitutions near the receptor binding site determine major antigenic change during influenza virus evolution. *Science*, 342(6161):976–979, 2013. ISSN 0036-8075. doi:10.1126/science.1244730. URL <https://science.scienmag.org/content/342/6161/976>.
- [6] Dylan H Morris, Katelyn M Gostic, Simone Pompei, Trevor Bedford, Marta Luksza, Richard A Neher, Bryan T Grenfell, Michael Lässig, and John W McCauley. Predictive modeling of influenza shows the promise of applied evolutionary biology. *Trends in microbiology*, 26(2):102–118, 2018.
- [7] Thorsten R. Klingen, Susanne Reimering, Carlos A. Guzmán, and Alice C. McHardy. In Silico Vaccine Strain Prediction for Human Influenza Viruses. *Trends in Microbiology*, 26(2):119–131, February 2018. ISSN 0966-842X. doi:10.1016/j.tim.2017.09.001. URL <http://www.sciencedirect.com/science/article/pii/S0966842X17302068>.
- [8] Peter Bogner, Ilaria Capua, David J Lipman, and Nancy J Cox. A global initiative on sharing avian flu data. *Nature*, 442(7106):981–981, 2006.
- [9] Yuelong Shu and John McCauley. Gisaid: Global initiative on sharing all influenza data—from vision to reality. *Eurosurveillance*, 22(13), 2017.
- [10] Andrew Rambaut, Oliver G. Pybus, Martha I. Nelson, Cecile Viboud, Jeffery K. Taubenberger, and Edward C. Holmes. The genomic and epidemiological dynamics of human influenza A virus. *Nature*, 453(7195):615–619, May 2008. ISSN 1476-4687. doi:10.1038/nature06945. URL <https://www.nature.com/articles/nature06945>.
- [11] Marta Luksza and Michael Lässig. A predictive fitness model for influenza. *Nature*, 507(7490):57–61, March 2014. ISSN 1476-4687. doi:10.1038/nature13087. URL <https://www.nature.com/articles/nature13087>. Number: 7490 Publisher: Nature Publishing Group.
- [12] L. Steinbrück, T. R. Klingen, and A. C. McHardy. Computational prediction of vaccine strains for human influenza a (h3n2) viruses. *Journal of Virology*, 88(20):12123–12132, 2014. ISSN 0022-538X. doi:10.1128/JVI.01861-14. URL <https://jvi.asm.org/content/88/20/12123>.

- [13] Richard A. Neher, Trevor Bedford, Rodney S. Daniels⁸⁷¹, Colin A. Russell, and Boris I. Shraiman. Prediction⁸⁷² dynamics, and visualization of antigenic phenotypes of⁸⁷³ seasonal influenza viruses. *Proceedings of the National⁸⁷⁴ Academy of Sciences of the United States of America*, 113⁸⁷⁵(12):E1701–1709, March 2016. ISSN 1091-6490 0027-8424⁸⁷⁶. doi:10.1073/pnas.1525578113.⁸⁷⁷
- [14] Richard A Neher, Colin A Russell, and Boris I Shraiman⁸⁷⁸. Predicting evolution from the shape of genealogical⁸⁷⁹ trees. *eLife*, 3, November 2014. ISSN 2050-084X. doi:⁸⁸⁰ 10.7554/eLife.03568. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4227306/>.⁸⁸²
- [15] Natalja Strelkowa and Michael Lässig. Clonal Inter-⁸⁸³ ference in the Evolution of Influenza. *Genetics*, 192⁸⁸⁴(2):671–682, October 2012. ISSN 0016-6731, 1943-⁸⁸⁵ 2631. doi:10.1534/genetics.112.143396. URL <http://www.genetics.org/content/192/2/671>.⁸⁸⁷
- [16] Yuri I Wolf, Cecile Viboud, Edward C Holmes, Eugene V⁸⁸⁸ Koonin, and David J Lipman. Long intervals of stasis⁸⁸⁹ punctuated by bursts of positive selection in the sea-⁸⁹⁰ sonal evolution of influenza A virus. *Biology Direct*, 1⁸⁹¹ 34, October 2006. ISSN 1745-6150. doi:10.1186/1745-⁸⁹² 6150-1-34. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1647279/>.⁸⁹⁴
- [17] Juhye M Lee, Rachel Eguia, Seth J Zost, Saket Choudhary,⁸⁹⁵ Patrick C Wilson, Trevor Bedford, Terry Stevens⁸⁹⁶, Ayers, Michael Boeckh, Aeron C Hurt, Seema S Lakdawala,⁸⁹⁷ Scott E Hensley, and Jesse D Bloom. Mapping⁸⁹⁸ person-to-person variation in viral mutations that escape⁸⁹⁹ polyclonal serum targeting influenza hemagglutinin.⁹⁰⁰ *eLife*, 8:e49324, August 2019. ISSN 2050-084X. doi:⁹⁰¹ 10.7554/eLife.49324. URL <https://doi.org/10.7554/eLife.49324>. Publisher: eLife Sciences Publications, Ltd.⁹⁰³
- [18] Motoo Kimura. Diffusion Models in Population Genetics. *Journal of Applied Probability*, 1(2):177–232,⁹⁰⁴ 1964. ISSN 0021-9002. doi:10.2307/3211856. URL <http://www.jstor.org/stable/3211856>.⁹⁰⁷
- [19] Le Yan, Richard A Neher, and Boris I Shraiman. Phylogodynamic theory of persistence, extinction and speciation of rapidly adapting pathogens. *eLife*, 8:e44205, September 2019. ISSN 2050-084X. doi:10.7554/eLife.44205.⁹¹¹ URL <https://doi.org/10.7554/eLife.44205>. Publisher: eLife Sciences Publications, Ltd.⁹¹³
- [20] R. A. Neher and B. I. Shraiman. Genetic draft and quasi-neutrality in large facultatively sexual populations. *Genetics*, 188(4):975–996, August 2011. ISSN 1943-2631.⁹¹⁶ doi:10.1534/genetics.111.128876.⁹¹⁷
- [21] Fabio Zanini and Richard A. Neher. FFPopSim: an efficient forward simulation package for the evolution of large populations. *Bioinformatics*, 28(24):3332–3333, 10 2012.⁹²⁰ ISSN 1367-4803. doi:10.1093/bioinformatics/bts633. URL <https://doi.org/10.1093/bioinformatics/bts633>.⁹²²
- [22] R. M. Bush, C. A. Bender, K. Subbarao, N. J. Cox,⁹²³ and W. M. Fitch. Predicting the evolution of human influenza A. *Science (New York, N.Y.)*, 286(5446):1921–⁹²⁴ 1925, December 1999. ISSN 0036-8075.⁹²⁶
- [23] Lars Steinbrück and Alice Carolyn McHardy. Inference⁹²⁷ of Genotype–Phenotype Relationships in the Antigenic⁹²⁸ Evolution of Human Influenza A (H3N2) Viruses. *PLOS⁹²⁹ Computational Biology*, 8(4):e1002492, April 2012. ISSN 1553-7358. doi:10.1371/journal.pcbi.1002492. URL <http://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1002492>.
- [24] Thorsten R. Klingen, Susanne Reimering, Jens Loers, Kyra Mooren, Frank Klawonn, Thomas Krey, Gülsah Gabriel, and Alice C. McHardy. Sweep Dynamics (SD) plots: Computational identification of selective sweeps to monitor the adaptation of influenza A viruses. *Scientific Reports*, 8(1):373, January 2018. ISSN 2045-2322. doi:10.1038/s41598-017-18791-z. URL <https://www.nature.com/articles/s41598-017-18791-z>.
- [25] Richard A. Neher and Oskar Hallatschek. Genealogies of rapidly adapting populations. *Proceedings of the National Academy of Sciences of the United States of America*, 110(2):437–442, January 2013. ISSN 1091-6490 0027-8424. doi:10.1073/pnas.1213113110.
- [26] Michael M. Desai, Aleksandra M. Walczak, and Daniel S. Fisher. Genetic Diversity and the Structure of Genealogies in Rapidly Adapting Populations. *Genetics*, 193(2):565–585, February 2013. ISSN 0016-6731, 1943-2631. doi:10.1534/genetics.112.147157. URL <http://www.genetics.org/content/193/2/565>.
- [27] Jason Schweinsberg. Coalescent processes obtained from supercritical Galton–Watson processes. *Stochastic Processes and their Applications*, 106(1):107–139, July 2003. ISSN 0304-4149. doi:10.1016/S0304-4149(03)00028-0. URL <http://www.sciencedirect.com/science/article/pii/S0304414903000280>.
- [28] Dylan H. Morris, Velislava Petrova, Fernando W. Rossine, Edyth Parker, Bryan Grenfell, Richard Neher, Simon Levin, and Colin Russell. Asynchrony between virus diversity and antibody selection limits influenza virus evolution. preprint, Open Science Framework, April 2020. URL <https://osf.io/847p2>.
- [29] James Hadfield, Colin Megill, Sidney M Bell, John Huddleston, Barney Potter, Charlton Callender, Pavel Sagunenko, Trevor Bedford, and Richard A Neher. Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics*, 34(23):4121–4123, 05 2018. ISSN 1367-4803. doi:10.1093/bioinformatics/bty407. URL <https://doi.org/10.1093/bioinformatics/bty407>.
- [30] Y. Rubner, C. Tomasi, and L. J. Guibas. A metric for distributions with applications to image databases. In *Sixth International Conference on Computer Vision (IEEE Cat. No.98CH36271)*, pages 59–66, Jan 1998. doi:10.1109/ICCV.1998.710701.
- [31] Matt J. Kusner, Yu Sun, Nicholas I. Kolkin, and Kilian Q. Weinberger. From word embeddings to document distances. In *Proceedings of the 32nd International Conference on International Conference on Machine Learning - Volume 37*, ICML’15, page 957–966. JMLR.org, 2015.
- [32] Iain Dunning, Joey Huchette, and Miles Lubin. Jump: A modeling language for mathematical optimization. *SIAM Review*, 59(2):295–320, 2017. doi:10.1137/15M1020575.
- [33] Julia Sigwart. Gene Genealogies, Variation and Evolution: A Primer in Coalescent Theory.—Jotun Hein, Mikkel H. Schierup, and Carsten Wiuf. 2004. Oxford University Press, Oxford. xiii + 276 pp. ISBN 0-19-852996-1, £29.95 (paperback); ISBN 0-19-852995-3, £65.00 (hardback). *Systematic Biology*, 54(6):986–987, 12 2005. ISSN 1063-5157. doi:10.1080/10635150500354860. URL <https://doi.org/10.1080/10635150500354860>.

930

SUPPLEMENTARY MATERIAL

931

1. Consensus sequence as a predictor for neutrally evolving populations

932 We consider the case of a neutrally evolving and structure-less population, such as the one in the Wright-Fisher
 933 model of evolution [33]. At an initial time $t = 0$, the population consists of N individuals with genomes $(\sigma^1 \dots \sigma^N)$ of
 934 length L (not necessarily distinct).

935 We make two hypotheses about this population. We first suppose that *no* mutations occur during the evolution of this
 936 population. This may seem surprising and is of course not true in the case of influenza. This assumption is however in
 937 line with the fact that the object of this work is to predict the outcome of *already existing* mutations in the influenza
 938 population. The prediction of mutations that we have not yet seen is not in its scope. Thus, assuming that no new
 939 mutations take place can be seen as a simple way to model the fact that we have no information about such events.
 940 The second assumption is that the population evolves in a completely neutral way, meaning that the average number
 941 of descendants of each genome σ^n is the same. Let us now consider the population after it has evolved for a long time
 942 $t \gg T$ where T is the typical coalescence time (for the Wright-Fisher model, $T = 2N$). At this point, all individuals in
 943 the future population will descend from a unique individual n_0 in the $t = 0$ population. Our two hypotheses now allow
 944 us to make two statements. First, since no new mutations are allowed, the population at $t \gg T$ will be clonal, with all
 945 individuals having genome σ^{n_0} . Second, since the evolution is neutral and does not favour any genome in particular,
 946 the probability that σ^{n_0} is equal to a given genome σ is $1/N$. In other words, the probability that a genome at $t = 0$
 947 ultimately becomes the ancestor of all the future population is equal to its frequency in the $t = 0$ population.

948 We now try to find the genome σ that best predicts the future population on the long run, that is for $t \gg T$. Here,
 949 we take best to mean that the predictor minimizes $H(\sigma, \sigma^{n_0})$ where H is the Hamming distance defined by

$$H(\sigma^a, \sigma^b) = \sum_{i=1}^L (1 - \delta_{\sigma_i^a, \sigma_i^b}), \quad (1)$$

951 with σ_i being the character appearing at position i of genome σ and δ the Kronecker delta. Since we do not know n_0 ,
 952 we have to average over all its possible values. σ must thus minimize the following quantity:

$$\begin{aligned} \langle H(\sigma, \sigma^{n_0}) \rangle_{n_0} &= \sum_{n=1}^N H(\sigma, \sigma^n) \\ &= \sum_{i=1}^L \sum_{n=1}^N (1 - \delta_{\sigma_i, \sigma_i^n}) \end{aligned} \quad (2)$$

by using the definition of the Hamming distance. We now assume that characters at each positions of the genomes can
 be indexed by an integer a running from 1 to q . For instance, if these were amino acid sequences, we could index the
 20 amino acids by a running from 1 to $q = 20$. We rewrite the Kronecker delta in the previous expression using this
 indexation:

$$\delta_{\sigma_i, \sigma_i^n} = \sum_{a=1}^q \delta_{\sigma_i, a} \delta_{\sigma_i^n, a}.$$

953 We also introduce the *profile* frequencies $p_i(a)$ of the population at time $t = 0$:

$$p_i(a) = \sum_{n=1}^N \delta_{\sigma_i^n, a}. \quad (3)$$

954 $p_i(a)$ represents the frequency at which character a appears at position i in genomes of the initial population.
 955 Equation 2 now becomes

$$\begin{aligned} \langle H(\sigma, \sigma^{n_0}) \rangle_{n_0} &= \sum_{i=1}^L \sum_{n=1}^N \left(1 - \sum_{a=1}^q \delta_{\sigma_i, a} \delta_{\sigma_i^n, a} \right) \\ &= \sum_{i=1}^L \left(1 - \sum_{a=1}^q \delta_{\sigma_i, a} p_i(a) \right) \\ &= \sum_{i=1}^L (1 - p_i(\sigma_i)) \end{aligned} \quad (4)$$

956 This means that the genome $\sigma = (\sigma_1 \dots \sigma_L)$ which best predicts the future population according to our definition is
 957 the one that minimizes the quantity $(1 - p_i(\sigma_i))$ for all positions i . This obviously implies that each σ_i must be chosen
 958 as to maximize $p_i(a)$, that is σ_i must be the character that appears the most frequently at position i . Thus, σ must be
 959 the *consensus* sequence of the initial population.

960

2. Predictor based on the local LBI maxima

In figure 15, we use several sequences as a predictor of the future population. Distance between two sets of sequences, *i.e.* the predictor sequences and the ones of the future population, is defined as the Earth Mover's Distance (EMD). Here, we show that for a population evolving under the same hypotheses as in section 1, the best *multiple* sequence long term predictor is again the consensus sequence with weight 1.

Let the predictor be a set of weighted sequences $\{(s^\alpha, q_\alpha)\}$. We again use the fact that in the long term, a unique sequence σ^{n_0} from the present will be the ancestor of the entire population. We want to compute the EMD from the predictor to σ^{n_0} , that is the EMD between the sets $\mathcal{X} = \{(s^\alpha, q_\alpha)\}$ and $\mathcal{Y} = \{\sigma^{n_0}, 1\}$. Applying the definition of the Methods section, it follows that the weights \mathbf{w} are in this case equal to the q_α s. By averaging over all values of n_0 , we now obtain

$$\langle \text{EMD}(\{(s^\alpha, q_\alpha)\}) \rangle_{n_0} = \sum_{n=1}^N \sum_{\alpha} H(s^\alpha, \sigma^n) \cdot q_\alpha.$$

By the same calculation procedure as in the previous section, this expression simplifies to

$$\langle \text{EMD}(\{(s^\alpha, q_\alpha)\}) \rangle_{n_0} = \sum_{i=1}^L \left(1 - \sum_{a=1}^q p_i(a) q_i(a) \right),$$

where the profile of the present population $p_i(a)$ has already been defined, and $q_i(a)$ stands for the profile of the predictor, that is

$$q_i(a) = \sum_{\alpha} \delta_{s_i^\alpha, a} q_\alpha.$$

961 To minimize this distance, we find a profile $q_i(a)$ that maximizes the quantity $\sum_{\alpha} \delta_{s_i^\alpha, a} q_\alpha$ for each position i . It is
 962 clear that this is done by assigning a value $q_i(a) = 1$ if a maximizes $p_i(a)$, and $q_i(a) = 0$ otherwise. Thus, the profile of
 963 the predictor must be that of the consensus sequence, which is only possible if the predictor becomes $\{\sigma^{cons}, 1\}$.

964

3. Biases in frequency estimations

965 The frequency of mutations in a given time-bin is simply performed by computing their frequency in sequences
 966 sampled in that time bin. This leads to potential biases in estimating frequencies, that arise for two reasons:

- 967 (i) A mutation present at frequency p in the population might be observed at another frequency $f \neq p$ if f is
 968 estimated using a sub-sample of the population.
- 969 (ii) For a neutrally evolving population, the distribution of frequencies of alleles is of the form $P(p) \propto 1/p$. This
 970 means that the amount of alleles at frequency p is lower when p is higher.

971 To illustrate (i), let us compute the probability that a mutation present at “real” frequency p in the population is
 972 found to be in a given frequency bin $[f_1, f_2]$ when p is estimated from a sample of size n . The sample consists of n
 973 observations $\{x_i\}$ with $1 \leq i \leq n$, with $x_i = 1$ if sequence i of the sample bears the mutation, and $x_i = 0$ if
 974 not. If n is small with regard to the total population size, we can consider the x_i as random variables with a binomial
 975 distribution, meaning that $P(x_i = 1) = p$ and $P(x_i = 0) = 1 - p$. The empirical frequency f is then estimated by
 976 taking the average of the x_i variables, that is $f = (x_1 + \dots + x_n)/n$. If those are independently sampled and n is large
 977 enough, the probability of measuring value f is given by the Central Limit Theorem:

$$P_{n,p}(f) \propto e^{(f-p)^2/2\sigma^2}, \quad \text{where } \sigma^2 = \frac{p(1-p)}{n}. \quad (5)$$




Figure S 1. **Left:** For a mutation present at frequency p in the population, probability of being observed in the frequency bin $[0.1, 0.2]$ as a function of p and for different sample sizes n . The dashed black line sketches the (non-normalized) background distribution $P_b(p)$. **Right:** Expected “real” average frequency of mutations found in frequency bin $[f_1, f_2]$ as a function of the centre of the bin $(f_1 + f_2)/2$, for different sample sizes.

988 To compute the probability that this mutation is found in a given frequency bin $[f_1, f_2]$, we integrate this distribution:

$$P_{f_1, f_2}(p, n) = \int_{f_1}^{f_2} dx P_{n,p}(x). \quad (6)$$

989 Function $P_{f_1, f_2}(p, n)$ is shown as a function of p for a fixed interval and for different values of n in the first panel of
990 figure S1. Note the asymmetry of it: the variance of a binomial distribution of parameter p is small when p is close to
991 0 or 1, and goes through a maximum at $p = 0.5$. For this reason, mutations present at frequency p close to 0.5 have a
992 higher probability of being observed in other frequency bins. On the contrary, this is unlikely for very rare or very
993 frequent mutations.

994 We now try to estimate biases in frequency estimation due this phenomenon. Given a set of mutations that have
995 been measured in frequency bin $[f_1, f_2]$, what is the average *real* frequency of these mutations? To compute this, we
996 need to sum $P_{f_1, f_2}(p, n)$ over all possible real frequencies p , giving us the amount of mutations that are observed in
997 interval $[f_1, f_2]$, and weigh this sum by the frequency value p as well as by the background distribution of frequencies
998 $P_b(p) \propto 1/p$. This last quantity represents the expected amount of mutations that are present at frequency p in the
999 population. Note that there is no divergence problem as the smallest non zero frequency is $1/N$, where N is the
999 population size. This leads us to the following expression for the average of “real” frequencies:

$$\begin{aligned} \langle p \rangle(f_1, f_2, n) &= \int_{1/N}^{1-1/N} dp P_{f_1, f_2}(p, n) P_b(p) p \\ &= \int_{1/N}^{1-1/N} dp P_{f_1, f_2}(p, n). \end{aligned} \quad (7)$$

993 We have not made normalization explicit in these equations. It is simply achieved by dividing the above expression by
994 $\int dp P_{f_1, f_2}(p, n) P_b(p)$.

995 In the second panel of figure S1, $\langle p \rangle(f_1, f_2, n)$ is plotted as a function of the centre of the interval $[f_1, f_2]$ and for
996 different values of n . For sample sizes $n > 100$, the biases due to this effect are almost non existent. For smaller
997 samples, for instance $n = 10$, they are small but non negligible. However, we argue that this is not a significant problem
998 with respect to the main results presented in this article. First, figure S6 shows that sample sizes of the order of $n = 10$
999 are only the case for a few months in the period going from year 2000 to 2018. From 2010 and onwards, more than a
1000 hundred sequences are available per month for most months. Secondly, even if most samples were in the $n = 10$ case,

1001 deviations shown in figure S1 are small enough that results shown in figures 2 and 3 would be *qualitatively* unchanged.
1002 Note that using the centre of the interval as a reference in figure S1, *i.e.* $(f_1 + f_2)/2$, would be correct in the case of
1003 a very large n and a flat background distribution $P_b(p)$. For figures 2 and 3 of the main text however, the average
1004 frequency of mutations found in an interval $[f_1, f_2]$ is computed by taking the average of the observed frequencies, and
1005 not the centre of the interval. This partially takes into account biases considered here, as the background distribution
1006 $P_b(p)$ is then accounted for, even though it is equivalent to assuming infinite sample sizes.

1007

4. Cutting off the HA1 159S branch




Figure S 2. Tree used for this study, based on a random selection of 100 strains per month from year 2002 to 2018. Nodes and branches are colored according to the amino acid found at position HA1:159. The HA1 159S mutation is visible as a thin but long light-greened color branch, coalescing with the “trunk” around 2013.

1008 The analysis of the main text is in a large part based on the probability of fixation of mutations. The motivation
 1009 underlying this choice is the relatively short coalescence time of the A/H3N2 influenza population, typically around
 1010 three years. This can be seen in figure 2 of the main text, which shows the typical lifetime of frequency trajectories,
 1011 ending in fixation or loss after at most 3 years in most cases. The tree in figure S2 is another illustration of this: for
 1012 the most part of it, a “trunk” is clearly identifiable, and lineages that depart from it have a relatively short lifetime.
 1013 This is no longer the case since the year ~ 2013: two clades have been competing since then, with no definite way to
 1014 identify a trunk in the tree. The clade defined by the HA1 159S mutation, colored in light green on figure S2, is one of
 1015 these two competing lineages. Because of this particular situation, the number of mutations fixating in the population
 1016 is strongly reduced, as a mutation must appear in both clades to reach a frequency of 1. This is a potential flaw in our
 1017 analysis, which concentrates on mutations fixating.

1018 For this reason, we decided to re-run our analysis after having cut off the HA1 159S clade. In other words, we remove
 1019 from the set of sequences those that carry the HA1 159S mutation. Results are shown in figures , equivalent to figures 2
 1020 and 3 of the main text. It is clear that qualitative results are left unchanged when this competing clade is removed.
 1021 This can be surprising, as almost no complete fixation of an amino acid mutation has occurred since 2013. Cutting off
 1022 the HA1 159S branch should thus result in many new fixations, changing the analysis. The reason for the similarity
 1023 of results can be explained: fixation (resp. loss) of a mutation are defined here as the frequency of this mutation
 1024 being measured above 95% (resp. 5%) frequency for two months in a row. As the HA1 159S clade is rather sparsely
 1025 populated, it reaches frequencies lower than 5% two times (in 2015 and 2017), allowing mutations in the competing
 1026 clade to “fix” as defined here. Thus, removing strains carrying HA1 159S does not introduce a significant amount of
 1027 “new” fixation events.




Figure S 3. Equivalent to figure 2 of the main text, but with strains carrying the HA1 159S mutation removed.




Figure S 4. Equivalent to figure 3 of the main text, but with strains carrying the HA1 159S mutation removed.

1028

5. Probability of fixation in single locus model of evolution

1029 In [18], Kimura investigates a simple model of evolution with a single locus and a population of size N . In this
 1030 framework, a mutation at this locus with fitness effect s and observed at frequency f has the following probability of
 1031 fixation:

$$P_{fix}(f|s, N) = \frac{1 - e^{-sNf}}{1 - e^{-sN}}. \quad (8)$$

1032 Expanding this formula for $sN \ll 1$, that is in the weak selection regime, yields at the first order

$$P_{fix}(f|s, N) = f + f(1 - f) \frac{sN}{2}. \quad (9)$$

1033 Equation 9 tells us two things. First, when the mutation is neutral, that is $s = 0$, we have $P_{fix}(f) = f$. This naturally
 1034 confirms the result obtained for a neutral model of evolution. Seconds, when $sN \neq 0$, we can expect deviations from
 1035 the diagonal in a P_{fix} against f plot. The sign of these deviations is determined by the sign of s , with beneficial
 1036 mutations being found above diagonal while deleterious one are found below. The amplitude of these deviations
 1037 depends on the strength of selection sN , as well as on the frequency through the $f(1 - f)$ term, making them larger
 1038 for $f \sim 0.5$.

6. Mutation tables

Gene	Position	AA	Start date	End date	Shih	Luksza	Koel	Tree counts
HA1	144	D	2001-06-09	2002-02-04	true	true	false	0
HA1	189	N	2003-07-29	2004-05-24	false	true	true	2
HA1	159	F	2003-08-28	2004-05-24	false	true	true	2
HA1	226	I	2003-09-27	2004-09-21	true	true	false	3
HA1	145	N	2003-12-26	2004-11-20	false	true	true	2
HA1	227	P	2003-05-30	2005-04-19	false	true	false	2
HA2	32	I	2004-06-23	2005-07-18	false	false	false	1
HA1	193	F	2004-12-20	2006-03-15	false	true	true	1
HA2	46	D	2006-06-13	2007-05-09	false	false	false	2
HA2	121	K	2006-06-13	2007-06-08	false	false	false	1
HA1	50	E	2006-09-11	2007-06-08	false	true	false	2
HA1	140	I	2006-11-10	2007-11-05	true	false	false	1
HA1	173	Q	2007-07-08	2009-01-28	true	true	false	2
HA2	32	R	2007-07-08	2009-01-28	false	false	false	1
HA1	158	N	2009-01-28	2009-07-27	true	true	true	2
HA1	189	K	2009-01-28	2009-07-27	false	true	true	2
HA1	212	A	2009-03-29	2011-01-18	false	false	false	2
HA1	45	N	2010-03-24	2013-02-06	false	false	false	3
HA1	223	I	2010-12-19	2013-02-06	false	false	false	2
HA1	48	I	2011-03-19	2013-02-06	false	false	false	1
HA1	198	S	2011-03-19	2013-02-06	false	false	false	1
HA1	312	S	2009-08-26	2013-03-08	false	false	false	3
HA1	278	K	2011-06-17	2013-03-08	false	true	false	1
HA1	145	S	2011-04-18	2013-04-07	false	true	true	4
HA1	33	R	2011-06-17	2013-06-06	false	false	false	2
HA2	160	N	2012-07-11	2015-09-24	false	false	false	3
HA1	225	D	2013-08-05	2015-09-24	false	false	false	3
HA1	3	I	2013-08-05	2016-11-17	false	false	false	2
HA1	159	Y	2014-02-01	2016-11-17	false	true	true	2
HA1	160	T	2014-01-02	2017-07-15	false	true	false	2

Table S I. The 30 trajectories that took place between year 2000 and year 2018 and resulted in fixation. Columns **Shih**, **Luksza** and **Koe1** respectively indicate whether the position is found in the epitopes lists in (respectively) [3], [11] and [5]. The **Tree counts** column indicates the number of times the mutation corresponding to the trajectory can be found in the phylogenetic tree. Note that a trajectory is only shown in the table if the sequenced population counts more than 10 strains at its time of fixation. This explains that only 30 trajectories are displayed, whereas more mutations did fix in this period of time.

Gene	Position	AA	Start date	End date	Fixation	Max. freq.
HA1	106	A	2001-02-09	2002-02-04	lost	1.0
HA1	144	D	2001-06-09	2002-02-04	fixed	1.0
HA1	105	H	2003-04-30	2003-10-27	lost	1.0
HA1	126	D	2003-04-30	2004-05-24	lost	1.0
HA1	140	Q	2004-01-25	2004-06-23	lost	0.31
HA1	226	I	2003-09-27	2004-09-21	fixed	1.0
HA1	173	E	2004-12-20	2006-03-15	lost	0.63
HA1	142	G	2006-06-13	2007-05-09	lost	0.71
HA1	144	D	2006-07-13	2007-05-09	lost	0.67
HA1	128	A	2006-09-11	2007-05-09	lost	0.25
HA1	157	S	2006-09-11	2007-05-09	lost	0.59
HA1	140	I	2006-11-10	2007-11-05	fixed	1.0
HA1	173	N	2007-12-05	2008-07-02	lost	0.3
HA1	157	S	2007-12-05	2008-09-30	lost	0.31
HA1	173	E	2006-06-13	2008-12-29	lost	0.67
HA1	173	Q	2007-07-08	2009-01-28	fixed	0.96
HA1	158	N	2009-01-28	2009-07-27	fixed	0.96
HA1	62	K	2009-01-28	2011-05-18	lost	0.73
HA1	144	K	2009-01-28	2011-05-18	lost	0.75
HA1	62	V	2011-04-18	2011-09-15	lost	0.34
HA1	157	S	2013-05-07	2015-09-24	lost	0.35
HA1	128	A	2012-08-10	2016-11-17	lost	0.81
HA1	197	K	2015-11-23	2016-11-17	lost	0.27
HA1	142	R	2018-05-11	2018-10-08	lost	0.38
HA1	142	G	2012-03-13		poly	0.86
HA1	144	S	2013-12-03		poly	0.96
HA1	121	K	2015-12-23		poly	0.82
HA1	142	K	2016-05-21		poly	0.77
HA1	62	G	2017-03-17		poly	0.75
HA1	128	A	2018-01-11		poly	0.56

Table S II. Trajectories of mutations at epitope positions in [3] (*Shih et. al.*) that have been observed at least once above frequency 0.25. The **Fixation** column indicates whether the mutation has fixed, disappeared, or is still polymorphic as of October 2018. The **Max.freq.** column indicates the maximum frequency reached by the trajectory. A maximum frequency of 1 for mutations that finally disappear is explained by trajectories reaching frequency 1 for one time bin and going back to lower values for following ones (a frequency above 0.95 for two time bins in a row defines fixation).

Gene	Position	AA	Start date	End date	Fixation	Max. freq.
HA1	50	G	2001-02-09	2002-02-04	lost	1.0
HA1	144	D	2001-06-09	2002-02-04	fixed	1.0
HA1	126	D	2003-04-30	2004-05-24	lost	1.0
HA1	189	N	2003-07-29	2004-05-24	fixed	1.0
HA1	159	F	2003-08-28	2004-05-24	fixed	1.0
HA1	226	I	2003-09-27	2004-09-21	fixed	1.0
HA1	145	N	2003-12-26	2004-11-20	fixed	1.0
HA1	188	N	2004-07-23	2005-02-18	lost	0.36
HA1	227	P	2003-05-30	2005-04-19	fixed	1.0
HA1	173	E	2004-12-20	2006-03-15	lost	0.63
HA1	193	F	2004-12-20	2006-03-15	fixed	0.97
HA1	142	G	2006-06-13	2007-05-09	lost	0.71
HA1	144	D	2006-07-13	2007-05-09	lost	0.67
HA1	157	S	2006-09-11	2007-05-09	lost	0.59
HA1	50	E	2006-09-11	2007-06-08	fixed	0.95
HA1	173	N	2007-12-05	2008-07-02	lost	0.3
HA1	157	S	2007-12-05	2008-09-30	lost	0.31
HA1	173	E	2006-06-13	2008-12-29	lost	0.67
HA1	173	Q	2007-07-08	2009-01-28	fixed	0.96
HA1	158	N	2009-01-28	2009-07-27	fixed	0.96
HA1	189	K	2009-01-28	2009-07-27	fixed	0.96
HA1	213	A	2009-01-28	2010-02-22	lost	0.68
HA1	144	K	2009-01-28	2011-05-18	lost	0.75
HA1	53	N	2009-11-24	2013-02-06	lost	0.72
HA1	278	K	2011-06-17	2013-03-08	fixed	0.98
HA1	145	S	2011-04-18	2013-04-07	fixed	0.99
HA1	159	S	2013-11-03	2015-08-25	lost	0.46
HA1	157	S	2013-05-07	2015-09-24	lost	0.35
HA1	159	Y	2014-02-01	2016-11-17	fixed	0.97
HA1	159	S	2015-10-24	2016-11-17	lost	0.4
HA1	197	K	2015-11-23	2016-11-17	lost	0.27
HA1	160	T	2014-01-02	2017-07-15	fixed	0.96
HA1	142	R	2018-05-11	2018-10-08	lost	0.38
HA1	135	N	2018-06-10	2018-10-08	lost	0.38
HA1	142	G	2012-03-13		poly	0.86
HA1	144	S	2013-12-03		poly	0.96
HA1	121	K	2015-12-23		poly	0.82
HA1	142	K	2016-05-21		poly	0.77
HA1	131	K	2016-09-18		poly	0.77
HA1	135	K	2016-11-17		poly	0.47

Table S III. Same as table SII, for [11] (*Luksza et. al.*).

Gene	Position	AA	Start date	End date	Fixation	Max. freq.
HA1	189	N	2003-07-29	2004-05-24	fixed	1.0
HA1	159	F	2003-08-28	2004-05-24	fixed	1.0
HA1	145	N	2003-12-26	2004-11-20	fixed	1.0
HA1	193	F	2004-12-20	2006-03-15	fixed	0.97
HA1	158	N	2009-01-28	2009-07-27	fixed	0.96
HA1	189	K	2009-01-28	2009-07-27	fixed	0.96
HA1	145	S	2011-04-18	2013-04-07	fixed	0.99
HA1	159	S	2013-11-03	2015-08-25	lost	0.46
HA1	159	Y	2014-02-01	2016-11-17	fixed	0.97
HA1	159	S	2015-10-24	2016-11-17	lost	0.4

Table S IV. Same as table SII, for [5] (*Koel et. al.*).

7. Supplementary figures




Figure S 5. Number of A/H3N2 HA sequences per year from year 1990.




Figure S 6. Number of H3N2 HA and NA sequences per month from year 2000.




Figure S 7. Frequency trajectories for the 9 most entropic positions in the A/H3N2 HA protein.




Figure S 8. Equivalent to panel **B** of figure 1 of the main text for A/H3N2, with f_0 equal to 0.5 in **A** (76 trajectories), and 0.7 in **B** (63 trajectories).




Figure S 9. Equivalent to panel **B** of figure 1 of the main text for A/H1N1pdm influenza. 89 trajectories are shown and participate to the mean (thick black line).




Figure S 10. Equivalent to panel **B** of figure 1 of the main text for A/H1N1pdm, with f_0 equal 0.5 in **A** (50 trajectories), and 0.7 in **B** (41 trajectories).




Figure S 11. Equivalent of figure 3 of the main text for the HA gene of A/H1N1pdm influenza. Fixation probability $P_{fix}(f)$ as a function of frequency. **A:** Mutation with higher or lower LBI values, based on their position with respect to the median LBI value. **B:** Different lists of epitope positions in the HA protein. The authors and the number of positions is indicated in the legend. **C:** Mutations for binary positions, *i.e.* positions for which we never see more than two amino acids in the same time bin. **D:** Mutations that appear once or more than once in the tree for a given time bin.




Figure S 12. Based on A/H3N2 HA and NA. **A:** Mutations with a higher or lower geographical spread, based on the median value of the score used (see Methods). *Note:* the words *local* and *global* only reflect the position of the geographic spread of the mutation relative to the median value computed for all mutations found at this frequency. As this median value may change with the considered frequency bin, so does the definition of local and global mutations. **B:** Mutations whose trajectories are older or more recent, based on the median age of trajectories when reaching the considered frequency f .




Figure S 13. Based on A/H1N1pdm HA and NA. **A:** Mutations with a higher or lower geographical spread, based on the median value of the score used (see Methods). *Note:* the words *local* and *global* only reflect the position of the geographic spread of the mutation relative to the median value computed for all mutations found at this frequency. As this median value may change with the considered frequency bin, so does the definition of local and global mutations. **B:** Mutations whose trajectories are older or more recent, based on the median age of trajectories when reaching the considered frequency f .




Figure S 14. Geographic spread of mutations as a function of the time for which they have been present in the population above a frequency of 5%. Points represent individual mutations and for a population in a given time bin. The line is the average of dots for a given value on the x -axis. Based on data for A/H3N2 HA.




Figure S 15. Earth mover's distance to the future population for different predictors. A present population consists of all A/H3N2 HA sequences sampled in a 4 months time window. Quantities are averaged over all possible "present" populations from the year 2002. Predictors are: **Global consensus**: Consensus sequence of the present population. Best long-term predictor for a structure-less neutrally evolving population. **All present population**: All sequences in the present population. Perfect predictor if the population does not change at all through time. **Cluster-wise consensus**: Consensus sequence for each cluster in the present population. Clusters are based on local maxima of the LBI. Sequences are assigned to a given cluster based on their tree branch-length distance to the corresponding local maximum.




Figure S 16. Fixation probability as a function of frequency for the simulations discussed in the main text. **Top:** Simulation without antibodies. The three colored curves reflect different rate of change for the fitness landscape. Visual inspection of the frequency trajectories indicates a typical sweep time of ~ 400 generations. **Bottom:** Simulation with antibodies. The different colored curves indicate the rate at which antibodies are introduced.




Figure S 17. Fixation probability as a function of frequency for the simulations discussed in the main text, with trajectories stratified according to real fitness values. “High” and “low” fitness classes are defined with respect to the median value. **Top:** Simulation without antibodies and with changes to the fitness landscape every $dt = 10$ generations. **Bottom:** Simulation with antibodies, with a new antibody every $dt = 10$ generations.