

THÈSE DE DOCTORAT
DE SORBONNE UNIVERSITÉ

Spécialité : Physique

École doctorale n°564: Physique en Île-de-France

réalisée

au Laboratoire Kastler Brossel

sous la direction de Pierre-François Cohadon

présentée par

Pierre-Edouard Jacquet

pour obtenir le grade de :

DOCTEUR DE SORBONNE UNIVERSITÉ

Sujet de la thèse :

Progress towards cryogenic squeezed light optomechanics

soutenue le ?? ??? 2025

devant le jury composé de :

M.	Mr.	Rapporteur
M.	Mr.	Rapporteur
M.	Mr.	Examineur
Mme.	coucou	Examinatrice
Mme.	salut	Présidente du Jury
M.	aurevoir	Directeur de thèse

Contents

I	Introduction	1
I.1	Historical background	1
I.2	State of the art	1
I.3	Relevance of this work	1
II	Theory: background	3
II.1	Quantum Optics Concepts	3
II.1.1	Quantum Description of Light	3
II.1.2	*Coherent States	4
II.1.3	Squeezed States	5
II.1.4	Field Operators	5
II.1.5	Optical Field Modulations	5
II.1.6	Quantum Noise and Uncertainty	5
II.1.7	Sideband Representation	5
II.2	Optical Cavities : Basics	5
II.2.1	Cavity types and Resonance Conditions	5
II.2.2	Spatial and Longitudinal Modes	5
II.2.3	Static and Dynamical effects	5
II.3	Optical Cavities : Three Mirror Cavities	6
II.3.1	6
II.4	Cavity Optomechanics	6
II.4.1	Radiation Pressure Coupling	6
II.4.2	Quantum Langevin Equations	6
II.4.3	Mechanical Resonators	6
II.4.4	Noise spectra	6
II.4.5	Three Mirror Cavities as Novel Optomechanical Systems	6
II.5	Squeezed Light Theory	6
II.5.1	Single-mode Squeezing	6
II.5.2	Noise Spectra	6
II.5.3	Frequency-dependent Squeezing and its use	6
II.6	Numerical Simulations of FDS Optomechanical Experiments	6
III	Theory: MATE systems with squeezed light	7

IV Experiments : A MATE system in vacuum	9
V Experiments : a high frequency squeezed light source	11
VI Un appendice	15

Chapter I

Introduction

- I.1 Historical background**
- I.2 State of the art**
- I.3 Relevance of this work**

Chapter II

Theory: background

This chapter will cover the elementary concepts required to describe an membrane based optomechanical system in a quantum regime. We will first recall basics on optical field quantization as well describing coherent and squeezed light field, to then turn to the more specific frequency dependent squeezed light field. Secondly, we will cover the mathematical description of a mechanical resonator interacting with a generic coherent optical field, highlighting the differences with the seminal optomechanical system of a mirror on a spring. Finally, we will derive the equations of motions of a membrane based optomechanical system with frequency dependent squeezed optical fields.

II.1 Quantum Optics Concepts

II.1.1 Quantum Description of Light

We introduce briefly field quantization concepts needed to describe monochromatic field propagation and measurements. key words : eigenmodes of the field, quantization, annihilation operators, quadratures, phase space, displacement operators, squeezing operators, coherent states, generic squeezed states,

Quantised Electromagnetic Field

We consider the quantised electromagnetic field in volume V . The electric field operator can be expressed as:

$$\hat{\mathbf{E}}(\mathbf{r}, t) = i \sum_{\ell} \mathcal{E}_{\ell} \left[\hat{a}_{\ell}(t) \mathbf{f}_{\ell}(\mathbf{r}) - \hat{a}_{\ell}^{\dagger}(t) \mathbf{f}_{\ell}^*(\mathbf{r}) \right] \quad (\text{II.1})$$

where $\mathcal{E}_{\ell} = \sqrt{\frac{\hbar \omega_{\ell}}{2 \varepsilon_0 V}}$ is the field per photon in mode ℓ , \hbar the reduced Planck constant, ε_0 the vacuum permittivity, $\mathbf{f}_{\ell}(\mathbf{r})$ are spatial mode functions satisfying orthonormality, and $\hat{a}_{\ell}(t)$, $\hat{a}_{\ell}^{\dagger}(t)$ are the time dependent annihilation and creation operators associated with each mode (ℓ). Here, we describe

These operators satisfy the canonical commutation relations:

$$[\hat{a}_{\omega, \ell}(t), \hat{a}_{\omega', \ell'}^{\dagger}(t)] = \delta_{\omega, \omega'} \delta_{\ell, \ell'}, \quad [\hat{a}_{\omega, \ell}(t), \hat{a}_{\omega', \ell'}(t)] = 0 \quad (\text{II.2})$$

The Hamiltonian for the electromagnetic field becomes a sum of harmonic oscillator energies:

$$\hat{H} = \sum_{\omega, \ell} \hbar \omega \left(\hat{a}_{\omega, \ell}^\dagger \hat{a}_{\omega, \ell} + \frac{1}{2} \right) \quad (\text{II.3})$$

Quadrature Operators

To describe the phase space properties of a field mode, we define the quadrature operators:

$$\hat{q}_{\omega, \ell} = \frac{1}{\sqrt{2}} (\hat{a}_{\omega, \ell} + \hat{a}_{\omega, \ell}^\dagger) \quad (\text{II.4})$$

$$\hat{p}_{\omega, \ell} = \frac{1}{\sqrt{2}i} (\hat{a}_{\omega, \ell} - \hat{a}_{\omega, \ell}^\dagger) \quad (\text{II.5})$$

These are Hermitian operators corresponding to measurable observables and satisfy:

$$[\hat{q}_{\omega, \ell}, \hat{p}_{\omega, \ell}] = i \quad (\text{II.6})$$

Generalised quadratures are defined via:

$$\hat{q}_\theta^{\omega, \ell} = \hat{q}_{\omega, \ell} \cos \theta + \hat{p}_{\omega, \ell} \sin \theta \quad (\text{II.7})$$

Uncertainty Principle and Quantum Noise

From the commutation relation, the uncertainty principle follows:

$$\Delta q_{\omega, \ell} \Delta p_{\omega, \ell} \geq \frac{1}{2} \quad (\text{II.8})$$

This defines the minimum amount of quantum noise (vacuum fluctuations) in the electromagnetic field.

II.1.2 *Coherent States

Coherent states $|\alpha_{\omega, \ell}\rangle$ are eigenstates of the annihilation operator:

$$\hat{a}_{\omega, \ell} |\alpha_{\omega, \ell}\rangle = \alpha_{\omega, \ell} |\alpha_{\omega, \ell}\rangle \quad (\text{II.9})$$

They can be generated by displacing the vacuum:

$$|\alpha_{\omega, \ell}\rangle = \hat{D}(\alpha_{\omega, \ell}) |0\rangle, \quad \hat{D}(\alpha) = \exp(\alpha \hat{a}^\dagger - \alpha^* \hat{a}) \quad (\text{II.10})$$

They exhibit:

- Minimum uncertainty: $\Delta q = \Delta p = 1/\sqrt{2}$
- Classical-like dynamics
- Poissonian photon statistics

II.1.3 Squeezed States

Squeezed states reduce the variance of one quadrature below vacuum level:

$$|\xi_{\omega,\ell}\rangle = \hat{S}(\xi_{\omega,\ell})|0\rangle \quad (\text{II.11})$$

$$\hat{S}(\xi) = \exp \left[\frac{1}{2}(\xi^* \hat{a}^2 - \xi \hat{a}^{\dagger 2}) \right], \quad \xi = r e^{i\phi} \quad (\text{II.12})$$

For phase quadrature squeezing ($\phi = 0$):

$$\Delta q_{\omega,\ell} = e^{-r}/\sqrt{2}, \quad \Delta p_{\omega,\ell} = e^r/\sqrt{2} \quad (\text{II.13})$$

Squeezed light is a key resource for precision metrology and quantum information.

II.1.4 Field Operators

The quantised vector potential and electric field can be expressed as:

$$\hat{\mathbf{A}}(\mathbf{r}) = \sum_{\omega,\ell} \sqrt{\frac{\hbar}{2\varepsilon_0\omega V}} \left[\hat{a}_{\omega,\ell} \mathbf{u}_{\omega,\ell}(\mathbf{r}) + \hat{a}_{\omega,\ell}^\dagger \mathbf{u}_{\omega,\ell}^*(\mathbf{r}) \right] \quad (\text{II.14})$$

$$\hat{\mathbf{E}}(\mathbf{r}) = i \sum_{\omega,\ell} \sqrt{\frac{\hbar\omega}{2\varepsilon_0 V}} \left[\hat{a}_{\omega,\ell} \mathbf{u}_{\omega,\ell}(\mathbf{r}) - \hat{a}_{\omega,\ell}^\dagger \mathbf{u}_{\omega,\ell}^*(\mathbf{r}) \right] \quad (\text{II.15})$$

These operators are central to describing interactions between light and matter in cavity QED, optomechanics, and other quantum platforms.

This concludes our introduction to the quantum description of light, setting the stage for modelling interactions between quantum optical fields and mechanical resonators.

II.1.5 Optical Field Modulations

II.1.6 Quantum Noise and Uncertainty

II.1.7 Sideband Representation

II.2 Optical Cavities : Basics

II.2.1 Cavity types and Resonance Conditions

II.2.2 Spatial and Longitudinal Modes

II.2.3 Static and Dynamical effects

II.3 Optical Cavities : Three Mirror Cavities

II.3.1

II.4 Cavity Optomechanics

II.4.1 Radiation Pressure Coupling

II.4.2 Quantum Langevin Equations

II.4.3 Mechanical Resonators

II.4.4 Noise spectra

II.4.5 Three Mirror Cavities as Novel Optomechanical Systems

II.5 Squeezed Light Theory

II.5.1 Single-mode Squeezing

II.5.2 Noise Spectra

II.5.3 Frequency-dependent Squeezing and its use

II.6 Numerical Simulations of FDS Optomechanical Experiments

lets write smth here

Chapter III

Theory: MATE systems with squeezed light

Chapter IV

Experiments : A MATE system in vacuum

Chapter V

Experiments : a high frequency squeezed light source

Summary, conclusion and perspectives

Chapter VI

Un appendice

I'll write something here.

Sujet : Progress towards cryogenic squeezed light optomechanics

Résumé : .

Mots clés : Optomecanique, Lumière comprimée, Cavit  de grande Finesse, Interferom trie, Bruit thermique, Bruit de grenaille quantique, Resonateur de grand facteur de Qualit  , Interf rom tres pour la detection d'ondes gravitationnelles, Bruit de pression de radiation quantique

Subject : Optomechanics and squeezed light

Abstract:

Keywords : Optomechanics, Squeezing, High-Finesse cavity, Interferometry, Thermal Noise, Quantum Shot Noise, High-Q Resonator, Gravitational wave Interferometer, Quantum Radiation Pressure Noise