How Did We Get Here? Retracing the Lives of Binary Stars with POSYDON

A collection of work regarding the channels of binary star evolution

Pierson Lipschultz[1]

[1]Department of Astronomy, UIUC

Abstract

This poster serves as an introduction to **POSYDON** (**PO**pulation **SY**nthesis with **D**etailed binary-evolution simulati**ON**s), a tool used for population synthesis of binary stars [1]. Additionally, it is a collection of finished and work-in-process projects which utilize POSYDON as the backbone of the research process.

Current Work

It is known that the progenitors to stellar mass BHs are high mass stars, with many of these stars being in binaries with stellar type stars. However, there is a large amount of uncertainty when it comes to formation and survival rate of these systems. As seen in [2], due to the 'quiet' nature of these systems visual detection and observation is very difficult. Currently, I am hoping to find a formation rate and common evolutionary channel of low P_{orb} BH-Sol systems via population synthesis with POSYDON.

Previous Work

This previous work was completed during the fall 2025 semester at the College of DuPage as a part of an honors independent study. It served as a general introduction to POSYDON, its general data structure, and the various processes driving binary evolution. I broke down into simplified 'stages' of binary evolution, which I defined to be detached, semi-detached, and contact systems, comparing simulated populations with the 'poster children' of said stage. These reference stars included Vela X-1, V404 Cygni, and W Ursae Majoris respectively. From this comparison I found that these systems are largely outsiders in their respective stages, and furthermore present evidence that V404 Cygni must have evolved as a tertiary system with all three stars playing a respective roll in the evolution.

Introduction

Why Does Binary Star Evolution Matter?

Binary star evolution is a complicated and nuanced field of investigation that is deeply critical to our understanding of the universe. Binaries are critical for our understanding of cosmology due to type la SN and BH mergers are massively driving our understanding of GR. These binary systems make a large up the majority of the stars in our galaxy, with estimated 60%-80% of all stars in our MW being in a binary pair [3]. While a large portion of those will evolve with a wide enough separation to effectively evolve independently, not interacting during their lifetimes, there is a sizable fraction which do interact. Due to the already moderately complicated process of single star evolution, the possibility for binary evolution in interacting systems is extremely large and nuanced (see fig 5). This process poses a great problem when it comes to accurately modeling these systems from formation to end, something that POSYDON hopes to solve.



Figure 1: Detailed flowchart of binary star evolution. Reprint from [4].

What is POSYDON?

POSYDON is an open source tool for simulation, interpretation, and analyzation of binary systems. While POSYDON is heavily reliant on MESA grids, it takes a largely different approach. POSYDON takes an initial zero-age main sequence (ZAMS) system and evolves it through a series of grids and solvers, recording the properties at the start and end of each stage. By taking this approach, it allows for systems with vast, complicated evolutionary paths to be recorded from start to end. POSYDON provides the framework to do this process in a high performance computing environment, allowing massive and rapid analysis [1]. See table 1 for an example of data structure.

Binary ID	System State	Orbital Period (days)	$\log_{10}$ Mass Transfer Rate	Donor State	Donor Mass $M_\odot$	Accretor State	Accretor Mass $M_\odot$
54	Detached	0.047520	-99.000	NS	1.196033	stripped He Core He burning	≈ 1.002
183	Detached	0.0429883	≈ -80.8	NS	1.196033	stripped He Core He burning	$\approx .9957$

Table 1: Example of POSYDON data, heavily modified for readability

Previous Work

The goal with this piece of work was to compare how observed systems, and their respective populations, may differ from the 'true' distribution of their populations. There are many

My process for analyzing these systems consisted of first finding said 'poster children' for the systems, and then comparing said systems to their simulated population counterparts. To select my reference systems, mainly relying on historic means, picking the systems that helped drive the research forward in their respective categories forward. From this initial process I settled on Vela X-1, V404 Cygni, and W Ursae Majoris.

For my reference populations I made the choice to evolve a grid with as minimal of initial biasing. In order to do this, I sourced a finished $1Z_\odot$ POSYDON grid of a million systems where all the systems were initially binaries, but with random mass ratios, eccentricities, and periods. This allowed for a dataset which contained binaries in all stages of evolution, at a wide range of masses, allowing for more detailed analysis.

Current Work

As outlined in [2], the formation rate and evolutionary channels of BH-Sol binaries are useful for deeper understanding of XrBs. However, these systems pose great difficultly when trying to use conventional observational data. This is because, in contrast to XrBs, they are 'quiet', with the only sign of a BH companion being in slight variations to the sun-like stars spectra. [2] puts an upper limit via non-detection of the formation rate of these systems at 1 in 10^5.

The end goal of this project is to both propose an expected formation (and possibly detection rate) of BH-Sol systems, as well as investigating the evolutionary channels of BH-Sol systems

Even when investigating a very specific type of system, it is best practice to start with as generalized of a grid as possible. In this case, I started with a grid which utilized initial mass distributions from [5], with the initial masses being constrained to $4 < M_{1,\odot} < 150$, $.1 < M_{2,\odot} < 150$. This 'wide net' approach is crucial, as the evolutionary channels can be very diverse, with many 'paths' leading to the same destination.

Methods

Previous Work

My general workflow for analysis relied on a custom-written HR Diagram script, which allowed easy graphing of the donor star within the evolved population. Furthermore, this script utilized the Python package Bokeh, which allowed for interactive html plots, allowing for more efficient analysis. A simple use example can be found at `https://piersonlip.github.io/Posydon_HR_Graphing_Script/` This script was then applied to each observed system (see fig 2).



Figure 2: Vela X-1's observed temperature and luminosity in comparison to similar detached HMXBs. Note that the 'black box' is a representation of the error range

From this process it quickly became clear that both Vela X-1 and WUMa were extremes in their respective populations. This is predicatable, as the more extreme nature is directly correlated to ease of observation.



Figure 3: W Ursae Majoris observed temperature and luminosity in comparison to evolved contact binaries. Note that the CE solver has been altered since this population.

However, it quickly became clear that V404 Cygni was more than just an extreme case, it was radically different from the evolved population. This is discussed further in results.

Current Work

After evolving the initial grid I found a dominate channel of evolution emerged, this being `ZAMS_oCE1_CC1_oRLO2_CC2`. Via inspection of the evolution history, it is evident that the systems evolve detached, then into post-MS unstable RLO via S1 $\rightarrow$ S2. This period decrease leads to common envelope, which then *greatly* decreases the orbital period. The envelope is then ejected, leaving a depleted stripped He core. The core then collapses in a couple of thousand years, leaving the system in a BH-Sol state with low p_{orb}/separation. Furthermore, I found that the systems were quite constrained with their initial properties (see table 2)

	orbital_period_i
count	6.000000
mean	4022.716197
std	151.522384
min	3873.118288
25%	3908.300983
50%	3978.305326
75%	4124.397101
max	4248.507680

	S1_mass_i
count	6.000000
mean	17.515761
std	0.520269
min	16.815111
25%	17.190036
50%	17.510176
75%	17.818155
max	18.254965

	S2_mass_i
count	6.000000
mean	2.296788
std	0.320702
min	1.797809
25%	2.120753
50%	2.353828
75%	2.547210
max	2.620529

Table 2: Distributions of initial conditions in BH-Sol progenitors in Grid 1.

After the discovery of this channel in the initial population, I decided to focus on trying to understand it more in depth. From this I moderately constrained the initial grids to the distribution of previously evolved BH-Sol systems. This allowed for a much greater quantity of BH-Sol systems to be evolved, allowing for further analysis. Previous work from [2] placed an upper limit on the orbital period of these systems when filtering, however, when investigating the population, it became clear that the period of the systems was dependent on the stage of CE evolution, with little distinction between systems with $P_{orb} < 300$d. As a side effect of the evolutionary channel of BH-Sol systems it has become key to understand the survivability of the CE stage, which can start to be seen in fig 4.



Figure 4: Plot of the initial properties in BH-Sol progenitors. Includes a grid using a mass distribution from [5] and a sampled distribution.

As of now, 5 more grids have been simulated with various initial properties and metallicities. These grids are all still in the process of investigation.

Previous Results and Current Questions

Previous Work

With W UMa and Vela X-1 the results, while on the more extreme side, were expected, however, with V404 Cygni the results seemed to hint at a much larger phenomenon



Figure 5: V404 Cygni plotted against all other finished binaries.

Next Steps and Current Questions

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Discussion

Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetuer a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetuer. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus.

Acknowledgments

This work was supported by Ilia Kiato, Contact: `yourname@institution.edu`

References

[1] Tassos Fragos, Jeff J. Andrews, Simone S. Bavera, et al. "POSYDON: A General-Purpose Population Synthesis Code Based on Detailed Binary-Evolution Grids". In: *The Astrophysical Journal Supplement Series* 264.2 (2023), p. 45. DOI: 10.3847/1538-4365/ac90c1.

[2] Matthew J. Green et al. "An upper limit on the frequency of short-period black hole companions to Sun-like stars". In: *Astronomy and Astrophysics* 695 (Mar. 2025). ADS Bibcode: 2025A&A...695A.210G, A210. ISSN: 0004-6361. DOI: 10.1051/0004-6361/202453271. URL: https://ui.adsabs.harvard.edu/abs/2025A&A...695A.210G (visited on 02/12/2026).

[3] A. Author and B. Collaborator. "Title of the Second Reference". In: *Monthly Notices of the Royal Astronomical Society* 000 (2023), pp. 1–10. DOI: 10.1000/placeholder.

[4] Xuefei Chen, Zhengwei Liu, and Zhanwen Han. "Binary stars in the new millennium". In: *Progress in Particle and Nuclear Physics* 134 (2024), p. 104083. ISSN: 0146-6410. DOI: https://doi.org/10.1016/j.ppnp.2023.104083. URL: https://www.sciencedirect.com/science/article/pii/S0146641023000649.

[5] Maxwell Moe and Rosanne Di Stefano. "Mind Your Ps and Qs: The Interrelation between Period (P) and Mass-ratio (Q) Distributions of Binary Stars". In: *The Astrophysical Journal Supplement Series* 230 (June 2017). ADS Bibcode: 2017ApJS..230...15M. p. 15. ISSN: 0067-0049. DOI: 10.3847/1538-4365/aa6fb6. URL: https://ui.adsabs.harvard.edu/abs/2017ApJS..230...15M (visited on 03/31/2026).