

# Astro HW 8

Pierson Lipschultz \*

October 28, 2025

## 8.1 Stellar Properties

Given two stars A and B with properties ...

Name	Spectral type	$m_B$ (mag)	$m_V$ (mag)	Diameter ( $\times 10^{-3}$ " )	Parallax (")	Mass ( $M_\odot$ )
Star A	M2 Ia	2.27	0.42	43.43	0.00451	19.6
Star B	M2 V	8.96	7.52	1.43	0.393	0.46

### (a) Distances

$$d(\text{pc}) = \frac{1}{\theta_{\text{parallax}}(")}$$

$$d_A = \frac{1}{0.00451} \boxed{\approx 221.72 \text{ pc}}, \quad d_B = \frac{1}{0.393} \boxed{\approx 2.544 \text{ pc}}$$

### (b) Radii

Given the small-angle relation ...

$$\theta_{\text{diam}} = \frac{2R}{d} \quad \Rightarrow \quad R = \frac{\theta_{\text{diam}}}{2} d$$

We find ...

$$R_A = \frac{43.43 \times 10^{-3}}{2 \times 206265} (221.72)(3.086 \times 10^{16}) \boxed{\approx 7.20 \times 10^{11} \text{ m} \approx 1.035 \times 10^3 R_\odot}$$

$$R_B = \frac{1.43 \times 10^{-3}}{2 \times 206265} (2.544)(3.086 \times 10^{16}) \boxed{\approx 2.72 \times 10^8 \text{ m} \approx 0.391 R_\odot}$$

\*If there's anything I can do to make grading easier please let me know :)

**(c) Surface Gravity**

$$g = \frac{GM}{R^2}$$

$$M_A = 19.6 M_\odot = 3.89 \times 10^{31} \text{ kg}, \quad M_B = 0.46 M_\odot = 9.15 \times 10^{29} \text{ kg}$$

$$g_A = \frac{6.67 \times 10^{-11} (3.89 \times 10^{31})}{(7.20 \times 10^{11})^2} \boxed{\approx 5.0 \times 10^{-3} \text{ m/s}^2}$$

$$g_B = \frac{6.67 \times 10^{-11} (9.15 \times 10^{29})}{(2.72 \times 10^8)^2} \boxed{\approx 8.2 \times 10^2 \text{ m/s}^2}$$

**(d) Luminosity Class Comparison**

We expect the main-sequence dwarf (V) to have much higher surface gravity than the supergiant (Ia). Indeed,  $g_B \gg g_A$ , consistent with their classifications.

**(e) Temperatures**

Using

$$B - V = m_B - m_V \quad (13.35)$$

and

$$T \approx \frac{9000 \text{ K}}{(B - V) + 0.93} \quad (13.36)$$

$$T_A = \frac{9000}{(2.27 - 0.42) + 0.93} \boxed{\approx 3.24 \times 10^3 \text{ K}}, \quad T_B = \frac{9000}{(8.96 - 7.52) + 0.93} \boxed{\approx 3.80 \times 10^3 \text{ K}}$$

**(f) Luminosities**

From the Stefan-Boltzmann law:

$$L = 4\pi R^2 \sigma_{\text{SB}} T^4$$

$$L_A = 4\pi (7.20 \times 10^{11})^2 (5.67 \times 10^{-8}) (3.24 \times 10^3)^4 \boxed{\approx 4.06 \times 10^{31} \text{ W}}$$

$$L_B = 4\pi (2.72 \times 10^8)^2 (5.67 \times 10^{-8}) (3.80 \times 10^3)^4 \boxed{\approx 1.10 \times 10^{25} \text{ W}}$$

### (g) H-R Diagram Location <sup>1</sup>

Star A: very cool, extremely luminous → upper right of the H–R diagram (red supergiant). Star B: cooler main-sequence dwarf → lower right region (red dwarf). Both positions are consistent with their spectral classifications.

Name	Distance (pc)	Radius (m)	Radius ( $R_\odot$ )	$g$ (m/s <sup>2</sup> )	Temp (K)	$L_{\text{tot}}$ (W)
Star A	221.72	$7.20 \times 10^{11}$	$1.04 \times 10^3$	$5.0 \times 10^{-3}$	$3.24 \times 10^3$	$4.06 \times 10^{31}$
Star B	2.544	$2.72 \times 10^8$	$3.91 \times 10^{-1}$	$8.2 \times 10^2$	$3.80 \times 10^3$	$1.10 \times 10^{25}$

## 8.2 Stellar main sequence lifetimes

Given ...

$$N_H = \frac{M}{m_p} \quad (15.52)$$

$$E_{fus} = \frac{N_h}{4} \Delta E \quad (15.53)$$

$$t_{fus} = \frac{E_{fus}}{L} \quad (15.54)$$

We can derive ...

$$t_{fus} = \frac{1}{L} \frac{1}{4} \frac{\Delta E}{1} \frac{M}{m_p}$$

$$t_{fus} = \frac{\Delta E \times M}{4L \times m_p}$$

Where ...

$$\Delta E = 4.1 \times 10^{-12} \text{J}, \quad m_p = 1.67 \times 10^{-27} \text{kg}$$

Additionally, we are given ...

$$\tau = 10 \text{Gyr} \left( \frac{M}{M_\odot} \right)^{-3} \quad (15.55)$$

### a) High-Mass

Given ...

$$M = 100 M_\odot \approx 1.989 \times 10^{32} \text{kg}, \quad L = 10^6 L_\odot \approx 3.83 \times 10^{32} \text{W}$$

We find that ...

$$t_{fus} = \frac{4.1 \times 10^{-12} \text{J} \times 1.989 \times 10^{32} \text{kg}}{4(3.83 \times 10^{32} \text{W}) \times 1.67 \times 10^{-27} \text{kg}} \boxed{\approx 3.187 \times 10^{14} \text{s} \approx 1 \times 10^7 \text{yr}}$$

<sup>1</sup>If whoever is grading this is horribly bored, and has nothing better to do, check out the script I wrote to streamline H–R diagram graphing of POSYDON Data using matplotlib and bokeh :) [https://github.com/PiersonLip/Posydon\\_HR\\_Graphing\\_Script](https://github.com/PiersonLip/Posydon_HR_Graphing_Script)

Using equation 15.55 ...

$$\tau = 10\text{Gyr} \left( \frac{100M_{\odot}}{M_{\odot}} \right)^{-3} = \frac{1}{100000} \times 10\text{Gyr} = 1 \times 10^4\text{yr}$$

Thus, we find that...

$$\frac{t_{fus}}{\tau} = 10^3$$

This means our approximation is off by 3 orders of magnitude

### b) Low-mass

Given ...

$$M = .5M_{\odot} \approx 9.945 \times 10^{29}\text{kg}, \quad L = .1L_{\odot} \approx 3.83 \times 10^{25}\text{W}$$

We find that ...

$$t_{fus} = \frac{4.1 \times 10^{-12}\text{J} \times 9.945 \times 10^{29}\text{kg}}{4(3.83 \times 10^{25}\text{W}) \times 1.67 \times 10^{-27}\text{kg}} \approx \boxed{1.58 \times 10^{19}\text{s} \approx 5.006 \times 10^{11}\text{yr}}$$

Using equation 15.55 we find ...

$$\tau = 10\text{Gyr} \left( \frac{.5M_{\odot}}{M_{\odot}} \right)^{-3} = 8 \times 10\text{Gyr} = \boxed{8 \times 10^{10}\text{yr}}$$

Thus, we find that ...

$$\frac{t_{fus}}{\tau} = \frac{5.06 \times 10^{11}}{8 \times 10^{10}} \approx 6.25$$

Hence, our approximation is only off by a factor of  $\approx 6.25$

## 8.3 Stellar structure from opacity

Given ...

$$\kappa \propto \rho T^{-3.5}$$

$$\frac{dT}{dr} \approx \frac{T}{R}$$

$$\rho = M / \left( \frac{4}{3} \pi R^3 \right)$$

$$L = 4\pi R^2 \sigma_{sb} T^4 \rightarrow T = \left( \frac{L}{4\pi R^2 \sigma_{sb}} \right)^{1/4}$$

### a) Deriving $L \propto M^{5.5} R^{-.5}$

$$\kappa = \frac{3M}{4\pi R^2} \left( \frac{L}{4\pi R^2 \sigma_{sb}} \right)^{1/4} R$$

## 8.4 Extra Credit

a)

The parallax error would affect its location on the y-axis. This is because the parallax would affect the distance measured, which in turn effect the absolute luminosity, hence shifting the location on the y-axis.<sup>2</sup>

b)

Given ...

$$d = \frac{1}{\theta_{par}} \quad F = \frac{L_{abs}}{4\pi d^2}$$

We can find the difference as ...

$$\frac{L_{meas}}{L_{true}} = \left( \frac{d_{meas}}{d_{true}} \right)^2 = \left( \frac{1}{0.9} \right)^2 = 1.234 \dots$$

c)

$$\approx 23\%_{error}$$

This means that the L value on the y-axis of the HR diagram should be 23% brighter, meaning the star should be further **up** on the HR diagram.

---

<sup>2</sup>It could also affect the x-axis through redshift + Hubble const, but if a parallax error is that big you might have bigger problems.