

R1

Pierson Lipschultz

March 31, 2026

1 New Code / Devs

1.1 Python Terminal Script for Filtering

Because we're likely to need to filter quite a few systems with very similar methods I decided to make an actual script instead of just using a notebook repetably. The largest upside is that this can easily be run on a compute node, which makes filtering larger sims easier.

Takes a user input in the CLI (see below), filters for all BH_Sol systems, and returns the population file with said system as well as a .csv of just the BH_Sol rows. Furthermore, while it is running it provides feedback via the terminal stating how many BH_MS systems it found, how many are the expected S1:S2 configuration. If any are S2:S1, it prints a warning as well as the total count.

Need to further refine it, but there's also a SBATCH script.

Usage Example

```
python BH-Sol_Filter.py /Data/1e+00_Zsun_population.h5 -o ./testResults
```

Arguments

Argument	Use	Ex value
-o	output location	.results
-outputName	name for the outputed files ¹	BH_Sol_Systems
-overwrite	Whether to overwrite existing results, T/F bool. Defaults to F	True
--maxMass	Max mass used when filtering, defaults to $3M_{\odot}$	3.5
--maxPeriod	Max period used when filtering, defaults to 3.5 days	4

Can be found at https://github.com/PiersonLip/stellarBHs/blob/main/FilteringScripts/BH-Sol_Filter.py

¹If a name is not specified it will use the original filename, appending a specifier to the front

2 New Research

2.1 Pop 1

When filtering via notebook and script, both returned 6 systems that had $p_{orb} < 3.5, M_{\odot,2} < 3, t < 10e9$

These systems all evolved via the same channel, that being ZAMS_oCE1_CC1_oRL02_CC2. Via inspection of the history df and the oneline, it is evident that the systems evolve detached, then post-MS **unstable** RLO happens via $S1 \rightarrow S2$. This leads to common envelope, which *greatly* decreases the orbital period. This envelope is then ejected, leaving a depleted stripped He core. The core then collapses in a couple thousand years, leaving the system in a BH-MS state with low p_{orb} /seperation.

These systems then evolve in a semi-peculiar way, where it is highly likely that there will be RLO from S2 onto S1, and because $S1 > S2$, this is stable, and leads to *very* prolonged stable RLO (seemingly on the scale of $> 10Gyr$), which leaves S2 at low masses ($< .08M_{\odot}$).

Interestingly, the initial period for all the systems is very close to $4000d$, with mean of 4022 and a std of only 151. However, a larger sample size is needed to confirm this.

Additionally, the initial masses of S1 and S2 appear to be quite constrained as well.

	orbital_period.i
count	6.000000
mean	4022.716197
std	151.522384
min	3873.118288
25%	3908.300983
50%	3978.305326
75%	4124.397101
max	4248.507680

	S1_mass.i
count	6.000000
mean	17.515761
std	0.520269
min	16.815111
25%	17.190036
50%	17.510176
75%	17.818155
max	18.254965

	S2_mass.i
count	6.000000
mean	2.296788
std	0.320702
min	1.797809
25%	2.120753
50%	2.353828
75%	2.547210
max	2.620529

2.2 Pop 2

Ran a second grid including a eccentricity distro, as well as slightly constraining the dataset further, hoping to confirm some of the init conditions

	orbital_period.i
count	12.000000
mean	4136.066721
std	177.676729
min	3844.570082
25%	4024.214908
50%	4119.397535
75%	4273.345845
max	4370.857404

	S1_mass.i
count	12.000000
mean	2.240382
std	0.412075
min	1.551266
25%	1.903135
50%	2.300024
75%	2.579111
max	2.729828

	S2_mass.i
count	12.000000
mean	17.357478
std	1.073496
min	15.506937
25%	16.714083
50%	16.998229
75%	18.162633
max	19.435080

2.3 Full P_{orb}

When filtering pop 2 with a max $p_{orb} = 300$ days we find 46 BH-Sol systems, all of these (except one!) evolve via the same CE channel. There is one system which does *not* do RLO before CC1, and it also only reaches a solar-like mass after RLO2, so I'm excluding it for now. Probably important to keep in mind though.

3 To-Do

- Find out if it makes sense for the onset of un-stable rlo to happen at $P_{orb} \approx 4000\text{days}$

4 Questions

- Tried running a grid, seems to have just completely failed from trying to reference itself before it was fully initialized. Entire log was filled with

```
Traceback (most recent call last):
File "/gpfs/projects/b1095/bku2126/UCXBInvestigation/POSYDON/posydon/popsyn/binarypopulation.py", 1
    binary.evolve()
File "/gpfs/projects/b1095/bku2126/UCXBInvestigation/POSYDON/posydon/binary_evol/binarystar.py", 1
    self.run_step()
File "/gpfs/projects/b1095/bku2126/UCXBInvestigation/POSYDON/posydon/binary_evol/binarystar.py", 1
    next_step(self)
File "/gpfs/projects/b1095/bku2126/UCXBInvestigation/POSYDON/posydon/binary_evol/DT/step_merged.py", 1
    super().__call__(binary)
File "/gpfs/projects/b1095/bku2126/UCXBInvestigation/POSYDON/posydon/binary_evol/DT/step_isolated.py", 1
    super().__call__(binary)
File "/gpfs/projects/b1095/bku2126/UCXBInvestigation/POSYDON/posydon/binary_evol/DT/step_detached.py", 1
    self.update_after_evo(t, binary, primary, secondary)
File "/gpfs/projects/b1095/bku2126/UCXBInvestigation/POSYDON/posydon/binary_evol/DT/step_detached.py", 1
    current = history[-1]
                ~~~~~~
IndexError: index -1 is out of bounds for axis 0 with size 0
```

and from what I can tell, this happened to nearly all the simmed bianaries

- Should we include an initial eccentricity distribution? I imagine it would impact formation with unstable RLO and CE.
- Does it make sense to still be filtering for $p_{orb} < 3\text{days}$? The simulations seem to be suggesting that *all* BH-Sol systems evolve through CE, and that the main thing effecting p_{orb} post CE is the kick strength of the BH. So, maybe we should be initially filterting for all BH-Sol, and if we want the formation rate of specifically low p_{orb} ones, just take the subpop.
- Cool if i present an assortment of general POSYDON things ive worked on at astrofest?