

HPV DNA Genotyping Test & Risk

The Fight Against Cervical Cancer – Cervical Cancer is the cancer we can eradicate

Promo RPP @ **RM180** (Single), **RM166** (Tag-on)



How many HPV genotype caused infection?

There are quite a variety of HPV types. At present, more than **200 types** have been found. About **40 types** can infect the skin or mucous membrane of the lower genital tract. Sexual contact is the main route of infection, but it may also be caused by genital-skin contact.

What cancer related to HPV infection?

Carcinogenic (High Risk) HPV can cause a variety of cancers, such as **cervical cancer, vaginal cancer, vulvar cancer, anal cancer**, etc. Among them, cervical cancer is the most common. Carcinogenic HPV is a necessary cause of cervical cancer.

HPV genotype classification?

HPV that can infect the lower genital tract can be classified according to its degree of correlation with cervical cancer, our test able to detect:

19 High-risk Type (A carcinogenic virus highly associated with cervical cancer, CINII, and III.)

16, 18, 26, 31, 33, 35, 39, 45, 51, 52, 53, 56, 58, 59, 66, 68, 69, 73, 82.

In cervical cancer tissues, the 16 and 18 types are the most common.

9 Low-risk Type (low-degree association with cervical cancer, CINII, and III)

6, 11, 40, 42, 43, 44, 54, 61, 70.

Among them, types 6 and 11 are often found in the genital warts (commonly known as cauliflower).

How is the prevalence of HPV infection?

According to the National Bureau of Health, the prevalence of high-risk HPV in women is about 10-15%, and more than half of the HPV of infected people disappear within one year. Sustained high-risk HPV infection can cause cervical cell variability and even lead to cancer.

HPV/PAP Referral Guideline

- These recommendations do not apply to women who have received a diagnosis of high grade precancerous cervical lesion or cervical cancer, women with in-utero exposure to diethylstilbestrol, or women who are immunocompromised such as those who are HIV positive.
- Pap Smear Screening begins at age 21 regardless of when sexual activity starts. Adolescents/young women 20 and below are not recommended to have a Pap test or HPV testing. Pap screening may end at age 65 if the Pap history is unremarkable and the patient is low risk.
- Screening recommended every 3 years for women 21-29. Women 30-65 and older who have had 3 consecutive negative Pap test and who have no history of CIN2 or 3, etc. OR low risk women 30 and above may go every 3 years if Pap only; or 5 years if co-testing. Risk factors that should be considered are HPV infection, STD's, HIV, smoking, new sexual partners, previous SIL or immunosuppression.
- When "absent/insufficient endocervical/transformation zone cells" is reported in a negative Pap smear in ages 21-29, follow routine screening. For ages 30 or greater, perform HPV. If HPV negative, do routine screening. If HPV positive, do PAP + HPV in 1 year or genotype, then manage per ASCCP guidelines.
- After hysterectomy for benign causes, women need not undergo routine Pap smears unless symptomatic, history of "SIL," or has associated risk factors as above.
- HPV testing (high risk types) is the preferred method for triage of ASCUS results using liquid cytology for ages 25-65. If 21-24 years, repeat PAP in 12 months.
- Screening practice should not change on the basis of HPV vaccination.
- Primary HPV testing can be considered for women starting at age 25 (2015 ASCP and SGO interim guideline.)

References:

1. ACS/ASCCP/ASCP Cervical Cancer Screening Recommendations, CA: A Cancer Journal of Clinicians Vol 62, May/June 2012, 147-172.
2. ACOG Practice Bulletin 109, December 2009.
3. American Society for Colposcopy and Cervical Pathology (<http://www.ascp.org>): Journal of Lower Genital Tract Disease, Vol 11, Number 4, 2007, 201-222.
4. AM J Ob Gyn 2007; 346-355
5. ACOG guidelines for cervical cancer screening, Obstet Gynecol 2006; 107(4) 963-8.
6. ASCP Guidelines 2009) 330-337.
7. Use of Primary High-Risk human papillomavirus testing for cervical cancer screening, Obstet Gynecol 2015; 125(2) 330-337.

Test Code HPV DNA

RPP

Specimen Type

: HPV D

: **Promo RM 180** ~~Normal RM240~~

: Female - FLOQ Swab (Self-collection or by doctor).

Male - Urethral Swab by doctor

Test Code HPV DNA + Liquid PAP

RPP

Specimen Type

: CXC

: **Promo RM 230** ~~Normal RM280~~

: Cervical Brush in collection vial

TAT

Promo validity until

: 7-10 working days

: 30/6/2026